



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Population density and developmental stress in the Neolithic: A diachronic study of dental fluctuating asymmetry at Çatalhöyük**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Population density and developmental stress in the Neolithic: A diachronic study of dental fluctuating asymmetry at Çatalhöyük (Turkey, 7,100–5,950 BC) / Milella, Marco\*; Betz, Barbara J.; Knüsel, Christopher J.; Larsen, Clark Spencer; Dori, Irene. - In: AMERICAN JOURNAL OF PHYSICAL ANTHROPOLOGY. - ISSN 0002-9483. - STAMPA. - 167:(2018), pp. 737-749. [10.1002/ajpa.23700]

*Availability:*

This version is available at: 2158/1151323 since: 2019-03-19T11:23:30Z

*Published version:*

DOI: 10.1002/ajpa.23700

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)



111 RIVER STREET, HOBOKEN, NJ 07030

**\*\*\*IMMEDIATE RESPONSE REQUIRED\*\*\***

Your article will be published online via Wiley's EarlyView® service ([wileyonlinelibrary.com](http://wileyonlinelibrary.com)) shortly after receipt of corrections. EarlyView® is Wiley's online publication of individual articles in full text HTML and/or pdf format before release of the compiled print issue of the journal. Articles posted online in EarlyView® are peer-reviewed, copyedited, author corrected, and fully citable via the article DOI (for further information, visit [www.doi.org](http://www.doi.org)). EarlyView® means you benefit from the best of two worlds--fast online availability as well as traditional, issue-based archiving.

Please follow these instructions to avoid delay of publication.

**READ PROOFS CAREFULLY**

- This will be your only chance to review these proofs. **Please note that once your corrected article is posted online, it is considered legally published, and cannot be removed from the Web site for further corrections.**
- Please note that the volume and page numbers shown on the proofs are for position only.

**ANSWER ALL QUERIES ON PROOFS** (If there are queries they will be found on the last page of the PDF file.)

- In order to speed the proofing process, we strongly encourage authors to correct proofs by annotating PDF files. Please see the instructions on the Annotation of PDF files. If unable to annotate the PDF file, please print out and mark changes directly on the page proofs.

**CHECK FIGURES AND TABLES CAREFULLY**

- Check size, numbering, and orientation of figures.
- All images in the PDF are downsampled (reduced to lower resolution and file size) to facilitate Internet delivery. These images will appear at higher resolution and sharpness in the final, published article.
- Review figure legends to ensure that they are complete.
- Check all tables. Review layout, title, and footnotes.

RETURN  **PROOFS**

**Other forms, as needed**

**Return corrections immediately via email to [DL-AJPA@wiley.com](mailto:DL-AJPA@wiley.com)**

**QUESTIONS**

Production Editor, AJPA

E-mail: [DL-AJPA@wiley.com](mailto:DL-AJPA@wiley.com)

Please refer to journal acronym and article production number


USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required software to e-annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 11 or above). (Note that this document uses screenshots from Adobe Reader DC.)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab (right-hand panel or under the Tools menu).

This will open up a ribbon panel at the top of the document. Using a tool will place a comment in the right-hand panel. The tools you will use for annotating your proof are shown below:




1. **Replace (Ins) Tool – for replacing text.**

 Strikes a line through text and opens up a text box where replacement text can be entered.

**How to use it:**

- Highlight a word or sentence.
- Click on .
- Type the replacement text into the blue box that appears.



2. **Strikethrough (Del) Tool – for deleting text.**

 Strikes a red line through text that is to be deleted.

**How to use it:**

- Highlight a word or sentence.
- Click on .
- The text will be struck out in red.

experimental data if available. For ORFs to be had to meet all of the following criteria:


1. Small size (35-250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminus terminus with another coding feature; over both ends; or ORF containing a tRNA.

3. **Commenting Tool – for highlighting a section to be changed to bold or italic or for general comments.**

  Use these 2 tools to highlight the text where a comment is then made.

**How to use it:**

- Click on .
- Click and drag over the text you need to highlight for the comment you will add.
- Click on .
- Click close to the text you just highlighted.
- Type any instructions regarding the text to be altered into the box that appears.




4. **Insert Tool – for inserting missing text at specific points in the text.**

 Marks an insertion point in the text and opens up a text box where comments can be entered.

**How to use it:**

- Click on .
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the box that appears.



**5. Attach File Tool – for inserting large amounts of text or replacement figures.**

 Inserts an icon linking to the attached file in the appropriate place in the text.

**How to use it:**

- Click on .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

The attachment appears in the right-hand panel.

chondrial preparator  
ative damage injury  
re extent of membra  
i, malondialdehyde (TBARS) formation.  
used by high perform


**6. Add stamp Tool – for approving a proof if no corrections are required.**

 Inserts a selected stamp onto an appropriate place in the proof.

**How to use it:**

- Click on .
- Select the stamp you want to use. (The **Approved** stamp is usually available directly in the menu that appears. Others are shown under *Dynamic*, *Sign Here*, *Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

on the business cycle, starting with the  
on perfect competition, constant ret  
production. In this environment goods  
extra costs and be a source of market  
he market is determined by the model. The New-Key  
otaki (1987), has introduced produc  
general equilibrium models with nomin  
and downward sloping. Most of this literat




**How to use it:**


- Click on one of the shapes in the **Drawing Markups** section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select *Open Pop-up Note*.
- Type any text in the red box that appears.

**7. Drawing Markups Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.**

Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.



For further information on how to annotate proofs, click on the **Help** menu to reveal a list of further options:





### **Additional reprint purchases**

Should you wish to purchase additional copies of your article, please click on the link and follow the instructions provided:

<https://caesar.sheridan.com/reprints/redir.php?pub=10089&acro=AJPA>

Corresponding authors are invited to inform their co-authors of the reprint options available.

Please note that regardless of the form in which they are acquired, reprints should not be resold, nor further disseminated in electronic form, nor deployed in part or in whole in any marketing, promotional or educational contexts without authorization from Wiley. Permissions requests should be directed to mail to: [permissionsus@wiley.com](mailto:permissionsus@wiley.com)

For information about 'Pay-Per-View and Article Select' click on the following link: [wileyonlinelibrary.com/aboutus/ppv-articleselect.html](http://wileyonlinelibrary.com/aboutus/ppv-articleselect.html)

# AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.

Query References	Query	Remarks
<b>Q1</b>	Please confirm that given names (blue) and surnames/family names (vermilion) have been identified and spelled correctly.	
<b>Q2</b>	Please check if link to ORCID is correct.	
<b>Q3</b>	Reference "Stock and Pinhasi (2011)" has not been included in the Reference List, please supply full publication details.	
<b>Q4</b>	Please provide exact date along with citation "I. Hodder, personal communication."	
<b>Q5</b>	Parts in Tables have been removed and renumbered according to citation order. Please check.	

## Funding Info Query Form

Please confirm that the funding sponsor list below was correctly extracted from your article: that it includes all funders and that the text has been matched to the correct FundRef Registry organization names. If no FundRef Registry organization name has been identified, it may be that the funder was not found in the FundRef registry, or there are multiple funders matched in the FundRef registry. If a name was not found in the FundRef registry, it may not be the canonical name form, it may be a program name rather than an organization name, or it may be an organization not yet included in FundRef Registry. If you know of another name form or a parent organization name for a “not found” item on this list below, please share that information.

<b>Funding Agency</b>	<b>FundRef Organization Name</b>
Agence Nationale de la Recherche	Agence Nationale de la Recherche;
H2020 Marie Skłodowska-Curie Actions	H2020 Marie Skłodowska-Curie Actions;
National Geographic Society	National Geographic Society

RESEARCH ARTICLE

# Population density and developmental stress in the Neolithic: A diachronic study of dental fluctuating asymmetry at Çatalhöyük (Turkey, 7,100–5,950 BC)

Marco Milella<sup>1</sup>  | Barbara J. Betz<sup>2</sup> | Christopher J. Knüsel<sup>3</sup> | Clark Spencer Larsen<sup>2</sup>  | Irene Dori<sup>3,4</sup>

<sup>1</sup>Department of Anthropology, University of Zurich, Zurich, Switzerland

<sup>2</sup>Department of Anthropology, 4034 Smith Laboratory, The Ohio State University, Columbus, Ohio

<sup>3</sup>Université de Bordeaux, Pessac, France

<sup>4</sup>Department of Biology, Laboratory of Anthropology, University of Florence, Florence, Italy

**Correspondence**

Marco Milella, Department of Anthropology, University of Zurich, Winterthurerstrasse 190, 8057, Zurich, Switzerland.  
Email: marco.milella@aim.uzh.ch

**Funding information**

Agence Nationale de la Recherche, Grant/Award Number: ANR-10-IDEX-03-02; H2020 Marie Skłodowska-Curie Actions, Grant/Award Number: 752626; National Geographic Society

**Abstract**

**Objectives:** The transition from foraging to farming is usually associated with unprecedented population densities coupled with an increase in fertility and population growth. However, little is known about the biological effects of such demographic changes during the Neolithic. In the present work, we test the relationship between diachronic changes in population size, relative exposure to developmental stressors, and patterns of dental fluctuating asymmetry in the Neolithic population of Çatalhöyük (Turkey, 7,100–5,950 cal BC).

**Materials and Methods:** We calculate fluctuating asymmetry of mesio-distal and bucco-lingual diameters of upper and lower permanent canines and first and second molars on a large (N = 259) sample representing adults of both sexes and various age classes.

**Results:** Results show only a moderate decrease of fluctuating asymmetry during the late phase of occupation of the site, possibly linked to a decrease in population density, and no differences in asymmetry between sexes.

**Discussion:** Though preliminary, our data reflect the presence of developmental stressors throughout the occupation of the site, albeit with a slight improvement in living conditions during the latest periods of occupation. At the same time, these data confirm the key role of diet as buffer against the detrimental effects of fluctuating demographic pressures on the biology of prehistoric human populations.

**KEYWORDS**

developmental stress, fluctuating asymmetry, neolithic, population density

## 1 | INTRODUCTION

The transition from foraging to farming is associated with the “...first human experiment in unprecedented population concentrations” (Bocquet-Appel, 2008). High energetic inputs due to calorie-rich food availability and a concomitant decrease in energy expenditure prefigured a shift in the energetic equilibrium of fertile women, resulting in an increase in fertility and birthrates, with unprecedented demographic effects (Bocquet-Appel, 2002, 2008, 2011). On the other hand, while these increases in fertility are today widely recognized due to convincing bioarchaeological and ethnological data, there is also growing evidence of an increase in infant mortality, and of

alternating patterns of boom-and-bust demographic changes during the early-mid Holocene of the Near East and Western Europe (Bocquet-Appel, 2008; Shennan et al., 2013).

Higher population densities resulted in increasingly complex social networks, increased social tensions, and the unprecedented flourishing of ritual and symbolism (Garfinkel, 1987; Kuijt & Goring-Morris, 2002; Pearson & Meskell, 2015; Rollefson, 2002; Wright, 2014). These patterns, in addition to their specific bearing on prehistoric archaeology and palaeodemography, have wider consequences, forming the basis of later phenomena such as social differentiation, social stratification, and the rise and development of social inequality

1 (Powers & Lehmann, 2014; Powers, van Schaik, & Lehmann, 2016;  
2 Price, 1995).

3 The transition to farming is associated by various authors with a  
4 decrease in health, differential patterns and sex-based division of  
5 physical activity, and a variety of micro-evolutionary processes  
6 (González-José et al., 2005; Katz, Grote, & Weaver, 2017; Larsen,  
7 1995; O'Brien et al., 2012; Paschetta et al., 2010; Pinhasi, Eshed, &  
8 Shaw, 2008; Pinhasi, Eshed, & von Cramon-Taubadel, 2015; von  
9 Cramon-Taubadel, 2011, 2017). The greater proximity of people to  
10 animals that this transition brought and the closeness of living quar-  
11 ters to waste disposal areas are recognized as the causes of the so-  
12 called "first epidemiological transition," generally characterized by the  
13 earliest appearance of infectious and parasitic diseases among human  
14 populations (Armélagos, Brown, & Turner, 2005; Larsen, 1995). All  
15 these factors, together with nutritional (especially protein) deficiencies  
16 stemming from a narrower dietary niche, are the basis of well-  
17 documented social changes coinciding with the Palaeolithic to Neo-  
18 lithic transition. Previous bioarchaeological comparisons between  
19 hunter-gatherer and farming communities highlight a general worsen-  
20 ing of skeletal and dental health, a pattern consistent with an increase  
21 in socio-environmental stressors with the adoption of farming  
22 (Cohen & Armélagos, 1984; Larsen, 1995). It should be stressed, how-  
23 ever, that the above pattern was probably not ubiquitous but rather  
24 characterized by a certain diachronic and regional heterogeneity  
25 (Cohen & Crane-Kramer, 2007; Starling & Stock, 2007; Stock & Pin-  
26 hasi, 2011). Also, biocultural reconstructions based on paleopathologi-  
27 cal patterns need to be considered with caution, given the possible  
28 mismatch between observed frequencies of skeletal lesions and actual  
29 health of the individual/population (Wood, Milner, Harpending, &  
30 Weiss, 1992—but see Goodman, 1993).

31 The above features raise the question of the types of biocultural  
32 adaptive strategies adopted by humans during the Neolithic transition,  
33 and, more specifically, of the socio-cultural and biological evolutionary  
34 processes triggered by the new environmental challenges associated  
35 with sedentism (Hawks, Wang, Cochran, Harpending, & Moyzis, 2007;  
36 Holden & Mace 1997; Laland, Odling-Smee, & Myles, 2010; Mace,  
37 2009; Naugler, 2008; O'Brien et al., 2012). An interesting angle from  
38 which to approach this issue is the exploration of possible relation-  
39 ships between changes in population densities, deviations in human  
40 developmental trajectories, and the evolution of developmental  
41 and/or behavioral buffers to such stressors over time.

42 Developmental stressors consist of any genetic and/or environ-  
43 mental factor hampering a specific genotype to reach its target pheno-  
44 type, given a particular growth environment (Palmer & Strobeck,  
45 2003). Examples include (among others) inbreeding, nutritional defi-  
46 ciencies, and infectious and metabolic diseases (Parsons, 1990). Inde-  
47 pendently from the type of stressors, threats to physiological  
48 homeostasis may result in deviations from the expected developmen-  
49 tal trajectories under given environmental conditions. Accordingly,  
50 such deviations are potentially informative on two levels: about the  
51 relative developmental stability (or instability—DI) of an organism  
52 (Møller & Swaddle, 1997), and about the number of stressors to which  
53 the organism is exposed during growth and development.

54 Studies of stress in past populations are most often based on a  
55 suite of skeletal and dental features (e.g., enamel hypoplasia, *cribra*

*orbitalia*, periostotic lesions, and long bone dimensions) associated 56  
either with developmental disruptions, non-specific bone reactive pro- 57  
cesses, or nutritional deficiencies. An additional proxy of developmen- 58  
tal stress, widely used in biology, is fluctuating asymmetry (FA). FA is 59  
defined as subtle, random deviations from symmetry in bilateral traits 60  
and is usually adopted in developmental biology as a proxy for devel- 61  
opmental instability (Klingenberg, 2003). FA differs from the two 62  
other types of bilateral asymmetry, anti-symmetry (AS), and direc- 63  
tional asymmetry (DA). In AS a population shows a mean difference 64  
between sides for bilateral traits, but it is composed of both left- and 65  
right-biased individuals. In DA, on the other hand, individuals present 66  
a bias in bilateral traits consistently favoring one side over the other. 67  
Mathematically, FA, AS, and DA are defined, respectively, by a normal 68  
and a platykurtic or bimodal distribution about a mean of zero, and by 69  
a normal distribution about a mean other than zero. The rationale for 70  
the use of FA in studies of developmental stress is that random, local- 71  
ized perturbations differently affect the development of the two sides 72  
of symmetric traits otherwise sharing the same genotype, resulting in 73  
asymmetry (Van Valen, 1962). Differences between sides can there- 74  
fore be interpreted as proxy for developmental instability, and corre- 75  
lated with the presence of stress-induced developmental noise. 76

77 Though largely under-represented in the bioarchaeological litera- 78  
ture, previous studies have already explored the use of dental, cranio- 79  
facial, and skeletal FA in testing patterns of developmental stress in 80  
both living and past populations, with results that overall confirm this 81  
parameter as a useful tool when investigating developmental stress in 82  
past populations (Albert & Greene 1999; Barrett, Guatelli-Steinberg, & 83  
Sciulli, 2012; Costa, 1986; DeLeon, 2007; Doyle & Johnston, 1977;  
84 Gawlikowska-Sroka, Dabrowski P. Szczurowski, Dzieciolowska-  
85 Baran, & Staniowski, 2017; Gawlikowska-Sroka, Dąbrowski, Szczur-  
86 owski, & Staniowski, 2013; Greene, 1984; Hoover, Corruccini, Bon-  
87 dioli, & Macchiarelli, 2005; Hoover & Matsumura, 2008; Kieser,  
88 Groeneveld, & Preston, 1986; Kujanova, Bigoni, Velemínska, & Vele-  
89 minsky, 2008; Perzigian, 1977). FA offers advantages over classic  
90 osteological markers of stress, namely that FA is a quantitative vari-  
91 able, and it is not linked to episodes of stress (as is the case with  
92 enamel hypoplasia); but represents the outcome of continuous alter-  
93 ations in physiological homeostasis. Collectively, these attributes  
94 make analyses of FA less biased by the analytical problems typically  
95 affecting qualitative features and are more suited to discussions of  
96 long-term patterns of developmental stress.

97 Many bioarchaeological studies of stress in past populations com- 98  
pare skeletal assemblages representing different social statuses and 99  
contrasting socio-economic strategies, with others focused on the  
100 analysis of differences in stature and/or skeletal and dental morpho-  
101 logical changes between different social groups and between foragers  
102 and farmers (Bigoni, Krajčec, Sladek, Velemínsky, & Velemínska,  
103 2013; Cardoso & Gomes, 2009; Cohen & Armélagos, 1984; Cohen &  
104 Crane-Kramer, 2007; Larsen, 1995; Pinhasi & Stock, 2011; Sakashita,  
105 Inoue, Inoue, & Zhu, 1997; Starling & Stock, 2007; Temple & Larsen,  
106 2007). In addition to demonstrating the usefulness of dental and skel-  
107 etal changes when testing biocultural hypotheses, results of these  
108 studies are, in general, consistent in highlighting a general decrease in  
109 quality of life and increase in environmental stressors associated with  
110 the adoption of a sedentary lifestyle and farming economy. On the

other hand, much less is known about the biological effects of fluctuating population densities in Neolithic communities after the transition to farming (but see Larsen et al., 2015). This study applies a more fine-grained approach by exploring the relationship between changes in population density and patterns of developmental stress in a single archaeological setting—Neolithic Çatalhöyük in Central Anatolia—characterized by changes in population size in the context of a millennium-long occupation of a large village setting.

## 1.1 | Biocultural context

Neolithic Çatalhöyük (7100–5,950 cal BC—Bayliss et al., 2015) is one of the most important settlements of the Near Eastern Pre-Pottery or Aceramic Neolithic, being characterized by a large occupation footprint (13 ha), long-standing occupation (ca., 1,150 years) and archaeological features consistent with the presence of complex sociocultural traditions.

The exceptional archaeological and bioarchaeological contextualization of this site (e.g., Hodder, 2014a), unmatched by any other Neolithic context in the Old World, permits a unique opportunity to test central hypotheses about the biocultural changes associated with large population agglomerations in a key geographic area for the study of subsequent Neolithic processes.

Two main excavation areas are represented at Çatalhöyük (North and South—including TP and TPC areas) (Hodder, 2014b). Stratigraphic levels at Çatalhöyük, from the start of the occupation to the abandonment of the site, are subdivided into four diachronic phases: early (7,100–6,700 BC), middle (6,700–6,500 BC), late (6,500–6,300 BC), and final (6,300–5,950 BC) (I. Hodder, personal communication, and forthcoming). Current estimates based on architectural features (Cessford, 2005) indicate a population of 3,500–8,000 individuals, while previous archaeological and paleodemographic data (number of buildings, number of buried individuals per building, juvenility indices) suggested the presence of salient demographic fluctuations throughout the occupation of the site (Cessford, 2005; Düring, 2001; Larsen et al., 2015). Similarly, estimates of percentage of open spaces and number of building per excavated area, suggest a progressive increase of building dispersal throughout time (Düring, 2001: Table 1; Hodder, 2014a).

As for paleodemographic patterns, previous calculation of juvenility indices (used as proxy for fertility) (Larsen et al., 2015) describe in

clear fashion an increase in population size and fertility throughout the early and middle phases of occupation, with a peak occurring around 6,610–6,250 BC, followed by a significant decrease (Early = .29, Middle = .46, Late = .29). This pattern, taken together with archaeological sites in the Konya plain pre-dating the site at Çatalhöyük (Baird, 2005; Baird et al., 2018) represented by typically small and rather dispersed settlements fits a scenario of population agglomeration, with demographic growth mainly driven by increases in fertility and birthrate. Conversely, the demographic decline estimated for the later phases could be related to the progressive abandonment of the site, likely due to a complex mosaic of factors, possibly including environmental changes driven by human activity and over-exploitation of resources (Doherty, 2013, Orton et al. 2018).

The presence of demographic fluctuations in a relatively narrow chronological interval raises the question of their possible impact on the members of the Çatalhöyük community, the question being which kind (if any) of effects these changes had on population density (and its biocultural correlates) of the settlement. Previous research highlighted diachronic patterns in the frequency of periostotic lesions mirroring the postulated changes in population size at the site, a result interpreted as a result of differential exposures to pathogens through time due to variable population densities (Larsen et al., 2015). This hypothesis seems supported by both isotopic and biomechanical data, which converge in suggesting the presence during the later phase of occupation of a more mobile and dispersed population, a factor that would have led to a reduced exposure to pathogens.

Conversely, reconstructed growth trajectories of both stature and body mass seem to contradict such a scenario, equating to an overall “healthy” population, a result interpreted as due to the buffering role of an adequate nutrition against the detrimental effects of environmental stressors (e.g., parasitic diseases, infections) on development (Larsen et al., 2015). These contrasting results stress the need for further research into the type of demographic changes (population size and density) characterizing the occupation of Çatalhöyük, and on their biological and cultural effects on the occupants of the settlement.

Here, we use Çatalhöyük as a natural experiment to explore the biological correlates of population changes associated with the Neolithic transition. More specifically, we focus on the developmental effects of such changes using dental fluctuating asymmetry (DFA) as a proxy for differential exposure to environmental stressors to test three main hypotheses:

**TABLE 1** Sample size by sex, age, and occupation phase

	Early			Middle			Late/final			Unstratified Neolithic			Total
	F	M	NA	F	M	NA	F	M	NA	F	M	NA	
Infant (2 months–3 y)	0	0	0	0	0	3	0	0	1	0	0	0	4
Child (3–12 y)	0	0	5	0	0	58	0	0	13	0	0	5	81
Adolescent (12–19 y)	0	1	1	3	2	17	2	1	5	0	1	3	36
Young adult (20–34 y)	2	2	1	13	12	1	5	4	0	2	0	2	44
Mature adult (35–50 y)	2	2	0	13	10	0	7	7	1	2	1	1	46
Older adult (>50 y)	2	2	0	7	2	1	0	3	0	0	0	0	17
Adult (>20 y)	1	0	0	6	6	2	3	6	3	1	1	2	31
Total	7	7	7	42	32	82	17	21	23	5	3	13	259

Y = years; M = males; F = females; NA = undetermined sex.

1 Hp1: Given a correlation between population density (and fertil-  
2 ity) and relative exposure to environmental stressors, mean DFA  
3 values will correlate with fluctuations in population size (and fertility),  
4 and will be characterized by an increase throughout the early and mid-  
5 dle phases followed by a decrease.

6 Hp2: Given the hypothesized higher developmental buffering and  
7 greater genetic control of odontogenesis in females (Garn, Lewis, &  
8 Kerewsky, 1967; Garn, Lewis, Kerewsky, & Jegart, 1965; Stinson,  
9 1985), males will exceed females in expressing developmental  
10 disturbance.

11 Hp3: Because of the expected correlation between FA and devel-  
12 opmental stress, there will be an inverse correlation between asym-  
13 metry and age-at-death in this setting.

## 16 2 | MATERIALS AND METHODS

17 Before proceeding with the analysis of asymmetry, we determined if  
18 inclusion of the skeletal remains excavated in 2016 and 2017 altered  
19 the juvenility index estimates proposed by Larsen et al. (2015), and  
20 recalculated the number of buildings and buried individuals per build-  
21 ing using data not available for previous estimates. For the latter  
22 purpose, only primary and primary disturbed burials were considered.  
23 Note that the use of the term buried individuals rather than burials is  
24 preferred here since burials at Çatalhöyük may represent both single  
25 and multiple interments.

26 The juvenility index D3-19/D3+ (the ratio between individuals  
27 aged between 3 and 19 years and the total of individuals above  
28 3 years of age) was calculated on a sample of 324 individuals from pri-  
29 mary and primary disturbed burials from all occupation levels at Çatal-  
30 höyük. In this and the following age-at-death and sex were  
31 determined according to standard protocols (Buikstra & Ubel-  
32 ker, 1994).

33 Analyses of DFA were performed on a sample representing all  
34 stratigraphic phases of Çatalhöyük and composed of individuals  
35 selected on the basis of the presence of the following antimeric per-  
36 manent teeth: canines (C), first molars (M1), and second molars  
37 (M2) of both maxillae and mandibles. These teeth were chosen in  
38 order to be able to study developmental stress for a relatively long  
39 period of individual growth and development. The choice to consider  
40 both molars was suggested by the possible differences in develop-  
41 mental stability between polar versus non-polar teeth of a dental field  
42 (Dahlberg, 1945; Goodman, 1989; Goodman & Armelagos, 1985). On  
43 each tooth, bucco-lingual (BL) and mesio-distal (MD) diameters were  
44 taken according to Hillson, Fitzgerald, & Flinn (2005) with a Mitutoyo  
45 digital calipers (accuracy = .01). Teeth with incomplete crown forma-  
46 tion, caries, intra-vitam and post-mortem damage, high degrees of cal-  
47 culus (calcified plaque) deposition, and wear were excluded from the  
48 study. In total, the sample is composed of 259 individuals representing  
49 both sexes and various age classes (Table 1).

50 Because of the need to maximize the sample size in the context  
51 of time constraints, data collection was subdivided between two  
52 observers (MM and ID). Both observers measured each tooth two  
53 times, with an interval of a week between observations. This strategy,  
54 while unavoidable, exposes the resulting dataset to possible biases  
55

56 due to inter-observer error. To control for this error, both observers  
57 independently measured 51 specimens taken from the identified col-  
58 lection of "Fiorentini" (individuals of both sexes who died in the Flor-  
59 entine area of Italy) during the second half of the 19th century AD,  
60 housed at the Museum of Natural History of the University of Flor-  
61 ence (Moggi-Cecchi, Pacciani, & Pinto-Cisternas, 1994) (Table 2). This  
62 sample was chosen due to the lack of available time for performing  
63 the same analysis in the field at Çatalhöyük, and on the basis of its  
64 excellent preservation, which maximized the sample size for error  
65 analysis. The relative amount of variance due to FA versus interob-  
66 server error was estimated by means of a two-way, mixed-model analysis  
67 of variance (ANOVA) with individuals as the random factor and sides as  
68 fixed. Traits showing nonsignificant FA when controlling for interob-  
69 server error were excluded from the dataset.

70 Further analyses were developed according to the following  
71 steps, based on the work of Palmer and Strobeck (2003):

- 72 1. The presence of outliers possibly inflating FA values was first  
73 assessed by visual inspection by R-L scatterplots, and tested by  
74 means of a Grubb's test.
- 75 2. Because FA analyses may be biased by the presence of antisym-  
76 metry or directional asymmetry, we determined the presence of  
77 deviations from normality of the difference (R-L) in each trait,  
78 after computing the average of two observation sessions. Direc-  
79 tional asymmetry and antisymmetry were tested by means of a  
80 D'Agostino-Pearson omnibus test (cf., Barrett et al., 2012).
- 81 3. We performed a two-way mixed-model analysis of variance  
82 (ANOVA) with individuals as the random factor and sides as fixed to  
83 assess the relative amount of variance due to FA versus measure-  
84 ment error (ME).
- 85 4. We used a Spearman test for testing the correlation between  
86  $|R - L|$  and  $|(R + L)/2|$  and possible allometric patterns in asymmetry.
- 87 5. To assess diachronic and demographic patterns in fluctuating  
88 asymmetry, we first calculated a trait-specific index  $|(R - L)|$   
89 (Palmer & Strobeck, 2003) for each individual. Owing to missing  
90 data, we did not attempt to calculate a composite index of  
91 FA. Because calculations of inter-observer error are influenced by  
92 the relative weight of the latter versus asymmetry, we also com-  
93 pared the asymmetry of each trait between the Fiorentini and  
94 Çatalhöyük samples by means of a Wilcoxon test. Owing to the  
95 small size of the Fiorentini sample, we did not attempt to test for  
96 differences controlling for age and/or sex.

97 Potential differences in FA between periods, age groups, and  
98 sexes were tested by means of Kruskal Wallis and Kendall Tau cor-  
99 relation tests. Because of unbalanced sample sizes, the Late and Final  
100

101 **TABLE 2** Size and age distribution of the sample used for the  
102 calculation of the inter-observer error

	F	M	NA	
≥18 years	23	24	0	
<18 years	2	1	1	
Total	25	25	1	51

103 F = females; M = males; NA = undetermined sex.

56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110

1 phases were combined into one subsample. Correlation tests were  
2 undertaken after chronologically ordering each phase from 1 (earliest)  
3 to 3 (latest).


4 The possible association between FA and age-at-death was  
5 tested using both the original age classes and after splitting the sam-  
6 ple into two groups: subadults (up to 19 years of age) and adults (from  
7 20 years of age). The latter strategy was dictated by the small sample  
8 sizes of individual age classes, and in order to include in our calcula-  
9 tion 31 adults in this study who were not assigned to specific age clas-  
10 ses due to their poor preservation. Correlation between FA and age  
11 classes was tested with a Kendall Tau correlation test, whereas possi-  
12 ble differences between adults and subadults were checked with a  
13 Wilcoxon test. Difference in asymmetry among traits was tested by  
14 means of a Wilcoxon test. Comparisons were made only between  
15 asymmetry values calculated using the same procedure  
16 (i.e., standardized or unstandardized by size). All statistical analyses  
17 were performed in R version 3.4.1, setting alpha at .05.

### 20 3 | RESULTS

#### 22 3.1 | Diachronic patterns of fertility, number of 23 buildings and individuals buried per building

24 The new juvenility indices calculated as part of this study (early = 9/29  
25 [.31], middle = 96/218 [.44], late/final = 20/77 [.26]) are consistent  
26 with those presented by Larsen and collaborators in their earlier study  
27 (Larsen et al., 2015) (early = .29, middle = 0.46, late = .29). These find-  
28 ings confirm a diachronic pattern of increased fertility from the early  
29 to middle phases of occupation of the site, followed by a decrease in  
30 the late-final phase.

31 The number of buildings and of individuals buried per building for  
32 the North, South, TP, and TPC areas of the site increase from the early  
33 occupation phase, reach a peak in the middle, and decrease during the  
34 late and final phases (Figure 1, Supporting Information Table S1), a  
35 pattern in overall agreement with previous studies and confirming the  
36



37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
**FIGURE 1** Temporal changes in number of buildings and buried individuals per building at Çatalhöyük. Data include all main excavation areas (north, south, TPC and TP), and only primary and primary disturbed burials

56 postulated increase in population size followed by a decline until the  
57 final abandonment of the site. The possibility of the above trend being  
58 a by-product of differences in the extent of excavation or differential  
59 preservation can be dismissed since excavation of the site is by strati-  
60 graphic layers (i.e., following the contours of the layered deposits and  
61 not arbitrary layers of similar thicknesses), and skeletal preservation is  
62 uniformly good to excellent throughout the stratigraphic layers of Çat-  
63 alhöyük. This means that buried individuals coming from the same  
64 phase are of the same relative date across the excavated area of the  
65 site. Note also that the perimeter of the excavated areas did not  
66 change throughout the field seasons.

#### 68 3.2 | Interobserver error, deviation from normality, 69 measurement error, and size effect

70 On the basis of the ANOVA results, five traits were excluded due to  
71 their high inter-observer error: UM1 (MD and BL), UCMD, LM1BL,  
72 and LCBL (Supporting Information Table S2). LM2MD and UCBL were  
73 further excluded due to the significant deviation from normality of  
74 their asymmetry values. In addition, the test for outliers by means of  
75 Grubb's test suggested exclusion of a number of observations (max:  
76 31, min: 7) from each trait-specific set.

77 All the remaining traits showed significant FA after controlling for  
78 measurement error (Table 3: Interaction Individual × Side), therefore  
79 ensuring that differences between sides in each trait were not due to  
80 directional asymmetry, antisymmetry, or observer error. Regression of  
81 asymmetry on size (see methods section) points to only two cases  
82 (LM2BL and LCMD) exhibiting a significant correlation between these  
83 variables (Table 4). Thus, we decided to use two different FA indices  
84 in the following analyses:  $|(R - L)|$  for traits showing no correlation  
85 between asymmetry and size (UM2MD, UM2BL, and LM1MD) and  
86  $|R - L| / |(R + L)/2|$  for traits with asymmetry and size correlations  
87 (LM2BL and LCMD).

88 A comparison of these indices between the Fiorentini sample and  
89 the Çatalhöyük dataset (Supporting Information Table S3) provides  
90 only one (LM1MD) significant difference.

#### 92 3.3 | Temporality, age-at-death, sex, and trait

93 A comparison between chronological phases indicates a general  
94 decrease in asymmetry with time, a pattern that reaches significance,  
95 however, only when considering sexes separately and only for LM2BL  
96 in females (Tables 5–7). As for age, the Kendall's Tau test on FA versus  
97 age classes highlights one trait (UM2MD) showing a positive correla-  
98 tion between asymmetry and age-at-death, and two traits (LM2BL,  
99 LCMD) characterized by a negative correlation (Table 8, Figure 2).  
100 When grouping the individuals into two age groups, subadults show  
101 significant higher asymmetry values for LM2BL, LM1MD, and LCMD  
102 (Table 9, Figure 3). As for sex, no trait shows significant differences  
103 between males and females, both considering each period separately  
104 and without chronological subdivisions (Table 10). Finally, when com-  
105 paring traits to each other, average asymmetry is significantly differ-  
106 ent between UM2MD versus LM1MD and between UM2BL and  
107 LM1MD (Table 11). Interestingly, these differences favor maxillary  
108 teeth in both cases.  
109  
110

**TABLE 3** Results of two-way mixed-model ANOVA (two repeated measurements)

		Df	SumSq	MeanSq	F	p
UM2MD	Individual	115.00	218.20	1.90	42.18	<.001
	Side	1.00	0.24	0.24	5.28	.023
	Individual x side	101.00	11.01	0.11	2.42	<.001
	Residuals	226.00	10.17	0.05		
UM2BL	Individual	109.00	226.39	2.08	76.88	<.001
	Side	1.00	0.28	0.28	10.40	.001
	Individual x side	94.00	7.00	0.07	2.76	<.001
	Residuals	213.00	5.75	0.03		
LM2BL	Individual	99.00	105.56	1.07	185.68	<.001
	Side	1.00	0.01	0.01	0.99	.321
	Individual x side	81.00	3.94	0.05	8.47	<.001
	Residuals	183.00	1.05	0.01		
LM1MD	Individual	102.00	218.59	2.14	99.54	<.001
	Side	1.00	0.01	0.01	0.52	.471
	Individual x side	89.00	4.64	0.05	2.42	<.001
	Residuals	202.00	4.35	0.02		
LCMD	Individual	102.00	60.61	0.59	175.01	<.001
	Side	1.00	0.07	0.07	20.33	<.001
	Individual x side	79.00	2.09	0.03	7.79	<.001
	Residuals	183.00	0.62	0.00		

UM2MD; second upper molar; LM2: second lower molar; LM1: first lower molar; LC: lower canine; MD = mesio-distal diameter; BL = bucco-lingual diameter.

## 4 | DISCUSSION

Before discussing the results of this study, we first address some theoretical and methodological issues regarding FA studies. A straightforward interpretation of FA as a measure of stress has been recognized as inadequate (Dongen, 2006; Leamy & Klingenberg, 2005), especially due to the current poor understanding of the genetic background of developmental instability, and the difficulty of estimating the latter from single traits. However, for this study, the risk of a genetic bias should be reduced by focusing on a single population (note also that patterns of nonmetric dental traits suggest a substantial lack of gene flow at Çatalhöyük (Pilloud & Larsen, 2011). Another issue, possibly more serious, is represented by the sample used for the test of inter-observer error. Although the use of a population different from that of Çatalhöyük was unavoidable, it is obvious that the use of a single observer, and the test for any measurement error on a single sample would have been ideal. To better accommodate these potential sources of bias inter-observer error as measured in this study is quantified with respect to the amount of variance due to asymmetry. In this sense, our error estimates appear rather solid in light of the large overlap in asymmetry values between the Çatalhöyük and Fiorentini samples.

Also, the exclusion of a number of traits characterized by unacceptable levels of inter-observer error led to a sensible decrease of sample size, already constrained by the need to exclude those teeth not preserving maximum crown diameters. Dental pathological conditions and loss of dental hard tissue due to wear (e.g., due to attrition and abrasion), represent two of the principal causes that affect the dimensions of a tooth. Mastication and the use of teeth as tools in

working of materials (paramastication), are among the major factors of abrasion (e.g., Larsen, 1985; Molnar, 1972).

Both issues (exclusion of traits due to interobserver error and exclusion of individuals due to missing data) pose the question of the real representativeness of our data for the whole population. This is especially relevant for the first and last chronological phases of Çatalhöyük, represented in some cases by relatively few individuals, but demographically extremely relevant (representing the beginning and end of occupation). In addition, the small and unbalanced sample sizes prevented the use of a composite index, with a resulting decrease in our ability to capture nuanced but possibly relevant asymmetry patterns. Finally, the reduced sample sizes also hindered detailed analysis of developmental trajectories of asymmetry, and of polar versus anti-polar teeth sensitivity to stress. Because of these (unavoidable) problems, this work must be considered as a preliminary test of our starting hypotheses, to be integrated in the future by the inclusion of additional data (e.g., enamel hypoplasia, Harris lines). On the other hand, when considered with the caution and criticism required by the

**TABLE 4** Correlation between asymmetry and size

	n	Spearman $\rho$	p
LM1MD	108	.00	.977
LCMD	89	.49	<.001
LM2BL	87	.51	<.001
UM2BL	111	.08	.426
UM2MD	114	.13	.163

UM2MD; second upper molar; LM2: second lower molar; LM1: first lower molar; LC: lower canine; MD = mesio-distal diameter; BL = bucco-lingual diameter.

**TABLE 5** Results of Kruskal-Wallis test on asymmetry values between periods

	F							M						
	Early		Middle		Late-final		p	Early		Middle		Late-final		p
	n	Median	n	Median	n	Median		n	Median	n	Median	n	Median	
UM2MD	1	0.430	21	0.255	5	0.070	0.247	4	0.540	18	0.368	4	0.155	0.158
UM2BL	1	0.325	21	0.135	4	0.283	0.477	4	0.203	18	0.143	3	0.135	0.536
LM2BL	1	0.054	17	0.034	7	0.007	0.017 (0.011 2vs3)	3	0.043	13	0.028	7	0.011	0.307
LM1MD	1	1.170	16	0.250	7	0.350	0.307	0		14	0.185	7	0.155	0.279
LCMD	3	0.046	18	0.025	7	0.007	0.425	2	0.033	15	0.011	6	0.012	0.811

M = males; F = females (NA individuals not included).

mentioned issues, we believe that our results are nonetheless able to identify some general patterns. These can be summarized as:

1. Homogenous levels of asymmetry throughout the occupation of the site, with the only diachronic change (i.e., decrease over time) observed in females for a single trait only.
2. Higher asymmetry in subadults versus adults.
3. No differences in asymmetry between the sexes.

Our first hypothesis postulated a decrease in dental asymmetry across chronological phases, and, specifically, an inverse correlation between developmental stress and postulated population density at the site. Results of both the Kruskal-Wallis and the Kendall tests, while indicating a slight decrease in FA with time, point to only a single case, LM2BL in females, for which this trend is significant (Table 5). Among the factors underpinning these results, three may play a relatively important role: (1) variable developmental buffering among different teeth, (2) a complex relationship between population density and developmental stress, and (3) a lack of correlation between calculated population size and inferred population density at the site.

Differences in asymmetry between tooth classes have been previously reported and their developmental background extensively discussed (Bailit, Workman, Niswander, & Maclean, 1970; Garn & Bailey, 1977; Garn, Lewis, & Kerewsky, 1966; Garn et al., 1967; Guatelli-Steinberg, Sciulli, & Edgar, 2006; Harris & Nweeia, 1980; Hershkovitz Livshits, Moskona, Arensburg, & Kobylansky, 1993; Perzigian, 1977; Sofaer, Bailit, & MacLean, 1971). Among the causes of such variability, one can mention differences in developmental stability of polar versus distal teeth (Hershkovitz et al., 1993; Townsend, Brook, Yong, & Hughes, 2015), upper versus lower dentitions (see Guatelli-Steinberg et al., 2006; Hershkovitz et al., 1993), and between MD and BL dimensions on the same tooth (Kolakowski & Bailit, 1981; Potter & Nance, 1976).

An additional possibility to consider may also be a lack of marked differences in stressors between periods. Thus, different degrees of developmental stability contributed to obscure possible diachronic patterns of developmental disruption. However, acknowledging such differences among traits, our data clearly indicate, overall, an absence of chronological differences in FA at Çatalhöyük. This result may shed some light on the biological correlates of local demographic changes in the prehistoric community. Bioarchaeologists postulate a link between the increase in population densities and worsening life

conditions (e.g., unprecedented levels of zoonotic diseases, infections, and malnutrition) characterizing the Neolithic transition (Armélagos et al., 2005; Larsen, 1995). Investigations on a smaller scale focused on local shifts in population densities are fewer, and their results less clear. From this perspective, skeletal and biochemical data from Çatalhöyük depict a complex picture. That is, diachronic patterns of periostosis show a sudden decrease in skeletal lesions in the late phase of occupation of the site, whereas estimates of stature and body mass fail to highlight changes in these parameters through time. At the same time, juvenile  $\delta^{15}\text{N}$  values are consistent with a relatively "premature" start of weaning (at about one year and-a-half—Larsen et al., 2015). Previous interpretations of these data postulated a diachronic worsening of life conditions due to increasing population densities, coupled with a weak immune response to environmental stressors due to early weaning. The parallel lack of abrupt deviations in body size, body mass, and cortical bone mass from those expected from a "healthy" population was further interpreted as the result of the access to cereal grains and animal (caprine) proteins, which would have acted as a buffer against these potential sources of stress (Larsen et al., 2015). Our results confirm the lack of marked temporal changes in developmental disruption at the site (therefore in agreement with data on stature), and are in only slight agreement with previous hypotheses about a decrease of exposure to stressors during the late phase of occupation (as evidenced by diachronic frequencies of periostosis). This result may strengthen the hypothesis of Larsen et al. (2015) about the buffering effect of diet versus the potential developmental effects of population crowding during the early-mid occupation of Çatalhöyük. Results on FA suggest a complex scenario, namely the possibility of a moderate amount of environmental stressors throughout the entire occupation of the site (therefore even in a situation of decreased population density), though mitigated by the effects of diet. A comparison of our data with those from previous work on dental FA is unwarranted due to methodological and

**TABLE 6** Results of Kendall's test on asymmetry versus periods

	M			F		
	n	Tau	p	n	Tau	p
UM2MD	26	-.312	.053	27	-.259	.110
UM2BL	25	-.109	.511	26	.011	.948
LM2BL	23	-.230	.178	25	-.488	.004
LM1MD	21	-.203	.279	24	-.068	.693
LCMD	23	-.066	.704	28	-.171	.270

M = males; F = females (NA individuals not included).

**TABLE 7** Results of Kruskal-Wallis and Kendall's test on asymmetry versus periods

	Early		Middle		Late-final		<i>p</i> (Wilcoxon)	Tau	<i>P</i> Kendall
	<i>n</i>	Median	<i>n</i>	Median	<i>n</i>	Median			
UM2MD	9	.155	78	.240	20	.225	.530	-.088	.276
UM2BL	9	.130	77	.185	18	.225	.567	.075	.368
LM2BL	8	.044	48	.036	22	.014	.181	-.150	.121
LM1MD	8	.335	66	.250	24	.283	.397	-.042	.626
LCMD	9	.053	53	.029	20	.015	.387	-.054	.571

Pooled sexes—M, F, and NA. UM2MD, second upper molar; LM2: second lower molar; LM1: first lower molar; LC: lower canine; MD = mesio-distal diameter; BL = bucco-lingual diameter. UM2MD, UM2BL, LM1MD asymmetry index = |(R-L)|. LM2BL, LCMD asymmetry index = |R-L| / |(R + L)/2|.

biological discrepancies. First, the comparability between results would be hampered by the use of different statistical protocols from the one adopted here. Moreover, comparing FA values between populations representing different gene pools (and developmental stability) and environments (and environmental stressors) would be problematic.

A better approach would be the analysis of dental FA on samples predating the occupation of Çatalhöyük and theoretically representing the same biological population (the ideal candidate being the nearby but the earlier site of Boncuklu—see Baird et al., 2018). Such a study would provide essential data for further testing our interpretations.

A potential issue affecting all these reconstructions is the hypothesized correspondence between population size and population density at Çatalhöyük. It is indeed likely that the extent of occupation at the community changed through time (cf., Cessford, 2005), and these fluctuations had significant consequences for population densities and their biological correlates. Furthermore, the observed lack of correspondence between the frequency of periostosis and the temporal trends in stature and FA is hardly unexpected, given both the multifaceted nature of physiological stress (Reitsemá & McIlvaine, 2014; Temple & Goodman, 2014), and the complex and not necessarily co-occurring series of variables influencing these features (e.g., time and duration of disruption, type of stress, genetic and epigenetic factors).


Our second and third hypotheses postulated, respectively, the presence of sexual differences in FA, and a negative correlation between FA and age-at-death. Comparisons between males and females fail to highlight any significant contrast in asymmetry values, a result that contradicts our expectations based on a suspected greater developmental buffering (Stinson, 1985) and more robust genetic control of odontogenesis (Garn, Lewis, & Kerewsky, 1965; Garn et al., 1967) in females, and on clinical and epidemiological data

**TABLE 8** Results of Kendall's test on asymmetry versus age classes (unspecified adults >20 not included)

	<i>n</i>	Tau	<i>p</i>
UM2MD	104	.16	.0348
UM2BL	102	.03	.7324
LM2BL	77	-.24	.0043
LM1MD	97	-.14	.0778
LCMD	79	-.25	.0033

indicating a lower immunocompetence in males (Guerra-Silveira & Abad-Franch, 2013; Jansen, Stark, Schneider, & Schoneberg, 2007; Klein, 2000, 2004; Leone et al., 2004; Markle & Fish, 2014; Muenchhoff & Goulder, 2014; Owens, 2002; Pennell, Galligan, & Fish, 2012). Though apparently counter-intuitive, our results are nonetheless consistent with those from other studies (e.g., Guatelli-Steinberg et al., 2006; Kieser et al., 1986). In a recent study of dental fluctuating asymmetry in a contemporary African-American population, Guatelli-Steinberg et al. (2006) found that the only significant difference between sexes was due to higher FA values for the permanent mandibular canine in females, with no apparent sexual dimorphism for the maxillary canine. In discussing these contradictory results, these authors postulated that sexual differences in developmental buffering might have obscured by the presence of relatively high levels of developmental noise (and the possible presence of sex-biased childcare practices).

In the present analysis, we previously mentioned the possibility of a relatively homogenous exposure to developmental stressors throughout the occupation of the site as the main factor responsible of the weak diachronic decrease in FA. Isotope data also indicates a lack of differences in diet between the sexes at Çatalhöyük (females:



**FIGURE 2** Results of Kendall's test of dental fluctuating asymmetry vs. age classes. Significant correlations are all negative (see also Table 5)


**TABLE 9** Results of Wilcoxon test on asymmetry among adults and subadults

	Adult		Subadult		p (Wilcoxon)
	n	Median	n	Median	
UM2MD	56	.270	58	.213	.0874
UM2BL	54	.178	57	.210	.6904
LM2BL	57	.020	31	.053	.0076
LM1MD	53	.220	55	.330	.0097
LCMD	55	.012	35	.050	.0014

UM2MD: second upper molar; LM2: second lower molar; LM1: first lower molar; LC: lower canine; MD = mesio-distal diameter; BL = bucco-lingual diameter. UM2MD, UM2BL, LM1MD asymmetry index = |(R-L)|. LM2BL, LCMD asymmetry index = |R-L| / |(R + L)/2|.

$\delta^{13}\text{C} = -18.8\%$ ,  $\delta^{15}\text{N} = 12.6\%$ ; males:  $\delta^{13}\text{C} = -18.6\%$ ,  $\delta^{15}\text{N} = 12.7\%$  (Larsen et al., 2015), therefore suggesting the absence of sex-biased childcare practices in this community. Altogether, these results strengthen the hypothesis of a weak influence of population size on actual exposure to developmental stress. Rather, applying the argument of Guatelli-Steinberg et al. (2006) we suggest that continuous exposure to relatively high developmental stressors obscured possible sexual differences in immunocompetence and fragility.

With regard to our third hypothesis, patterns of FA across age classes and between adults and subadults tend to confirm the expected correlation between high FA and premature mortality. Several studies identify an association between FA and various health variables, such as parasitism, chromosomal abnormalities, and genetic diseases (Kieser, Groeneveld, & Da Silva, 1997; Møller, 2006; Thornhill & Møller, 1997), and the link between assaults on the immune system and the development of asymmetrical phenotypes (Fair, Hansen, & Ricklefs, 1999; Møller, 2006; Whitaker & Fair, 2002). In addition, age-at-death profiles of individuals showing dental effects of

**FIGURE 3** Differences between medians (DBM) of adults and subadults, and p values of the Wilcoxon tests. Note that all significant differences show higher FA in subadults**TABLE 10** Results of Wilcoxon test on asymmetry between sexes

	F		M		p (Wilcoxon)
	n	Median	n	Median	
UM2MD	29	.255	27	.345	.1584
UM2BL	28	.210	26	.153	.9931
LM2BL	30	.014	25	.021	.7353
LM1MD	27	.250	23	.175	.0643
LCMD	29	.016	25	.011	.4932

M = males; F = females (NA individuals not included)

high exposure to early life stressors (i.e., enamel defects) are overall consistent with a link between developmental disruptions and higher fragility (Amoroso, Garcia, & Cardoso, 2014; Armelagos, Goodman, Harper, & Blakey, 2009). The observed contrast between adults and subadults in FA at Çatalhöyük is relevant since it supports the interpretation of this parameter as a marker of developmental stress. Accordingly, it justifies the socio-ecological interpretation of the data proposed in the present investigation. Also, note that this result is in agreement with the age distribution and relative prevalence of periostitis at Çatalhöyük (higher incidence in subadults), as well as with biochemical reconstructions suggesting a condition of weakened immunocompetence in juveniles (Larsen et al., 2015).

## 5 | CONCLUSION

We examined the relationship between diachronic changes in population size, relative exposure to developmental stressors, and patterns of dental fluctuating asymmetry in the Neolithic population of Çatalhöyük (Turkey, 7,100–5,950 cal BC). Though preliminary, our results are consistent with the presence of developmental stressors throughout the occupation of the site, though with a slight improvement in living conditions during the latest periods of occupation. Moreover, our data agree with the results of previous research suggesting the important role diet played as a developmental buffer at Çatalhöyük, and overall demonstrates the usefulness of dental fluctuating asymmetry as an analytical tool in biocultural reconstructions of health and living conditions. More work is needed in order to further test our hypotheses (e.g., combine data on dental fluctuating asymmetry with patterns of enamel hypoplasia and Harris lines). On a more general level, our study highlights the complex and multifaceted nature of developmental stress and the usefulness of a multipronged approach

**TABLE 11** Results of Wilcoxon test on asymmetry between traits

		Z	p
UM2MD	vs. UM2BL	1.41	.159
UM2MD	vs. LM1MD	-2.05	.040
UM2BL	vs. LM1MD	-3.39	.001
LM2BL	vs. LCMD	-.21	.833

UM2MD: second upper molar; LM2: second lower molar; LM1: first lower molar; LC: lower canine; MD = mesio-distal diameter; BL = bucco-lingual diameter. UM2MD, UM2BL, LM1MD asymmetry index = |(R-L)|. LM2BL, LCMD asymmetry index = |R-L| / |(R + L)/2|. Only asymmetry indices calculated with the same formula are compared.

1 to developmental studies, while stressing the need for new palaeode-  
2 mographic models that permit more informed discussions of the links  
3 between environment, growth, and development in the past.

#### 6 ACKNOWLEDGMENTS

7 The authors thank the Editor in Chief and the two anonymous  
8 reviewers for their detailed comments and suggestions, which  
9 improved the quality of this manuscript. The authors also thank  
10 Jacopo Moggi-Cecchi (Università degli Studi di Firenze) for his com-  
11 ments on the manuscript and Monica Zavattaro for granting access to  
12 the identified skeletal collection of the Museum of Natural History  
13 (section of Anthropology and Ethnology), Università degli Studi di  
14 Firenze. This study has received financial support from the National  
15 Geographic Society, the European Union's Horizon 2020 research and  
16 innovation program under the Marie Skłodowska-Curie grant agree-  
17 ment No 752626 (ID), and from the French State within the frame-  
18 work of the "Investments for the future" Program, IdEx Bordeaux,  
19 reference ANR-10-IDEX-03-02 (CJK).

#### 21 ORCID

22 Marco Milella  <http://orcid.org/0000-0003-1027-6601>

23 Clark Spencer Larsen  <http://orcid.org/0000-0002-6905-8417>

#### 26 REFERENCES

27 Albert, A., & Greene, D. (1999). Bilateral asymmetry in skeletal growth and  
28 maturation as an indicator of environmental stress. *American Journal of*  
29 *Physical Anthropology*, 110, 341–349. [https://doi.org/10.1002/\(SICI\)  
30 \*1096-8644\(199911\)110:3<341::AID-AJPA6>3.0.CO;2-8\*  
31 Amoroso, A., Garcia, S. J., & Cardoso, H. F. V. \(2014\). Age at death and lin-  
32 ear enamel hypoplasias: Testing the effects of childhood stress and  
33 adult socioeconomic circumstances in premature mortality. \*American\*  
34 \*Journal of Human Biology\*, 26, 461–468. \[https://doi.org/10.1002/ajhb.  
35 \\*22547\\*  
36 Armelagos, G. J., Brown, P. J., & Turner, B. \\(2005\\). Evolutionary, historical  
37 and political economic perspectives on health and disease. \\*Social Sci-\\*  
38 \\*ence and Medicine\\*, 61, 755–765. \\[https://doi.org/10.1016/j.socscimed.  
39 \\\*2004.08.066\\\*  
40 Armelagos, G. J., Goodman, A. H., Harper, K. N., & Blakey, M. L. \\\(2009\\\).  
41 Enamel hypoplasia and early mortality: Bioarchaeological support for  
42 the barker hypothesis. \\\*Evolutionary Anthropology\\\*, 18, 261–271.  
43 <https://doi.org/10.1002/evan.20239>  
44 Bailit, H. L., Workman, P. L., Niswander, J. D., & Maclean, C. J. \\\(1970\\\). Den-  
45 tal asymmetry as an indicator of genetic and environmental conditions  
46 in human populations. \\\*Human Biology\\\*, 42, 626–638.  
47 Baird, D. \\\(2005\\\). The history of settlement and social landscapes in the  
48 early Holocene in the Çatalhöyük area. In I. Hodder \\\(Ed.\\\), \\\*Çatalhöyük\\\*  
49 \\\*Perspectives: Reports from the 1995–1999 Seasons\\\* \\\(pp. 55–74\\\). Cam-  
50 bridge, UK: McDonald Institute for Archaeological Research.  
51 Baird, D., Fairbairn, A., Jenkins, E., Martin, L., Middleton, C., Pearson, J., ...  
52 Elliott, S. \\\(2018\\\). Agricultural origins on the Anatolian plateau. \\\*Proceed-\\\*  
53 \\\*ings of the National Academy of Sciences of the United States of America\\\*,  
54 \\\*115\\\(14\\\)\\\*, 3077–3086.  
55 Barrett, C. K., Guatelli-Steinberg, D., & Scullin, P. W. \\\(2012\\\). Revisiting den-  
56 tal fluctuating asymmetry in Neandertals and modern humans. \\\*Ameri-\\\*  
57 \\\*can Journal of Physical Anthropology\\\*, 149, 193–204. \\\[https://doi.org/10.\\\]\\\(https://doi.org/10.1002/ajpa.22107\\\)  
58 \\\*1002/ajpa.22107\\\*  
59 Bayliss, A., Brock, F., Farid, S., Hodder, I., Southon, J., & Taylor, R. E.  
60 \\\(2015\\\). Getting to the bottom of it all: A Bayesian approach to dating  
61 the start of Catalhöyük. \\\*Journal of World Prehistory\\\*, 28, 1–26. \\\[https://\\\]\\\(https://doi.org/10.1007/s10963-015-9083-7\\\)  
62 \\\*doi.org/10.1007/s10963-015-9083-7\\\*\\]\\(https://doi.org/10.1016/j.socscimed.2004.08.066\\)\]\(https://doi.org/10.1002/ajhb.22547\)](https://doi.org/10.1002/(SICI)1096-8644(199911)110:3<341::AID-AJPA6>3.0.CO;2-8)

63 Bigoni, L., Krajcicek, V., Sladek, V., Veleminsky, P., & Veleminska, J. (2013). 56  
64 Skull shape asymmetry and the socioeconomic structure of an early 57  
65 medieval central European society. *American Journal of Physical Anthro-*  
66 *pology*, 150, 349–364. <https://doi.org/10.1002/ajpa.22210>  
67 Bocquet-Appel, J.-P. (2002). Paleoanthropological traces of a neolithic 59  
60 demographic transition. *Current Anthropology*, 43, 637–650.  
61 Bocquet-Appel, J.-P. (2008). Explaining the Neolithic demographic transi- 62  
63 tion. In J.-P. Bocquet-Appel & O. Bar-Yosef (Eds.), *The Neolithic demo-*  
64 *graphic transition and its consequences* (pp. 35–55). New York, NY:  
65 Springer.  
66 Bocquet-Appel, J.-P. (2011). When the world's population took off: The 64  
65 springboard of the Neolithic demographic transition. *Science*, 333,  
66 560–561. <https://doi.org/10.1126/science.1208880>  
67 Buikstra, J. E., & Ubelaker, D. H. (1994). *Standards for data collection from*  
68 *human skeletal remains*. Fayetteville, AR: Arkansas Archaeological  
69 Survey.  
70 Cardoso, H., & Gomes, J. (2009). Trends in adult stature of peoples who 69  
71 inhabited the modern Portuguese territory from the Mesolithic to the  
72 late 20th century. *International Journal of Osteoarchaeology*, 19,  
73 711–725. <https://doi.org/10.1002/oa.991>  
74 Cessford, C. (2005). Estimating the Neolithic population of Çatalhöyük. In 72  
73 I. Hodder (Ed.), *Inhabiting Çatalhöyük: Reports from the 1995–1999 sea-*  
74 *sons* (pp. 323–328). Cambridge, UK: McDonald Institute for Archaeo-  
75 logical Research.  
76 Cohen, M. N., & Armelagos, G. J. (1984). *Paleopathology at the origins of*  
77 *agriculture*. Orlando, FL: Academic Press.  
78 Cohen, M. N., & Crane-Kramer, G. M. M. (2007). *Ancient health: Skeletal*  
79 *indicators of agricultural and economic intensification*. Gainesville, FL:  
80 University Press.  
81 Costa, R. L., Jr. (1986). Asymmetry of the mandibular condyle in Haida 79  
82 Indians. *American Journal of Physical Anthropology*, 70, 119–123.  
83 <https://doi.org/10.1002/ajpa.1330700116>  
84 Dahlberg, A. A. (1945). The changing dentition of man. *The Journal of the*  
85 *American Dental Association*, 32, 676–690. doi:10.14219/jada.  
86 archive.1945.0112  
87 De Leon, V. B. (2007). Fluctuating asymmetry and stress in a medieval 83  
84 Nubian population. *American Journal of Physical Anthropology*, 132,  
85 520–534. <https://doi.org/10.1002/ajpa.20549>  
86 Doherty, C. (2013). Sourcing Çatalhöyük's clays. In I. Hodder (Ed.), *Substan-*  
87 *tive Technologies at Çatalhöyük: Reports from the 2000–2008 seasons*  
88 (pp. 51–66). Los Angeles, CA: Cotsen Institute for Archaeology.  
89 ~~Dongen, S. V. (2006). Fluctuating asymmetry and developmental instability 88~~  
90 ~~in evolutionary biology: Past, present and future. *Journal of Evolution-*~~  
91 ~~*ary Biology*, 19, 1727–1743. [https://doi.org/10.1111/j.1420-9101.](https://doi.org/10.1111/j.1420-9101.2006.01175.x)~~  
92 ~~*2006.01175.x*~~  
93 Doyle, W. J., & Johnston, O. (1977). On the meaning of increased fluctuat- 92  
94 ing dental asymmetry: A cross populational study. *American Journal of*  
95 *Physical Anthropology*, 46, 127–134.  
96 Düring, B. S. (2001). Social dimensions in the architecture of Neolithic Çat- 93  
94 alhöyük. *Anatolian Studies*, 51, 1–18. [https://doi.org/10.](https://doi.org/10.2307/3643025)  
95 *2307/3643025*  
96 Fair, J. M., Hansen, E. S., & Ricklefs, R. E. (1999). Growth, developmental 96  
97 stability and immune response in juvenile Japanese quails (*Coturnix*  
98 *coturnix japonica*). *Proceedings of the Royal Society of London. Series B,*  
99 *Biological Sciences*, 266, 1735–1742. [https://doi.org/10.1098/rspb.  
100 \*1999.0840\*  
101 Garfinkel, Y. \(1987\). Burnt lime products and social implications in the 100  
102 pre-pottery Neolithic B villages of the near east. \*Paléorient\*, 13, 69–76.  
103 Garn, S. M., & Bailey, S. M. \(1977\). The symmetrical nature of bilateral 101  
104 asymmetry \( \$\delta\$ \) of deciduous and permanent teeth. \*Journal of Dental\*  
105 \*Research\*, 56, 1422–1422.  
106 Garn, S. M., Lewis, A. B., & Kerewsky, R. S. \(1965\). Genetic, nutritional, and 104  
105 maturational correlates of dental development. \*Journal of Dental\*  
106 \*Research\*, 44, 228–242.  
107 Garn, S. M., Lewis, A. B., & Kerewsky, R. S. \(1966\). The meaning of bilateral 106  
108 asymmetry in the permanent dentition. \*The Angle Orthodontist\*, 36,  
109 55–62.  
110 Garn, S. M., Lewis, A. B., & Kerewsky, R. S. \(1967\). Buccolingual size asym- 107  
108 metry and its developmental meaning. \*The Angle Orthodontist\*, 37,  
109 186–193.](https://doi.org/10.1098/rspb.1999.0840)

- Garn, S. M., Lewis, A. B., Kerewsky, R. S., & Jegart, K. (1965). Sex differences in Intraindividual tooth-size communalities. *Journal of Dental Research*, 44, 476–479.
- Gawlikowska-Sroka, A., Dabrowski, P., Szczurowski, J., Dzieciolowska-Baran, E., & Staniowski, T. (2017). Influence of physiological stress on the presence of hypoplasia and fluctuating asymmetry in a medieval population from the village of Sypniewo. *International Journal of Paleopathology*, 19, 43–52. <https://doi.org/10.1016/j.ijpp.2017.10.002>
- Gawlikowska-Sroka, A., Dabrowski, P., Szczurowski, J., & Staniowski, T. (2013). Analysis of interaction between nutritional and developmental instability in mediaeval population in Wrocław. *Anthropological Review*, 76, 51–62. <https://doi.org/10.2478/anre-2013-0009>
- González-José, R., Ramírez-Rozzi, F., Sardi, M., Martínez-Abadías, N., Hernández, M., & Pucciarelli, H. M. (2005). Functional-cranial approach to the influence of economic strategy on skull morphology. *American Journal of Physical Anthropology*, 128, 757–771. <https://doi.org/10.1002/ajpa.20161>
- Goodman, A. H. (1989). Dental enamel hypoplasias in prehistoric populations. *Advances in Dental Research*, 3, 265–271.
- Goodman, A. H. (1993). On the interpretation of health from skeletal remains. *Current Anthropology*, 34, 281–288. <https://doi.org/10.1086/204170>
- Goodman, A. H., & Armelagos, G. J. (1985). Factors affecting the distribution of enamel hypoplasias within the human permanent dentition. *American Journal of Physical Anthropology*, 68, 479–493. <https://doi.org/10.1002/ajpa.1330680404>
- Greene, D. L. (1984). Fluctuating dental asymmetry and measurement error. *American Journal of Physical Anthropology*, 65, 283–289. <https://doi.org/10.1002/ajpa.1330650308>
- Guatelli-Steinberg, D., Sciuilli, P. W., & Edgar, H. H. (2006). Dental fluctuating asymmetry in the Gullah: Tests of hypotheses regarding developmental stability in deciduous vs. permanent and male vs. female teeth. *American Journal of Physical Anthropology*, 129, 427–434. <https://doi.org/10.1002/ajpa.20237>
- Guerra-Silveira, F., & Abad-Franch, F. (2013). Sex bias in infectious disease epidemiology: Patterns and processes. *PLoS One*, 8, e62390. <https://doi.org/10.1371/journal.pone.0062390>
- Harris, E. F., & Nweeia, M. T. (1980). Dental asymmetry as a measure of environmental-stress in the Ticuna Indians of Colombia. *American Journal of Physical Anthropology*, 53, 133–142. <https://doi.org/10.1002/ajpa.1330530118>
- Hawks, J., Wang, E. T., Cochran, G. M., Harpending, H. C., & Moyzis, R. K. (2007). Recent acceleration of human adaptive evolution. *Proceedings of the National Academy of Sciences of the United State of America*, 104, 20753–20758. <https://doi.org/10.1073/pnas.0707650104>
- Hershkovitz, I., Livshits, G., Moskona, D., Arensburg, B., & Kobylansky, E. (1993). Variables affecting dental fluctuating asymmetry in human isolates. *American Journal of Physical Anthropology*, 91, 349–365. <https://doi.org/10.1002/ajpa.1330910308>
- Hillson, S., Fitzgerald, C., & Flinn, H. (2005). Alternative dental measurements: Proposals and relationships with other measurements. *American Journal of Physical Anthropology*, 126, 413–426. <https://doi.org/10.1002/ajpa.10430>
- Hodder, I. (2014a). Mosaics and networks: The social geography at Çatalhöyük. In I. Hodder (Ed.), *Integrating Çatalhöyük: Themes from the 2000–2008 seasons* (pp. 149–167). Los Angeles, CA: Cotsen Institute of Archaeology.
- Hodder, I. (2014b). Çatalhöyük: The leopard changes its spots. A summary of recent work. *Anatolia Studies*, 64, 1–22. <https://doi.org/10.1017/S0066154614000027>
- Holden, C., & Mace, R. (1997). Phylogenetic analysis of the evolution of lactose digestion in adults. *Human Biology*, 69, 605–628.
- Hoover, K. C., Corruccini, R. S., Bondioli, L., & Macchiarelli, R. (2005). Exploring the relationship between hypoplasia and odontometric asymmetry in Isola sacra, an imperial roman necropolis. *American Journal of Human Biology*, 17, 752–764. <https://doi.org/10.1002/ajhb.20436>
- Hoover, K. C., & Matsumura, H. (2008). Temporal variation and interaction between nutritional and developmental instability in prehistoric Japanese populations. *American Journal of Physical Anthropology*, 137, 469–478. <https://doi.org/10.1002/ajpa.20892>
- Jansen, A., Stark, K., Schneider, T., & Schoneberg, I. (2007). Sex differences in clinical leptospirosis in Germany: 1997–2005. *Clinical Infectious Diseases*, 44, e69–e72. <https://doi.org/10.1086/513431>
- Katz, D. C., Grote, M. N., & Weaver, T. D. (2017). Changes in human skull morphology across the agricultural transition are consistent with softer diets in preindustrial farming groups. *Proceedings of the National Academy of Sciences of the United State of America*, 114, 9050–9055. <https://doi.org/10.1073/pnas.1702586114>
- Kieser, J. A., Groeneveld, H. T., & Da Silva, P. C. (1997). Dental asymmetry, maternal obesity, and smoking. *American Journal of Physical Anthropology*, 102, 133–139. [https://doi.org/10.1002/\(SICI\)1096-8644\(199701\)102:1<133::AID-AJPA11>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1096-8644(199701)102:1<133::AID-AJPA11>3.0.CO;2-1)
- Kieser, J. A., Groeneveld, H. T., & Preston, C. B. (1986). Fluctuating dental asymmetry as a measure of odontogenic canalization in man. *American Journal of Physical Anthropology*, 71, 437–444. <https://doi.org/10.1002/ajpa.1330710407>
- Klein, S. L. (2000). The effects of hormones on sex differences in infection: From genes to behavior. *Neuroscience and Biobehavioral Reviews*, 24, 627–638.
- Klein, S. L. (2004). Hormonal and immunological mechanisms mediating sex differences in parasite infection. *Parasite Immunology*, 26, 247–264. <https://doi.org/10.1111/j.0141-9838.2004.00710.x>
- Klingenberg, C. P. (2003). A developmental perspective on developmental instability: Theory, models and mechanisms. In M. Polak (Ed.), *Developmental instability: Causes and consequences* (pp. 14–34). Oxford, UK: Oxford University Press.
- Kolakowski, D., & Bailit, H. L. (1981). A differential environmental effect on human anterior tooth size. *American Journal of Physical Anthropology*, 54, 377–381. <https://doi.org/10.1002/ajpa.1330540311>
- Kuijt, I., & Goring-Morris, N. (2002). Foraging, farming, and social complexity in the pre-pottery Neolithic of the southern Levant: A review and synthesis. *Journal of World Prehistory*, 16, 361–440. <https://doi.org/10.1023/A:1022973114090>
- Kujanova, M., Bigoni, L., Velemínska, J., & Velemínský, P. (2008). Limb bones asymmetry and stress in medieval and recent populations of Central Europe. *International Journal of Osteoarchaeology*, 18, 476–491. <https://doi.org/10.1002/oa.958>
- Laland, K. N., Odling-Smee, J., & Myles, S. (2010). How culture shaped the human genome: Bringing genetics and the human sciences together. *Nature Reviews. Genetics*, 11, 137–148. <https://doi.org/10.1038/nrg2734>
- Larsen, C. S. (1985). Dental modification and tool use in the western Great Basin. *American Journal of Physical Anthropology*, 67, 393–402. <https://doi.org/10.1002/ajpa.1330670411>
- Larsen, C. S. (1995). Biological changes in human-populations with agriculture. *Annual Review of Anthropology*, 24, 185–213. <https://doi.org/10.1146/annurev.an.24.100195.001153>
- Larsen, C. S., Hillson, S. W., Boz, B., Pilloud, M. A., Sadvari, J. W., Agarwal, S. C., ... Knüsel, C. J. (2015). Bioarchaeology of Neolithic Çatalhöyük: Lives and lifestyles of an early farming society in transition. *Journal of World Prehistory*, 28, 27–68. <https://doi.org/10.1007/s10963-015-9084-6>
- Leamy, L. J., & Klingenberg, C. P. (2005). The genetics and evolution of fluctuating asymmetry. *Annual Review of Ecology, Evolution, and Systematics*, 36, 1–21. <https://doi.org/10.1146/annurev.ecolsys.36.102003.152640>
- Leone, M., Honstetter, A., Lepidi, H., Capo, C., Bayard, F., Raoult, D., & Mege, J.-L. (2004). Effect of sex on *Coxiella burnetii* infection: Protective role of 17 $\beta$ -estradiol. *The Journal of Infectious Diseases*, 189, 339–345. <https://doi.org/10.1086/380798>
- Mace, R. (2009). Update to Holden and Mace's "phylogenetic analysis of the evolution of lactose digestion in adults" (1997): Revisiting the coevolution of human cultural and biological diversity. *Human Biology*, 81, 621–624. <https://doi.org/10.3378/027.081.0610>
- Markle, J. G., & Fish, E. N. (2014). SexX matters in immunity. *Trends in Immunology*, 35, 97–104. <https://doi.org/10.1016/j.it.2013.10.006>
- Moggi-Cecchi, J., Pacciani, E., & Pinto-Cisternas, J. (1994). Enamel hypoplasia and age at weaning in 19th-century Florence, Italy. *American*

- 1 *Journal of Physical Anthropology*, 93, 299–306. <https://doi.org/10.1002/ajpa.1330930303>
- 2 Møller, A. P. (2006). A review of developmental instability, parasitism and  
3 disease: Infection, genetics and evolution. *Infection, Genetics, and Evolution: Journal of Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases*, 6, 133–140. <https://doi.org/10.1016/j.meegid.2005.03.005>
- 4 Møller, A. P., & Swaddle, J. P. (1997). *Asymmetry, developmental stability and evolution*. Oxford, UK: Oxford University Press.
- 5 Molnar, S. (1972). Tooth wear and culture: A survey of tooth functions  
6 among some prehistoric population. *Current Anthropology*, 13, 511–526. <https://doi.org/10.1086/201284>
- 7 Muenchhoff, M., & Goulder, P. J. (2014). Sex differences in pediatric infectious diseases. *The Journal of Infectious Diseases*, 209(S3), S120–S126. <https://doi.org/10.1093/infdis/jiu232>
- 8 Naugler, C. (2008). Hemochromatosis: A Neolithic adaptation to cereal grain diets. *Medical Hypotheses*, 70, 691–692. <https://doi.org/10.1016/j.mehy.2007.06.020>
- 9 O'Brien, M. J., Laland, K. N., Broughton, J. M., Cannon, M. D., Fuentes, A., Gerbault, P., ... Mackinnon, M. J. (2012). Genes, culture, and agriculture: An example of human niche construction. *Current Anthropology*, 53, 434–470.
- 10 Orton, D., Anvari, J., Gibson, C., Last, J., Bogaard, A., Rosenstock, E., & Biehl, P. F. (2018). A tale of two tells: Dating the Çatalhöyük west mound. *Antiquity*, 92(363), 620–639. [doi.org/10.15184/aqy.2018.91](https://doi.org/10.15184/aqy.2018.91)
- 11 Owens, I. P. (2002). Ecology and evolution. *Sex differences in mortality rate. Science*, 297, 2008–2009. <https://doi.org/10.1126/science.1076813>
- 12 Palmer, A. R., & Strobeck, C. (2003). Fluctuating asymmetry analyses revisited. In M. Polak (Ed.), *Developmental instability: Causes and consequences* (pp. 279–319). Oxford, UK: Oxford University Press.
- 13 Parsons, P. A. (1990). Fluctuating asymmetry: An epigenetic measure of stress. *Biological Reviews of the Cambridge Philosophical Society*, 65, 131–145.
- 14 Paschetta, C., de Azevedo, S., Castillo, L., Martinez-Abadias, N., Hernandez, M., Lieberman, D. E., & González-José, R. (2010). The influence of masticatory loading on craniofacial morphology: A test case across technological transitions in the Ohio valley. *American Journal of Physical Anthropology*, 141, 297–314. <https://doi.org/10.1002/ajpa.21151>
- 15 Pearson, J., & Meskell, L. (2015). Isotopes and images: Fleshing out bodies at Çatalhöyük. *Journal of Archaeological Methods and Theory*, 22, 461–482. <https://doi.org/10.1007/s10816-013-9184-5>
- 16 Pennell, L. M., Galligan, C. L., & Fish, E. N. (2012). Sex affects immunity. *Journal of Autoimmunity*, 38, J282–J291. <https://doi.org/10.1016/j.jaut.2011.11.013>
- 17 Perzigian, A. J. (1977). Fluctuating dental asymmetry—Variation among skeletal populations. *American Journal of Physical Anthropology*, 47, 81–88. <https://doi.org/10.1002/ajpa.1330470114>
- 18 Pilloud, M. A., & Larsen, C. S. (2011). "official" and "practical" kin: Inferring social and community structure from dental phenotype at Neolithic Catalhöyük, Turkey. *American Journal of Physical Anthropology*, 145, 519–530. <https://doi.org/10.1002/ajpa.21520>
- 19 Pinhasi, R., Eshed, V., & Shaw, P. (2008). Evolutionary changes in the masticatory complex following the transition to farming in the southern Levant. *American Journal of Physical Anthropology*, 135, 136–148. <https://doi.org/10.1002/ajpa.20715>
- 20 Pinhasi, R., Eshed, V., & von Cramon-Taubadel, N. (2015). Incongruity between affinity patterns based on mandibular and lower dental dimensions following the transition to agriculture in the near east, Anatolia and Europe. *PLoS One*, 10, e0117301. <https://doi.org/10.1371/journal.pone.0117301>
- 21 Pinhasi, R., & Stock, J. T. (2011). *Human bioarchaeology of the transition to agriculture*. Chichester, UK: Wiley-Blackwell.
- 22 Potter, R. H., & Nance, W. E. (1976). A twin study of dental dimension. I. Discordance, asymmetry, and mirror imagery. *American Journal of Physical Anthropology*, 44, 391–395.
- 23 Powers, S. T., & Lehmann, L. (2014). An evolutionary model explaining the Neolithic transition from egalitarianism to leadership and despotism. *Proceedings of the Royal Society of London. Series B, Biological Sciences*, 281, 20141349. <https://doi.org/10.1098/rspb.2014.1349>
- 24 Powers, S. T., van Schaik, C. P., & Lehmann, L. (2016). How institutions shaped the last major evolutionary transition to large-scale human societies. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 371, 20150098. <https://doi.org/10.1098/rstb.2015.0098>
- 25 Price, T. D. (1995). Social inequality at the origins of agriculture. In T. D. Price & G. M. Feinman (Eds.), *Foundations of social inequality* (pp. 129–151). New York, NY: Springer.
- 26 Reitsem, L. J., & McIlvaine, B. K. (2014). Reconciling "stress" and "health" in physical anthropology: What can bioarchaeologists learn from the other subdisciplines? *American Journal of Physical Anthropology*, 155, 181–185. <https://doi.org/10.1002/ajpa.22596>
- 27 Rollefson, G. O. (2002). Ritual and social structure at Neolithic Ain Ghazal. In I. Kuijt (Ed.), *Life in Neolithic farming communities* (pp. 165–190). Boston, MA: Springer.
- 28 Sakashita, R., Inoue, M., Inoue, N., Pan, Q. F., & Zhu, H. (1997). Dental disease in the Chinese yin-Shang period with respect to relationships between citizens and slaves. *American Journal of Physical Anthropology*, 103, 401–408. [https://doi.org/10.1002/\(SICI\)1096-8644\(199707\)103:3<401::AID-AJPA9>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1096-8644(199707)103:3<401::AID-AJPA9>3.0.CO;2-S)
- 29 Shennan, S., Downey, S. S., Timpson, A., Edinborough, K., Colledge, S., Kerig, T., ... Thomas, M. G. (2013). Regional population collapse followed initial agriculture booms in mid-Holocene Europe. *Nature Communications*, 4, 2486. <https://doi.org/10.1038/ncomms3486>
- 30 Sofaer, J. A., Bailit, H. L., & MacLean, C. J. (1971). A developmental basis for differential tooth reduction during hominid evolution. *Evolution: International Journal of Organic Evolution*, 25, 509–517. <https://doi.org/10.1111/j.1558-5646.1971.tb01910.x>
- 31 Starling, A. P., & Stock, J. T. (2007). Dental indicators of health and stress in early Egyptian and Nubian agriculturalists: A difficult transition and gradual recovery. *American Journal of Physical Anthropology*, 134, 520–528. <https://doi.org/10.1002/ajpa.20700>
- 32 Stinson, S. (1985). Sex differences in environmental sensitivity during growth and development. *Yearbook of Physical Anthropology*, 28, 123–147. <https://doi.org/10.1002/ajpa.1330280507>
- 33 Temple, D. H., & Goodman, A. H. (2014). Bioarchaeology has a "health" problem: Conceptualizing "stress" and "health" in bioarchaeological research. *American Journal of Physical Anthropology*, 155, 186–191. <https://doi.org/10.1002/ajpa.22602>
- 34 Temple, D. H., & Larsen, C. S. (2007). Dental caries prevalence as evidence for agriculture and subsistence variation during the Yayoi period in prehistoric Japan: Biocultural interpretations of an economy in transition. *American Journal of Physical Anthropology*, 134, 501–512. <https://doi.org/10.1002/ajpa.20694>
- 35 Thornhill, R., & Møller, A. P. (1997). Developmental stability, disease and medicine. *Biological Reviews of the Cambridge Philosophical Society*, 72, 497–548.
- 36 Townsend, G., Brook, A., Yong, R., & Hughes, T. (2015). Tooth classes, field concepts, and symmetry. In J. D. Irish & R. A. Scott (Eds.), *A companion to dental anthropology* (pp. 171–201). London, UK: Wiley Blackwell.
- 37 Valen, L. (1962). A study of fluctuating asymmetry. *Evolution: International Journal of Organic Evolution*, 16, 125–142. <https://doi.org/10.2307/2406192>
- 38 von Cramon-Taubadel, N. (2011). Global human mandibular variation reflects differences in agricultural and hunter-gatherer subsistence strategies. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 19546–19551. <https://doi.org/10.1073/pnas.1113050108>
- 39 von Cramon-Taubadel, N. (2017). Measuring the effects of farming on human skull morphology. *Proceedings of the National Academy of Sciences of the United States of America*, 114, 201711475–201718919. <https://doi.org/10.1073/pnas.1711475114>
- 40 Whitaker, S., & Fair, J. (2002). The costs of immunological challenge to developing mountain chickadees, *Poecile gambeli*, in the wild. *Oikos*, 99, 161–165. <https://doi.org/10.1034/j.1600-0706.2002.990116.x>
- 41 Wood, J. W., Milner, G. R., Harpending, H. C., & Weiss, K. M. (1992). The osteological paradox: Problems of inferring prehistoric health from skeletal samples. *Current Anthropology*, 33, 343–358. <https://doi.org/10.1086/204084>

- 1 Wright, K. I. K. (2014). Domestication and inequality? Households, corpo-  
2 rate groups and food processing tools at Neolithic Çatalhöyük. *Journal*  
3 *of Anthropological Archaeology*, 33, 1–33. [https://doi.org/10.1016/j.jaa.](https://doi.org/10.1016/j.jaa.2013.09.007)  
4 2013.09.007

#### 5 SUPPORTING INFORMATION

6 Additional supporting information may be found online in the Sup-  
7 porting Information section at the end of the article.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

**How to cite this article:** Milella M, Betz BJ, Knüsel CJ, Larsen CS, Dori I. Population density and developmental stress in the Neolithic: A diachronic study of dental fluctuating asymmetry at Çatalhöyük (Turkey, 7,100–5,950 BC). *Am J Phys Anthropol.* 2018;1–13. <https://doi.org/10.1002/ajpa.23700>

56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110

Uncorrected Proofs