



Article scientifique

Article

2010

Accepted version

Open Access

This is an author manuscript post-peer-reviewing (accepted version) of the original publication. The layout of the published version may differ .

A Chiral and Colorful Redox Switch: Enhanced π Acidity in Action

Misek, Jiri; Vargas Jentzsch, Andreas; Sakurai, Shin-ichiro; Emery, Daniel; Mareda, Jiri; Matile, Stefan

How to cite

MISEK, Jiri et al. A Chiral and Colorful Redox Switch: Enhanced π Acidity in Action. In: Angewandte Chemie, 2010, vol. 49, n° 42, p. 7680–7683. doi: 10.1002/anie.201003722

This publication URL: <https://archive-ouverte.unige.ch//unige:12065>

Publication DOI: [10.1002/anie.201003722](https://doi.org/10.1002/anie.201003722)

© The author(s). This work is licensed under a Other Open Access license

<https://www.unige.ch/biblio/aou/fr/guide/info/references/licences/>

Enhanced π -Acidity in Action, with a Chiral and Colorful Redox Switch**

Jiří Mišek, Andreas Vargas Jentzsch, Shin-ichiro Sakurai, Daniel Emery, Jiri Mareda and Stefan Matile*

In this report, we introduce simple sulfur redox chemistry^[1,2] to generate enhanced π -acidity,^[3-5] demonstrate functional relevance with anion transport experiments,^[2,4-7] and create the switchable chiral π -acidic surfaces that are needed for future applications toward asymmetric anion- π catalysis. Efforts to expand the number of non-covalent interactions available to construct new supramolecular systems with significant functions such as sensors, optoelectronic devices, catalysts or drugs, are of paramount importance.^[4] To contribute to this adventure, we have explored the use of potential-macrodipole,^[2,7] anion-macrodipole,^[7] aromatic electron donor-acceptor^[7] as well as anion- π interactions^[4,5] to build stimuli-responsive transport systems^[4-7] in lipid bilayer membranes. Core-substituted naphthalenediimides (cNDIs)^[8] are ideal to explore the usefulness of anion- π interactions to create function because the quadrupole moment of the parent NDI **1** is already $Q_{ZZ} = +19$ B (Figures 1a and b).^[3-5] This intrinsic π -acidity is very high, as high as that of the explosive TNT.^[3-5] The introduction of two cyano π -acceptors in the cNDI core doubles the quadrupole moment to $Q_{ZZ} = +39$ B, the highest value known today.^[4] This increase in π -acidity is reflected in decreasing LUMO energy levels from -4.31 eV to -4.62 eV, and increasing anion binding affinity, charge mobility and anion transport activity.^[4,8,9] Synthetic efforts to further lower LUMO energies and generate enhanced π -acidity by adding more cyano π -acceptors into the cNDI core failed so far despite much interest in academia and industry.^[4,8,9]

Looking for synthetic routes to “super- π -acidity,” we considered that oxidation of sulfide donors^[10] in the core of cNDIs could reversibly yield chiral sulfoxide and powerful sulfone acceptors (Figure 1). Aromatic sulfur compounds, particularly thiophenes, are ubiquitous in materials sciences because of the high conductivity of their π -stacks.^[11] Their reversible oxidation has been much less explored to generate function, although elegant examples exist for fluorescent sensors, p - to n -charge mobility inversion (oligothiophene oxidation), switchable secondary peptide structures,

amyloid disease prevention and voltage-gated synthetic ion channels.^[1,2] In the following, we report synthesis, optoelectronic properties, anion transport activity and computational studies of cNDIs **2-9**. Tetrasulfone **9** emerges as the most π -acidic cNDI known today, the resulting anion transport activity is naturally outstanding.




Figure 1. Structures, frontier orbital energies, absorption maxima and favorable (a) and unfavorable (b) anion complexes of anion- π redox switches. HOMO (bold) and LUMO (dashed) energies are in eV against vacuum compared to native NDI **1** (grey), absorption maxima in nm (dashed arrows).

The cNDI **2** with two sulfide donors in the core and two diisopropylphenyl (Dip) substituents in the periphery was prepared as described in the literature (Figure S1).^[10,11] To enable anion- π interactions on the most electron-deficient pyridinedione heterocycles, the obstructing Dip substituents (Figure 1b) were replaced by mesityl (Mes) substituents which allow for a practical compromise between acceptable anion- π interactions and reasonable solubility (Figure 1a).^[4] Reactivity and optoelectronic properties of the Mes- and Dip-cNDIs were roughly the same. Namely, conversion of the π -donating sulfides **4** into π -accepting sulfoxides **5** with stoichiometric and sulfones **6** with excess mCPBA caused the

[*] Dr. J. Mišek, A. Vargas Jentzsch, Dr. S. Sakurai, D. Emery, Dr. J. Mareda, Prof. S. Matile
Department of Organic Chemistry
University of Geneva
Geneva, Switzerland
Fax: (+) 41 22 379 3215
E-mail: stefan.matile@unige.ch
Homepage: www.unige.ch/sciences/chiorg/matile/

[**] We thank D.-H. Tran for contributions to synthesis, D. Jeannerat, A. Pinto and S. Grass for NMR measurements, the Sciences Mass Spectrometry (SMS) platform at the Faculty of Sciences, University of Geneva, for mass spectrometry services, the Swiss National Supercomputing Center (CSCS) in Manno for CPU time, and the University of Geneva and the Swiss NSF for financial support. J. Mišek is a fellow of the Roche Research Foundation.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

peaks for the protons in position 3 and 7 of the naphthyl core to shift downfield from 8.81 ppm to 9.66 ppm in the ^1H NMR spectra (Figures 2l-2n). The absorption maxima suffered a hypsochromic shift from a strong, broad charge-transfer (CT) band at 528 nm for sulfide cNDIs to weakened CT at 448 nm for sulfoxide cNDIs and complete CT-silencing for sulfone cNDIs (Figures 2e-2g, Table S1). The π,π^* -transition responded to sulfur oxidation with a weak bathochromic shift from 382 nm for sulfide cNDIs to 398 nm for sulfone cNDIs. None of the sulfur containing cNDIs was fluorescent.




Figure 2. Cyclic voltammograms (a-d), absorption spectra (e-k) and ^1H NMR spectra (l-r) of cNDIs 4-9. This covers sulfide 4 (a, e, h, l, o), sulfoxide 5 (b, f, h, m) and sulfone 6 (c, g, n) compared to tetrasulfide 7 (i), tetrasulfoxide 8 (j) and tetrasulfone 9 (d, k), colorimetric redox switching with 4 and 5 (h), 4 as product of the reduction of 5 (o), and racemic 5 (p), enantioenriched 5 (q) and achiral 4 (r) in the presence of the shift reagent (-)-(R)-TFAE (250 eq, CDCl_3 , 50 $^\circ\text{C}$, Figure S5).

Cyclic voltammetry (CV) revealed a significant increase in π -acidity in response to the oxidation of the sulfur substituents in the core (Figures 2a-c). LUMO energies against vacuum were obtained by subtraction of -5.1 eV for Fc/Fc^+ from the first redox potential at -1.17 V for sulfide cNDIs, -0.76 V for sulfoxide cNDIs and -0.64 V for sulfone cNDIs (Figure 1, Table S1). HOMO/LUMO gaps were calculated from the onset of the lowest energy absorptions to approximate the HOMO energies (Figure 1).

The reversibility of sulfide oxidation was confirmed by reduction of 5 with triflic anhydride (TiF_2O) and KI (Figure 2o vs 2l). Colorimetric redox switching was readily demonstrated for repeated reduction and oxidation cycles for 4 and 5 (Figures 2h, S4). In the presence of the chiral shift reagent (-)-(R)-1-(9-anthryl)-2,2,2-trifluoroethanol (TFAE, or Pirkle's alcohol),^[12] the NMR signals for the protons in position 3 and 7 of the naphthyl core in chiral

sulfoxide 5 (but not achiral sulfide 4) splitted into four equal resonances (Figure 2p vs 2r). Asymmetric oxidation of 4 with *tert*-butylhydroperoxide (TBHP) catalyzed by (*R*)-(+)-1,1'-bi(2-naphthol) ((*R*)-BINOL, 20%) and $\text{Ti}(\text{O}i\text{Pr})_4$ (10%) changed the magnitude of the split resonances of 5 (Figure 2q).^[11,13] The upfield signals for the pair of enantiomers integrated for an enantiomeric excess of 40% *ee*. Their 1:1 ratio compared to the *meso* diastereomer suggested that the second sulfide oxidation occurs without stereoselectivity (Figure S5). The circular dichroism (CD) spectrum of enantioenriched 5 showed a weakly negative Cotton effect (CE) for the CT band at 449 nm and a stronger positive CE at 303 nm (Figure S6).

Today, the use of standard intermolecular interactions to build functional systems that can transport ions across lipid bilayers is almost routine.^[5-8] In a recent inversion of paradigm, activity to transport anions was considered as a tool to identify more exotic interactions and assess their functional relevance for translocation and catalysis, where much weaker interactions are of interest than for simple binding.^[4,14] For this purpose, the ability of cNDIs 2-6 to use anion- π interactions to transport anions across lipid bilayer membranes was evaluated with the 8-hydroxy-1,3,6-pyrenetrisulfonate (HPTS) assay. In this assay, EYPC-LUVsDHPTS are exposed to a transmembrane pH gradient. These are large unilamellar vesicles composed of egg yolk phosphatidylcholine and loaded with the pH-sensitive fluorophore HPTS. The internal pH probes report the ability of externally added cNDIs to mediate the dissipation of transmembrane pH gradients (Figure S8). Dose response curves obtained for cNDIs 2-6 were then subjected to Hill analysis to give the effective monomer concentrations (EC_{50}) needed to reach 50% activity. For sulfide 4, $EC_{50} = 13.9 \pm 1.2 \mu\text{M}$ was found (Figure 3a, $\square$, Table S2). This value improved with sulfide oxidation to $EC_{50} = 2.1 \pm 0.2 \mu\text{M}$ for sulfoxide 5 and $EC_{50} = 1.8 \pm 0.4 \mu\text{M}$ for sulfone 6 (Figure 3a, $\triangle$ and $\circ$). Increasing activity with increasing π -acidity was in support of operational anion- π interactions in the Mes-cNDI series. In the Dip-cNDI series, activity decreased with π -acidity from $EC_{50} = 0.9 \pm 0.1 \mu\text{M}$ for 2 to $EC_{50} = 3.5 \pm 0.4 \mu\text{M}$ for 3, probably due to decreasing partitioning of more hydrophilic transporters (Table S2). This dichotomic behavior with obstructed anion- π interactions in the Dip-series was essential to corroborate the functional relevance of the operational anion- π interactions in the Mes-cNDI series 4-6.^[7c]

The ion selectivity of cNDIs 2-6 was determined by external ion exchange. The high sensitivity to external anion exchange and the negligible sensitivity to external cation exchange (except for Li^+) implied selectivity for anions (Figures 3b and 3c). The global anti-Hofmeister behavior in the halide series, preference for oxyanions, occasional nitrate recognition and fluoride/acetate rejection with increasing π -acidity were roughly as described for dicyano cNDIs and can be interpreted as support for strong, π - π enhanced anion- π interactions at work (Table S2).^[4] Anion selectivity sequences were partially obscured by a lithium anomaly that gained in importance with increasing sulfide oxidation (Figures 3b, 3c). This effect was attributed to the binding of lithium to the pocket formed by sulfone and diimide oxygens and considered as attractive approach to Lewis-acid enhanced π -acidity.

Quantum chemical computations suggested that the π -acidity of cNDIs with two sulfones in the core is comparable to that of previously investigated dicyano cNDIs.^[5a,11] Comparison of the electrostatic surface potentials (ESP) of disulfone 6 and dicyano cNDIs revealed only slightly less electron-deficient surfaces for the former (Figures S21).^[11] This suggested that oxidation of the four sulfides in the core of cNDI 7 should provide access to the enhanced π -acidity that couldn't be reached with tetracyano cNDIs (Figure

1).^[4] Tetrasulfide **7** was readily prepared by adapting procedures reported previously in the literature,^[10b] oxidation to tetrasulfoxide **8** with stoichiometric and tetrasulfone **9** with excess mCPBA was unproblematic.^[11] Solutions of the super- π -acid **9** in MeOH or DMSO turned black within hours, as did spots on thin layer chromatograms.




Figure 3. Transport activity (a) and selectivity (b, c) of **4** ($\square$), **5** ($\triangle$), **6** ($\circ$), **7** ($\blacksquare$), **8** ($\blacktriangle$) and **9** ($\bullet$). a) Dependence of fractional activity Y on cNDI concentration with Hill analysis. (b, c) Dependence of Y on the dehydration energy of external anions (b, 100 mM NaX, inside 100 mM NaCl) and cations (c, 100 mM XCl, inside 100 mM NaCl), normalized for $Y = 1$ for Na^+ and Cl^- .^[11]

The transport activity of tetrasulfone **9** was outstanding ($EC_{50} = 239 \pm 24$ nM, Figure 3a, $\bullet$), the best known so far for anion- π transporters.^[4] The Hill coefficient $n = 2.5 \pm 0.7$ confirmed that the active structure is an unstable supramolecule composed of at least three monomers. The anti-Hofmeister selectivity topology with AcO^-/F^- exclusion and significantly enhanced nitrate recognition found for **8** and **9** supported strong anion recognition (Figure 3b, $\bullet$, $\blacktriangle$). Compared to disubstituted cNDIs (Figures 2e-g), the absorption spectra of tetrasubstituted cNDIs were broadened, showed a stronger, more bathochromic CT band for tetrasulfoxide **8** and a stronger hypsochromic effect for the π, π^* -bands (Figures 2i-k). The origin of these differences is unknown, chromophore twisting due to topological crowding with four substituents in the core is likely to contribute (Figure 4). The cyclic voltammograms of di- and tetrasubstituted cNDIs were identical in appearance (Figures 2a-d, S7). From the first redox potential at -0.36 V, a LUMO energy of -4.74 eV was obtained for **9** (Figure 1). This finding demonstrated that with tetrasulfone cNDI **9**, the so far elusive “super- π -acidity” was finally realized.

Molecular models confirmed the validity of this interpretation. The exceptional ESP of super- π -acid **9** with a deeply blue aromatic surface contrasted nicely to the greenish surfaces of the less π -acidic tetrasulfide **7** (Figure 4, top). Increasing peripheral crowding with increasing sulfur oxidation slightly twisted the pyridinedione heterocycles in tetrasulfone **9** out of the naphthyl plane. This unprecedented NDI deplanarization is presumably subject to

extensive atropisomerism of the ethylsulfonyl side chains (Figure 4, bottom).

In summary, we report facile access to aromatic surfaces that unify “super- π -acidity,” operational anion- π interactions, chirality and red color in a switchable manner. π -Acidic cNDIs with sulfur substituents in the core are of fundamental interest in materials science (e.g., air-stable n -semiconductors with high charge mobility)^[1,9] and organocatalysis. In view of the stabilization of cationic transition states by cation- π interactions as classical motif in catalysis, particularly carbocation chemistry,^[14] anion- π interactions have the potential to stabilize the complementary anionic reactive intermediates such as enolates.




Figure 4. Electrostatic surface potentials computed for optimized structures viewed from the top (blue +1.24, green 0, red -1.24 eV, PBE1PBE/6-311G**) and side views of ball and stick representations along the N, N' axes (atom color coding) of tetrasulfide cNDI **7** (left) and tetrasulfone cNDI **9** (right).

- [1] a) E. L. Dane, S. B. King, T. M. Swager, *J. Am. Chem. Soc.* **2010**, *132*, 7758-7768; b) M. Moreno Oliva, J. Casado, J. T. López Navarrete, S. Patchkovskii, T. Goodson III, M. R. Harpham, J. S. Seixas de Melo, E. Amir, S. Rozen, *J. Am. Chem. Soc.* **2010**, *132*, 6231-6242; c) C. Wolschner, A. Giese, H. A. Kretzschmar, R. Huber, L. Moroder, N. Budisa, *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 7756-7761; d) G. P. Dado, S. H. Gellman, *J. Am. Chem. Soc.* **1993**, *115*, 12609-12610.
- [2] a) N. Sakai, S. Matile, *J. Am. Chem. Soc.* **2002**, *124*, 1184-1185; b) N. Sakai, D. Gerard, S. Matile, *J. Am. Chem. Soc.* **2001**, *123*, 2517-2524; c) F. Robert, J.-Y. Winum, N. Sakai, D.

- Gerard, S. Matile, *Org. Lett.* **2000**, *2*, 37-39; d) N. Sakai, S. Matile, *Chem. Eur. J.* **2000**, *6*, 1731-1737.
- [3] a) B. L. Schottel, H. T. Chifotides, K. R. Dunbar, *Chem. Soc. Rev.* **2008**, *37*, 68-83; b) B. P. Hay, V. S. Bryantsev, *Chem. Commun.* **2008**, 2417-2428; c) G. Gil-Ramírez, E. C. Escudero-Adán, J. Benet-Buchholz, P. Ballester, *Angew. Chem. Int. Ed.* **2008**, *47*, 4114-4118; d) P. Gamez, T. J. Mooibroek, J. S. Teat, J. Reedijk, *Acc. Chem. Res.* **2007**, *40*, 435-444; e) D. Quinonero, C. Garau, C. Rotger, A. Frontera, P. Ballester, A. Costa, P. M. Deya, *Angew. Chem. Int. Ed.* **2002**, *41*, 3389-3392; f) M. Mascal, A. Armstrong, M. D. Bartberger, *J. Am. Chem. Soc.* **2002**, *124*, 6274-6276; g) I. Alkorta, I. Rozas, J. Elguero, *J. Am. Chem. Soc.* **2002**, *124*, 8593-8598.
- [4] R. E. Dawson, A. Hennig, D. P. Weimann, D. Emery, S. Gabutti, J. Montenegro, V. Ravikumar, M. Mayor, J. Mareda, C. A. Schalley, S. Matile, *Nat. Chem.* **2010**, *2*, 533-538.
- [5] a) J. Mareda, S. Matile, *Chem. Eur. J.* **2009**, *15*, 28-37; b) A. Perez-Velasco, V. Gorteau, S. Matile, *Angew. Chem. Int. Ed.* **2008**, *47*, 921-923.
- [6] a) A. P. Davis, D. N. Sheppard, B. D. Smith, *Chem. Soc. Rev.* **2007**, *36*, 348-357; b) C. P. Wilson, S. J. Webb, *Chem. Commun.* **2008**, 4007-4009; c) J. T. Davis, P. A. Gale, O. A. Okunola, P. Prados, J. C. Iglesias-Sanchez, T. Torroba, R. Quesada, *Nat. Chem.* **2009**, *1*, 138-144; d) X. Li, B. Shen, X. Q. Yao, D. Yang, *J. Am. Chem. Soc.* **2009**, *131*, 13676-13680; e) G. W. Gokel, N. Barkey, *New J. Chem.* **2009**, *33*, 947-963; f) J. K. W. Chui, T. M. Fyles, *Chem. Commun.* **2010**, *46*, 4169-4171.
- [7] a) N. Sakai, N. Sordé, S. Matile, *J. Am. Chem. Soc.* **2003**, *125*, 7776-7777; b) A. Hennig, L. Fischer, G. Guichard, S. Matile, *J. Am. Chem. Soc.* **2009**, *131*, 16889-16895; c) P. Talukdar, G. Bollob, J. Mareda, N. Sakai, S. Matile, *Chem. Eur. J.* **2005**, *11*, 6525-6532; d) S. Bhosale, A. L. Sisson, P. Talukdar, A. Fürstenberg, N. Banerji, E. Vauthey, G. Bollob, J. Mareda, C. Röger, F. Würthner, N. Sakai, S. Matile, *Science* **2006**, *313*, 84-86; e) S. Hagihara, H. Tanaka, S. Matile, *J. Am. Chem. Soc.* **2008**, *130*, 5656-5657.
- [8] N. Sakai, J. Mareda, E. Vauthey, S. Matile, *Chem. Commun.* **2010**, *46*, 4225-4237.
- [9] a) S. Chopin, F. Chaignon, E. Blart, F. Odobel, *J. Mater. Chem.* **2007**, *17*, 4139-4146; b) X. Gao, C. Di, Y. Hu, X. Yang, H. Fan, F. Zhang, Y. Liu, H. Li, D. Zhu, *J. Am. Chem. Soc.* **2010**, *132*, 3697-3699; c) B. A. Jones, A. Facchetti, M. R. Wasielewski, T. J. Marks, *J. Am. Chem. Soc.* **2007**, *129*, 15259-15278; d) M. Könemann, PCT Int. Appl. WO 2007074137, **2007**.
- [10] a) A. Blaszczyk, M. Fischer, C. von Hänisch, M. Mayor, *Helv. Chim. Acta* **2006**, *89*, 1986-2005; b) C. Röger, F. Würthner, *J. Org. Chem.* **2007**, *72*, 8070-8075.
- [11] See Supplementary Information.
- [12] a) W. H. Pirkle, D. L. Sikkenga, M. S. Pavlin, *J. Org. Chem.* **1977**, *42*, 384-387; b) D. Parker, *Chem. Rev.* **1991**, *91*, 1441-1457.
- [13] a) N. Komatsu, M. Hashizume, T. Sugita, S. Uemura, *J. Org. Chem.* **1993**, *58*, 4529-4533; b) I. Fernández, N. Khiar, *Chem. Rev.* **2003**, *103*, 3651-3705.
- [14] R. R. Knowles, S. Lin, E. N. Jacobsen, *J. Am. Chem. Soc.* **2010**, *132*, 5030-5032.