

## REVIEW

**Leaf senescence and activities of the antioxidant enzymes**

D. PROCHÁZKOVÁ\* and N. WILHELMOVÁ

*Institute of Experimental Botany, Academy of Sciences of the Czech Republic,  
Na Karlovce 1a, Prague 6, CZ-16000, Czech Republic***Abstract**

Senescence is a genetically regulated process that involves decomposition of cellular structures and distribution of the products of this degradation to other plant parts. Reactions involving reactive oxygen species are the intrinsic features of these processes and their role in senescence is suggested. The malfunction of protection against destruction induced by reactive oxygen species could be the starting point of senescence. This article reviews biochemical changes during senescence in relation to reactive oxygen species and changes in antioxidant protection.

*Additional key words:* ageing, enzymatic antioxidants, lipid peroxidation, non-enzymatic antioxidants, oxidative stress.

**Introduction**

Leaf senescence – a final developmental phase between maturity and death of a plant or a plant part - has been defined as a highly organized, energy-requiring, genetically programmed process.

The nature of the inducer(s), initiating a natural senescence program, has not been fully explained yet. Among the various possibilities, such as source-sink competitions during redirection of nutrients, changing hormonal control, or changes of the energy status of the cell (Thimann 1980, Thomas and Stoddart 1980, Kar and Feierabend 1984, Buchanan-Wollaston 1997), the free-radical theory has attracted attention. A hypothesis has been developed that ageing results from an accumulation of harmful free radicals (Harman 1981) and that the onset of senescence is mainly due to the uncontrolled strong enhancement in the generation of reactive oxygen species (ROS), especially superoxide, singlet oxygen, hydroxyl radical and hydrogen peroxide (Thompson and Barber 1987).

Three phases may be distinguished in a typical senescent process (Peñarrubia and Moreno 1995):

1) Storage mobilization. A phase of selective degradation of certain molecules. This degradation does not cause a major impairment of the physiological functions. The

mobilized molecules may be considered as nutrient storage materials. In some cases senescence may be reversed during this phase by certain changes in environmental conditions.

2) Generalized breakdown. Extension and generalization of breakdown to components which are central in maintaining physiological function. As a result of this breakdown a physiological function is consequently lost. During this phase the senescence process becomes irreversible and death of the cells becomes inevitable.

3) Abscission. The final phase of senescence is abscission and death.

Leaf senescence is accompanied by changes of many organelles. Chloroplasts of senescing leaves show reduced volume, their shape is spherical and the thylakoid system is reduced (Matile 1992). The final developmental stage of chloroplasts – gerontoplasts – shows an increase in the number and diameter of plastoglobuli, a reduction of the thylakoid system, a loosening of the stacking of thylakoids and a swelling of intrathylakoid spaces. Mitochondria remain intact until late senescence, however, in later stages some swelling or distortion of cristae becomes apparent. Nuclei do not show substantial structural changes until relatively late stage of senescence

*Received 14 June 2006, accepted 22 January 2007.*

*Abbreviations:* APX - ascorbate peroxidase; Asc - ascorbate; CAT - catalase; Chl - chlorophyll; DHA - dehydroascorbate; DHAR - dehydroascorbate reductase; GPX - glutathione peroxidase; GR - glutathione reductase; GSH - reduced glutathione; GSSG - oxidized glutathione; MDHA - monodehydroascorbate radical; MDHAR - monodehydroascorbate reductase; ROS - reactive oxygen species; SOD - superoxide dismutase;  $\alpha$ -toc -  $\alpha$ -tocopherol;  $\beta$ -car -  $\beta$ -carotene.

*Acknowledgements:* This work was supported by the Grant Agency of the Czech Republic, project No. 522/05/P558.

\* Corresponding author; fax: (+420) 224310113, e-mail: prochazkova@ueb.cas.cz

(Noodén *et al.* 1997), nevertheless, there are some reports of chromatin condensation and other changes (Kuran

1993, Biradar and Rayburn 1994, Noodén *et al.* 1997).

### Biochemical changes during senescence

**Lipids:** One of the most characteristic features of senescence is the initiation of membrane leakiness and a consequent loss of intracellular compartmentalization predominantly due to lipid damage. Lipid peroxidation is commonly used as an indicator of prevalence of free radicals in tissues (Smirnoff 1993). Lipid peroxidation not only threatens the integrity and function of membranes and membranous proteins but also produces a variety of toxic aldehydes and ketones (Valentine *et al.* 1998, Wilhelmová *et al.* 2006). Some of such products of lipid peroxidation, malondialdehyde and 4-hydroxynonenal, cause protein damage by means of reactions with lysine amino groups, cysteine sulfhydryl groups and histidine imidazole groups (Esterbauer *et al.* 1991, Uchida and Stadtman 1992, Friguet *et al.* 1994, Brunner *et al.* 1995, Refsgaard *et al.* 2000).

The striking chemical change in the membrane lipids is a dramatic increase in the sterol/phospholipid ratio (Borochoy *et al.* 1978, McKersie *et al.* 1978, Lees and Thompson 1980), however, content of sterols declines with physiological ageing (Paliyath and Droillard 1992).

**Photosynthetic pigments:** Breakdown of chlorophylls (Chl) may be one of the earliest symptoms of senescence. All final catabolites are not derived only from chlorophyll *a* (Chl *a*) but represent the breakdown products of chlorophyll *b* (Chl *b*) as well (Curty and Engel 1996, Ginsburg and Matile 1993, Matile and Hörtensteiner 1999). The first step in the degradation of Chl *a* is hydrolysis of the phytol ester bond catalyzed by chlorophyllase, which forms chlorophyllide *a* and phytol (Tsuchiya *et al.* 1997, Suzuki and Shioi 1999). In the next step magnesium is released from the macrocyclic ring and pheophorbide *a* forms with the action of Mg-dechelataze substance (Shioi *et al.* 1996a, Suzuki and Shioi 1999). The final step of macrocycle ring modification is the conversion of pheophorbide *a* to pyropheophorbide *a* (Shioi *et al.* 1996b, Watanabe *et al.* 1999, Suzuki and Shioi 1999). The next reaction, oxidative cleavage of the tetrapyrrole ring, is probably catalyzed by pheophorbide *a* oxygenase (Matile *et al.* 1996, Rodoni *et al.* 1997, Suzuki and Shioi 1999).

The main catabolites of degraded chlorophylls are derivatives of bilin (Engel *et al.* 1991, Iturraspe and Moyano 1995, Doi *et al.* 1997, Suzuki and Shioi 1999). The final products have been also termed nonfluorescent Chl catabolites (Matile and Hörtensteiner 1999). These linear tetrapyrroles are accumulated in the vacuoles of senescent mesophyll cells (Matile *et al.* 1988, Kräutler 2003, Hörtensteiner 2004).

**Nucleic acids:** The content of DNA remains relatively constant during senescence (Makrides and Goldthwaite

1981, Buchanan-Wollaston 1997). It has been shown that repeated sequences are selectively degraded while coding regions of nuclear DNA remain to a large extent intact (Abeles and Dunn 1990).

Total RNA content decreases during senescence, with a completely yellow leaf having about 10-times less RNA than a green one (Lohman *et al.* 1994). The expression of many genes is switched off (Bate *et al.* 1991, Hensel *et al.* 1993, Lohman *et al.* 1994). New transcripts were originally detected as a changed pattern of products observed in *in vitro* translation experiments using mRNA from senescing leaves when compared to green leaves (Malik 1987, Thomas *et al.* 1992).

Transfer RNA synthetase activities are greatly reduced during senescence (Jaybaskaran *et al.* 1990). A general increase in RNase activities has been described during senescence (Green 1994, Buchanan-Wollaston 1997). Nevertheless, qualitative changes in mRNA are probably more relevant in the senescence process.

The fate of the purines and pyrimidines resulting from nucleic acid degradation is not clear, but they are presumably broken down further, to keep their nitrogen and carbon components for further use in the plant. The increased levels of uricase and xanthine oxidase observed in the peroxisomes of senescent leaves indicate that purine catabolism takes place in this organelle (Vicentini and Matile 1993, Pastori and del Río 1994).

The nucleic acids, in particular the rRNA, are used as an important source of carbon, nitrogen and especially phosphorus within a senescing cell.

**Proteins:** Leaf senescence is characterized by a progressive decrease of total protein content (Brady 1988). The patterns of protein loss are characteristic and independent of the cause of senescence. Many specific proteins are degraded while others remain intact.

Probably both reduced synthesis and enhanced proteolysis are responsible for protein loss observed during senescence. In this regard, synthesis of all thylakoid proteins is known to be severely curtailed in senescing bean leaves except for the D1 protein of photosystem 2 (Droillard *et al.* 1992). Increased protein breakdown may result from different mechanisms: *de novo* synthesis of proteolytic enzymes, activation of pre-existing proteases, decompartmentalization of proteases and their substrates for degradation.

Reaction of most ROS with proteins results in the initial formation of carbon-centered radicals at either the protein backbone or side chains (Hawkins and Davies 2001).

**Oxidation of the protein backbone:** The carbon-centered radical reacts rapidly with O<sub>2</sub> to form an

alkylperoxyl radical intermediate, which can give rise to the alkylperoxide, followed by a formation of an alkoxy radical, which may be converted to a hydroxyl protein derivative. The alkyl, alkylperoxyl and alkoxy radical intermediates in this pathway may undergo side reactions with other amino acid residues in the same or a different protein molecule to generate a new carbon-centered radical (Berlett and Stadtman 1997).

**Oxidation of amino acid side chains:** Carbon-centered radicals react rapidly with O<sub>2</sub> to give peroxyl species. Peroxyl radicals undergo a number of reactions that result in the formation of carbonyl groups (aldehydes or ketones), alcohols and hydroperoxides. Alkoxy radicals can be generated from peroxyl radicals. They can undergo rapid addition and hydrogen abstraction reactions, as well as facile unimolecular fragmentation and rearrangement reactions (Hawkins and Davies 2001).

### Protection against reactive oxygen species during senescence

ROS are accumulated with age in a progressive manner. Their most powerful source is chloroplast (Foyer and Noctor 2000). Other sources are electron transport chain in mitochondria, oxidation in microbodies (Dat *et al.*

2000) or cell wall peroxidases (Eltner 1982). Plants have evolved sophisticated mechanisms to maintain ROS concentrations under the control (see Table 1).

Leaf senescence is often associated with increased

Table 1. Compendium of the most important enzymatic and non-enzymatic antioxidants.

Antioxidant	Reaction
Superoxide dismutase (SOD)	$2\text{O}_2^{\cdot-} + 2\text{H}^+ \Rightarrow \text{O}_2 + \text{H}_2\text{O}_2$
Ascorbate peroxidase (APX)	$\text{H}_2\text{O}_2 + 2\text{Asc} \Rightarrow 2\text{H}_2\text{O} + 2\text{MDHA}$
Monodehydroascorbate reductase (MDHAR)	$2\text{MDHA} + \text{NAD(P)H} + \text{H}^+ \Rightarrow 2\text{Asc} + \text{NAD(P)}^+$
Dehydroascorbate reductase (DHAR)	$\text{DHA} + 2\text{GSH} \Rightarrow \text{Asc} + \text{GSSG}$
Glutathione reductase (GR)	$\text{GSSG} + \text{NAD(P)H} + \text{H}^+ \Rightarrow 2\text{GSH} + \text{NAD(P)}^+$
Catalase (CAT)	$2\text{H}_2\text{O}_2 \Rightarrow 2\text{H}_2\text{O} + \text{O}_2$
Glutathione peroxidase (GPX)	$\text{H}_2\text{O}_2 + 2\text{GSH} \Rightarrow 2\text{H}_2\text{O} + \text{GSSG}$
Glutathione (GSH)	$\text{H}_2\text{O}_2 + 2\text{GSH} \Rightarrow 2\text{H}_2\text{O} + \text{GSSG}$
	$\text{GSH} + \text{OH}^{\cdot} \Rightarrow \text{H}_2\text{O} + \text{GS}^{\cdot}$
Ascorbate (Asc)	$\text{Asc} + \text{H}_2\text{O}_2 \Rightarrow 2\text{H}_2\text{O} + \text{DHA}$
β-carotene (β-car)	$\beta\text{-car} + \text{ROO}^{\cdot} \Rightarrow \beta\text{-car} + \text{ROOH}$
	$^3\text{Chl}^* + \beta\text{-car} \Rightarrow ^1\text{Chl} + ^3\beta\text{-car}^*$
α-tocopherol (α-toc)	$\alpha\text{-toc} + \text{LOO}^{\cdot} \Rightarrow \alpha\text{-toc} + \text{LOOH}$

oxidative damage to cellular macromolecules by ROS. Since ROS levels and their damage products in many plants are known to increase during senescence, it is possible that these changes are due to a decline in the activity of certain antioxidant enzymes. There is superiority of evidence from number of laboratories that antioxidant protection declines at the later stages of leaf senescence. However, it is still impossible to draw any explicit conclusion. For example, Scebba *et al.* (2001) described entirely different courses of the activities on two different cultivars of *Prunus armeniaca* during senescence. Another illustration of a surprising course of antioxidant activity was described in *Arabidopsis* where APX activity dramatically decreased after pre-bolt stage and the lowest was in medium bolting stage. Its activity rebounded with further development (reproduction and seed setting) and this increase precedes the onset of visible senescence symptoms (Ye *et al.* 2000).

The course of activity in various experiments depends

on too many factors: time of sampling, length of monitored time course, a kind of a method employed, plant organ, plant species or cultivar. A selection of reference unit is very important. Almost all of them undergo quantitative change to various extents. The tendency of studied characteristics can achieve even contradictory course on dependence of such reference parameter (per fresh mass, dry mass, proteins, per volume of enzyme extract or per one leaf, one cotyledon, *etc.*).

The antioxidant activity in senescence varies on the level of organelles as well. For example a notable increase in the activity of peroxisomal MnSOD and a decline in the activity of mitochondrial MnSOD from pea leaf extract were determined at the same time (Pastori and del Río 1994). On the other hand a decrease in the activity of whole pea leaf extract MnSOD was described in another experiment (Jiménez *et al.* 1998). Moreover differences between natural and artificially induced senescence would deserve a separate article.

Table 2. The antioxidant enzyme activity and content of non-enzymatic antioxidant during plant senescence. Symbol ↑ describes continuous increase of activity or concentration during monitored time course; symbol ↓ describes initiative increase, followed by a decrease of activity or concentration; symbol ↓ describes continuous decline of activity or concentration during monitored time course.

Antioxidant	Plant	Organ	Course	Reference
Total SOD	clusterbean	cotyledons	↓	Deo <i>et al.</i> 2006
	tobacco	leaves	↑↓	Dertinger <i>et al.</i> 2003
	bean	cotyledons	↑↓	Procházková and Wilhelmová 2007a
	maize	leaves	↑↓	Procházková <i>et al.</i> 2001
	wheat	leaves	↓	Srivalli and Khanna-Chopra 2001
CuZnSOD	bean	cotyledons	↑↓	Procházková and Wilhelmová 2007a
MnSOD	bean	cotyledons	↑↓	Procházková and Wilhelmová 2007a
	pea	leaves	↓	Jiménez <i>et al.</i> 1998
APX	pea	leaves	↓	Jiménez <i>et al.</i> 1998
	cucumber	cotyledons	↑	Kanazawa <i>et al.</i> 2000
	bean	cotyledons	↑↓	Procházková and Wilhelmová 2004
	maize	leaves	↑↓	Procházková <i>et al.</i> 2001
CAT	cucumber	cotyledons	↓	Kanazawa <i>et al.</i> 2000
	soybean	leaves	↑↓	Fu <i>et al.</i> 2000
	maize	leaves	↑↓	Procházková <i>et al.</i> 2001
	bean	cotyledons	↓	Procházková and Wilhelmová 2004
	<i>Arabidopsis</i>	leaves	↓	Zimmermann <i>et al.</i> 2006
GR	cucumber	cotyledons	↑↓	Kanazawa <i>et al.</i> 2000
	bean	cotyledons	↑↓	Procházková and Wilhelmová 2004
	tobacco	leaves	↓	Dertinger <i>et al.</i> 2003
DHAR	cucumber	cotyledons	↓	Kanazawa <i>et al.</i> 2000
GPX	cucumber	cotyledons	↑	Kanazawa <i>et al.</i> 2000
GSH	bean	cotyledons	↓	Procházková and Wilhelmová 2007b
α-toc	sage	leaves	↓	Munné-Bosch <i>et al.</i> 2001
	tobacco	leaves	↑	Dertinger <i>et al.</i> 2003
Asc	tobacco	leaves	↓	Dertinger <i>et al.</i> 2003
	bean	cotyledons	↓	Procházková and Wilhelmová 2007b

## Conclusion

Recent studies unravelled many questions concerning senescence and ROS, but many other questions arise. The focus of the current research is aimed rather at increasing our knowledge of leaf senescence at the molecular level. The identification and the characterization of genes

involved in the complex degradation events that occur during leaf senescence and the elucidation of their mechanisms of regulation will help in increasing of yield, stress resistance and shelf life of agricultural products.

## References

- Abeles, F.B., Dunn, L.J.: Restriction fragment length polymorphism analysis of DNA from senescing cotyledon tissue. - *Plant Sci.* **72**: 13-17, 1990.
- Bate, N.J., Rothstein, S.J., Thompson, J.E.: Expression of nuclear and chloroplast photosynthesis-specific genes during leaf senescence. - *Planta* **189**: 74-79, 1991.
- Berlett, B.S., Stadtman, E.R.: Protein oxidation in aging, disease and oxidative stress. - *J. biol. Chem.* **272**: 20313-20316, 1997.
- Biradar, D.P., Rayburn, A.L.: Flow cytometric probing of chromatin condensation in maize diploid nuclei. - *New Phytol.* **126**: 31-35, 1994.
- Borochoy, A., Halevy, A.H., Borochoy, H., Shinitzky, M.: Microviscosity of plasmalemmas in rose petals as affected by age and environmental factory. - *Plant Physiol.* **61**: 812-815, 1978.
- Brady, C.J.: Nucleic acid and protein synthesis. - In: Noodén, L.D., Leopold, A.C. (ed.): *Senescence and Ageing in Plants*. Pp. 147 - 181. Academic Press, London 1988.
- Brunner, B.A., Jones, A.D., German, J.B.: Direct characterization of protein adduct of the lipid peroxidation product 4-hydroxy-2-nonenal using electrospray mass spectrometry. - *Chem. Res. Toxicol.* **8**: 552 - 559, 1995.
- Buchanan-Wollaston, V.: The molecular biology of leaf senescence. - *J. exp. Bot.* **307**: 181-199, 1997.
- Curty, C., Engel, N.: Detection, isolation and structure elucidation of a chlorophyll a catabolite from autumnal senescent leaves of *Cercidiphyllum japonicum*. - *Phytochemistry* **42**: 1531-1536, 1996.
- Dat, J., Vandenabeele, S., Vranová, E., Van Montagu, M.,

- Inzé, D., Van Breusegem, F.: Dual action of the active oxygen species during plant stress responses. - *Cell. mol. Life Sci.* **57**: 779-795, 2000.
- Deo, P.M., Biswal, U.C., Biswal, B.: Water stress-sensitized photoinhibition in senescing cotyledons of clusterbean: Changes in thylakoid structures and inactivation of photosystem 2. - *Photosynthetica* **44**: 187-192, 2006.
- Dertinger, U., Schaz, U., Schulze, E.-D.: Age-dependence of the antioxidative system in tobacco with enhanced glutathione reductase activity or senescence-induced production of cytokinins. - *Physiol. Plant* **119**: 19-29, 2003.
- Doi, M., Shima, S., Egashira, T., Nakamura, K., Okayama, S.: New bile pigment extracted by a *Chlamydomonas reinhardtii* mutant a possible breakdown catabolite of chlorophyll *a*. - *J. Plant Physiol.* **150**: 504-508, 1997.
- Droillard, M.J., Bate, N.J., Rothstein, S.J., Thompson, J.E.: Active translation of the D1 protein of photosystem II in senescing leaves. - *Plant Physiol.* **99**: 589-594, 1992.
- Elstner, E.F.: Oxygen activation and oxygen toxicity. - *Annu. Rev. Plant Physiol.* **33**: 73-96, 1982.
- Engel, N., Jenny, T.A., Mooser, V., Gossauer, A.: Chlorophyll catabolism in *Chlorella protothecoides*. Isolation and structure elucidation of a red bilin derivatives. - *FEBS Lett.* **293**: 131-133, 1991.
- Esterbauer, H., Schaur, R.J., Zollner, H.: Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. - *Free Radical Biol. Med.* **11**: 81-128, 1991.
- Foyer, C.H., Noctor, G.: Oxygen processing in photosynthesis: a molecular approach. - *New Phytol.* **146**: 359-392, 2000.
- Friguet, B., Szveda, L.I., Stadtman, E.R.: Susceptibility of glucose-6-phosphate dehydrogenase modified by 4-hydroxy-2-nonenal and metal-catalyzed oxidation to proteolysis by the multicatalytic protease. - *Arch. Biochem. Biophys.* **311**: 168-173, 1994.
- Fu, J., Huang, B., Zhang, G.: Physiological and biochemical changes during seed filling in relation to leaf senescence in soybean. - *Biol. Plant.* **43**: 545-548, 2000.
- Ginsburg, S., Matile, P.: Identification of catabolites of chlorophyllporphyrin in senescent rape cotyledons. - *Plant Physiol.* **102**: 521-527, 1993.
- Green, P.J.: The ribonucleases of higher plants. - *Annu. Rev. Plant Physiol. Plant mol. Biol.* **45**: 421-445, 1994.
- Harman, D.: The aging process. - *Proc. nat. Acad. Sci. USA* **78**: 7124-7128, 1981.
- Hawkins, C.L., Davies, M.J.: Generation and propagation of radical reactions on proteins. - *Biochim. biophys. Acta* **1504**: 196-219, 2001.
- Hensel, L.L., Grbic, V., Baumgarten, D.A., Bleeker, A.B.: Developmental and age-related processes that influence the longevity and senescence of photosynthetic tissues in *Arabidopsis*. - *Plant Cell* **5**: 553-564, 1993.
- Hörtensteiner, S.: The loss of green color during chlorophyll degradation – a prerequisite to prevent cell death? - *Planta* **219**: 191-194, 2004.
- Iturraspe, J., Moyano, N.: A new 5-formylbilinone as a major chlorophyll *a* catabolite in tree senescent leaves. - *J. org. Chem.* **60**: 6664-6665, 1995.
- Jaybaskaran, C., Kuntz, M., Guillemaut, P., Weil, J.H.: Variations in the leaves of chloroplast transfer RNA and aminoacyl transfer RNA synthetases in senescing leaves of *Phaseolus vulgaris*. - *Plant Physiol.* **92**: 136-140, 1990.
- Jiménez, A., Hernandez, J.A., Pastori, G., Del Río, L.A., Sevilla, F.: Role of ascorbate-glutathione cycle of mitochondria and peroxisomes in the senescence of pea leaves. - *Plant Physiol.* **118**: 1327-1335, 1998.
- Kanazawa, S., Sano, S., Koshiba, T., Ushimaru, T.: Changes in antioxidative enzymes in cucumber cotyledons during natural senescence: comparison with those during dark-induced senescence. - *Physiol. Plant.* **109**: 211-216, 2000.
- Kar, M., Feierabend, J.: Metabolism of activated oxygen in detached wheat and rye leaves and its relevance to the initiation of senescence. - *Planta* **160**: 385-391, 1984.
- Kräutler, B.: Chlorophyll breakdown and chlorophyll catabolites. - In: Kadish, K.M., Smith, K.M., Guillard, R. (ed.): *The Porphyrin Handbook*. Vol. 13. Pp. 183-209. Elsevier, San Diego 2003.
- Kuran, H.: Changes in DNA, dry mass and protein content of leaf epidermic nuclei during aging of perennial monocotyledonous plants. - *Acta Soc. Bot. Pol.* **62**: 149-154, 1993.
- Lees, G.L., Thompson, J.E.: Lipid composition and molecular organization in plasma membrane-enriched fractions from senescing cotyledons. - *Physiol. Plant.* **49**: 215-221, 1980.
- Lohman, K.N., Gan, S., John, M.C., Amasino, R.M.: Molecular analysis of natural leaf senescence in *Arabidopsis thaliana*. - *Physiol. Plant.* **92**: 322-328, 1994.
- Makrides, S.C., Goldthwaite, J.: Biochemical changes during bean leaf growth, maturity and senescence. Contents of DNA, polyribosomes, ribosomal RNA, protein and chlorophyll. - *J. exp. Bot.* **32**: 725-735, 1981.
- Malik, N.S.A.: Senescence in oat leaves: changes translatable mRNAs. - *Physiol. Plant.* **70**: 438-446, 1987.
- Matile, P.: Chloroplast senescence. - In: Baker, N.R., Thomas, H. (ed.): *Crop Photosynthesis: Spatial and Temporal Determinants*. Pp. 413-440. Elsevier, Amsterdam 1992.
- Matile, P., Ginsburg, S., Schellenberg, M., Thomas, H.: Catabolites of chlorophyll in senescing leaves are localized in the vacuoles of mesophyll cell. - *Proc. nat. Acad. Sci. USA* **85**: 9529-9532, 1988.
- Matile, P., Hörtensteiner, S.: Chlorophyll degradation. - *Annu. Rev. Plant Physiol. Plant mol. Biol.* **50**: 67-95, 1999.
- Matile, P., Hörtensteiner, S., Thomas, H., Kräutler, B.: Chlorophyll breakdown in senescent leaves. - *Plant Physiol.* **112**: 1403-1409, 1996.
- McKersie, B.D., Lepock, J.R., Kruuv, J., Thompson, J.E.: The effects of cotyledon senescence on the composition and physical properties of membrane lipid. - *Biochim. biophys. Acta* **508**: 197-212, 1978.
- Munné-Bosch, S., Schwarz, K., Alegre, L.: Water deficit in combination with high solar radiation leads to midday depression of  $\alpha$ -tocopherol in field-grown lavender (*Lavandula stoechas*) plants. - *Aust. J. Plant Physiol.* **28**: 315-321, 2001.
- Noodén, L.N., Guimét, J.J., John, I.: Senescence mechanisms. - *Physiol. Plant.* **101**: 746-753, 1997.
- Paliyath, G., Droillard, M.J.: The mechanisms of membrane deterioration and disassembly during senescence. - *Plant Physiol. Biochem.* **30**: 789-812, 1992.
- Pastori, G.M., Del Río, L.A.: An activated-oxygen-mediated role for peroxisomes in the mechanism of senescence of *Pisum sativum* L. leaves. - *Planta* **193**: 385-391, 1994.
- Peñarrubia, L., Moreno, J.: Senescence in Plants and Crops. - In: Pessarakli, M. (ed.): *Handbook of Plant and Crop Physiology*. Pp. 461 - 481. Marcel Dekker, New York - Basel - Hong Kong 1995.
- Procházková, D., Sairam, R.K., Srivastava, G.C., Singh, D.V.: Oxidative stress and antioxidant activity as the basis of senescence in maize leaves. - *Plant Sci.* **161**: 765-771, 2001.
- Procházková, D., Wilhelmová, N.: Changes in antioxidative

- protection in bean cotyledons during natural and continuous irradiation-accelerated senescence. - *Biol. Plant.* **48**: 33-39, 2004.
- Procházková, D., Wilhelmová, N.: The capacity of antioxidant protection during modulated ageing of bean (*Phaseolus vulgaris*) cotyledons. 1. The antioxidant enzyme activities. - *Cell Biochem. Funct.* **25**: 87-95, 2007a.
- Procházková, D., Wilhelmová, N.: The capacity of antioxidant protection during modulated ageing of bean (*Phaseolus vulgaris*) cotyledons. 2. The low-molecular weight antioxidants. - *Cell Biochem. Funct.* **25**: 97-102, 2007b.
- Refsgaard, H.H., Tsai, L., Stadtman, E.R.: Modifications of proteins by polyunsaturated fatty acid peroxidation products. - *Proc. nat. Acad. Sci. USA* **97**: 611-616, 2000.
- Rodoni, S., Mühlecker, W., Anderl, W., Kräutler, B., Moser, D., Thomas, H., Matile, P., Hörtensteiner, S.: Chlorophyll breakdown in senescent chloroplasts. Cleavage of pheophorbide *a* in two enzymatic steps. - *Plant Physiol.* **115**: 669-676, 1997.
- Scebba, F., Sebastiani, L., Vitagliano, C.: Activities of antioxidant enzymes during senescence of *Prunus armeniaca* leaves. - *Biol. Plant.* **44**: 41-46, 2001.
- Shioi, Y., Tomita, N., Tsuchiya, T., Takamiya, K.: Conversion of chlorophyllide to pheophorbide by Mg-dechelating substance in extracts of *Chenopodium album*. - *Plant Physiol. Biochem.* **34**: 41-47, 1996a.
- Shioi, Y., Watanabe, K., Takamiya, K.: Enzymatic conversion of pheophorbide *a* to the precursor of pyropheophorbide *a* in leaves of *Chenopodium album*. - *Plant Cell Physiol.* **37**: 1143-1149, 1996b.
- Smirnoff, N.: The role of active oxygen in the response of plants to water deficit and desiccation. - *New Phytol.* **125**: 27-58, 1993.
- Srivalli, B., Khanna-Chopra, R.: Induction of new isoforms of superoxide dismutase and catalase enzymes in the flag leaf of wheat during monocarpic senescence. - *Biochem. biophys. Res. Commun.* **288**: 1037-1042, 2001.
- Suzuki, Y., Shioi, Y.: Detection of chlorophyll breakdown products in the senescent leaves of higher plants. - *Plant Cell Physiol.* **40**: 909-915, 1999.
- Thimann, K.V.: The senescence of leaves. - In: Thimann, K.V. (ed.): *Senescence in Plants*. Pp. 86-115. CRC Press, Boca Raton 1980.
- Thomas, H., Ougham, H.J., Davies, T.G.E.: Leaf senescence in a non-yellowing mutant of *Festuca pratensis* – transcripts and translation products. - *J. Plant Physiol.* **139**: 403-412, 1992.
- Thomas, H., Stoddard, J.L.: Leaf senescence. - *Annu. Rev. Plant Physiol.* **31**: 83-111, 1980.
- Thompson, J.E., Barber, R.F.: The role of free radicals in senescence and wounding. - *New Phytol.* **105**: 317-344, 1987.
- Tsuchiya, T., Ohta, H., Masuda, T., Mikami, B., Kita, N., Shioi, Y., Takamiya, K.: Purification and characterization of two isozymes of chlorophyllase from mature leaves of *Chenopodium album*. - *Plant Cell Physiol.* **38**: 1026-1031, 1997.
- Uchida, K., Stadtman, E.R.: Modification of histidine residues in proteins by reactions with 4-hydroxynonenal. - *Proc. nat. Acad. Sci. USA* **89**: 4544-4548, 1992.
- Valentine, J.S., Wertz, D.L., Lyons, T.J., Liou, L.L., Goto, J.J., Gralla, E.B.: The dark side of dioxygen biochemistry. - *Curr. Opin. Chem. Biol.* **2**: 253-262, 1998.
- Vicentini, F., Matile, P.: Gerontosomes, a multi-functional type of peroxisome in senescent leaves. - *J. Plant. Physiol.* **142**: 50-56, 1993.
- Watanabe, K., Ohta, H., Tsuchiya, T., Mikami, B., Masuda, T., Shioi, Y., Takamiya, K.: Purification and some properties of pheophorbidease in *Chenopodium album*. - *Plant Cell Physiol.* **40**: 104-108, 1999.
- Wilhelmová, N., Dominques, P.M.D.N., Srbová, M., Fuksová, H., Wilhelm, J.: Changes in nonpolar aldehydes in bean cotyledons during ageing. - *Biol. Plant.* **50**: 559-564, 2006.
- Ye, Z.Z., Rodriguez, R., Tran, A., Hoang, H., De los Santos, D., Brown, S., Vellanoweth, R.L.: The developmental transition to flowering represses ascorbate peroxidase activity and induces enzymatic lipid peroxidation in leaf tissue in *Arabidopsis thaliana*. - *Plant Sci.* **158**: 115-127, 2000.
- Zimmermann, P., Heinlein, C., Orendi, G., Zentgraf, U.: Senescence-specific regulation of catalases in *Arabidopsis thaliana* (L.) Heynh. - *Plant Cell Environ.* **29**: 1049-1060, 2006.