ORIGINAL ARTICLE

Biosorption of Pb^{2+} and Zn^{2+} by Non-Living Biomass of *Spirulina* sp.

Rajiv Kumar Aneja · Gunjan Chaudhary · Sarabjeet Singh Ahluwalia · Dinesh Goyal

Received: 9 March 2009/Accepted: 28 January 2010/Published online: 26 January 2011 © Association of Microbiologists of India 2011

Abstract Removal of heavy metals (Pb^{2+} , Zn^{2+}) from aqueous solution by dried biomass of *Spirulina* sp. was investigated. *Spirulina* rapidly adsorbed appreciable amount of lead and zinc from the aqueous solutions within 15 min of initial contact with the metal solution and exhibited high sequestration of lead and zinc at low equilibrium concentrations. The specific adsorption of both Pb^{2+} and Zn^{2+} increased at low concentration and decreased when biomass concentration exceeded 0.1 g 1^{-1} . The binding of lead followed Freundlich model of kinetics where as zinc supported Langmuir isotherm for adsorption with their r^2 values of 0.9659 and 0.8723 respectively. The adsorption was strongly pH dependent as the maximum lead biosorption occurred at pH 4 and 10 whereas Zn^{2+} adsorption was at pH 8 and 10.

Keywords Biosorption · Spirulina sp. · Lead · Zinc

List of symbols

Variables

- Q Metal ion adsorbed per unit mass of *Spirulina* (mg g^{-1}), at equilibrium
- V Volume of solution (1)
- C_i Initial concentration of metal ions in solution (mg l^{-1})
- C_f Final concentration of metal ions in solution (mg l^{-1})
- M Mass of biomass (g)

- q_{max} Langmuir parameter, maximum theoretical adsorption upon complete saturation of the surface (mg g⁻¹)
- b Langmuir constant related to the energy of adsorption desorption (1 mg^{-1})
- k Freundlich constant related to the strength of the adsorptive bond $(mg^{1-n}/g l^n)$
- n Freundlich constant related to bond distribution

Introduction

Environmental pollution due to toxic heavy metals is a significant worldwide problem due to their incremental accumulation in the food chain and continued persistence in the ecosystem [1]. Efforts have been made to remove toxic heavy metal contamination from the environment by using conventional technologies such as ion exchange or limit precipitation, which are sometimes inefficient and expensive, particularly for removal of heavy metal ions lower than 100 mg 1^{-1} , and also leads to generation of toxic sludge which further adds burden on the technoeconomic feasibility of the treatment process [2, 3].

In recent years, role of microbes in removal of toxic metal ions from the polluted effluent have taken more importance in this area as microorganisms due to their small size have a high surface area-to-volume ratio and therefore provide a large contact area for metal binding [4, 5]. In this race bacteria, algae, fungi, actinomycetes, yeast, activated sludge and various other biopolymers have been well recognized for heavy metal removal directly or indirectly [4, 6–12]. In this context the role of algae is known for a few decades, but has received increased attention only in recent years because of its potential for application in environmental protection or recovery of precious or strategic metals [13–16]. The affinity

R. K. Aneja · G. Chaudhary · S. S. Ahluwalia · D. Goyal (⊠) Department of Biotechnology and Environmental Sciences, Thapar University, Patiala 147004, Punjab, India e-mail: dineshgoyal63@gmail.com

of various algal species for binding of metal ions shows different hierarchies whereas, in general, metal ions with greater electronegativity and smaller ionic radii are preferably sorbed by algal biomass [16]. Metal accumulation capacity of algal biomass is either comparable or sometimes higher than chemical sorbents therefore algal biomass may serve as an economically feasible and efficient alternative to the existing physicochemical methods for metal removal and recovery from wastewaters. The major challenge in biosorption studies is to select the most promising types of biomass from an extremely large pool of readily available and inexpensive biomaterials [17]. Contribution of *Spirulina* (family Oscillatoriaceae) in the metal sorption is of considerable importance with adherent advantage of mass cultivation [18, 19].

In the present work, non-living biomass of *Spirulina* sp. was characterized for its Zn^{2+} and Pb^{2+} removal potential from synthetic metal solution whereas, the effect of pH, biomass concentration, initial metal concentration and time was studied for metal sorption process along with equilibrium sorption kinetics.

Materials and Methods

Biosorbent Material

The dry biomass of *Spirulina* sp. was procured from Global Green Company, Bangalore (India).

Lead and Zinc Sorption Experiments

Metal ion solutions were prepared by diluting 1.0 g l^{-1} of stock solutions, which were obtained by dissolving weight quantity of zinc sulphate heptahydrate (4.39 gm) and lead nitrate (1.59 gm) of analytical grade in double distilled water. Metal sorption studies were carried out to evaluate the capacity of non-living dry biomass of Spirulina to adsorb metal ions form solution. In batch process 100 ml of synthetic metal solution having different concentrations (20-100 mg l^{-1}) of zinc or lead were taken in 250 ml Erlenmeyer flasks with a range of biomass concentrations (0.1, 0.4, 0.8, 1.6 and 3.2 g) as biosorbent. Erlenmeyer flasks were kept under shaking condition at 100 rpm at ambient temperature. Samples were drawn after 5, 15, 30, 60, 120 and 240 min and filtered and analysed for residual metal concentration $(C_{\rm f})$ using atomic absorption spectrophotometer (GBC 932 AA; GBC Scientific Equipment Pvt. Ltd. Australia).

To see the effect of pH on Pb²⁺ and Zn²⁺ removal, a range of pH (2, 4, 6, 8 and 10) was adjusted with 0.1 M NaOH and 0.1 M H₂SO₄ in 100 ml metal synthetic solutions containing fixed concentration of metal at 50 mg 1^{-1} and biomass of 0.1 g followed by contact time of 30 min at 100 rpm. The metal adsorption (q) by the alga and bioremoval efficiency (R) were calculated by the following formulae [20, 21].

$$q = \frac{(C_{\rm i} - C_{\rm f})V}{M} \tag{1}$$

$$R(\%) = \frac{(C_{\rm i} - C_{\rm f})}{C_{\rm i}} \times 100$$
(2)

Where q = metal adsorption (mg/g); M = dry mass of alga (g); V = volume of initial metal ion solution used (L); R = bioremoval efficiency (%); $C_i =$ initial concentration of metal in aqueous solution (mg l⁻¹); $C_f =$ final concentration of metal in aqueous solution (mg l⁻¹).

Adsorption Isotherm

During biosorption, a rapid equilibrium is established between absorbed metal ion on the algal cell (q) and unabsorbed metal ions in the solution (C_f). This equilibrium represented by Langmuir and Freundlich adsorption isotherms, are widely used to analyse data for wastewater treatment application [22, 23]. Langmuir equation, which is valid for monolayer sorption onto a surface, a finite number of identical sites are given by Eq. 3.

$$q = q_{\max} \frac{bC_{\rm f}}{1 + bC_{\rm f}} \tag{3}$$

Where q_{max} is the maximum amount of the metal ion per unit weight of alga to form a complete monolayer on the surface bound at high C_{f} (mg l⁻¹), and b is a constant related to the affinity of the binding sites (l mg⁻¹), q_{max} represents a practical limiting adsorption capacity when the surface is fully covered with metal ions and assists in the comparison of adsorption performance, particularly in cases where the sorbent did not reach its full saturation in experiments. q_{max} and b can be determined from the liner plot of C_{f}/q versus C_{f} . The empirical Freundlich equation based on sorption on a heterogeneous surface is given below by Eq. 4.

$$q = kC_{\rm f}^{(1/n)} \tag{4}$$

where k and n are the constants, characteristic of the system. k and n are indicators of adsorption capacity and adsorption intensity, respectively Eq. 4 can be linearized in logarithmic form and Freundlich constants can be determined. Freundlich isotherm is also more widely used but provides no information on the monolayer adsorption capacity, in contrast to the Langmuir model.

Results

pH is one of the important parameters in metal sorption by *Spirulina* sp. [24] therefore metal sorption studies were



Fig. 1 pH-dependent sorption of Pb²⁺ and Zn²⁺ by *Spirulina* biomass (C_i Pb²⁺: 20 mg l⁻¹, Zn²⁺: 20 mg l⁻¹)

carried out at different pH values. Results revealed maximum biosorption of Pb^{2+} at pH 4 (82%) and 10 (75%) whereas, Zn^{2+} demonstrated the same at pH of 8 (90%) and 10 (89%) from synthetic solution containing 50 mg l^{-1} of metal concentration (Fig. 1). It was observed that Pb^{2+} and Zn^{2+} adsorption was less than 5% at pH 2. Biosorption studies carried out for both Pb^{2+} and Zn^{2+} in 100 ml solutions containing metals varying from 20 to 100 mg l^{-1} by 3.2 g of biomass exhibited effective role of initial metal concentration on metal removal. A consistent decrease in metal removal was observed by increasing external metal concentration (Fig. 2) where 90-91% Pb²⁺ removal efficiency was reported from 20 to 40 mg l^{-1} Pb²⁺ containing solution followed by 5% decline in solution containing 100 mg l^{-1} Pb²⁺ (Fig. 2). Similarly 35% decrease in Zn²⁺ removal efficiency was observed by increasing external metal concentration from 40 to 100 mg l^{-1} Zn²⁺ with 90% removal from solution containing 20 mg l^{-1} of Zn^{2+} (Fig. 2).

Rapid metal adsorption profile of Spirulina sp. was obtained for both Pb^{2+} and Zn^{2+} , which is important when the material is to be used for bioremediation. It exhibited rapid biosorption in first 0–15 min by removing 90% Pb^{2+} and 89% of Zn²⁺ from metal solutions thereafter increase in metal removal was marginal. Equilibrium was established between absorbed metal ions and the metal ions in the solution after 30 min with maximum removal of 90 and 93% of Pb^{2+} and Zn^{2+} (Fig. 3). Increase in metal removal efficiency from 20 mg l^{-1} synthetic metal solution for both Pb^{2+} and Zn^{2+} was observed on increasing biomass concentration (Fig. 4). Pb^{2+} adsorption was increased by 76.71% by increasing biomass concentration from 0.1 to $3.2 \text{ g } \text{l}^{-1}$ from 20 mg l^{-1} of lead containing synthetic solution whereas, same trend was observed in case of Zn^{2+} with the increase of 76.73%. Spirulina biomass displayed its equilibrium for Pb^{2+} removal at 0.8 g 1^{-1} of biomass concentration whereas, a continuous increase in Zn²⁺



Fig. 2 Initial concentration dependent sorption of Pb^{+2} and Zn^{+2} by *Spirulina* biomass



Fig. 3 Time dependent removal of Pb²⁺ and Zn⁺² by *Spirulina* biomass (C_i Pb²⁺: 20 mg l⁻¹, Zn⁺²: 20 mg l⁻¹)



Fig. 4 Effect of *Spirulina* biomass concentration on Pb²⁺ and Zn²⁺ removal. (C_i Pb²⁺: 20 mg l⁻¹, Zn²⁺: 20 mg l⁻¹)

removal on increasing biomass concentration was observed till 3.2 g l⁻¹. To the contrary of metal removal, metal adsorption studies from solution mass balance revealed a decline by 75 and 80% in sorption of Pb²⁺ and Zn²⁺ respectively by increasing biomass from 0.1 to 3.2 g l⁻¹ after 15 min of contact time. The results showed that amount of metal ion adsorbed per unit mass of *Spirulina* (mg g⁻¹), at equilibrium was 48.21 mg g⁻¹ Pb²⁺ and 8.75 mg g⁻¹ Zn²⁺ by 0.1 g l⁻¹ of biomass (Table 1).

 Table 1
 Langmuir and Freundlich parameters for the sorption of the test metals by Spirulina sp

Metal	Langmuir isotherm			Freundlich isotherm		
	<i>b</i> (1 mg ⁻¹)	r^2	$q_{\max} \pmod{(\mathrm{mg g}^{-1})}$	$\frac{k(\mathrm{mg}^{1-n})}{\mathrm{g}\;\mathrm{l}^n}$	n	r^2
Pb^{2+}	1.237	0.649	48.21	2.738	4.11	0.965
Zn ²⁺	1.57	0.872	8.75	1.073	7.31	0.743

A linear regression of the experimental results for Pb²⁺ and Zn²⁺ differed in terms of Pb²⁺ sorption fitted better to Freundlich isotherm and Zn²⁺ to Langmuir isotherms with r^2 values of 0.9659 and 0.8723 respectively (Table 1). The maximum adsorption per unit mass of *Spirulina* (mg g⁻¹), at equilibrium (q_{max}) for Pb²⁺ and Zn²⁺ were calculated to be 48.21 and 8.75 mg g⁻¹ respectively (Table 1).

Discussion

Enhanced adsorption of Pb^{2+} and Zn^{2+} ions at higher pH was observed which coincides with earlier findings where in most cases the removal efficiency increased steadily with rise in pH [9]. The adsorption of metal ions was lower at low pH because of high concentration of protons in the solution which competed with metal ions in forming a bond with the active sites on the surface of the algal biomass [25]. Selective sorption of specific metals due to distinct pH optima for their sorption may be due to differences in chemical composition of cell surface [16, 26]. A distinct relationship between pH of aqueous metal solution and involvement of functional group in binding of Pb²⁺ onto Spirulina maxima was observed at pH range of 2-5, 5-9 and 9-12 with the involvement of functional groups such as carboxyl, carboxyl and phosphate and carboxyl, phosphate and hydroxyl respectively [27]. Rapid metal adsorption profile of Spirulina sp. was obtained for both Pb²⁺ and Zn²⁺, which is important when the material is to be used for bioremediation. It exhibited rapid biosorption in first 0-15 min by removing 90% Pb²⁺ and 89% of Zn²⁺ from metal solutions thereafter increase in metal removal was marginal [19]. Decrease in metal removal on increasing initial metal concentration was supported by the findings of Mehta and Gaur [28] who observed that the removal of metal generally decreases with increasing concentration of metals in the solution. Algal cell surface has different functional groups with varying affinity for ionic species, where low and high affinity of functional groups in sorption of metal ions also depends upon concentration of metal ions therefore decline in metal removal is largely attributed to saturation of adsorption sites [19].

Spirulina biomass showed rapid biosorption in first 15 min, similar to the findings reported by Incharoensakdi and Kitjaharn [6] for the rapid adsorption of Zn^{+2} by Aphanotheca halophytica from aqueous solution. It has been reported that the sorption of heavy metal ions by algae followed a two-step mechanism where the metal ion is physically or chemically taken up onto the surface of the algal cell before being taken up biologically into the cell [29, 30]. The first step, known as a passive uptake which occurs rapidly, while the second biological step or active transport could take much longer time to complete. In this case, since the alga was dried and biological functions were no longer active, the sorption could only take place on the cell surface. The increase in Pb^{2+} adsorbed by increasing biomass was also expected as a result of increase in available area-to-volume ratio and therefore providing a large contact area for metal binding [4, 5]. In a similar study, 82.64% decrease in Pb²⁺ sorption capacity of Spirulina maxima with increasing biomass from 0.1 to 20 g 1^{-1} was reported although this is generally attributed to a shift in the sorption equilibrium [18]. The other probable explanation for such a relationship between biomass concentration and adsorption may be limited availability of metal, increased electrostatic interactions between binding sites and reduced mixing at higher biomass concentration [16].

The sorption isotherm is the relationship between equilibrium concentration of solute in the solution and equilibrium concentration of solute in the sorbent at constant temperature where either Freundlich or Langmuir model can describe the passive biosorption equilibrium of zinc and lead [22, 23]. An extremely high r^2 value of Freundlich isotherm for Pb²⁺ sorption indicated that ion exchange interaction takes place between metal ion and the biosorbent where as, Zn²⁺ follows Langmuir isotherm and thus supported physico–chemical interactions [12]. q_{max} for Pb²⁺ and Zn²⁺ adsorption by *Spirulina* sp. were 48.21 and 8.75 mg g⁻¹ respectively (Table 1), which is higher than earlier report of Sandau and his co-workers who observed 16.98 and 9.58 mg g⁻¹ of Pb²⁺ and Zn²⁺ adsorption respectively by *Spirulina platensis* [31].

Acknowledgments The authors thank the Director, Thapar University, Patiala, India, for providing the infrastructure and facilities and to University Grant Commission for providing the financial support vide Approval F-3-16/2002 (SR-II), dated March 16, 2002.

References

- Dudka S, Miller WP (1999) Accumulation of potentially toxic elements in plant and their transfer to human food chain. J Environ Sci Heal B 34:681–708
- Volesky B (1994) Advances in biosorption of metals: selection of biomass types. FEMS Microbiol Rev 14:291–302

- Ahluwalia SS, Goyal D (2007) Microbial and plant derived biomass for removal of heavy metals from wastewater. Bioresour Technol 98:2243–2257
- 4. Kim DW, Che DK, Wang J, Huang CP (2002) Heavy metal removal by activated sludge: influence of *Nocardia amarae*. Chemosphere 46:137–142
- Zouboulis AI, Loukidou MX, Matis KA (2004) Biosorption of toxic metals from aqueous solution by bacterial strains isolated from metal-polluted soils. Process Biochem 39:909–916
- Incharoensakdi A, Kitjaharn P (2002) Zinc biosorption from aqueous solution by halotolerant cyanobacterium *Aphanotheca halophytica*. Curr Microbiol 45:261–264
- 7. Yan G, Viraraghavan T (2003) Heavy-metal removal from aqueous solution by fungus *Mucor rouxii*. Water Res 37:4486–4496
- Oswald WJ (2003) My sixty years in applied algology. J Appl Phycol 15:99–106
- Pavasant P, Apiratikul R, Sungkhum V, Suthiparinyanont P, Wattanachira S, Marhaba TF (2006) Biosorption of Cu²⁺, Cd²⁺, Pb²⁺, and Zn²⁺ using dried marine green macroalga *Caulerpa lentillifera*. Bioresour Technol 97:2321–2329
- Polti MA, Amaroso MJ, Abate CM (2007) Chromium(IV) resistance and removal by actinomycete strains isolated from sediments. Chemosphere 67:660–667
- Xiong J, He Z, Liu D, Mahmood Q, Yang X (2008) The role of bacteria in the heavy metals removal and growth of *Sedum alfredii* Hance in an aqueous medium. Chemosphere 70:489–494
- Ahluwalia SS, Goyal D (2007) Microbial and plant derived biomass for removal of heavy metals from wastewater. Bioresour Technol 98:2243–2257
- Dönmez G, Aksu Z (2002) Removal of chromium(VI) from saline wastewaters by *Dunaliella* species. Process Biochem 38: 751–762
- Davis TA, Volesky B, Mucci A (2003) A review of the biochemistry of heavy metal biosorption by brown algae. Water Res 37:4311–4330
- Chojnacka K, Chojnacki A, Górecka H (2004) Trace element removal by *Spirulina* sp. from copper smelter and refinery effluents. Hydrometallurgy 73:137–153
- Mehta SK, Gaur JP (2005) Use of algae for removing heavy metal ions from wastewater: progress and prospects. Crit Rev Biotechnol 25:113–152
- Kratochvil D, Volesky B (1998) Advances in the biosorption of heavy metals. Trends Biotechnol 16:291–300

- Gong R, Ding Y, Liu H, Chen Q, Liu Z (2001) Lead biosorption and desorption by intact and pretreated *Spirulina maxima* biomass. Chemosphere 58:125–130
- Hong C, Shan-shan P (2005) Bioremediation potential of *Spirulina* toxicity and biosorption studies of lead. J Zhejiang Univ Sci B 6:171–174
- Volesky B (1992) Removal of heavy metals by biosorption. In: Ladisch MR, Bose A (eds) Harnessing biotechnology for the 21st Century. Amer Chem Soc, Washington DC, pp 462–466
- Zhang L, Zhao L, Yu Y, Chen C (1998) Removal of Pb²⁺ from aqueous solution by non-living *Rhizopus nigricans*. Water Res 32:1437-1444
- Langmuir I (1918) The adsorption of gases in plain surface of glass, mica, and platinum. J Am Chem Soc 40:1361–1403
- Freundlich H (1907) Uber die adsorption in lousumgen. Z F Phys Chem 57:385–471
- Solisio C, Lodi A, Soletto D, Converti A (2008) Cadmium biosorption on *Spirulina platensis* biomass. Bioresour Technol 99:5933–5937
- Parent L, Campbell PGC (1994) Aluminium bioavailability to the green alga *Chlorella pyrenoidosa* in acidified synthetic soft water. Environ Toxicol Chem 13:587–598
- Aksu Z, Kutsal T (1990) Comparative study for biosorption characteristic of Zn ion with C. vulgaris. Environ Technol 11: 979–987
- 27. Chojnacka K, Chojnacki A, Górecka H (2005) Biosorption of Cr³⁺, Cd²⁺ and Cu²⁺ ions by blue-green algae *Spirulina* sp: kinetics, equilibrium and the mechanism of the process. Chemosphere 59:75–84
- Mehta SK, Gaur JP (2001) Removal of Ni and Cu from single and binary metal solution by free and immobilized *Chlorella vulgaris*. Eur J Protistol 37:261–271
- Volesky B (1990) Removal and recovery of heavy metals by biosorption. In: Volesky B (ed) Biosorption of heavy metals. CRC Press, Boca Raton, pp 7–43
- 30. Kojima H, Lee KY (2001) Photosynthetic microorganisms in environmental biotechnology. Springer-Verlag, Hong Kong
- Sandau E, Sandau P, Pulz O (1996) Heavy metal sorption by microalgae. Acta Biotechnol 16:227–235