

Erratum to: Infestation Rates and Tiller Morphology Effects by the Switchgrass Moth on Six Cultivars of Switchgrass

Veronica Calles Torrez · Paul J. Johnson · Arvid Boe

Published online: 22 March 2013
© Springer Science+Business Media New York 2013

Erratum to: Bioenerg Res DOI 10.1007/s12155-013-9306-7

In Table 3, of the original version, the column headings were incorrect. The correct version is shown here.

Table 3 Weights of healthy tillers and tillers infested by the larva of the switchgrass moth at Brookings, SD, USA

Tiller type	Year	
	2011	2012
	(mg tiller ⁻¹)	
Healthy	607 ^a	657 ^a
Infested	206	185

^a Comparison of healthy and infested tiller significant at 0.01 level by *t* test

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12155-013-9306-7>.

V. Calles Torrez · P. J. Johnson (✉)
Insect Biodiversity Lab, Department of Plant Science, South
Dakota State University, Brookings, SD, Box 2207A, 57007, USA
e-mail: paul.johnson@sdstate.edu

V. Calles Torrez
e-mail: veronica.callestorrez@sdstate.edu

A. Boe
Department of Plant Science, South Dakota State University,
Brookings, SD, Box 2140C, 57007, USA
e-mail: arvid.boe@sdstate.edu