



In vitro and in vivo toxicity assessment of nanoparticles

Vinay Kumar¹ · Neha Sharma² · S. S. Maitra¹

Received: 26 November 2016 / Accepted: 13 November 2017 / Published online: 25 November 2017

© The Author(s) 2017. This article is an open access publication

Abstract

Nanotechnology has revolutionized gene therapy, diagnostics and environmental remediation. Their bulk production, uses and disposal have posed threat to the environment. With the appearance of these nanoparticles in the environment, their toxicity assessment is an immediate concern. This review is an attempt to summarize the major techniques used in cytotoxicity determination. The review also presents a detailed and elaborative discussion on the toxicity imposed by different types of nanoparticles including carbon nanotubes, gold nanoparticles, silver nanoparticles, quantum dots, fullerenes, aluminium nanoparticles, zinc nanoparticles, iron nanoparticles, titanium nanoparticles and silica nanoparticles. It discusses the in vitro and in vivo toxicological effects of nanoparticles on bacteria, microalgae, zebrafish, crustacean, fish, rat, mouse, pig, guinea pig, human cell lines and human. It also discusses toxicological effects on organs such as liver, kidney, spleen, sperm, neural tissues, liver lysosomes, spleen macrophages, glioblastoma cells, hematoma cells and various mammalian cell lines. It provides information about the effects of nanoparticles on the gene-expression, growth and reproduction of the organisms.

Keywords Nanoparticles · Toxicity assessment · In-vivo toxicity · In-vitro toxicity · Diagnostics

Abbreviations

CdSe/ZnS	Cadmium selenide/zinc sulphide
CdTe	Cadmium telluride
CNTs	Carbon nanotubes
DCFDA	2',7'-Dichlorofluorescein diacetate
DTNB	5,5'-Dithiobis-(2-nitrobenzoic acid)
EPR	Electron paramagnetic resonance
HBMVECs	Human brain microvascular endothelial cells
HEK	Human epidermal keratinocyte
HMSC	Human bone marrow derived mesenchymal stem cells
LD50	Lethal dose 50
LDH	Lactate dehydrogenase
MTT	3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
MW-CNTs	Multi-walled carbon nanotubes
PI	Propidium iodide
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
SCGE	Single cell gel electrophoresis assay

SOD	Superoxide dismutase
SW-CNTs	Single-walled carbon nanotubes
TEMP	2,2,6,6-Tetramethylpiperidine
TUNEL	TDT-mediated dUTP-biotin nick-end labelling
WTS-1	Water soluble tetrazolium salts
XTT	2,3-Bis-(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5-carboxanilide

Introduction

Nanoparticles are produced in the size range 1–100 nm [1]. With the advancement in the technology, there has been a tremendous growth in their applications [2, 3]. Nanoparticles are used in additives for paints, ceramics, foods, paper, packaging, drug delivery, biosensor and cancer therapy [4]. They are also used as tumour detector [5], paclitaxel [6] and radiotherapy dose enhancer [7, 8]. They are in demand due to their small size and greater surface area to volume ratio [9]. These properties result in higher chemical reactivity and increased reactive oxygen (ROS) production [10, 11]. Nanoparticles have received much attention due to their toxicity imposed on the environment during production and disposal of consumer products [12] (Fig. 1). They impose toxicity by various processes. Nanoparticles can easily cross the cell


✉ Vinay Kumar
vkmjnu@gmail.com

¹ Lab No. 117, School of Biotechnology, Jawaharlal Nehru University, New Delhi 110067, India

² Arbro Pharmaceuticals Pvt Ltd., New Delhi 110015, India



Fig. 1 Applications of nanoparticles in different occupation



membranes and interact with intracellular metabolism [13]. ROS formation is one of the mechanisms for nanoparticle toxicity [14, 15]. Interaction of nanoparticles with cells induces pro-oxidant effects leading to ROS generation, mitochondrial respiration and NADPH-dependent enzyme systems [16–18]. Upon internalization of nanoparticles, phagocytosis induces the production of reactive oxygen species ROS [17–19].

The reviews published so far on nanoparticle toxicology provide information about toxicity of engineered nanoparticles to environmental microorganisms [20], titanium oxide nanoparticle toxicity [21], gold nanoparticle toxicity [21], risk management of inhaled nanoparticles [22], induced mitochondrial toxicity of silver nanoparticles [23] and single-wall carbon nanotube toxicity [24] and multi-wall carbon nanotube toxicity [25]. The published reviews are either too general or very specific (focusing on a single nanoparticle or toxicological effects on a particular organ) to explain the toxicity. Second, these reviews are unable to provide a comprehensive and detailed information on the toxicity assessment on the higher organisms and cell lines.

This review presents a detailed and elaborative discussion on the toxicity imposed by the nanoparticles on rat, mouse, pig, guinea pig, human cell lines and human. The review also focuses on (1) summarizing the techniques useful in determining the toxicity of nanoparticles. (2) Determining the toxic effects of nanoparticles (carbon nanotubes, gold nanoparticles, silver nanoparticles, aluminium nanoparticles and quantum dots, etc.) both in vitro and in vivo and (3) evaluating the effect of nanoparticles on the gene-expression, growth, behaviour and reproduction of organisms.

Assessment of nanoparticle toxicity

Various methods are available for the toxicity assessment imposed by nanoparticles on the organisms. Figure 2 presents the types of nanoparticles, experimental models and

toxic effects imposed by nanoparticles. The methods for toxicity assessment can be categorized as in vitro and in vivo.

In vitro assessment methods

In vitro nanoparticle toxicity assessment is one of the important methods. The advantages include lower cost, faster and minimum ethical concerns. Assessment can be subdivided into proliferation assay, apoptosis assay, necrosis assay, oxidative stress assay and DNA damage assays.

Proliferation assays

This assay is used to measure the cellular metabolism by assessment of metabolically active cells. 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) is the most commonly used tetrazolium salt for in vitro toxicity assessment of nanoparticles [26]. The technique is advantageous due to quick yields, reproducible results and minimum manipulation of the model cells [27]. The assay is based on the measurement of tetrazolium salt and it can sometimes lead to altered measurement due to changes in the culture media additives [28], media pH [29], ascorbate [30] and cholesterol [31]. The MTT assay also produces formazan; therefore, the assays such as such XTT or WST-1 which produce soluble dyes are preferred. [³H] thymidine incorporation is a method used for assessment of cellular proliferation, but this method is avoided due to toxicity and relatively high cost [32]. Alamar Blue is used to measure the cellular redox potential and advantageous as compared to MTT assay due to simpler sample preparation [33]. But the success of the Alamar Blue is hindered due to unavailability of the biochemical mechanisms of the assay and reaction of non-porous silicon with Alamar Blue in the absence of the cells [34]. Another assay is known as cologenic assay where the proliferating cells are counted by visual inspection after nanoparticle exposure is also used [35].



Types	Experimental models	Toxic effects
Carbon nanoparticles	<ul style="list-style-type: none"> Bacteria, microalgae, crustacean, zebrafish, <i>Drosophila melanogaster</i>, <i>Caenorhabditis elegans</i> Fish, oysters, Pigs, guinea pig, mouse, rat, 	<ul style="list-style-type: none"> Metabolic activity, membrane activity Cell viability Mitochondrial damage Mitochondrial integrity Reactive oxygen species production
Gold nanoparticles		
Silver nanoparticles		
Quantum dots	<ul style="list-style-type: none"> Astrocyte Breast cancer cells Human epidermal keratinocyte cells Hippocampal neuronal cells Human hepatoma cells Human hepatocytes Human lymphocytes Carcinoma cells Human bronchial epithelial cells Mesenchymal stem cells Mouse lung epithelial cells 	<ul style="list-style-type: none"> Oxidative stress DNA damage Apoptosis Actin filament integrity Blood brain barrier destruction Alteration of gene expression
Fullerenes		
Aluminum nanoparticles		
Zinc nanoparticles		
Iron nanoparticles		
Titanium nanoparticles		
Silicon nanoparticles		
	<ul style="list-style-type: none"> Liver, spleen, colon mucosa, kidney Sperms, lung, gill, pulmonary organs Human skin 	

Fig. 2 Nanoparticles types, experimental models used for the studies and toxic effects of nanoparticles

Apoptosis assay

Apoptosis is one of the major markers observed in the in vitro assessment of nanoparticle toxicity. Generation of excessive free radical is considered the cause of apoptosis and DNA damage [36, 37]. Evidence suggested that apoptosis and DNA damage can be caused by oxidative stress in cell culture systems [36]. Many studies have reported the apoptosis induced by nanoparticles. In vitro studies indicated that silver nanoparticles caused apoptosis in mouse embryonic stem cells [38]. In another investigation, the release of apoptosis markers viz. caspase-3 and caspase-9 were examined on the treatment of larval tissues of *Drosophila melanogaster* with silver nanoparticles at concentrations of 50 and 100 µg/ml for 24 and 48 h. The results suggested the involvement of silver nanoparticles in the apoptotic pathway of *D. melanogaster* [39]. Up-regulation of p38 protein expression was also demonstrated in the exposure of silver nanoparticles in *D. melanogaster* in time- and dose-dependent manner. Up-regulation of genes and extensive DNA insult is responsible for inducing cell death and a cascade of apoptosis pathway [40, 41].

There are a number of methods for assessment of apoptosis. These include Annexin-V assay [42], Comet assay [43], TdT-mediated dUTP-biotin nick end labelling (TUNEL) assay [44] and inspection of morphological changes [45]. The DNA laddering technique is a technique which is used to visualize the endonuclease cleavage products of apoptosis

[46]. An irregular reduction in the size of cells and by DNA fragmentation confirms the induction of apoptosis. Agarose gel electrophoresis can easily discriminate between apoptotic and necrotic modes of cell death [47, 48]. Genomic fragments obtained during electrophoresis of irregular sizes are typical of necrotic cells and a ladder-like electrophoretic pattern indicates apoptotic internucleosomal DNA fragmentation [49]. Annexin-V and propidium iodide (PI) are typical cell death markers used in toxicity assessment. The assay works on the principle that when Annexin-V binds to phosphatidylserine, it shows increased fluorescence and hence indicates the externalization of the plasma membrane. This externalization of the plasma membrane is induced by activation of the caspase-dependent pathway. PI is an impermeable dye which stains the nucleus only when the integrity of the cell membrane is lost, which can be related to the late stage of apoptosis [21, 50]. When human HepG2 hepatoma cells were treated with silica nanoparticles, morphological changes in the nucleus and induction of apoptosis was observed [51]. The investigation demonstrated the induction of apoptosis measured by Annexin V/PI in HeLa cell lines treated with gold nanoparticle [52]. Single cell gel electrophoresis assay (SCGE), or comet assay, is a sensitive tool for the detection of the mutagenic potential of a test material [53]. It is used to detect single- and double-stranded DNA breaks in individual cells, both in vitro and in vivo [54, 55]. It is also used to quantify oxidative DNA damage, alkali-labile sites, DNA–DNA or DNA–protein cross-links



and abasic sites [56–58]. The assay is based on the principle that the damaged DNA fragments will migrate out of the cell when an electric current is applied, whereas the undamaged DNA will remain in the cell nucleus. In assay, damaged DNA resemble the tail and the intact DNA resemble the head. The extent of DNA damage is correlated with the size and shape of the tail and the distribution of DNA within the comet [59–61]. In the procedure, the cells are lysed to remove cellular protein and the damaged DNA is allowed to migrate away from the nucleus by undergoing electrophoresis. DNA-specific fluorescent dye is used to stain the samples. The gel is then analysed for the amount of fluorescence in the head and tail and the tail length [62–64]. Comet assay was used to assess the toxicity of zinc oxide nanoparticles at 25 mg Zn/L on *D. tertiolecta* which resulted in 55% nuclei damage [65]. In another investigation, comet assay was used to measure the toxicity imposed by SiO₂ nanoparticles on *D. tertiolecta* at 125 mg/L, which resulted in increasing genotoxic effects after 72 h [66]. In the similar study on *D. tertiolecta*, using TiO₂ nanoparticles for 24 h resulted in more than 70% damage to the nuclei after 72 h [66].

TUNEL assay is one of the most widely used methods for detecting DNA damage in situ TUNEL staining [67]. IT was initially described as a method for staining cells that have undergone apoptosis or programmed cell death and internucleosomal DNA fragmentation [47, 68–70]. TUNEL assay is based on the ability of the enzyme terminal deoxynucleotidyl transferase to incorporate labelled dUTP into free 3'-hydroxyl termini generated by the fragmentation of DNA [67]. IT is not limited to the detection of apoptotic cells only. It can be used to detect DNA damage associated with non-apoptotic events including necrotic cell death which is induced by exposure to toxic compounds [71]. TUNEL assay was reported to stain cells undergoing active DNA repair [72]. The pancreatic islet function of Goto Kakizaki rats was estimated by TUNEL assay of pancreatic β -cells after treatment with insulin-loaded selenium nanoparticles (25 IU/kg) daily for 2 weeks. In TUNEL assay, apoptotic cells declined to 2.3 % from initial 17.6 % in the treatment [73].

Necrosis assay

Necrosis is measured by the integrity of the membrane and it is commonly used to determine the viability of the cells. Membrane integrity is measured by uptake of the dye such as Neutral Red [74] and Trypan Blue [75]. The need for a reliable, rapid, inexpensive and reproducible quantitative in vitro assay for screening of nanoparticles is generally acknowledged. Neutral red (2-amino-3 methyl-7-dimethyl-aminophenazoniumchloride) is a weakly cationic supravital dye which at slightly acid pH yields a deep red colour. The Neutral red readily diffuses through the plasma membrane. It concentrates in the lysosomes and binds by electrostatic

hydrophobic bonds with anionic sites in the lysosomal matrix [76, 77]. Alterations of the cell surface lead to lysosomal fragility [76]. Such changes brought about by the action of xenobiotics or nanoparticles [77, 78] can result in a decreased uptake and binding of Neutral red. It is thus possible to distinguish between viable and dead cells [79, 80]. In a study, endosome–lysosome system stability which was measured by neutral red assay decreased after exposure to the silver nanoparticle (30%) [81]. Another method is called Trypan blue exclusion test. The dye trypan blue enters dead cells and is excluded from living cells [82]. A trypan blue exclusion assay was performed for the evaluation of cell membrane stability. The medium was then replaced with zinc nanoparticles at 12, 61, 123, 184, 369 and 737 μ M. The results demonstrated that Zn compounds exerted considerable cytotoxicity at 369 μ M and higher [83].

Oxidative stress assay

Exposure of nanoparticles leads to the production of reactive ROS and reactive nitrogen species (RNS) [84]. The method for detection of ROS and RNS involves the reaction of 2,2,6,6-tetramethylpiperidine (TEMP) with O₂^{•−} stable radical which can be detected using X-band electron paramagnetic resonance (EPR) [85]. The application of these methods is hindered due to their high cost. Fluorescent probe molecules have emerged as an alternative and cost-effective approach [86]. But there are limitations with fluorescent probes as they are inefficient due to their ability to react with a variety of reactive species. This property leads to misleading results sometimes [87]. The above problems can be solved by the use of 2',7'-dichlorofluorescein diacetate (DCFDA), a non-fluorescent probe. DCFDA is reactive to HO[•], RO[•], ROO[•] and H₂O₂ in the presence of cellular peroxidases [88]. Oxidative stress can also be assessed by measuring lipid peroxidation C11-BIODIPY assay and TBA assay for malondialdehyde [89]. Availability of numerous other assays makes the assessment much convenient. These assays include lipid hydro peroxide's measurement using Amplex Red assay, antioxidant depletion measurement by 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB) and superoxide dismutase (SOD) activity by Nitro blue tetrazolium assay [90].

In vivo toxicity assessment methods

The in vivo toxicity assessment is normally performed on animal models such as mice and rat. The assessment methods for in vivo toxicity include bio distribution, clearance, haematology, serum chemistry and histopathology. Biodistribution studies examine the localization route of nanoparticles to the tissue or organ. Nanoparticles are detected in the killed or live animals through radiolabels [91]. Clearance of



nanoparticles is performed by the examination of excretion and metabolism of nanoparticles at various time points after exposure [92]. Another method for in vivo toxicity assessment is the examination of changes in the serum chemistry and cell type after exposure of nanoparticles [93]. Histopathology of the cell, tissue or organ after exposure is used to determine the toxicity level caused by a nanoparticle [94]. Histopathology examination has been used to nanoparticles' exposed tissues such as lung, eyes, brain, liver, kidneys, heart and spleen [93, 95]. The advancement of toxicity assessment includes use of micro-electrochemistry and microfluidics [96].

In-vitro toxicity of nanoparticles

Cell viability and lethality

Cell viability and lethality are the two parameters which are used to measure the toxicity caused by the nanoparticles. Among the various nanoparticles, carbon nanoparticles (CNTs) are used most frequently for assessment of viability and lethality of cells. They are widely used in chemical, industrial and biomedical applications due to their unique properties [97, 98]. They are synthesized as single-walled carbon nanotubes (SW-CNTs) and multi-walled carbon nanotubes (MW-CNTs) [99]. The anti-microbial properties of CNTs have been observed by studies in various bacteria due to the mechanical damage caused by the nanotubes [100–102]. A recent study has indicated that functionalized CNTs affect soil bacterial diversity [25]. The toxicity studies on a micro crustacean (*Daphnia magna*), freshwater microalgae (*Raphidocelis subcapitata* and *Chlorella vulgaris*) and a fish (*Oryzias latipes*) revealed inhibited the growth of the algae *C. vulgaris* and *R. subcapitata* with effective SW-CNT concentration 30.96 mg/mL and 29.99 mg/mL, respectively [103]. The nanoparticles synthesized in the form of iron oxide were also reported toxic in murine macrophage cells, human macrophages, human hepatocellular carcinoma cells and rat mesenchymal stem cells. Iron oxide nanoparticles reported toxicity at 25–200 µg/mL for 2 h exposure on murine macrophage cells. The study observed effects include the decrease in cell viability [104]. Another study reported a reduction in the cell viability when murine macrophage cells were treated with 0.1 mg/mL iron oxide nanoparticles for 7 days [105]. Another toxicity study which was performed on rat mesenchymal stem cells at 0.1 mg/mL for 2 days reported a decrease in cell viability [106]. Silica nanoparticles were also reported for toxicity to human keratinocytes. In a study, the toxicity of the silica nanoparticles at 30–300 µg/mL was evaluated using CHK (human keratinocytes). The results suggested a decrease in cell viability [107].

Effects on cell lines

The toxicity of nanoparticles was evaluated on various cell lines. The effect of SW-CNTs was observed by various researchers on human cell lines, including human HEL 293 cells, HEK cells, A549 cells, human macrophage cells and human epithelial-like Hela cells [108–110]. In a study lung fibroblast cells were treated with CNTs to evaluate the toxicity [53]. In another study when A549 cells were exposed to SW-CNTs at 250–500 µg/ml for 72 h, it resulted in oxidative response and membrane damage, induced by inflammatory response [111]. Another study reported suppression of inflammatory mediators including IL-6, IL-8 and MCP-1 in vitro [112]. The effects of multi-wall carbon nanotube were also evaluated on human epidermal keratinocytes [74]. It was suggested that toxicity induced by multi-wall carbon nanotubes is mediated by pro-inflammatory effects which are facilitated by NF-κB and ROS [113]. In vitro studies reported various toxicological effects of MW-CNTs including oxidative stress, DNA damage and apoptosis in mammalian cells lines. Other effects include VE-cadherin distribution and actin filament integrity in human aortic endothelial cells [114–118].

Gold nanoparticles (AuNPs) in MRC-5 human lung fibroblasts induced autophagy with oxidative stress [119]. In a study, the cellular motility was used to demonstrate the cytotoxicity of metal and semiconductor nanoparticles on animal cells. The study utilizes the potential of electrical cell-substrate impedance analysis as a highly suitable method to quantify the in vitro cytotoxicity of gold nanorods and quantum dots. The method was validated by fluorescence and dark field microscopy [120]. Toxicity of starch-coated silver nanoparticles was studied on human glioblastoma cells (U251) and human lung fibroblast cells (IMR-90). The study resulted in dose-dependent reduction in ATP content and DNA damage. The study demonstrated that the DNA damage is due to silver nanoparticle deposition and interaction with DNA followed by cell cycle arrest in G₂/M phase [121]. Cytotoxic study on fibroblast cells NIH3T3 showed that silver nanoparticles induced mitochondria-dependent apoptosis associated with JNK activation and reactive oxygen species [122]. Toxicity of silver nanoparticles was also evaluated on human hematoma cell line HepG2 using micronucleus test, viability assay and DNA microarray analysis [123]. A study reported the toxicity of silver nanoparticles in HeLa cells. Exposure resulted in upregulation of ho-1, mt-2A and oxidative stress genes [124]. The results of three different types of silver nanoparticle treatment on *E. coli* suggested compromised replication fidelity of the risk gene [125].

Long-term exposure of CdTe quantum dots was observed on the live cells. The study was focused on assessment of intracellular Cd²⁺ concentration in human breast cancer cells (MCF-7) which were treated with cadmium telluride



(CdTe), cadmium selenide/zinc sulphide (CdSe/ZnS) nanoparticles capped with cysteamine, mercaptopropionic acid and N-acetyl cysteine conjugated to cysteamine. The study demonstrated that CdTe quantum dots were cytotoxic with significant lysosomal damage and production of ROS [126]. The effects of different surface coatings on quantum dots were observed with relation to the toxicity on the human epidermal keratinocytes (HEKs). The results suggested that carboxylic acid-coated quantum dots enhanced the release of IL-1 β , IL-6 and IL-8. It was determined that surface coating is the primary determinant of immunotoxicity and cytotoxicity in HEKs [127]. The cytotoxicity of thiols stabilized CdTe, CdTe/CdS/ZnS core-shell-shell structured quantum dots and CdTe/CdS core shell structure was observed on cell lines including HEK293T and K562. The results demonstrated that CdTe QDs was highly toxic for cells [128]. A hippocampal neuronal culture model was used for the neurotoxicity investigation of cadmium selenium quantum dots. The study focused on voltage-gated sodium channel and cytoplasmic calcium level. The results evidenced induced neuron death and elevated cytoplasmic calcium levels [129]. The exposure of CdSe quantum dot on enterocyte-like Caco-2 cells as a model for intestine epithelium was investigated. Results suggested that acid treatment of PEG-coated quantum dots increased the toxicity [130]. Another study investigated the in vitro and in vivo toxicity of CdTe nanoparticles on human hepatoma HepG2 cells [131].

The aluminium nanoparticles in the size range 1–10 μm were used for 24 h on human brain microvascular endothelial cells (HBMVECs). The study demonstrated that treatment resulted in the decrease in mitochondrial function, cell viability and an increase in oxidative stress [132]. Mammalian cells were treated with 10–400 $\mu\text{g/mL}$ aluminium nanoparticles to examine the toxic effects. The observed results suggested that no significant toxicity was observed on cell viability in the study [133]. In another study cell viability was determined with respect to the interaction of human bone marrow derived mesenchymal stem cells (HMSC) with aluminium nanoparticles in the range 25–40 $\mu\text{g/mL}$. The results from the study indicated a decrease in the cell viability [134]. Another study investigated the effect of increasing concentration of aluminium nanoparticles on rat blood cells at 500–2000 mg/kg for 72 h. It was observed from the study that the toxicity was dose dependent [135]. A study conducted on mammalian cell lines suggested that at 0–5000 $\mu\text{g/mL}$, aluminium nanoparticles were responsible for DNA damage after treatment for 2 h [136].

Mechanistic studies

Mechanistic studies were performed to assess the effects of nanoparticles in vitro. In vitro studies have revealed CNTs disrupt the membrane potential, membrane integrity,

metabolic activity and cellular reproduction [137, 138]. Gold nanoparticles are responsible for mitochondrial damage, affecting cellular micro mobility, autophagy and oxidative stress [119, 139]. In vitro studies on silver nanoparticles' toxicity have suggested that interference with DNA replication, fidelity, apoptosis, oxidative stress, cytotoxicity, chromosome instability, intracellular calcium transients, JNK activation and cell cycle arrest in mammalian cells [121–125, 140, 141]. Fullerenes are responsible for DNA damage and oxidative stress in mammalian cells lines [142–144]. In a study on FE1-MutaTM mouse lung, epithelial cells were investigated to observe the effects of C₆₀ fullerenes and SW-CNTs for cytotoxicity, genotoxicity and ROS production [142].

In-vivo toxicity of nanoparticles

Dose and LD50

The toxicity of nanoparticles is determined by the exposure conditions, exposure duration and dose. In a study, 3T3 cells were treated with 1–10 mm size SWCNTs with an exposure time of 1 h. The treatment resulted in 20% viability [145]. In another study immortalized epidermal keratinocytes with 80% confluency were treated with SWCNTs in concentration range 0.06–0.24 mg/mL for 2, 4, 6, and 8 h. The treatment resulted in decreased viability after 4 h and ~ 65% viability on exposure to nanoparticles at 0.24 mg/mL [146]. When human embryonic kidney cells were treated with SWCNTs at 0.7812–200 $\mu\text{g/mL}$ for 24–120 h; then it was observed that cytotoxicity was time and dose-dependent with G1 cell cycle arrest in 43.5% cells after day 1 [147]. But when mouse peritoneal macrophage-like cells were treated with SWCNTs in the range 0–7.3 $\mu\text{g/mL}$ for 4, 8, 12 and 18 h, the cells ingested the nanoparticles without toxic effects [148]. When guinea pig alveolar macrophages were treated with SWCNTs at 1.41–226 $\mu\text{g/cm}^2$ for 3 h, the study resulted in cytotoxicity at 0.38 $\mu\text{g/cm}^2$ and necrosis at 3.06 $\mu\text{g/cm}^2$ [149]. It was observed that the cytotoxicity of SWCNTs was dose dependent and 60–80% reduction in cell number was observed when rat alveolar epithelial cells, NR8383 and human alveolar epithelial cells, A549 were treated at 5–100 $\mu\text{g/mL}$ for 24–96 h [150]. Strongest adverse effect of SWCNTs was observed when HEKs were treated at 0.8–100 $\mu\text{g/mL}$ for 24–120 h. The study resulted in 79, 50 and 31% viability at 100 $\mu\text{g/mL}$ when treated for 1, 3 and 5 days, respectively [151]. The toxicity of MWCNTs was also evaluated on different cell lines. In a study when HEK cells were treated with MWCNTs with 80% confluency at 0.1–0.4 mg/mL for 1, 4, 8, 12, 24 and 48 h, the study resulted in ~ 73% viability at 0.4 mg/mL. When purified



MWCNTs were used to examine the toxicity on human skin fibroblasts with 70% confluency at 0.06–0.6 mg/mL for 24 and 48 h, it was observed that the toxicity was dose dependent [152].

Gold nanoparticles' exposure at 150 pm for 3 h on HeLa and 3T3/NIH cells resulted in cell viability reduction by 20 and 5%, respectively [153]. In another study, gold nanoparticles were exposed to macrophage cells with particle size range 10–100 μm for 24–72 h. The study demonstrated that the 100 μm nanoparticles decreased the cell viability to 85% in 72 h [154]. In a study on human dermal fibroblasts, cells were used to assess the toxicity imposed by gold nanoparticles at 0–0.8 mg/mL for 2–6 days. The results from the study demonstrated dose-dependent decrease in the cell area and density [155]. Gold Nano shell was also used to study the cytotoxic effects. A study conducted on Vero cells at 0.001–200 $\mu\text{g/mL}$ for 6 and 24 h resulted in decreased cell viability on exposure to gold Nano shells [156].

Fe_3O_4 nanoparticles' exposure to human fibroblast cells at 0–1000 $\mu\text{g/mL}$ for 24 h resulted in 25–50% reduction in cell viability [157]. In another study conducted on mouse macrophages, Fe_3O_4 nanoparticles were exposed at 0.2 mg/mL for 1 and 4 days. The study demonstrated that cytotoxicity was dose dependent [158]. When human breast carcinoma SK-BR-3 cells were exposed to Fe_3O_4 nanoparticles at 10–400 nm for 1–48 h, the study resulted in 91% viability [159].

Treatment of Hela cells with CdSe quantum dots with sizes 1, 10 and 100 nm for 2 h resulted in survival of 90% cells [160]. In another experiment human lymphoblastoid cells were treated with CdSe quantum dots at 0.2 μm size for 12 h; the study resulted in decreased cell activity [161]. When three cell lines viz. primary hepatocytes, Hela cells and Vero cells, were treated with 0–4 mg/mL CdSe quantum dots for 24 h, the study resulted in low damage at 0.1 mg/mL, while increased damage was observed at 0.2 mg/mL. HepG2 cells' and Wister mice cells' treatment with CdSe quantum dots at 10–400 ppm for 12–72 h resulted in more than 80% cell viability at up to 400 ppm [162]. CdTe quantum dots were also examined to observe the cytotoxic effects on the cells. Treatment of rat pheochromocytoma cells with 0.01–100 $\mu\text{g/mL}$ CdTe quantum dots for 24 h resulted in a decrease in metabolic activity by 50% [163]. Similarly, treatment of human hepatoma HepG2 cells with CdTe quantum dots at 0– 10^{-5} M for 24 h resulted in ~ 50% reduced viability at 10^{-5} M [164].

Effects on organ systems

In *in vivo* models, the effect of CNTs appears to be related to their method of administration. Various research groups have found that exposure of nanoparticles to the respiratory

system could result in asthma, bronchitis, emphysema and lung cancer. Entry of nanoparticles through the gastrointestinal tract could lead to Crohn's disease and colon cancer. Furthermore, it has been discovered that the nanoparticles' exposure to the circulatory system may result in blood clotting and heart disease (reviewed in [99, 165]). *In vivo* toxicity of carbon nanotubes toward animals was evaluated in a few studies which focused on organisms including guinea pig, mouse and rat. The effect of soot-containing carbon nanotubes containing Co/Ni was observed on guinea pig [166]. Effect of single-walled carbon nanotubes containing metals was evaluated in mouse and rat by various researchers [167–170]. In one of the above studies, acute lung toxicity was evaluated for intratracheally instilled SW-CNTs in rats [170]. In this study cell injury, multifocal granulomas and transient inflammatory reactions were observed. Another study reported lymphocyte and macrophage influx, early neutrophil accumulation, elevation of pro-inflammatory cytokines and fibrogenic transforming growth factor [169]. Effect of MW-CNTs was also evaluated in a study conducted on rat [171]. In the study, MW-CNTs were administered intratracheally in rats. The study evaluated the inflammation, lung persistence and fibrotic reactions both histologically and biochemically. Pulmonary lesions were observed in the bronchial lumen which was characterized by collagen rich granulomas. Stimulation for the production of $\text{TNF-}\alpha$ was also observed. Toxicity of carbon nanotubes was also observed on an aquatic organism such as rainbow trout [172]. In the study, rainbow trout were treated with SW-CNTs at 0.1–0.5 mg/L for 10 days. The study recognized the SW-CNTs as a respiratory toxicant, responsible for neurotoxicity and cell cycle defects. MW-CNTs are also known for increased micronuclei frequency and chromosomal aberrations, promotion of allergic response in mice, activation of cyclooxygenase enzymes through suppression of systemic immune function in spleen and altered gene expression in the liver [173–177]. Other effects of MW-CNTs include apoptosis, phenotypic defects, toxicity in bacteria and formation of abnormal spinal cords in zebrafish embryo [178–180].

In vivo exposure of gold nanoparticles reported apoptosis and acute inflammation in the liver, bioaccumulation in organs and penetration ability in sperm head and tail regions [181–183]. In a study 13 nm sized PEG coated gold nanoparticles were used to observe the toxicity in the liver. It was observed that the particles accumulated in liver and spleen up to 7 days after injection. These nanoparticles induced apoptosis and inflammation in the liver. The study also demonstrated the presence of PEG-coated gold nanoparticles in spleen macrophages and lysosome of liver [181]. *In vivo* studies have reported the astrocyte swelling, blood–brain barrier destruction, oxidative stress induced by free radicals, alteration of gene expression and neuronal degeneration on exposure to silver nanoparticles [184–186]. The distribution



and accumulation of silver nanoparticles were investigated in rats by subcutaneous injection. The results indicated that particles were distributed in liver, spleen, kidney, brain and lung [186]. The ability to form oedema and permeability to blood–brain barrier was investigated on rats by intraperitoneal, intracerebral and intravenous administration. The results suggested that nanoparticles were able to induce oedema formation in the brain by influencing blood brain barrier in vivo [185]. A study utilized silver nanoparticles (25 nm) to evaluate its effects on the gene expression in the different regions of the mouse brain. The study suggested that the silver nanoparticles were able to produce neurotoxicity by the generation of free radicals [184]. The results demonstrated uptake of silver nanoparticles in liver, gills and kidney. The study also evidenced induced expression of *cyp1a2* in the gills suggesting an increase in oxidative metabolism [187].

It was observed from the in vitro studies that quantum dots are able to cross the placental barrier and reached mice pups in pregnant mouse. They also affect mouse oocyte development and were able to penetrate the UV-radiation and compromise skin barrier [188–190]. Titanium oxide-containing sunscreen was used on the human skin to evaluate its toxicity [191]. The results revealed that nanoparticles penetrated the open part of the hair follicles. The penetration of the nanoparticles was reported in a study where the sunscreen was applied to relatively hairy skin [192]. Previous studies reported pulmonary toxicity where the silica nanoparticles' administration was intra-tracheal. The results from the study evidenced acute pulmonary inflammation and neutrophil infiltration to the lung tissues in a dose-dependent manner [193]. Similar studies using silica nanoparticles reported induction of anti-inflammatory mediators and reversibility of fibrotic changes [194, 195]. A few studies in lung tissues suspected translocation and diffusion of silica nanoparticles away from the lung tissue through systematic circulation and deposition in extra pulmonary organs [196–198].

Mechanistic studies

In vivo, mechanistic studies were performed to assess the toxicity caused by nanoparticles. Silver nanoparticles are also known for genotoxicity and cytotoxicity in fish including accumulation in gill tissues and lysosomal destabilization in adult oysters. They are also known for adverse effects on oyster embryonic development, oxidative stress and expression of p53 protein in zebrafish [187, 199–202]. In another study conducted on zebrafish, it was observed that silver nanoparticles caused induction of apoptosis and oxidative stress in the liver. After nanoparticles' treatment, there was upregulation of the p53-related pro-apoptotic genes Noxa, Bax and p21 [200]. Oyster embryo was

used to examine the toxic effects of silver nanoparticles on embryonic development. The study evidenced a significant increase in metallothionein gene expression in embryos [201]. Another study demonstrated the genotoxicity and cytotoxicity of silver nanoparticles on fish cells. The results evidenced the induced aneuploidy and chromosomal aberrations after treatment. The study demonstrated that nanoparticles are genotoxic and cytotoxic to fish cells [202]. In *D. melanogaster*, they are known as oxidative stress, heat shock stress, upregulation of p53 proteins [39]. In *Caenorhabditis elegans*, they are known for oxidative stress and a decrease in reproduction potential [203].

Cytotoxicity of quantum dots was also determined in another study where cadmium sulphate CdS quantum dots were synthesized. The study revealed that CdS quantum dots are more toxic as compared to micro-sized CdS and they elevate ROS production by 20–30%. The study proposed that CdS quantum dots' cytotoxicity is mediated by intracellular ROS production, cadmium ions (Cd^{2+}) release and GSH depletion [204]. Quantum dots are responsible for phototoxicity in *Daphnia magna* under UV-B light [205]. In vivo studies suggested that fullerenes are responsible for elevated gene expression of MHC class II molecules, increased pro-inflammatory cytokines and increased T cell distribution in lungs [142, 206–210]. Oxidative damage in the liver, colon mucosa and lung was observed in the study conducted on rat by oral exposure of C_{60} fullerenes and SW-CNTs. Doses of SW-CNTs increased the levels of 8-oxodG in lung and liver. C_{60} fullerenes administration increased the hepatic 8-oxodG level and high dose generated 8-oxodG in the lung [206]. In oysters they are known for affecting oyster embryonic development, cellular damage in the alimentary canal in *Daphnia magna*, growth inhibition in freshwater fish *Carassius auratus*, increase in mortality rates in gestating daphnids, nitric oxide production in *Mytilus hemocytes* [211–215].

The toxicity of zinc nanoparticles has been reported on human cervix carcinoma cell line (HEp-2), human hepatocyte HEK 293 cell line and human bronchial epithelial cells. Zinc nanoparticles in the range 10–100 $\mu\text{g/mL}$ were used for 24–48 h on HEp-2 cells. The observed results from the study demonstrated that the zinc oxide nanoparticles were toxic for the cells causing DNA damage and reduction in cell viability [216]. Another study focused on the assessment of in vivo toxicity of zinc nanoparticles at 14–20 $\mu\text{g/mL}$ for 12 h to determine the effects of treatment on cell viability, DNA damage, ROS production and apoptosis [217]. When HEK 293 cell line was treated with 0–100 $\mu\text{g/mL}$ zinc oxide nanoparticles for 24 h, it was observed that there was a reduction in cell viability. Results also demonstrated DNA damage, oxidative stress and mitochondrial damage [218]. Similarly, when human bronchial epithelial cells were treated with zinc oxide at 100 $\mu\text{g/mL}$, the observed effects include the release of LDH, decrease in cell viability and oxidative



stress [219]. The toxicity of titanium nanoparticles affects differentiation, cell proliferation, apoptosis and mobility [220, 221]. The toxicity of the titanium oxide nanoparticle was observed when keratinocyte cell line (HaCaT), human dermal fibroblasts and human immortalized sebaceous gland cell lines (SZ95) were used. The cytotoxicity affected cellular functions including differentiation, cell proliferation and mobility which resulted in apoptosis [220]. Upregulation of fibrogenic mediators including IL-4, IL-10 and IL-13 was also observed contributing to fibrotic changes [195].

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Nel, A., et al.: Toxic potential of materials at the nanolevel. *Science* **311**(5761), 622–627 (2006)
- Robertson, T.A., Sanchez, W.Y., Roberts, M.S.: Are commercially available nanoparticles safe when applied to the skin? *J. Biomed. Nanotechnol.* **6**(5), 452–468 (2010)
- Thomas, T.A.: Nanotechnology in consumer products. In *Nanotoxicology: Progress toward Nanomedicine*, pp. 97–112. CRC Press, Boca Raton. (2014)
- Yang, X., et al.: SiO₂ nanoparticles induce cytotoxicity and protein expression alteration in HaCaT cells. *Part. Fibre Toxicol.* **7**(1), 1 (2010)
- Qian, X., et al.: In vivo tumor targeting and spectroscopic detection with surface-enhanced Raman nanoparticle tags. *Nat. Biotechnol.* **26**(1), 83–90 (2008)
- Gibson, J.D., Khanal, B.P., Zubarev, E.R.: Paclitaxel-functionalized gold nanoparticles. *J. Am. Chem. Soc.* **129**(37), 11653–11661 (2007)
- Hainfeld, J.F., et al.: Gold nanoparticles enhance the radiation therapy of a murine squamous cell carcinoma. *Phys. Med. Biol.* **55**(11), 3045 (2010)
- McMahon, S.J., et al.: Biological consequences of nanoscale energy deposition near irradiated heavy atom nanoparticles. *Sci. Rep.* **1**, 18 (2011)
- Caruthers, S.D., Wickline, S.A., Lanza, G.M.: Nanotechnological applications in medicine. *Curr. Opin. Biotechnol.* **18**(1), 26–30 (2007)
- Choi, O., Hu, Z.: Size dependent and reactive oxygen species related nanosilver toxicity to nitrifying bacteria. *Environ. Sci. Technol.* **42**(12), 4583–4588 (2008)
- Zoroddu, M.A., et al.: Toxicity of nanoparticles. *Curr. Med. Chem.* **21**(33), 3837–3853 (2014)
- Behra, R., Krug, H.: Nanoecotoxicology: nanoparticles at large. *Nat. Nanotechnol.* **3**(5), 253–254 (2008)
- Hanley, C., et al.: The influences of cell type and ZnO nanoparticle size on immune cell cytotoxicity and cytokine induction. *Nanoscale Res. Lett.* **4**(12), 1409 (2009)
- Wang, F., et al.: The biomolecular corona is retained during nanoparticle uptake and protects the cells from the damage induced by cationic nanoparticles until degraded in the lysosomes. *Nanomed. Nanotechnol. Biol. Med.* **9**(8), 1159–1168 (2013)
- Elsaesser, A., Howard, C.V.: Toxicology of nanoparticles. *Adv. Drug Deliv. Rev.* **64**(2), 129–137 (2012)
- Regoli, F., Giuliani, M.E.: Oxidative pathways of chemical toxicity and oxidative stress biomarkers in marine organisms. *Mar. Environ. Res.* **93**, 106–117 (2014)
- Jomova, K., Baros, S., Valko, M.: Redox active metal-induced oxidative stress in biological systems. *Transit. Met. Chem.* **37**(2), 127–134 (2012)
- Chen, H., et al.: Oxidative stress in ischemic brain damage: mechanisms of cell death and potential molecular targets for neuroprotection. *Antioxid. Redox. Signal.* **14**(8), 1505–1517 (2011)
- Soenen, S.J., et al.: Cellular toxicity of inorganic nanoparticles: common aspects and guidelines for improved nanotoxicity evaluation. *Nano Today* **6**(5), 446–465 (2011)
- Hegde, K., et al.: Current understandings of toxicity, risks and regulations of engineered nanoparticles with respect to environmental microorganisms. *Nanotechnol. Environ. Eng.* **1**(1), 5 (2016)
- Berghe, T.V., et al.: Determination of apoptotic and necrotic cell death in vitro and in vivo. *Methods* **61**(2), 117–129 (2013)
- Bakand, S., Hayes, A.: Toxicological considerations, toxicity assessment, and risk management of inhaled nanoparticles. *Int. J. Mol. Sci.* **17**(6), 929 (2016)
- Maurer, L., Meyer, J.: A systematic review of evidence for silver nanoparticle-induced mitochondrial toxicity. *Environ. Sci. Nano* **3**(2), 311–322 (2016)
- Jain, S., Singh, S.R., Pillai, S.: Toxicity issues related to biomedical applications of carbon nanotubes. *J. Nanomed. Nanotechnol.* **3**(5), 1–15 (2012)
- Kerfahi, D., et al.: Effects of functionalized and raw multi-walled carbon nanotubes on soil bacterial community composition. *PLoS One* **10**(3), e0123042 (2015)
- Sayes, C.M., Reed, K.L., Warheit, D.B.: Assessing toxicity of fine and nanoparticles: comparing in vitro measurements to in vivo pulmonary toxicity profiles. *Toxicol. Sci.* **97**(1), 163–180 (2007)
- Marshall, N.J., Goodwin, C.J., Holt, S.J.: A critical assessment of the use of microculture tetrazolium assays to measure cell growth and function. *Growth Regul.* **5**(2), 69–84 (1995)
- Molinari, B.L., et al.: Cell-based quantitative evaluation of the MTT assay. *Anal. Quant. Cytol. Histol.* **25**(5), 254–262 (2003)
- Jabbar, S., Twentyman, P., Watson, J.: The MTT assay underestimates the growth inhibitory effects of interferons. *Br. J. Cancer* **60**(4), 523 (1989)
- Natarajan, M., et al.: Antioxidant compounds interfere with the 3. *Cancer Detect. Prev.* **24**(5), 405–414 (2000)
- Abe, K., Saito, H.: Cholesterol does not affect the toxicity of amyloid β fragment but mimics its effect on MTT formazan exocytosis in cultured rat hippocampal neurons. *Neurosci. Res.* **35**(3), 165–174 (1999)
- Hussain, S.M., Frazier, J.M.: Cellular toxicity of hydrazine in primary rat hepatocytes. *Toxicol. Sci.* **69**(2), 424–432 (2002)
- Punshon, G., et al.: Interactions between endothelial cells and a poly(carbonate-silsesquioxane-bridge-urea)urethane. *Biomaterials* **26**(32), 6271–6279 (2005)
- Low, S.P., et al.: Evaluation of mammalian cell adhesion on surface-modified porous silicon. *Biomaterials* **27**(26), 4538–4546 (2006)
- Casey, A., et al.: Spectroscopic analysis confirms the interactions between single walled carbon nanotubes and various dyes commonly used to assess cytotoxicity. *Carbon* **45**(7), 1425–1432 (2007)
- Ryter, S.W., et al.: Mechanisms of cell death in oxidative stress. *Antioxid. Redox Signal.* **9**(1), 49–89 (2007)
- Li, G.-Y., Osborne, N.N.: Oxidative-induced apoptosis to an immortalized ganglion cell line is caspase independent but



- involves the activation of poly (ADP-ribose) polymerase and apoptosis-inducing factor. *Brain Res.* **1188**, 35–43 (2008)
38. Ahamed, M., et al.: DNA damage response to different surface chemistry of silver nanoparticles in mammalian cells. *Toxicol. Appl. Pharmacol.* **233**(3), 404–410 (2008)
 39. Ahamed, M., et al.: Silver nanoparticles induced heat shock protein 70, oxidative stress and apoptosis in *Drosophila melanogaster*. *Toxicol. Appl. Pharmacol.* **242**(3), 263–269 (2010)
 40. Liang, S.H., Clarke, M.F.: Regulation of p53 localization. *FEBS J.* **268**(10), 2779–2783 (2001)
 41. Sherr, C.J.: Principles of tumor suppression. *Cell* **116**(2), 235–246 (2004)
 42. Lee, K.J., et al.: In vivo imaging of transport and biocompatibility of single silver nanoparticles in early development of zebrafish embryos. *ACS Nano* **1**(2), 133–143 (2007)
 43. Jin, Y., et al.: Toxicity of luminescent silica nanoparticles to living cells. *Chem. Res. Toxicol.* **20**(8), 1126–1133 (2007)
 44. Mo, Y., Lim, L.-Y.: Paclitaxel-loaded PLGA nanoparticles: potentiation of anticancer activity by surface conjugation with wheat germ agglutinin. *J. Control. Release* **108**(2–3), 244–262 (2005)
 45. Pan, Y., et al.: Size-dependent cytotoxicity of gold nanoparticles. *Small* **3**(11), 1941–1949 (2007)
 46. Wyllie, A.H.: Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature* **284**(5756), 555–556 (1980)
 47. Kerr, J.F., Wyllie, A.H., Currie, A.R.: Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer* **26**(4), 239 (1972)
 48. Suman, S., Pandey, A., Chandna, S.: An improved non-enzymatic “DNA ladder assay” for more sensitive and early detection of apoptosis. *Cytotechnology* **64**(1), 9–14 (2012)
 49. Abdel-Khalek, A.A.: Comparative evaluation of genotoxic effects induced by CuO bulk and nano-particles in *Nile tilapia*, *Oreochromis niloticus*. *Water Air Soil Pollut.* **227**(1), 35 (2016)
 50. Silva, M.T.: Secondary necrosis: the natural outcome of the complete apoptotic program. *FEBS Lett.* **584**(22), 4491–4499 (2010)
 51. Lu, X., et al.: In vitro cytotoxicity and induction of apoptosis by silica nanoparticles in human HepG2 hepatoma cells. *Int. J. Nanomed.* **6**, 1889 (2011)
 52. Baharara, J., et al.: Induction of apoptosis by green synthesized gold nanoparticles through activation of caspase-3 and 9 in human cervical cancer cells. *Avicenna J. Med. Biotechnol.* **8**(2), 75 (2016)
 53. Kisin, E.R., et al.: Single-walled carbon nanotubes: geno- and cytotoxic effects in lung fibroblast V79 cells. *J. Toxicol. Environ. Health Part A* **70**(24), 2071–2079 (2007)
 54. Bajpayee M., Kumar A., Dhawan A.: The Comet Assay: Assessment of In Vitro and In Vivo DNA Damage. In: Dhawan A., Bajpayee M. (eds) *Genotoxicity Assessment. Methods in Molecular Biology (Methods and Protocols)*, vol 1044, pp. 325–345. Humana Press, Totowa, NJ (2013)
 55. Kumar A., Sharma V., Dhawan A.: Methods for Detection of Oxidative Stress and Genotoxicity of Engineered Nanoparticles. In: Armstrong D., Bharali D. (eds) *Oxidative Stress and Nanotechnology. Methods in Molecular Biology (Methods and Protocols)*, vol 1028, pp. 231–246. Humana Press, Totowa, NJ (2013)
 56. Kumar, A., Dhawan, A.: Genotoxic and carcinogenic potential of engineered nanoparticles: an update. *Arch. Toxicol.* **87**(11), 1883–1900 (2013)
 57. Dusinska, M., Collins, A.R.: The comet assay in human biomonitoring: gene–environment interactions. *Mutagenesis* **23**(3), 191–205 (2008)
 58. Pavanello, S., Clonfero, E.: Biological indicators of genotoxic risk and metabolic polymorphisms. *Mutat. Res. Rev. Mutat. Res.* **463**(3), 285–308 (2000)
 59. Fairbairn, D.W., Olive, P.L., O'Neill, K.L.: The comet assay: a comprehensive review. *Mutat. Res. Rev. Genet. Toxicol.* **339**(1), 37–59 (1995)
 60. Lemay, M., Wood, K.A.: Detection of DNA damage and identification of UV-induced photoproducts using the CometAssay [TM] Kit. *Biotechniques* **27**, 846–848 (1999)
 61. Malyapa, R.S., et al.: Detection of DNA damage by the alkaline comet assay after exposure to low-dose gamma radiation. *Radiat. Res.* **149**(4), 396–400 (1998)
 62. Henderson, L., et al.: The ability of the Comet assay to discriminate between genotoxins and cytotoxins. *Mutagenesis* **13**(1), 89–94 (1998)
 63. Collins, A.R., Ai-Guo, M., Duthie, S.J.: The kinetics of repair of oxidative DNA damage (strand breaks and oxidised pyrimidines) in human cells. *Mutat. Res. DNA Repair* **336**(1), 69–77 (1995)
 64. Singh, N., Stephens, R.: Microgel electrophoresis: sensitivity, mechanisms, and DNA electrostretching. *Mutat. Res. DNA Repair* **383**(2), 167–175 (1997)
 65. Schiavo, S., et al.: Genotoxic and cytotoxic effects of ZnO nanoparticles for *Dunaliella tertiolecta* and comparison with SiO₂ and TiO₂ effects at population growth inhibition levels. *Sci. Total Environ.* **550**, 619–627 (2016)
 66. Handy, R.D., et al.: Practical considerations for conducting ecotoxicity test methods with manufactured nanomaterials: what have we learnt so far? *Ecotoxicology* **21**(4), 933–972 (2012)
 67. Gavrieli, Y., Sherman, Y., Ben-Sasson, S.A.: Identification of programmed cell death in situ via specific labelling of nuclear DNA fragmentation. *J. Cell Biol.* **119**, 493 (1992)
 68. Arends, M., Morris, R., Wyllie, A.: Apoptosis. The role of the endonuclease. *Am. J. Pathol.* **136**(3), 593 (1990)
 69. Bortner, C.D., Oldenburg, N.B., Cidlowski, J.A.: The role of DNA fragmentation in apoptosis. *Trends Cell Biol.* **5**(1), 21–26 (1995)
 70. Loo, D.T.: TUNEL assay: an overview of techniques. In: Didenko, Vladimir, V. (Eds) *Situ Detection of DNA Damage: Methods and Protocols*, pp. 21–30. Humana Press, New York, United States (2002)
 71. Ansari, B., et al.: In situ end-labelling detects DNA strand breaks in apoptosis and other physiological and pathological states. *J. Pathol.* **170**(1), 1–8 (1993)
 72. Kano, M., et al.: Significance of myocytes with positive DNA in situ nick end-labeling (TUNEL) in hearts with dilated cardiomyopathy. *Circulation* **99**(21), 2757–2764 (1999)
 73. Deng, W., et al.: Selenium nanoparticles as versatile carriers for oral delivery of insulin: insight into the synergic anti-diabetic effect and mechanism. *Nanomed. Nanotechnol. Biol. Med.* **13**(6), 1965–1974 (2017). doi: <https://doi.org/10.1016/j.nano.2017.05.002>
 74. Monteiro-Riviere, N.A., et al.: Multi-walled carbon nanotube interactions with human epidermal keratinocytes. *Toxicol. Lett.* **155**(3), 377–384 (2005)
 75. Huang, M., Khor, E., Lim, L.-Y.: Uptake and cytotoxicity of chitosan molecules and nanoparticles: effects of molecular weight and degree of deacetylation. *Pharm. Res.* **21**(2), 344–353 (2004)
 76. Bitensky, L.: The reversible activation of lysosomes in normal cells and the effects of pathological conditions. In: *Ciba Foundation Symposium—Lysosomes*, pp. 362–383. Wiley Online Library (1963)
 77. Nemes, Z., et al.: The pharmacological relevance of vital staining with neutral red. *Experientia* **35**(11), 1475–1476 (1979)
 78. Lüllmann-Rauch, R.: Drug-induced lysosomal storage disorders. *Front. Biol.* **48**, 49 (1979)
 79. Borenfreund, E., Puerner, J.A.: Toxicity determined in vitro by morphological alterations and neutral red absorption. *Toxicol. Lett.* **24**(2–3), 119–124 (1985)



80. Borenfreund, E., Shopsis, C.: Toxicity monitored with a correlated set of cell-culture assays. *Xenobiotica* **15**(8–9), 705–711 (1985)
81. Miranda, R.R., et al.: Toxicological interactions of silver nanoparticles and non-essential metals in human hepatocarcinoma cell line. *Toxicol. In Vitro* **40**, 134–143 (2017)
82. Strober, W.: Trypan blue exclusion test of cell viability. In: *Current protocols in immunology*, pp. A3. B.1–A3. B.3. John Wiley & Sons, Unites States (2001)
83. Kononenko, V., et al.: Comparative in vitro genotoxicity study of ZnO nanoparticles, ZnO macroparticles and ZnCl₂ to MDCK kidney cells: size matters. *Toxicol. In Vitro* **40**, 256–263 (2017)
84. Magder, S.: Reactive oxygen species: toxic molecules or spark of life? *Crit. Care* **10**(1), 208 (2006)
85. Zang, L.: F. vanKuijk and HP Misra. *Biochem. Mol. Biol. Int.* **37**, 1187–1195 (1995)
86. Gomes, A., Fernandes, E., Lima, J.L.: Fluorescence probes used for detection of reactive oxygen species. *J. Biochem. Biophys. Methods* **65**(2), 45–80 (2005)
87. Halliwell, B., Whiteman, M.: Measuring reactive species and oxidative damage in vivo and in cell culture: how should you do it and what do the results mean? *Br. J. Pharmacol.* **142**(2), 231–255 (2004)
88. Wagner, A.J., et al.: Cellular interaction of different forms of aluminum nanoparticles in rat alveolar macrophages. *J. Phys. Chem. B* **111**(25), 7353–7359 (2007)
89. Fantel, A.G.: Reactive oxygen species in developmental toxicity: review and hypothesis. *Teratology* **53**(3), 196–217 (1996)
90. Hussain, S.M., et al.: The interaction of manganese nanoparticles with PC-12 cells induces dopamine depletion. *Toxicol. Sci.* **92**(2), 456–463 (2006)
91. Kim, S.C., et al.: In vivo evaluation of polymeric micellar paclitaxel formulation: toxicity and efficacy. *J. Control. Release* **72**(1), 191–202 (2001)
92. Li, Y.-P., et al.: PEGylated PLGA nanoparticles as protein carriers: synthesis, preparation and biodistribution in rats. *J. Control. Release* **71**(2), 203–211 (2001)
93. Baker, G.L., et al.: Inhalation toxicity and lung toxicokinetics of C60 fullerene nanoparticles and microparticles. *Toxicol. Sci.* **101**(1), 122–131 (2008)
94. Lei, R., et al.: Integrated metabolomic analysis of the nano-sized copper particle-induced hepatotoxicity and nephrotoxicity in rats: a rapid in vivo screening method for nanotoxicity. *Toxicol. Appl. Pharmacol.* **232**(2), 292–301 (2008)
95. Zhu, M.-T., et al.: Comparative study of pulmonary responses to nano- and submicron-sized ferric oxide in rats. *Toxicology* **247**(2), 102–111 (2008)
96. Ewing, A.G., Bigelow, J.C., Wightman, R.M.: Direct in vivo monitoring of dopamine released from two striatal compartments in the rat. *Science* **221**(4606), 169–171 (1983)
97. Guo, N.L., et al.: Multiwalled carbon nanotube-induced gene signatures in the mouse lung: potential predictive value for human lung cancer risk and prognosis. *J. Toxicol. Environ. Health Part A* **75**(18), 1129–1153 (2012)
98. Sathyanarayana, S., Hübner, C.: Thermoplastic nanocomposites with carbon nanotubes. In: Njuguna J. (eds) *Structural Nanocomposites*. Engineering Materials, pp. 19–60. Springer, Berlin (2013)
99. Madani, S.Y., Mandel, A., Seifalian, A.M.: A concise review of carbon nanotube's toxicology. *Nano Rev. Exp.* **4**, 21521–21534 (2013)
100. Amarnath, S., Hussain, M. A., Nanjundiah, V., Sood A. K.: β -Galactosidase leakage from *Escherichia coli* points to mechanical damages likely cause of carbon nanotube toxicity. *Soft Nanosci. Let.* **2**(3), 41–45 (2012)
101. Krause, C.J.: Evaluating the toxicity of nitrogen-functionalized carbon nanotubes MS (Master of Science) thesis, University of Iowa (2014). <http://ir.uiowa.edu/etd/4667>. Accessed 15 Aug 2014
102. Pasquini, L.M., et al.: Impact of surface functionalization on bacterial cytotoxicity of single-walled carbon nanotubes. *Environ. Sci. Technol.* **46**(11), 6297–6305 (2012)
103. Sohn, E. K., Chung, Y. S., Johari, S. A., Kim, T. G., Kim, J. K., Lee, J. H., Lee, Y. H., Kang, S. W., Yu, I. J.: Acute toxicity comparison of single-walled carbon nanotubes in various freshwater organisms. *BioMed Res. Int.* **2015** (2015). <http://dx.doi.org/10.1155/2015/323090>
104. Naqvi, S., et al.: Concentration-dependent toxicity of iron oxide nanoparticles mediated by increased oxidative stress. *Int. J. Nanomed.* **5**, 983–989 (2009)
105. Jeng, H.A., Swanson, J.: Toxicity of metal oxide nanoparticles in mammalian cells. *J. Environ. Sci. Health Part A* **41**(12), 2699–2711 (2006)
106. Delcroix, G.J.-R., et al.: Mesenchymal and neural stem cells labeled with HEDP-coated SPIO nanoparticles: in vitro characterization and migration potential in rat brain. *Brain Res.* **1255**, 18–31 (2009)
107. Park, Y.-H., et al.: Assessment of dermal toxicity of nanosilica using cultured keratinocytes, a human skin equivalent model and an in vivo model. *Toxicology* **267**(1), 178–181 (2010)
108. Yehia, H.N., et al.: Single-walled carbon nanotube interactions with HeLa cells. *J. Nanobiotechnol.* **5**(1), 1 (2007)
109. Fiorito, S., et al.: Effects of fullerenes and single-wall carbon nanotubes on murine and human macrophages. *Carbon* **44**(6), 1100–1105 (2006)
110. Davoren, M., et al.: In vitro toxicity evaluation of single walled carbon nanotubes on human A549 lung cells. *Toxicol. In Vitro* **21**(3), 438–448 (2007)
111. Choi, S.-J., Oh, J.-M., Choy, J.-H.: Toxicological effects of inorganic nanoparticles on human lung cancer A549 cells. *J. Inorg. Biochem.* **103**(3), 463–471 (2009)
112. Herzog, E., et al.: SWCNT suppress inflammatory mediator responses in human lung epithelium in vitro. *Toxicol. Appl. Pharmacol.* **234**(3), 378–390 (2009)
113. Ye, S.-F., et al.: ROS and NF- κ B are involved in upregulation of IL-8 in A549 cells exposed to multi-walled carbon nanotubes. *Biochem. Biophys. Res. Commun.* **379**(2), 643–648 (2009)
114. Cveticanin, J., et al.: Using carbon nanotubes to induce micro-nuclei and double strand breaks of the DNA in human cells. *Nanotechnology* **21**(1), 015102 (2009)
115. Patlolla, A., Patlolla, B., Tchounwou, P.: Evaluation of cell viability, DNA damage, and cell death in normal human dermal fibroblast cells induced by functionalized multiwalled carbon nanotube. *Mol. Cell. Biochem.* **338**(1–2), 225–232 (2010)
116. Ravichandran, P., et al.: Induction of apoptosis in rat lung epithelial cells by multiwalled carbon nanotubes. *J. Biochem. Mol. Toxicol.* **27**(5), 286 (2013)
117. Reddy, A.R.N., et al.: Multi wall carbon nanotubes induce oxidative stress and cytotoxicity in human embryonic kidney (HEK293) cells. *Toxicology* **272**(1), 11–16 (2010)
118. Walker, V.G., et al.: Potential in vitro effects of carbon nanotubes on human aortic endothelial cells. *Toxicol. Appl. Pharmacol.* **236**(3), 319–328 (2009)
119. Li, J.J., et al.: Autophagy and oxidative stress associated with gold nanoparticles. *Biomaterials* **31**(23), 5996–6003 (2010)
120. Tarantola, M., et al.: Cytotoxicity of metal and semiconductor nanoparticles indicated by cellular micromotility. *ACS Nano* **3**(1), 213–222 (2008)
121. AshaRani, P., et al.: Cytotoxicity and genotoxicity of silver nanoparticles in human cells. *ACS Nano* **3**(2), 279–290 (2008)
122. Hsin, Y.-H., et al.: The apoptotic effect of nanosilver is mediated by a ROS- and JNK-dependent mechanism involving the



- mitochondrial pathway in NIH3T3 cells. *Toxicol. Lett.* **179**(3), 130–139 (2008)
123. Kawata, K., Osawa, M., Okabe, S.: In vitro toxicity of silver nanoparticles at noncytotoxic doses to HepG2 human hepatoma cells. *Environ. Sci. Technol.* **43**(15), 6046–6051 (2009)
 124. Miura, N., Shinohara, Y.: Cytotoxic effect and apoptosis induction by silver nanoparticles in HeLa cells. *Biochem. Biophys. Res. Commun.* **390**(3), 733–737 (2009)
 125. Yang, W., et al.: Food storage material silver nanoparticles interfere with DNA replication fidelity and bind with DNA. *Nanotechnology* **20**(8), 085102 (2009)
 126. Cho, S.J., et al.: Long-term exposure to CdTe quantum dots causes functional impairments in live cells. *Langmuir* **23**(4), 1974–1980 (2007)
 127. Ryman-Rasmussen, J.P., Riviere, J.E., Monteiro-Riviere, N.A.: Surface coatings determine cytotoxicity and irritation potential of quantum dot nanoparticles in epidermal keratinocytes. *J. Investig. Dermatol.* **127**(1), 143–153 (2007)
 128. Su, Y., et al.: The cytotoxicity of cadmium based, aqueous phase-synthesized, quantum dots and its modulation by surface coating. *Biomaterials* **30**(1), 19–25 (2009)
 129. Tang, M., et al.: Unmodified CdSe quantum dots induce elevation of cytoplasmic calcium levels and impairment of functional properties of sodium channels in rat primary cultured hippocampal neurons. *Environ. Health Perspect.* **116**(7), 915 (2008)
 130. Wang, L., et al.: Toxicity of CdSe nanoparticles in Caco-2 cell cultures. *J. Nanobiotechnol.* **6**(1), 1 (2008)
 131. Zhang, Y., et al.: In vitro and in vivo toxicity of CdTe nanoparticles. *J. Nanosci. Nanotechnol.* **7**(2), 497–503 (2007)
 132. Chen, L., et al.: Manufactured aluminum oxide nanoparticles decrease expression of tight junction proteins in brain vasculature. *J. Neuroimmune Pharmacol.* **3**(4), 286–295 (2008)
 133. Radziun, E., et al.: Assessment of the cytotoxicity of aluminum oxide nanoparticles on selected mammalian cells. *Toxicol. In Vitro* **25**(8), 1694–1700 (2011)
 134. Alshatwi, A.A., et al.: Al₂O₃ nanoparticles induce mitochondria-mediated cell death and upregulate the expression of signaling genes in human mesenchymal stem cells. *J. Biochem. Mol. Toxicol.* **26**(11), 469–476 (2012)
 135. Balasubramanyam, A., et al.: In vivo genotoxicity assessment of aluminium oxide nanomaterials in rat peripheral blood cells using the comet assay and micronucleus test. *Mutagenesis* **24**(3), 245–251 (2009)
 136. Kim, Y.-J., et al.: Genotoxicity of aluminum oxide (Al₂O₃) nanoparticle in mammalian cell lines. *Mol. Cell. Toxicol.* **5**(2), 172–178 (2009)
 137. Clark, K.A., et al.: Evaluation of the interactions between multi-walled carbon nanotubes and Caco-2 cells. *J. Toxicol. Environ. Health Part A* **75**(1), 25–35 (2012)
 138. Kim, J.S., et al.: Determination of cytotoxicity attributed to multiwall carbon nanotubes (MWCNT) in normal human embryonic lung cell (WI-38) line. *J. Toxicol. Environ. Health Part A* **73**(21–22), 1521–1529 (2010)
 139. Pan, Y., et al.: Gold nanoparticles of diameter 1.4 nm trigger necrosis by oxidative stress and mitochondrial damage. *Small* **5**(18), 2067–2076 (2009)
 140. Asharani, P., Hande, M.P., Valiyaveetil, S.: Anti-proliferative activity of silver nanoparticles. *BMC Cell Biol.* **10**(1), 1 (2009)
 141. Foldbjerg, R., Dang, D.A., Autrup, H.: Cytotoxicity and genotoxicity of silver nanoparticles in the human lung cancer cell line, A549. *Arch. Toxicol.* **85**(7), 743–750 (2011)
 142. Jacobsen, N.R., et al.: Genotoxicity, cytotoxicity, and reactive oxygen species induced by single-walled carbon nanotubes and C60 fullerenes in the FE1-Muta™ Mouse lung epithelial cells. *Environ. Mol. Mutagen.* **49**(6), 476–487 (2008)
 143. Zhang, L.W., et al.: Endocytic mechanisms and toxicity of a functionalized fullerene in human cells. *Toxicol. Lett.* **191**(2), 149–157 (2009)
 144. Dhawan, A., et al.: Stable colloidal dispersions of C60 fullerenes in water: evidence for genotoxicity. *Environ. Sci. Technol.* **40**(23), 7394–7401 (2006)
 145. Pantarotto, D., et al.: Translocation of bioactive peptides across cell membranes by carbon nanotubes. *Chem. Commun.* **1**, 16–17 (2004)
 146. Shvedova, A., et al.: Exposure to carbon nanotube material: assessment of nanotube cytotoxicity using human keratinocyte cells. *J. Toxicol. Environ. Health Part A* **66**(20), 1909–1926 (2003)
 147. Cui, D., et al.: Effect of single wall carbon nanotubes on human HEK293 cells. *Toxicol. Lett.* **155**(1), 73–85 (2005)
 148. Cherukuri, P., et al.: Near-infrared fluorescence microscopy of single-walled carbon nanotubes in phagocytic cells. *J. Am. Chem. Soc.* **126**(48), 15638–15639 (2004)
 149. Jia, G., et al.: Cytotoxicity of carbon nanomaterials: single-wall nanotube, multi-wall nanotube, and fullerene. *Environ. Sci. Technol.* **39**(5), 1378–1383 (2005)
 150. Pulskamp, K., Diabaté, S., Krug, H.F.: Carbon nanotubes show no sign of acute toxicity but induce intracellular reactive oxygen species in dependence on contaminants. *Toxicol. Lett.* **168**(1), 58–74 (2007)
 151. Tian, F., et al.: Cytotoxicity of single-wall carbon nanotubes on human fibroblasts. *Toxicol. In Vitro* **20**(7), 1202–1212 (2006)
 152. Ding, L., et al.: Molecular characterization of the cytotoxic mechanism of multiwall carbon nanotubes and nano-onions on human skin fibroblast. *Nano Lett.* **5**(12), 2448 (2005)
 153. Tkachenko, A.G., et al.: Cellular trajectories of peptide-modified gold particle complexes: comparison of nuclear localization signals and peptide transduction domains. *Bioconjug. Chem.* **15**(3), 482–490 (2004)
 154. Shukla, R., et al.: Biocompatibility of gold nanoparticles and their endocytotic fate inside the cellular compartment: a microscopic overview. *Langmuir* **21**(23), 10644–10654 (2005)
 155. Pernodet, N., et al.: Adverse effects of citrate/gold nanoparticles on human dermal fibroblasts. *Small* **2**(6), 766–773 (2006)
 156. Su, C.-H., et al.: Nanoshell magnetic resonance imaging contrast agents. In: *Microprocesses and Nanotechnology, 2007 Digest of Papers, IEEE* (2007)
 157. Gupta, A.K., Wells, S.: Surface-modified superparamagnetic nanoparticles for drug delivery: preparation, characterization, and cytotoxicity studies. *IEEE Trans. Nanobiosci.* **3**(1), 66–73 (2004)
 158. Hu, F., et al.: Cellular response to magnetic nanoparticles “PEGylated” via surface-initiated atom transfer radical polymerization. *Biomacromolecules* **7**(3), 809–816 (2006)
 159. William, W.Y., et al.: Aqueous dispersion of monodisperse magnetic iron oxide nanocrystals through phase transfer. *Nanotechnology* **17**(17), 4483 (2006)
 160. Chen, F., Gerion, D.: Fluorescent CdSe/ZnS nanocrystal-peptide conjugates for long-term, nontoxic imaging and nuclear targeting in living cells. *Nano Lett.* **4**(10), 1827–1832 (2004)
 161. Hoshino, A., et al.: Physicochemical properties and cellular toxicity of nanocrystal quantum dots depend on their surface modification. *Nano Lett.* **4**(11), 2163–2170 (2004)
 162. Guo, G., et al.: Probing the cytotoxicity of CdSe quantum dots with surface modification. *Mater. Lett.* **61**(8), 1641–1644 (2007)
 163. Lovrić, J., et al.: Unmodified cadmium telluride quantum dots induce reactive oxygen species formation leading to multiple organelle damage and cell death. *Chem. Biol.* **12**(11), 1227–1234 (2005)



164. Liu, Y., et al.: Comparison of water-soluble CdTe nanoparticles synthesized in air and in nitrogen. *J. Phys. Chem. B* **110**(34), 16992–17000 (2006)
165. Buzea, C., Pacheco, I.I., Robbie, K.: Nanomaterials and nanoparticles: sources and toxicity. *Biointerphases* **2**(4), MR7–MR71 (2007)
166. Huczko, A., et al.: Physiological testing of carbon nanotubes: are they asbestos-like? *Fuller. Sci. Technol.* **9**(2), 251–254 (2001)
167. Lam, C.-W., et al.: A review of carbon nanotube toxicity and assessment of potential occupational and environmental health risks. *Crit. Rev. Toxicol.* **36**(3), 189–217 (2006)
168. Lam, C.-W., et al.: Pulmonary toxicity of single-wall carbon nanotubes in mice 7 and 90 days after intratracheal instillation. *Toxicol. Sci.* **77**(1), 126–134 (2004)
169. Shvedova, A.A., et al.: Unusual inflammatory and fibrogenic pulmonary responses to single-walled carbon nanotubes in mice. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **289**(5), L698–L708 (2005)
170. Warheit, D.B., et al.: Comparative pulmonary toxicity assessment of single-wall carbon nanotubes in rats. *Toxicol. Sci.* **77**(1), 117–125 (2004)
171. Muller, J., et al.: Respiratory toxicity of multi-wall carbon nanotubes. *Toxicol. Appl. Pharmacol.* **207**(3), 221–231 (2005)
172. Smith, C.J., Shaw, B.J., Handy, R.D.: Toxicity of single walled carbon nanotubes to rainbow trout, (*Oncorhynchus mykiss*): respiratory toxicity, organ pathologies, and other physiological effects. *Aquat. Toxicol.* **82**(2), 94–109 (2007)
173. Ji, Z., et al.: The hepatotoxicity of multi-walled carbon nanotubes in mice. *Nanotechnology* **20**(44), 445101 (2009)
174. Mitchell, L., et al.: Mechanisms for how inhaled multiwalled carbon nanotubes suppress systemic immune function in mice. *Nat. Nanotechnol.* **4**(7), 451–456 (2009)
175. Nygaard, U. C., Hansen, J. S., Samuelsen, M., Alberg, T., Mariøara, C. D., Løvik, M.: Single-walled and multi-walled carbon nanotubes promote allergic immune responses in mice. *Toxicol. Sci.* **109**(1), 113–23 (2009)
176. Park, E.-J., et al.: Pro-inflammatory and potential allergic responses resulting from B cell activation in mice treated with multi-walled carbon nanotubes by intratracheal instillation. *Toxicology* **259**(3), 113–121 (2009)
177. Patlolla, A.K., et al.: Comparative study of the clastogenicity of functionalized and nonfunctionalized multiwalled carbon nanotubes in bone marrow cells of Swiss-Webster mice. *Environ. Toxicol.* **25**(6), 608–621 (2010)
178. Asharani, P., et al.: Impact of multi-walled carbon nanotubes on aquatic species. *J. Nanosci. Nanotechnol.* **8**(7), 3603–3609 (2008)
179. Cheng, J., et al.: Acute and long-term effects after single loading of functionalized multi-walled carbon nanotubes into zebrafish (*Danio rerio*). *Toxicol. Appl. Pharmacol.* **235**(2), 216–225 (2009)
180. Kang, S., Mauter, M.S., Elimelech, M.: Physicochemical determinants of multiwalled carbon nanotube bacterial cytotoxicity. *Environ. Sci. Technol.* **42**(19), 7528–7534 (2008)
181. Cho, W.-S., et al.: Acute toxicity and pharmacokinetics of 13 nm-sized PEG-coated gold nanoparticles. *Toxicol. Appl. Pharmacol.* **236**(1), 16–24 (2009)
182. Lasagna-Reeves, C., et al.: Bioaccumulation and toxicity of gold nanoparticles after repeated administration in mice. *Biochem. Biophys. Res. Commun.* **393**(4), 649–655 (2010)
183. Wiwanitkit, V., Sereemasun, A., Rojanathanes, R.: Effect of gold nanoparticles on spermatozoa: the first world report. *Fertil. Steril.* **91**(1), e7–e8 (2009)
184. Rahman, M., et al.: Expression of genes related to oxidative stress in the mouse brain after exposure to silver-25 nanoparticles. *Toxicol. Lett.* **187**(1), 15–21 (2009)
185. Sharma, H.S., et al.: Influence of nanoparticles on blood–brain barrier permeability and brain edema formation in rats. In: *Brain Edema XIV*, pp. 359–364. Springer, Berlin (2010)
186. Tang, J., et al.: Distribution, translocation and accumulation of silver nanoparticles in rats. *J. Nanosci. Nanotechnol.* **9**(8), 4924–4932 (2009)
187. Scown, T.M., et al.: Effects of aqueous exposure to silver nanoparticles of different sizes in rainbow trout. *Toxicol. Sci.* **115**(2), 521–534 (2010)
188. Chu, M., et al.: Transfer of quantum dots from pregnant mice to pups across the placental barrier. *Small* **6**(5), 670–678 (2010)
189. Hsieh, M.-S., Shiao, N.-H., Chan, W.-H.: Cytotoxic effects of CdSe quantum dots on maturation of mouse oocytes, fertilization, and fetal development. *Int. J. Mol. Sci.* **10**(5), 2122–2135 (2009)
190. Mortensen, L.J., et al.: In vivo skin penetration of quantum dot nanoparticles in the murine model: the effect of UVR. *Nano Lett.* **8**(9), 2779–2787 (2008)
191. Lademann, J., et al.: Penetration of titanium dioxide microparticles in a sunscreen formulation into the horny layer and the follicular orifice. *Skin Pharmacol. Physiol.* **12**(5), 247–256 (1999)
192. Bennat, C., Muller-Goymann, C.: Skin penetration and stabilization of formulations containing microfine titanium dioxide as physical UV filter. *Int. J. Cosmet. Sci.* **22**(4), 271–284 (2000)
193. Cho, W.-S., et al.: Inflammatory mediators induced by intratracheal instillation of ultrafine amorphous silica particles. *Toxicol. Lett.* **175**(1), 24–33 (2007)
194. Choi, M., et al.: Transient pulmonary fibrogenic effect induced by intratracheal instillation of ultrafine amorphous silica in A/J mice. *Toxicol. Lett.* **182**(1), 97–101 (2008)
195. Barbarin, V., et al.: Pulmonary overexpression of IL-10 augments lung fibrosis and Th2 responses induced by silica particles. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **288**(5), L841–L848 (2005)
196. Oberdörster, G., et al.: Extrapulmonary translocation of ultrafine carbon particles following whole-body inhalation exposure of rats. *J. Toxicol. Environ. Health Part A* **65**(20), 1531–1543 (2002)
197. Oberdörster, G., et al.: Translocation of inhaled ultrafine particles to the brain. *Inhalation Toxicol.* **16**(6–7), 437–445 (2004)
198. Chen, Y., et al.: Comparing study of the effect of nanosized silicon dioxide and microsized silicon dioxide on fibrogenesis in rats. *Toxicol. Ind. Health* **20**(1–5), 21–27 (2004)
199. Bar-Ilan, O., et al.: Toxicity assessments of multisized gold and silver nanoparticles in zebrafish embryos. *Small* **5**(16), 1897–1910 (2009)
200. Choi, J.E., et al.: Induction of oxidative stress and apoptosis by silver nanoparticles in the liver of adult zebrafish. *Aquat. Toxicol.* **100**(2), 151–159 (2010)
201. Ringwood, A.H., et al.: The effects of silver nanoparticles on oyster embryos. *Mar. Environ. Res.* **69**, S49–S51 (2010)
202. Wise, J.P., et al.: Silver nanospheres are cytotoxic and genotoxic to fish cells. *Aquat. Toxicol.* **97**(1), 34–41 (2010)
203. Roh, J.-Y., et al.: Ecotoxicity of silver nanoparticles on the soil nematode *Caenorhabditis elegans* using functional ecotoxicogenomics. *Environ. Sci. Technol.* **43**(10), 3933–3940 (2009)
204. Li, K., et al.: Intracellular oxidative stress and cadmium ions release induce cytotoxicity of unmodified cadmium sulfide quantum dots. *Toxicol. In Vitro* **23**(6), 1007–1013 (2009)
205. Kim, J., et al.: Phototoxicity of CdSe/ZnSe quantum dots with surface coatings of 3-mercaptopropionic acid or tri-n-octylphosphine oxide/gum arabic in *Daphnia magna* under environmentally relevant UV-B light. *Aquat. Toxicol.* **97**(2), 116–124 (2010)
206. Folkmann, J.K., et al.: Oxidatively damaged DNA in rats exposed by oral gavage to C₆₀ fullerenes and single-walled carbon nanotubes. *Environ. Health Perspect.* **117**(5), 703 (2009)



207. Yang, S.-T., et al.: Long-term accumulation and low toxicity of single-walled carbon nanotubes in intravenously exposed mice. *Toxicol. Lett.* **181**(3), 182–189 (2008)
208. Kang, S., Mauter, M.S., Elimelech, M.: Microbial cytotoxicity of carbon-based nanomaterials: implications for river water and wastewater effluent. *Environ. Sci. Technol.* **43**(7), 2648–2653 (2009)
209. Liu, X., et al.: Differential toxicity of carbon nanomaterials in *Drosophila*: larval dietary uptake is benign, but adult exposure causes locomotor impairment and mortality. *Environ. Sci. Technol.* **43**(16), 6357–6363 (2009)
210. Park, E.-J., et al.: Carbon fullerenes (C60s) can induce inflammatory responses in the lung of mice. *Toxicol. Appl. Pharmacol.* **244**(2), 226–233 (2010)
211. Canesi, L., et al.: In vitro effects of suspensions of selected nanoparticles (C₆₀ fullerene, TiO₂, SiO₂) on *Mytilus* hemocytes. *Aquat. Toxicol.* **96**(2), 151–158 (2010)
212. Ringwood, A.H., Levi-Polyachenko, N., Carroll, D.L.: Fullerene exposures with oysters: embryonic, adult, and cellular responses. *Environ. Sci. Technol.* **43**(18), 7136–7141 (2009)
213. Tao, X., et al.: Effects of aqueous stable fullerene nanocrystals (nC₆₀) on *Daphnia magna*: evaluation of sub-lethal reproductive responses and accumulation. *Chemosphere* **77**(11), 1482–1487 (2009)
214. Yang, X., Edelmann, R., Oris, J.: Suspended C₆₀ nanoparticles protect against short-term UV and fluoranthene photo-induced toxicity, but cause long-term cellular damage in *Daphnia magna*. *Aquat. Toxicol.* **100**(2), 202–210 (2010)
215. Zhu, X., et al.: Oxidative stress and growth inhibition in the freshwater fish *Carassius auratus* induced by chronic exposure to sublethal fullerene aggregates. *Environ. Toxicol. Chem.* **27**(9), 1979–1985 (2008)
216. Osman, I.F., et al.: Genotoxicity and cytotoxicity of zinc oxide and titanium dioxide in HEp-2 cells. *Nanomedicine* **5**(8), 1193–1203 (2010)
217. Sharma, V., et al.: Induction of oxidative stress, DNA damage and apoptosis in mouse liver after sub-acute oral exposure to zinc oxide nanoparticles. *Mut. Res. Genet. Toxicol. Environ. Mutagen.* **745**(1), 84–91 (2012)
218. Guan, R., et al.: Cytotoxicity, oxidative stress, and genotoxicity in human hepatocyte and embryonic kidney cells exposed to ZnO nanoparticles. *Nanoscale Res. Lett.* **7**(1), 1–7 (2012)
219. Huang, C.-C., et al.: Oxidative stress, calcium homeostasis, and altered gene expression in human lung epithelial cells exposed to ZnO nanoparticles. *Toxicol. In Vitro* **24**(1), 45–55 (2010)
220. Kiss, B., et al.: Investigation of micronized titanium dioxide penetration in human skin xenografts and its effect on cellular functions of human skin-derived cells. *Exp. Dermatol.* **17**(8), 659–667 (2008)
221. Pan, Z., et al.: Adverse effects of titanium dioxide nanoparticles on human dermal fibroblasts and how to protect cells. *Small* **5**(4), 511–520 (2009)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

