# UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

## Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements

Pant, P.; Harrison, R.M.

DOI: 10.1016/j.atmosenv.2013.04.028

Document Version Early version, also known as pre-print

Citation for published version (Harvard):

Pant, P & Harrison, RM 2013, 'Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements: A review', *Atmospheric Environment*, vol. 77, pp. 78-97. https://doi.org/10.1016/j.atmosenv.2013.04.028

Link to publication on Research at Birmingham portal

#### Publisher Rights Statement:

10.1016/j.atmosenv.2013.04.028 NOTICE: this is the author's version of a work that was accepted for publication in Atmospheric Chemistry. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Atmospheric Chemistry, [vol. 77, 2013] DOI: http://dx.doi.org/10.1016/j.atmosenv.2013.04.028

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

2	Estimation of the Contribution of Road Traffic
3	<b>Emissions to Particulate Matter Concentrations</b>
4	from Field Measurements: A Review
5	
6	Pallavi Pant and Roy M. Harrison <sup>*1</sup>
7	
8	Division of Environmental Health & Risk Management,
9	School of Geography, Earth & Environmental Sciences
10	University of Birmingham, Edgbaston
11	Birmingham B15 2TT
12	United Kingdom
13	

<sup>\*</sup> Corresponding Author: Tel: 00 44 121 414 3494; Fax: 00 44 121 414 3709; Email: r.m.harrison@bham.ac.uk

<sup>&</sup>lt;sup>1</sup> Also at: Centre of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203, Jeddah (21589) Saudi Arabia

#### 14 ABSTRACT

Road traffic is one of the main sources of particulate matter in the atmosphere. Despite its 15 importance, there are significant challenges in quantitative evaluation of its contribution to airborne 16 concentrations. This article first reviews the nature of the particle emissions from road vehicles 17 including both exhaust and non-exhaust (abrasion and resuspension sources). It then briefly 18 reviews the various methods available for quantification of the road traffic contribution. This 19 includes tunnel/roadway measurements, twin site studies, use of vehicle-specific tracers and other 20 21 methods. Finally, the application of receptor modelling methods is briefly described. Based on the review, it can be concluded that while traffic emissions continue to contribute substantially to 22 23 primary PM emissions in urban areas, quantitative knowledge of the contribution, especially of nonexhaust emissions to PM concentrations remain inadquate. 24

25

26 Keywords: Particulate matter; road traffic; exhaust emissions; non-exhaust emissions

#### 28 1. INTRODUCTION

29 Emissions due to road traffic are known to make a large contribution to total particulate matter (PM) 30 concentrations in urban areas and exposure to PM from vehicular emissions has been demonstrated 31 to have detrimental impacts on human health (Mauderly, 1994; Buckeridge et al., 2002; Fan et al., 32 2006; HEI, 2010; Masiol et al., 2012; Rissler et al., 2012). An improved understanding of the characteristics of emissions from different traffic-related sources is therefore vital for conducting 33 34 source apportionment and health effects studies (Schauer et al., 2006). PM emissions from road 35 vehicles include emissions from the tailpipe (exhaust emissions) and emissions due to wear and tear 36 of vehicle parts such as brake, tyre and clutch and re-suspension of dust (non-exhaust emissions). 37 Non-exhaust emissions contribute mainly to the coarse mode of PM (PM<sub>2.5-10</sub>) while exhaust emissions contribute predominantly to fine PM (aerodynamic diameter  $< 2.5 \,\mu$ m) (Abu-Allaban et 38 al., 2003; Tervahattu et al., 2006; Thorpe et al., 2007; Kam et al., 2012) but this is a far from clear 39 40 distinction. A number of different properties of traffic emissions are studied including physical shape and structure, particle size distributions, chemical composition, and temporal and spatial 41 42 variation (Watson and Chow, 2007). Most of the research as well as policy action in the last few 43 decades has largely focused on exhaust emissions, and stringent regulations and technological upgrades have resulted in a decline of the percentage contribution of vehicle tailpipe emissions to 44 total ambient PM concentrations (Allen et al., 2006; Thorpe and Harrison, 2008; Mathissen et al., 45 2011). As a result, the contribution of non-exhaust PM is becoming more important although 46 detailed information on non-exhaust PM emissions is relatively scarce (Kuhlbusch et al., 2009; 47 Amato et al., 2012; Harrison et al., 2011; 2012b; Denier van der Gon et al., 2013). It has been 48 shown that even with zero tail-pipe emissions, traffic will continue to contribute to fine and 49 50 ultrafine particles through non-exhaust emissions (Dahl et al., 2006; Kumar et al., 2013) and it is estimated that nearly 90% of the total emissions from road traffic will come from non-exhaust 51 sources by the end of the decade (Rexeis and Hausberger, 2009). 52

In order to understand the contribution of road traffic emissions to the environment, it is important 53 to understand various aspects including sources, chemical characteristics and quantitative 54 55 contributions. In the last three decades, a significant amount of research has been conducted on 56 characterization and analysis of exhaust emissions and as a result, there is a very strong body of knowledge in the field. Non-exhaust emissions, however, are becoming more important now, and 57 further research is anticipated in this field in the coming years. This review aims to reconcile the 58 59 various studies that have been reported around this issue, and tease out some common 60 characteristics, and highlight areas with future research potential.

This paper reviews the current information on primary road traffic emissions including exhaust and 61 62 non-exhaust emissions. Key characteristics of the major sources are discussed. Various 63 methodologies used for measurement and analysis of such emissions are also discussed with a brief 64 discussion on the merits and shortcomings of each method. In the final section, various source apportionment techniques useful for the estimation of traffic source contributions are discussed 65 together with a summary of some of the reported studies. A number of in-depth review articles have 66 been published in the last few years on various issues related to road traffic emissions on topics 67 such as properties and characteristics of non-exhaust emissions (Thorpe and Harrison, 2008), 68 toxicological impacts of tyre wear emissions (Wik and Dave, 2009), chemical composition of brake 69 70 pads (Chan and Stachowiak, 2004), development of emission factors (Franco et al., 2013), 71 nanoparticle emissions from non-exhaust sources (Kumar et. al., 2013). However, each of these 72 reviews focuses on one or a few specific aspects of road traffic emissions. This review builds on the information provided in those review articles, and repetition is avoided. In summary, the key 73 74 questions that we hope to address through this review article are as follows:

75 1. What are the key characteristics/sources/contributions of road traffic emissions? [What and
76 how much do we know about each of the road traffic emission sources?]

2. How can such emissions be characterized? [Various methods, pros and cons of the methods]

78 3. What have we learnt so far?

79 4. What are the key issues that are not resolved yet?

80

The literature search for this review was primarily performed using Web of Knowledge, Science Direct and Google Scholar using various combinations of keywords such as traffic emissions, nonexhaust/exhaust emissions, resuspension, road dust, traffic PM etc. However, the authors acknowledge that not all articles published on the topic have been included in the review.

85 2. NON-EXHAUST EMISSIONS

Non-exhaust PM comprises the various emissions that do not derive from the tailpipe of a vehicle 86 including particles generated due to brake and tyre wear, road surface abrasion, wear and 87 tear/corrosion of other vehicle components such as the clutch, and re-suspension of road surface 88 dusts (Table 1). Key reasons for needing to understand non-exhaust emissions include their inherent 89 toxicity including their tendency to act as carriers of heavy metals and carcinogenic components 90 91 (Adachi and Tainosho, 2004; Hjortenkrans et al., 2007; Johansson et al., 2009; Amato et al., 2011a) and their contribution to exceedences of air quality guidelines and standards (Amato et al., 2011a; 92 Denier van der Gon, 2013). 93

Harrison et al. (2012b) showed that the size distributions of trace metals were indicative of particle 94 95 sources. Keuken et al. (2010) concluded that re-suspension of accumulated PM and road wear related particles are the primary contributors to non-exhaust emissions, and tyre wear and brake 96 97 wear contribute to the fine and coarse fraction respectively. However, Narvaez et al. (2008) reported that although a majority of abrasion particles are in the coarse fraction, abrasion can 98 99 contribute significantly to the fine fraction of PM. In Taiwan, EFs for elements such as Ba and Se 100 were found to be higher in the  $PM_{2.5-10}$  fraction indicating mechanical friction and brake wear as 101 potential sources (Chiang and Huang, 2009). Road traffic was found to be a key contributor to 102 fine/ultrafine and nano mode particles of Ba, Zn and Pb (Lin et al., 2005) and in Barcelona (Spain), 103 62-96% of elements including Cu, Sb, Ba, Mn and Zn were found to be present in the  $PM_{1-10}$ 104 fraction (Perez et al., 2010). A detailed review of the nanoparticle emissions from non-exhaust 105 sources including traffic and non-traffic sources is presented in Kumar et al. (2013).

106 Non-exhaust emissions are typically characterized by trace metals (e.g. Cu, Zn, Ba, Sb, Mn) 107 although organic markers (e.g. polycyclic aromatic hydrocarbons (PAHs), n-alkanes) have been 108 used in some cases (Wahlin et al., 2006; Amato et al., 2009; El Haddad et al., 2009; Gietl et al., 2010; Oliviera et al., 2011; Kwon and Castaldi, 2012). A list of key tracers used for source 109 110 characterization of non-exhaust emissions is presented in Table 2. However, emissions of trace metal markers are reported to vary with the fleet composition, with higher emissions reported for 111 112 some of the elements for heavy duty vehicles (HDVs) (Grieshop et al., 2006; Mancilla and 113 Mendoza, 2012). In addition, the profile of trace metal concentrations in non-exhaust particulate 114 matter is unique for every region and varies based on parameters such as traffic volume and pattern, vehicle fleet characteristics, driving and traffic patterns and climate and geology of the region 115 116 (Omstedt et al., 2005; Amato et al., 2011a,b; Duong and Lee, 2011). Another important aspect is the variability of type and brake composition depending on the manufacturer which makes it very 117 difficult to ascertain fleet-wide composition other than from environmental measurements 118 119 (Canepari et al., 2008; Gietl et al., 2010; Denier van der Gon et al., 2013).

120

Several factors are reported to affect non-exhaust emissions including increase in vehicle speed 121 122 (Chen et al., 2006; Gustafsson et al., 2008; Hussein et al., 2008; Mathissen et al., 2011) although Keuken et al. (2010) reported a non-linear relationship between traffic volume and non-exhaust 123 emissions. Metal emissions due to road dust show a low correlation to metal emissions due to 124 125 abrasion/brake wear and combustion (Johansson et al., 2009). One of the major problems in analysis of non-exhaust PM using field data has been the difficulty in distinguishing between wear 126 and tear emissions and road dust since the chemical composition is often very similar (Bukowiecki 127 128 et al., 2010). This may, in any case, not be a clear distinction as wear emissions may deposit to the road surface, only to be re-suspended subsequently. Gehrig et al. (2010) used mobile load simulators to compare the contributions of road abrasion and resuspension emissions, and concluded that particle emissions due to abrasion are a function of the state of the pavement whereas resuspension is normally higher, and increases with higher dust loads on the road. Thorpe and Harrison (2008) present a more elaborate account of sources and properties of non-exhaust emissions.

Mixed results have been obtained for the impact of street washing activities on PM and while Amato et al. (2009) observed a reduction, albeit statistically non-significant, in the concentration of trace metals known to be contributed by non-exhaust traffic sources in Spain, Keuken et al. (2010) reported a lack of significant reduction in the non-exhaust emissions after street sweeping or washing in the Netherlands.

#### 140 **2.1 Tyre Wear**

141 Tyre wear particles are generated either by shear forces between the tyre tread and the road surface 142 and are predominantly coarse (PM<sub>2.5-10</sub>) or by volatilization (Rogge et al., 1993; Kupiainen et al., 143 2005; Allen et al., 2006; Thorpe and Harrison, 2008; Aatmeeyata and Sharma, 2010). Tyre wear 144 particles are reported to be generally elongated with rough surfaces based on microscopic analysis 145 (Gunawardana et al., 2011). Tyre tread, a source of airborne particles, contains natural rubber 146 copolymers such as styrene-butadiene rubber and polyisoprene rubber, and zinc (Zn) is added to 147 tyre tread as zinc oxide and organozinc compounds to facilitate the vulcanization process. Passenger 148 car tyres in EU are known to contain nearly 1% zinc oxide, and rubber, metals and carbon black 149 make up typically 47%, 16.5% and 21.5% respectively (Milani et al., 2004). The key tracer 150 components of tyre wear include n-alkanes, n-alkanoic acids, PAHs, benzothiazoles and trace 151 metals (Rogge et al., 1993; Reddy et al., 1997; Camatani et al., 2001; Adachi and Tainosho, 2004; 152 Boulter et al., 2005; Aatmeeyata and Sharma, 2010). Zn is reported to be nearly 1% by weight in rubber tyres (Councell et al., 2004) and tyre wear has been reported to be a significant source of Zn 153

154 (Adachi and Tainosho, 2004; Hjortenkrans et al., 2007; Ondracek et al., 2011). The concentration of Zn was found to be approximately 15 times higher in tyres compared to brakes while concentrations 155 156 of other heavy metals such as copper (Cu) and barium (Ba) were higher for brake materials in 157 Massachusetts (USA) (Apaegyei et al., 2011). Differences have also been observed between tyre tread composition (primarily Zn) and tyre wear (Al, Si, Ca, Fe, Zn, Ti) (Adachi and Tainosho, 158 2004). However, the spatial and temporal trend for Zn was found to be different from other roadside 159 160 tracers in New York (USA), and a very small portion of airborne Zn concentration was attributed to 161 tyre wear (Peltier et al., 2011). However, Zn is emitted from brake wear, motor oil and other 162 sources and cannot be used as the only tracer for tyre wear (Lough et al., 2005). Benzothiazoles are 163 also used as markers for tyre wear, particularly benzothiazole (BT), 2-hydroxy benzothiazole (HOBT), 2-(4-morpholinyl)benzothiazole (24MoBT) and N-cyclohexyl-2-benzothiazolamine 164 (NCBA) (Kumata et al., 2002; Allen et al., 2006; Wik and Dave, 2009). Among the PAHs, pyrene, 165 166 benzo(ghi)perylene, fluoranthene and phenanthrene are known to be emitted from tyres (Boonyatumanond et al., 2007; Kwon and Castaldi, 2012). Aatmeeyata and Sharma (2010) reported 167 168 that small cars tested in India emitted 378 ng of total PAH/tyre/km while in the USA, 200 µg/g of PAHs were found in tyre wear with pyrene, fluoranthene and phenanthrene being present in the 169 highest concentrations (Rogge et al., 1993). Tyre tread wear has also been found to contain an 170 average of 0.53  $\mu$ g/g of dibenzopyrenes, indicating that tyre dust may be significant source of 171 172 dibenzopyrenes in the environment (Sadiktis et al., 2012).

173

Average tread wear for tyres is reported to be between 0.006-0.009 g/km based on the road, tyre and vehicle conditions (Rogge et al., 1993) and tyres can lose up to 10% of their mass during their lifetime (Milani et al., 2004). Speed has been reported as an influential parameter for particle mass and number concentrations for tyre wear emissions (Gustafsson et al., 2008; Mathissen et al., 2011). Other important parameters for tyre tread wear include road surface type, and tyre and driving conditions (Allen et al., 2006; Gustafsson et al., 2008). For example, asphalt surfaces have been reported to cause less tyre wear than concrete pavements and in Arizona, USA, the emission rate for tyre wear was found to be 1.4-2 times lower for asphalt pavement compared to concrete pavement (Allen et al., 2006). The type of tyre also impacts the magnitude of tyre wear emissions and studded tyres are known to cause more emissions compared with summer and friction tyres (Kupiainen et al., 2005; Gustafsson et al., 2008; Hussein et al., 2008; Schaap et al., 2009). Unimodal (70-90 nm) and bimodal (< 10 and 30-60 nm) peaks in the nano size range have been reported for tyre particles by number under low and high speed conditions respectively (Mathissen et al., 2011).

187

#### 188 2.2 Brake Wear

Brake wear, including abrasion of brake lining material and brake discs, caused by grinding of 189 190 brake pad constituents (coarse range particles) or volatilization and condensation of brake pad 191 materials (fine range particles), is known to release PM directly into the atmosphere and to 192 contribute to the trace metal concentration in airborne PM, particularly less than 10 µm (Garg et al., 193 2000; Blau and Meyer, 2003; Ingo et al., 2004; Wahlin et al., 2006; Gietl et al., 2010; Varrica et al., 194 2012). Key components of brake pads include fillers, frictional additives, reinforcing fibres and binder (Chan and Stachowiak, 2004) and the key chemical species used include sulphides of metal, 195 abrasives (e.g. silica), barium silicate/sulphate (particularly in brake linings) and other metallic 196 197 particles (as filler material), carbon fibres and lubricant (e.g. graphite) (Ingo et al., 2004; Dongarra et al., 2009). Chan and Stachowiak (2004) present a detailed review of the chemical constituents of 198 199 brake pads. Garg et al. (2000) observed an average mass median diameter (median particle 200 diameter based on mass) of 1.49 µm for brake wear particles using tests on a brake dynamometer 201 whereas Sanders et al. (2003) reported a mass median diameter of 6  $\mu$ m for brake debris generated 202 during urban driving conditions. Higher brake wear related emissions have been reported during rush hour. A higher number of particles are released in the braking phase as compared to the 203 204 acceleration phase (Hussein et al., 2008; Mathissen et al., 2011) and various studies have corroborated this observations including Greishop et al. (2006) who reported a higher particle 205

number concentration during rush hour where the traffic often operates in the stop-and-go mode and
Abu-Allaban et al. (2003) who observed higher contribution of brake wear at freeway exit sites
compared to other types of roadside sites (Abu-Allaban et al., 2003).

Passenger cars have been estimated to emit nearly 44 g/car/year brake dust (Iijima et al., 2007). 209 Sanders et al. (2003) reported high concentrations of Fe, Ba and Cu in brake lining wear using 210 211 dynamometer and track tests whereas Adachi and Tainosho (2004) reported Fe to be the most abundant metal in brake dust along with other metals such as Ba, Cu, Sb, Zr and Zn, and 212 213 Hjortenkrans et al. (2007) reported Cu, Zn and Sb to be present in brake wear emissions. Varrica et 214 al. (2012) undertook a detailed study on Sb in brake dust and reported the most commonly released 215 forms of Sb due to brake abrasion to be Sb (III) and Sb (V). However, brake pads are often found to 216 have different composition based on the brand, particularly with respect to Cu and Sb (Hjortenkrans 217 et al., 2007; Canepari et al., 2008). Polyalkylene glycol ethers are also reported to be present in 218 brake wear particles with small concentrations of n-alkanes and n-alkanoic acids (Rogge et al., 219 1993).

220

Iijima et al. (2007) reported a unimodal number-based distribution for brake abrasion dust with the mode at 1-2  $\mu$ m while the mass-based distribution had a peak at 3-6  $\mu$ m. EFs for brake wear have been reported by several studies including Garg et al. (2000) who reported a brake wear EF of 3-9 mg/km for light duty gasoline vehicles (LDGV) and Abu-Allaban et al. (2003) who reported an EF of 0-80 mg/km. Wahlin et al. (2006), on the other hand, reported a Cu emission factor of 0.7± 0.2 mg/km/vehicle in Copenhagen.

Sternbeck et al. (2002) proposed the ratio of Cu:Sb ( $4.6 \pm 2.3$ ) as characteristic of brake wear particles and Table 3 lists the Cu: Sb ratios as reported in the literature. Differences in the ratios can be attributed to the difference in the brake pad composition, and contributions of metals from other sources. In some cases, concentrations in ambient air can also vary due to the site characteristics.

## 2.3 Road Dust and Road Surface Wear

Road dust, of which crustal dust is a key component, consists of primarily coarse-sized particles 232 derived from different sources such as traffic, industrial emissions, mineralogical dust etc. 233 234 (Kupianinen et al., 2005; Tanner et al., 2008). Composition of road dust shows spatial as well as 235 temporal variation and it is often difficult to classify dust into crustal/resuspended/direct emission 236 etc. In Monterrey (Mexico), re-suspended dust was found to be contributing nearly 20-25% to the PM<sub>2.5</sub> EF (Mancilla and Mendoza, 2012). The amount of re-suspended road dust particles depends 237 238 on a number of factors including vehicle movement (particularly traffic speed), street maintenance, 239 season and associated meteorological parameters and speed of traffic (Etyemezian et al., 2003; 240 Gertler et al., 2006; Thorpe et al., 2007; Bhaskar and Sharma, 2008; Kaunhaniemi et al., 2011; Laidlaw et al., 2012; Majumdar et al., 2012). Nicholson (1988) presents a detailed review of the 241 mechanisms of resuspension of road dust. Thorpe et al. (2007) reported a strong association 242 243 between heavy duty traffic and re-suspension in the UK with wind speed not found to be a strong 244 influence. In addition, precipitation was found to have no influence on the amount of re-suspension. 245 In somewhat related research, street-washing has been reported to be ineffective for PM control based on experiments in Spain (Karanasiou et al., 2012). However, in Sweden, road wetness was 246 247 found to be an important factor in the amount of re-suspension (Omstedt et al., 2005), but this 248 appears related to road sanding and the use of studded tyres in Sweden. It is also important to note 249 that the residence time of PM<sub>10</sub> on paved roads (travel lanes) has been estimated as a few hours (Etyemezian et al., 2003). In the case of studded tyres, interaction between tyres and the road 250 251 pavement can generate particles less than 0.1 µm (Gustafsson et al., 2008) and such tyres have been shown to increase  $PM_{10}$  emissions by a factor of 1.5 (Tervahattu et al., 2006). The type of pavement 252 253 is another important factor for road dust emissions. Granite pavements have been reported to emit 254 more PM compared to quartzite pavements (Kupiainen et al., 2005; Gustafsson et al., 2009). Tervahattu et al. (2006) also explained the higher PM<sub>10</sub> concentration attributed to use of anti-skid 255

aggregate using the *sandpaper effect*, where the aggregates used to prevent skidding generate  $PM_{10}$ particles and lead to further particle emissions from the pavement.

258

Pristane, phytane, hopanes, steranes in addition to unresolved complex mixture (UCM) were 259 260 reported in road dust particles which were found to be enriched in biogenic component by Omar et al. (2007). High concentrations of phenanthrene, fluoranthene and pyrene along with high 261 molecular weight (4-7 ring) PAHs were reported in road dust samples in Taiwan (Fang et al., 2004), 262 263 China (Liu et al., 2007; Han et al., 2009) and Egypt (Hassanien and Abdel-Latif, 2008) respectively. 264 The spatial distribution of PAHs was reported to be affected by wind direction Liu et al. (2007) and 265 total organic carbon content by Liu et al. (2007) and Han et al. (2009). Mono- ( $\alpha$ - and  $\beta$ -glucose) 266 and disaccharides (sucrose and mycose) were used as markers for road dust re-suspension for 267 source apportionment by Omar et al. (2007).

Han et al. (2007) analysed suspended dust in Beijing and reported concentrations of elements such 268 269 as Ca, S, Cu, Zn, Ni, Pb, and Cd to be much higher than the crustal abundances. Cu, Zn, Ni, and Pb were attributed to traffic emissions together with coal burning, and ions such as  $Ca^{2+}$ ,  $SO_4^{2-}$ ,  $Cl^-$ , 270  $NO_3^-$ , K<sup>+</sup>, and Na<sup>+</sup> were also found to be present in re-suspended dust. Luo et al. (2011) observed 271 272 that the concentrations of trace metals contributed by vehicle exhaust and tyre abrasion, i.e. Pb, Zn 273 and Cu have been reported to be much higher in roadside samples than the concentrations of these elements in background soils in China. Wei and Yang (2010) analysed urban soil, urban road dust 274 275 and agricultural soil in Chinese cities and found relatively higher concentrations of Cd, Cu, Pb and 276 Zn in urban soil samples and Cu, Cr, Pb, Zn, Ni and Cd in urban road dusts. The high concentrations of metals were attributed to anthropogenic sources - road traffic and industrial 277 278 emissions. Higher concentration levels for several trace elements were reported for roadside dust compared to dust collected at a distance of 100 m in Bosnia, and particularly high concentrations 279 280 were reported for Hg and Co and were attributed to vehicle exhaust and brake pads respectively. 281 Elements such as platinum, palladium and rhodium have also been reported to be present in road

282 dust and are attributed to the catalytic converters (Prichard and Fisher, 2012). Duong and Lee (2011) analyzed heavy metal contamination in road dust in high traffic areas in Korea and concluded that 283 284 the concentrations of heavy metals are much higher in high traffic areas in relation to a background 285 site. Their analysis also confirmed a high degree of correlation between the concentration of heavy metals at a specific location and the traffic throughput in the area. Martuzevicius et al. (2011) 286 analysed street dust and reported that it can be a major source for particle-bound PAHs together 287 288 with other PM. Han et al. (2009) reported a high correlation between PAHs and OC in road dust, 289 and traffic emissions were identified as a major contributor to PAHs in road dust using diagnostic 290 ratio analysis. Mathematical models have been proposed for estimation of PM emissions due to 291 road dust although they do not include emissions due to wear of brake/tyre materials (Omstedt et al., 292 2005; Ketzel et al., 2007; Berger and Denby, 2011; Kaunhaniemi et al., 2011).

Aatmeeyata et al. (2009) reported a bimodal number and mass distribution (0.3  $\mu$ m and 4-5  $\mu$ m) for PM<sub>10</sub> generated due to surface (concrete pavement) and tyre wear . Duong and Lee (2011) reported a multi-modal distribution for road dust particles (based on weight) collected from the roadside with a majority of the particles between 180-850  $\mu$ m in Ulsan, Republic of Korea. Chen et al. (2006) reported a bimodal (5-10  $\mu$ m and > 30  $\mu$ m) mass size distribution for road dust particles in Beijing (China).

#### 299 **3. EXHAUST EMISSIONS**

Motor vehicles are an important source for carbonaceous aerosols particularly for the particles in the fine size range (aerodynamic diameter  $< 2.5 \ \mu$ m) (El Haddad et al., 2009; Kam et al., 2012; Keuken et al., 2012). PM emissions from vehicles depend on the engine type and age and maintenance and contain carbon in the form of OC and EC with smaller amounts of trace metals and ions (Brook et al., 2007; Robert et al., 2007a; Fulper et al., 2010; Peltier et al., 2011). For example, Robert et al. (2007a,b) reported that that emission rates for ultrafine and fine particles for heavy-duty diesel vehicles (HDDVs) are one order of magnitude higher than LDGVs. Similarly, 307 Chen et al. (2013) reported 4-times higher diesel EFs compared to gasoline EFs in a tunnel environment. Higher emissions from the high-emitting (smoker) vehicles have previously been 308 309 attributed to lubricating oil (Fujita et al., 2006). Emissions from diesel and gasoline vehicles are 310 different in terms of composition as diesel engines emit both a greater mass of PM and a larger 311 number of ultrafine particles (UFPs) compared to gasoline vehicles (Rose et al., 2006). Gasoline engines are known to release a higher fraction of OC while diesel engines emit more EC (Watson et 312 313 al., 1994; Weingartner et al., 1997; Ntziachristos et al., 2007) although Sodeman et al. (2005) 314 demonstrated a dominance of EC in the ultrafine range in gasoline vehicles. Zhu et al. (2010) 315 reported PM<sub>2.5</sub> emissions from diesel and gasoline vehicles to be rich in different fraction of EC and 316 OC; while EC2 and OC2 fraction were higher in case of diesel vehicles, EC3 and OC3 were higher 317 in case of gasoline vehicle emissions. Vehicles are also a major source of n-alkanes, and diesel 318 engines are known to emit more n-alkanes compared to gasoline engines (Rogge et al., 1993). 319 Results from tunnel studies and chassis dynamometer analyses have shown that lower molecular weight PAHs such as phenanthrene and fluoranthene are characteristic of diesel-derived aerosols 320 321 whereas gasoline engines contributed more to higher molecular weight PAHs such as benzo(ghi)perylene (Smith and Harrison, 1996). PAH emissions from gasoline engines are reported 322 to be one order of magnitude higher than the diesel engine emissions integrated across the U.S. 323 324 vehicle fleet (Lough et al., 2005). A detailed description of diesel particulate matter is presented in 325 Maricq (2007). The situation in relation to exhaust emissions from traffic is constantly changing as abatement technologies develop. In particular the adoption of diesel particle filters is greatly 326 327 reducing mass emissions.

Both trace elements and organic compounds are used as source markers for vehicle emissions. However, since the removal of lead from gasoline, trace metals have proved far less useful as a tracer of engine exhaust (Harrison et al., 2003). Organic compounds released from vehicles are particularly useful markers for conducting receptor modelling analyses since such markers help in distinguishing the vehicular emissions from other sources. The most commonly used molecular markers for vehicular emissions are hopanes and steranes and PAHs such as benzo(ghi)perylene which are attributed to lubricating oil (Rogge et al., 1993; Schauer et al., 1996; Cass, 1998; Lough et al., 2007) and due to their source, the distribution of hopane emissions from vehicles is suggested to be independent of fleet composition (He et al., 2008). It has also been suggested that hopanes and steranes (derived from lubricating oil) and PAHs (reactions at high temperature and fuel combustion) are indicative of different processes during the organic carbon formation in vehicle engines (Lee et al., 1995; Fujita et al., 2006; Riddle et al., 2007; Hanedar et al., 2008).

340

341 A number of PAHs have been reported to be present in vehicle exhaust including benzo(e)pyrene, 342 benzo(a)pyrene, indeno(1,2,3-cd) pyrene, coronene and benzo(ghi)perylene (Rogge et al., 1993; 343 Lough et al., 2007; Riddle et al., 2007). However, Gao et al. (2011) attributed indeno(1,2,3cd)pyrene and benzo(ghi)pervlene to biomass combustion in Guangzhou (China). PAHs are found 344 predominantly in the smallest size fraction ( $<0.4 \mu m$ ), and most of the high molecular weight PAHs 345 are found in the fine particle fraction (Hien et al., 2007). Diagnostic ratio analysis including the 346 ratio between concentrations of indeno(1,2,3-cd) pyrene to the sum of indeno(1,2,3-cd) pyrene and 347 348 benzo(ghi)perylene is also used for distinguishing between gasoline and diesel emissions (Chellam et al., 2005; He et al., 2006; Shen et al., 2010; Ancelet et al., 2011; Kim et al., 2012). Ratios 349 350 between methylphenanthrenes/phenanthrene has also been used for estimation of the contribution of 351 traffic to PAH concentrations in ambient air (Lim et al., 1999). However, it is important to remember that PAHs can be altered by atmospheric chemical reactions (Wu et al., 2007). Also, 352 emission profiles for PAHs are difficult to generalize since they vary across the different vehicle 353 classes (Miguel et al., 1998).Congener profiles of PAH from road traffic have been derived from 354 355 tunnel studies (e.g. Smith and Harrison, 1996) and from roadside concentrations from which the 356 local background is subtracted (Mari et al., 2010). Nielsen (1996) also prepared a traffic PAH profile for Copenhagen (Denmark) using a roadside study. 357

Elemental markers which have been used for vehicular emissions include Cu, Mn, Fe, Zn, Ba, Sn, 359 Ni, Mo and Sb (Lough et al., 2005; Almeida et al., 2006; Birmili et al., 2006; Crawford et al., 2007; 360 Dongarra et al., 2009; Fabretti et al., 2009; Gietl et al., 2010; Amato et al., 2011a,b). Metals can be 361 362 emitted from various exhaust-related sources including fuel and lubricant combustion, catalytic converters, particulate filters and engine corrosion (Lough et al., 2005; Pulles et al., 2012; Sysalova 363 et al., 2012; Varrica et al., 2012) but many of these appear most likely to arise from non-exhaust 364 365 sources. Ni and V have also been reported to be present in emissions due to oil combustion (Pey et 366 al.,, 2010). Cheng et al. (2010) reported higher emission rates for elements such as Br, Ba, Sb and V 367 for gasoline and liquefied petroleum gas (LPG) engines compared to diesel engines in Hong Kong 368 and indicated potential use of these species as markers for gasoline/LPG engines. Ba, Cd, Zn, Sb and V in nanoparticles have been reported to be strongly associated with diesel fuel whereas Cu, 369 Mn and Sr in the particles <0.1 µm have been found to be associated with gasoline (Lin et al., 2005). 370 371 Metal concentrations from diesel and gasoline vehicles are reported to vary over two orders of magnitude (Pulles et al., 2012). In a traffic emissions study in New York (USA), Ba, Br and black 372 carbon (BC) were used as tracers for gasoline/diesel combustion (Peltier et al., 2011). However, the 373 374 levels of trace elements emitted in the exhaust are very low, and great care is needed to distinguish them from non-exhaust traffic emissions and other sources. In addition, different authors have 375 376 attributed metals to different sources which renders precise source attribution with trace metals alone very difficult unless detailed local information on source particle composition is available (e.g. 377 Pant and Harrison, 2012). 378

379

## 4. **PARTICLE SIZE DISTRIBUTIONS**

The size distribution of a particular element or compound not only influences the potential health impact (in terms of respiratory deposition – Harrison et al., 2010), but also influences the extent of atmospheric dispersion (Allen et al., 2001). Size distributions can be measured in terms of mass, number or surface area (Harrison et al., 2000). A number of factors can influence the size distribution 384 including, but not limited to vehicle fleet mix, road type and grade, meteorological conditions (e.g. season type) and distance from road (Zhu et al., 2006; Beddows et al., 2009; Hays et al, 2011; Song 385 and Gao, 2011). For example, differences were observed in the modal structure of particles close to 386 387 the freeway and at a crossroad in Prague and the difference was attributed to the higher concentration of UFPs in the freeway due to the high volume of traffic (Ondracek et al., 2011). 388 However, in a study in the USA, Padro-Martinez et al. (2012) concluded that distance from the 389 390 highway does not impact the particle size distribution. Many other studies (e.g. Shi et al., 1999) 391 have reported the opposite. The behaviour of nanoparticles after emission is discussed later in this 392 section. Size distributions can be useful in ascertaining the sources of exhaust and non-exhaust 393 particle emissions. For example, sampling on a busy London (UK) highway, Harrison et al. (2011) used Positive Matrix Factorization (PMF) to disaggregate the particle size distribution which 394 demonstrated a nucleation mode centered on around 20 nm and a solid particle mode at 50-60 nm in 395 396 the number distribution. However, in using such data, it should be borne in mind that the size distributions measured by cascade impactors may be modified from those in ambient air as an 397 398 artefact caused by the semi-volatile nature of the PAHs, hopanes and steranes.

399

#### 400 4.1 Mass-Size Distributions

401 Based on size distribution analysis for trace metals found in atmospheric aerosols at urban and 402 background sites in the UK, it was concluded that there are three key categories; metals with most 403 of the mass in the accumulation (fine) mode (e.g. Cs, Sn, Pb) with an additional minor mode, metals with mass distributed between fine, coarse and intermediate mode (e.g. Ni, Zn, Cu, Mn) and metals 404 405 with most of the mass in the coarse range (Fe, Ba, Sr) (Allen et al., 2001; Birmili et al., 2006). Song 406 and Gao (2011) reported similar results for size-segregated PM at a highway site in the USA with a 407 bimodal mass-based distribution with the two peaks at 0.32-0.56 µm and 3.2-5.6 µm Table 4 408 summarizes some of the published data for traffic-related elements.

409

410 Similar observations regarding mass-based size distributions for trace metals were reported by Samara and Voutsa (2005) for a traffic site in Greece. Metals such as Ba, Pb and Zn were found to 411 412 exhibit bimodal distributions whereas some others such as Cr and V showed trimodal distributions 413 in Southern Taiwan (Lin et al., 2005). The presence of an Aitken mode in the ultrafine range may 414 be indicative of a contribution from vehicular emissions, both diesel and gasoline. Hays et al. (2011) 415 reported erratic mass size distributions for various elements including Ni, Cu, Cd, and Zn and 416 attributed this to a range of emission sources near highways in North Carolina (USA). A peak was 417 observed at 80 nm for soot particles in a street canyon corresponding to diesel engine emissions in 418 Leipzig, Germany and the concentration of soot particles was reported to be influenced by traffic 419 volume (Rose et al., 2006)

420

421 4.2 Number-Size Distributions

Ultrafine particles are characterized as particles with diameter less than 100 nm diameter and are 422 423 typically measured as particle number concentration (PNC) (Charron and Harrison, 2009). Such 424 particles may have a short lifetime and as a result, the spatial variability of the particles differs based on proximity to the emission source (Puustinen et al., 2007; Cyrys et al., 2008; Charron and 425 426 Harrison, 2009). Traffic has been reported as one of the major sources of UFPs (Abu-Allaban et al., 427 2002; Nanzetta and Holmen, 2004; Westerdahl et al., 2009) and particle number is linked to fresh vehicle exhaust emissions (Rodriguez et al., 2007). Number size distributions are heavily influenced 428 429 by proximity to traffic (Rose et al., 2006; Kerminen et al., 2007; Beddows et al., 2009; Keuken et 430 al., 2012). As a consequence of this, concentrations of organic compounds have also been reported to be highly variable for near-highway sites (Sun et al., 2012. Highest PNC has been reported 431 during morning rush hours in Germany (Wehner et al., 2002) and Spain (Perez et al., 2010). A 432 433 significant difference has also been reported between weekdays and weekends, highlighting the influence of traffic (Morawska et al., 2002; Wehner et al., 2002; Van Poppel et al., 2012). A 434 435 significant difference was observed in the particle number size distribution in the Kingsway tunnel

in the UK (peak at 30-40 nm) and Plabutsch tunnel in Austria (80-100 nm) and this was attributed
to the differences in traffic intensity and the fleet composition, i.e. proportion of HDVs and LDVs
(Imhof et al., 2005a). It is important to note that although PNC is heavily influenced by road traffic,
it is also influenced by other sources including atmospheric nucleation processes (Imhof et al., 2005a; Reche et al., 2011).

441

The typical particle number distribution of a vehicle has 3 modes: nucleation mode (particles < 50442 443 nm), consisting of particles formed due to condensation of exhaust particles; accumulation mode (50-300 nm with a peak at ~ 80-100 nm) consisting of soot particles, and a coarse mode (1µm-444 445 10µm) consisting of particles generated due to abrasion and wear and tear of brake pads and tyres 446 and re-suspension of road dust (Kittleson et al., 2004; Imhof et al., 2005a). The number size distribution for diesel engine emissions has been reported as lognormal with mean diameter 447 between 60-120 nm whereas gasoline engine emissions have asymmetric distributions with a mean 448 diameter between 40-80 nm (Harris and Maricq, 2001). PNC shows higher variability compared to 449 450 particle mass (Weijers et al., 2004). While nuclei mode particles only constitute about 10% of the 451 total particle mass, they contribute more than 90% to PNC (Kittleson et al., 2004). Fushimi et al. (2008) reported the presence of a nuclei mode at 20 nm in the number distribution for PM at 452 453 roadside while no such mode was observed at the background site in Kawasaki, Japan. PNC was 454 reported to be significantly higher in a street canyon compared to an urban background location in a study in Leipzig, Germany (Wehner et al., 2002). 455

456

The particle number distribution for vehicular exhaust emissions measured in ambient air is typically bimodal with one peak at 10-40 nm and another one at 50-90 nm (Charron and Harrison, 2009). Gasoline exhaust particles fall in the size range of 20-60 nm whereas diesel exhaust particles are in the size range of 20-130 nm (accumulation mode between 50-90 nm and a smaller nucleation mode) (Zhu et al., 2002b; Rissler et al., 2012). Such bimodal distributions have been reported for vehicular emissions in many parts of the world including Korea (30-50 nm and 100-400 nm for 463 diesel engines) (Lim et al., 2008), Czech Republic (25-30 and 60-70 nm) (Ondracek et al., 2011) and United Kingdom (13.3 nm and 86.6 nm) (Kumar et al., 2008). On the other hand, multi-modal 464 465 distributions have been reported in several cities and the larger number of modes has been attributed 466 to mixing of traffic and background aerosol (Ondracek et al., 2011). Lingard et al. (2006) reported a four-mode particle number size distribution for an urban roadside site in Leeds (UK) with 467 particles in modes II and III to be highest during the rush hours indicating a traffic source. The four 468 469 modes were classified as sub-11 nm nucleation mode, super-11 nm nucleation mode, Aitken mode 470 and accumulation mode and were attributed to photolysis, diesel engine emissions and secondary 471 aerosols (Lingard et al., 2006). Rosenbohm et al. (2006) observed a bimodal (20 nm and 80 nm) 472 size distribution at a site downwind of the motorway whereas a unimodal distribution (80 nm) was observed for an upwind site. El Haddad et al. (2009) reported a single mode (50-90 nm) distribution 473 for PM in a road tunnel in Marseille. France and Abu-Allaban et al. (2002) observed the highest 474 475 PNC in the nucleation mode (11-17 nm) in a roadway tunnel in USA. A bimodal (11 nm and 80 nm) distribution was reported for  $PM_{0.5}$  emissions during a vehicle chase experiment (Wang et al., 2011). 476 477 Although the magnitude of PNC emissions from a gasoline spark ignition engine exhaust is smaller than from a diesel engine, emissions from gasoline engines are comparable to diesel engines in the 478 cases of acceleration and freeway cruising (Kittleson et al., 2004). Klems et al., (2010) also reported 479 480 higher PNC at the time of acceleration at traffic lights.

Significant differences have been reported between dynamometer and real-world studies on PNC. A mean diameter of 15 nm for PM emissions was measured in Leipzig, Germany in a street canyon though some lab-studies on gasoline and diesel engines have shown mean diameters between 40-70 nm (Wehner et al., 2002). It is important for dynamometer studies to reflect adequately the dilution properties of the atmosphere or they may suppress the nucleation mode which is formed during the initial dilution of the exhaust plume (Shi and Harrison, 1999).

488 For traffic emissions, distance from the road is also reported to cause changes in PNC due to processes other than dilution (Zhu et al., 2006). This feature is of considerable interest; particularly 489 490 from an epidemiological perspective as this would mean that exposure from near-roadway sites can 491 be much different from other sites. HEI (2010) reported the key exposure zone for traffic emissions 492 to be within 300-500 m of a roadway. In Los Angeles, both PNC and size distribution changed with an increase in distance from the road, particularly for particles with a diameter less than 30 nm and 493 494 it has been suggested that this drastic change in PNC is caused by coagulation and atmospheric 495 dilution (Zhu et al., 2002a). Padro-Martinez et al. (2012) found PNC to be at least two times higher 496 at 0-50 m from the highway compared to further distances while Zhu et al. (2002b) observed the 497 PNC to be 25 times higher near the freeway compared to background locations. Rose et al. (2006) also observed a decrease in soot particle number concentration with increasing distance from the 498 street canyon which was attributed to coagulation processes. Kittleson et al. (2004) observed a 499 500 decline in the nuclei mode particles as the distance from the highway increased and recently, Canagaratna et al. (2010) also reported a decrease in PNC with increasing distance away from the 501 502 roadway. Dilution processes will inevitably cause a reduction in PNC with distance from the road, 503 as they do for other traffic-generated pollutants (e.g. NO<sub>x</sub>, BC, etc.). Coagulation causes an increase in particle size, and a decrease in number, but is unlikely to be a significant influence close 504 505 to the road due to an insufficient number concentration to drive the process, and short travel times. An observed increase in particle diameters with distance from a road is most probably the result of 506 mixing of traffic emissions with a coarser background aerosol leading to a large modal diameter 507 508 (Shi et al., 1999). The nucleation mode traffic particles are semi-volatile and this can lead to shifts in size distribution (typically to smaller sizes) and particle loss due to evaporation (Dall'Osto et al., 509 510 2011; Harrison et al., 2012a).

511

512 Charron and Harrison (2003) reported that increasing wind speeds and lower temperatures can513 increase the abundance of smaller nucleation mode particles (diameter of sub-30 nm) and this could

514 be due to the greater dilution and lower condensation sink. Ruellan and Cachier (2001) also reported an influence of wind speed and boundary layer height on PNC. Kumar et al. (2008) 515 reported traffic volume and wind speed as the key variables influencing PNC in street canyon 516 517 conditions and they observed a linear dependence of PNC on traffic volume with a higher correlation in the case of smaller particles (diameter < 300 nm) and a weaker correlation in the case 518 of larger particles for which other sources are large relative contributors. Highest PNC was 519 520 observed for calm winds and lowest for wind speeds above 1.6 m/s in Somerville (USA) (Padro-521 Martinez et al., 2012). PNC was found to be correlated with traffic intensity in France (Ruellan and 522 Cachier, 2001; El Haddad et al., 2009; Perez et al., 2010) and in Los Angeles, PNC was reported to 523 be influenced by traffic density and wind speed and direction (Zhu et al., 2002b). In a study in London, Milan and Barcelona, Rodriguez et al. (2007) concluded that vehicular exhaust emissions 524 largely influence ultrafine particles and found a unimodal distribution for particle number with a 525 526 mode at 37 nm. Kittleson et al. (2004) observed a direct influence of increasing vehicle speed on nuclei-mode particles. Decreasing the engine load from 100% to about 60% was reported to 527 decrease the PNC as well as EC and OC emissions from diesel vehicles (Lim et al., 2008). 528 Morawska et al. (2002), on the other hand, indicated a 70% increase in PNC with a 50% increase in 529 530 traffic flow. While roadside PNC measurements have indicated an influence of temperature on the 531 concentration, similar studies in laboratory conditions using chassis dynamometers have not shown such influence (Abu-Allaban et al., 2002; Kittleson et al., 2004). Abu-Allaban et al. (2002) reported 532 an inverse relationship between ambient air temperature and the diameter of nucleation mode 533 534 particles. Using modelling analysis, Jayaratne et al. (2009) observed an increase in PNC at a signalized pedestrian crossing due to the stop and go nature of traffic at the signal and the increase 535 was found to be higher for HDVs compared to LDVs. This is similar to the observations from Abu-536 537 Allaban et al. (2002) and Wang et al. (2010) who observed higher concentrations arising from HDVs compared to LDVs in USA and Denmark respectively. Rosenbohm et al. (2006) reported a 538 correlation between PNC and NO in Heidelberg (Germany) although no correlation was observed 539

for PNC and  $PM_{10}$  concentration. Similarly, Padro-Martinez et al. (2012) also reported a similar spatial trend for PNC,  $NO_x$ , particulate PAHs and BC in the USA while a different trend was reported for  $PM_{2.5}$ .

543

Thus, key factors that influence the particle number concentration measurements include wind speed, traffic density and the distance from source at which the measurements are made. Nanzetta and Holmen (2004) observed differences in the PNC across different seasons and concluded that results for one season cannot be used for generalization of PNC.

548

#### 549 5. METHODOLOGIES FOR ASSESSMENT OF TRAFFIC EMISSIONS

There are several ways of evaluating emissions from road vehicles including tests with 550 dynamometers, measurements in tunnels/highways and near roadside measurements and road 551 552 simulator tests (Figure 1) (Mclaren et al., 1996; Phuleria et al., 2007; Handler et al., 2008; Gustafsson et al., 2008; El Haddad et al., 2009; Yan et al., 2009; Sjodin et al., 2010). Table 5 553 554 presents a qualitative description of measurements using dynamometers under lab conditions and 555 tunnel/roadway/twin-site studies under ambient conditions. In recent years, significant differences have been observed between laboratory-tested and real-world mixed source traffic emissions 556 (Gertler et al., 2002; Yan et al., 2009; Ancelet et al., 2011). Near roadside/kerbside, traffic tunnel 557 558 and highway measurements are reported to be more realistic since they represent mixed-fleet emissions under real-world driving conditions (Phuleria et al., 2007). Beddows and Harrison (2008) 559 560 compared particle number EFs measured in dynamometer studies with those estimated in roadside 561 studies, finding generally good agreement overall. However, Canagaratna et al. (2010) reported the EFs calculated using on-road measurements to be nearly 50% lower than EFs measured using a 562 563 dynamometer.

An emission factor is defined as "the mass (or number) of pollutant released per unit time/distance 565 travelled or mass of fuel used" and typically increases with vehicle speed for LDVs as well as 566 HDVs (Jones and Harrison, 2006). Characterization of emissions from road vehicles includes 567 568 analysis of size distributions and chemical composition (organic, inorganic and trace metal) and can lead to generation of EFs. EFs typically depend on conditions at the sampling site, traffic intensity 569 and modal shares of HDVs and LDVs (Staehelin et al., 1995; Imhof et al., 2005a). A recent review 570 571 article on development of emission factors delves into this issue in greater detail (Franco et al., 2013). 572

573

EFs can be measured using a number of different methods including dynamometer studies (Oanh et
al., 2010; Chiang et al., 2012), roadway tunnel studies (Gertler et al., 2002; Handler et al., 2008;
Mancilla and Mendoza, 2012), on-road chase experiments (Wang et al., 2011) and remote sensing
(Zhang et al., 1995). Phuleria et al. (2007) reported overall agreement between EFs determined at
tunnel and roadway sites in USA.

579

Two of the most common methods for EF expression are fuel-based EF (Ban-Weiss et al., 2010; 580 581 Wang et al., 2011) and vehicle kilometre travelled (VKT)-based calculation (Abu-Allaban et al., 582 2002; Kittleson et al., 2004; Jones and Harrison, 2006; Amato et al., 2012). Phuleria et al. (2006, 2007) and Bukowiecki et al. (2010) describe the methodology of EF derivation in greater detail. 583 584 Fuel-based EF can be defined as "grams of pollutant emitted for every kilogram of fuel burned" and can be calculated for various pollutants using CO<sub>2</sub> and CO as a measure of fuel burned (Liacos et al., 585 2012). The concentration of pollutant with respect to the concentration of  $CO_2 + CO$  can be used to 586 587 calculate the fuel burnt using the following formula (Miguel et al., 1998; Liacos et al., 2012):

$$EF_{\rm P} = 10^6 \times \left(\frac{\Delta P}{\Delta CO2 + \Delta CO}\right) w_{\rm c} \tag{1}$$

where  $EF_P$  is the emission factor for the pollutant ( $\mu g/kg$  fuel burned),  $\Delta P$  ( $ng/m^3$ ),  $\Delta CO_2$  and  $\Delta CO$ ( $\mu gC/m^3$ ) are the increments over background concentration and w<sub>c</sub> is the fraction of carbon weight in the fuel.

591

592 One of the other most commonly used methods is the derivation of PM EFs using NO<sub>x</sub> EFs, which 593 are calculated as described in the following equation (Gehrig et al., 2004; Imhof et al., 2005b). It is 594 assumed in this case that both NO<sub>x</sub> and PM have similar dilution.

595 
$$D = \frac{N_{\text{LDV}} \times EF NOx_{\text{LDV}} + N_{\text{HDV}} \times EF NOx_{\text{HDV}}}{\Delta NOx}$$
(2)

596 where D is the dilution rate  $(m^2 h^{-1})$ , n is the number of vehicles  $(h^{-1})$ , and EF NO<sub>x</sub> is the emission 597 factor for NOx in mg/km.

598599 Using the dilution rate, the EF (per hour) can be calculated per vehicle based on:

$$EF_{\rm x} = \frac{\Delta Concentration_{\rm x} \times D}{n_{\rm total}} \tag{3}$$

600 where  $\Delta$ Concentration<sub>x</sub> is the difference in concentration of species x between roadside and 601 background and n is the total number of vehicles crossing the sampling point per hour. The 602 respective contribution of HDVs and LDVs can then be estimated with multiple linear regression 603 (MLR) (Gertler and Pierson, 1996; Imhof et al., 2005b). Table 6 presents some EFs reported for 604 PM<sub>10</sub>, PM<sub>2.5</sub> and particle number concentration in the literature.

605

#### 606 6.1 Tunnel/Roadway Measurements

EF calculations from tunnel studies are typically based on calculation of the difference in concentration between entrance and exit (Weingartner et al., 1997; Abu-Allaban et al., 2002), and Gertler and Pierson (1996) describe the methodology for calculation of emission factors using tunnel measurements. Chang et al. (1981) developed a model relating air quality in the tunnel to 611 vehicular emissions which can also be used for calculation of emission factors using tunnel 612 measurements. For the tunnel studies, "the velocity and concentration are assumed to be the same 613 across the tunnel's cross-section and the difference between the influx and outflux is assumed to be 614 the sum of emission rates of vehicles in the tunnel" (Rogak et al., 1998). The following formula is 615 typically used for calculation of the EF (Gertler and Pierson, 1996; He et al., 2008):

$$EF = \frac{C_{\rm out}V_{\rm out} - C_{\rm in}V_{\rm in}}{NL} \tag{4}$$

616 where C is the concentration of the pollutant, V is the volume of air calculated using duration of 617 sampling, cross-sectional area of tunnel and wind speed; N is the number of vehicles and; L is the 618 distance between sampling locations at inlet and outlet of the tunnel.

619

Several studies have been conducted for measurement of PM emissions in roadway tunnels in
different parts of the world including the USA (Fraser et al., 1998; Rogak et al., 1998; Gillies et al.,
2001; Abu-Allaban et al., 2002; Gertler et al., 2002), China (He et al., 2006; He et al., 2008; Chiang
and Huang, 2009), Europe (Stechmann and Dannecker, 1990; Smith and Harrison, 1996;
Weingartner et al., 1997; Imhof et al., 2005a; Handler et al., 2008; El Haddad et al., 2009; Oliviera
et al. 2011), New Zealand (Ancelet et al., 2011), Japan (Funasaka et al., 1998), Taiwan (Chiang and
Huang, 2009), Chile (Caceres et al., 1998) and Mexico (Mancilla and Mendoza, 2012).

627

Although tunnel studies provide an opportunity to conduct measurements on a mixed vehicle fleet, it is important to remember that factors such as variations in speed, aerodynamic conditions in the tunnel and the fleet characteristics (i.e. proportion of HDVs and LDVs) can cause variability in measurements (Rogak et al., 1998; He et al., 2008). In addition, vehicles in the tunnel are often driving at a steady speed which does not happen under other road conditions where traffic follows a stop-and-go pattern (Gertler et al., 2002). Chirico et al. (2011) have shown that the distribution of organic compounds between particles and vapour is heavily affected by the high PM concentrations 635 in a road tunnel, and thus may influence the estimate of emission factors for semi-volatile636 components.

637

 $PM_3$  (particles with diameter < 3  $\mu$ m) was analysed in the Gubrist Tunnel in Zurich, Switzerland 638 639 and it was concluded that most of the particles are emitted from the tailpipe with a small 640 contribution from tyre wear and road dust (Weingartner et al., 1997). While BC accounted for 31% of the total emissions, particulate PAHs accounted for 0.86% of PM<sub>3</sub> (Weingartner et al., 1997). In 641 642 New Zealand, PAHs were reported to contribute nearly 0.10% of the total PM mass with a higher proportion of low molecular weight PAHs indicating a higher contribution of diesel emissions 643 644 (Ancelet et al., 2011). In roadway tunnel studies in China, EC was found to contribute 63% and 45% 645 to PM mass whereas OC contributed 34% and 31% (He et al., 2006; He et al., 2008). Ancelet et al. (2011), however, found OC to contribute between 32-46% to PM mass. Gillies et al. (2001) and He 646 et al. (2008) have attributed higher PM<sub>2.5</sub> emissions to HDVs compared to light duty vehicles 647 (LDVs) in USA and China respectively. Caceres et al. (1998) prepared a PAH source signature for 648 traffic using tunnel measurements in Santiago de Chile whereas Smith and Harrison (1996) 649 650 prepared a source profile fingerprint using tunnel measurements for traffic in Birmingham, UK. 651 Funasaka et al. (1998) analysed emissions in a traffic tunnel in Osaka, Japan and EC and NO<sub>x</sub> (NO as the main component) were found to be correlated with diesel traffic volume whereas OC did not 652 653 show good correlation. In addition, the concentration of EC and SPM was observed to increase linearly with an increase in traffic volume. In Lisbon, (Portugal), nearly 84% of the total PAH mass 654 in a road tunnel was reported to be less than 0.49 µm diameter by Oliviera et al. (2011). 655

656 Concentrations of trace elements were not found to be correlated with EC concentration in Zhujiang 657 Tunnel (China) and the authors concluded non-exhaust sources to be contributing to the trace 658 element mass (He et al., 2008). A similar conclusion was reported by Funasaka et al. (1998) and 659 Ancelet et al. (2011) who attributed coarse particles in a road tunnel in Osaka, Japan and inorganic 660 elements such as Fe, Zn etc. respectively to non-exhaust emissions, primarily re-suspension of road dust. It should be noted, however, that the contributors to road dust in a tunnel will differ from those in a normal roadway environment and that both chemical composition and emission factors will be different. Partitioning of the semi-volatile components between particles and vapour can also be influenced by the high concentrations present (Chirico et al., 2011).

665

#### 666 6.2 Twin-Site Studies

A number of papers have reported the estimation of the contribution of traffic emissions to total PM using twin-site studies in recent years (Yan et al., 2009; Bukowiecki et al., 2010; Gietl et al., 2010; Oliviera et al., 2010; Pey et al., 2010). With the assumption that all sources other than traffic (including any local or regional sources) have the same impact at both roadside and background sites, the increment at the roadside site obtained using the following equation is used as a local traffic increment estimate (Harrison, 2009; Yan et al., 2009; Wang et al., 2010).

673 Concentration of 
$$X_{\text{traffic}} = Concentration of X_{\text{roadside}} - Concentration of X_{\text{background}}$$
 (5)

674 Typically, the difference between rural and urban concentration observation provides an estimate 675 for the urban increment while the difference between roadside and urban background concentration 676 observations provides an estimate for the traffic increment (Harrison, 2009). However, results from such studies can be influenced by street geometry. Harrison et al. (2004) observed different 677 behaviour for particle mass based on the site geometry, where the traffic contribution increases for 678 679 larger daily background mass concentrations at open sites, whereas the magnitude of the traffic 680 mass increment is independent of daily background concentration for enclosed sites, thus making selection of a suitable sampling location critical for coherent results. They also found the pattern of 681 682 air circulation to be an important determinant of ambient PM concentration at an enclosed street site. Roadway studies are useful in providing a reasonable estimate of pollutant concentrations that 683 684 commuters are exposed to (Phuleria et al., 2007) but if the sampling is carried out only at roadside

sites, the data generated is representative only of that part of the total vehicle fleet that was sampled(Liacos et al., 2012).

687

Harrison et al. (2003) concluded that the trace element enrichment observed at roadside sites is 688 689 contributed largely by wear and tear of tyres and brakes. Significant roadside enrichment has been 690 reported for elements, ions and organic compounds in various cities including London (UK) (Gietl et al., 2010) Los Angeles (USA) (Ntziachristos et al., 20082007), Sicily, Rome and Milan (Italy) 691 692 (Manno et al., 2006; Canepari et al., 2008; Perrone et al., 2012), Barcelona (Spain) (Amato et al., 2011), Portugal (Oliviera et al., 2011) and Queensland (Australia) (Gunawardana et al., 2011). . In 693 New York City, Fe, Al and Si were found to contribute between 0.5- 1.0  $\mu$ g/m<sup>3</sup> to the ambient 694 695 PM<sub>2.5</sub> loading (Peltier et al., 2011) whereas Lin et al., 2008 reported Ca, Al and Fe to constitute between 65-77% of suspended particles in Taiwan. Nearly 90% of the road traffic emissions of Cu 696 were reported to be due to brake wear in Stockholm (Sweden) (Johansson et al., 2009). Amato et al. 697 (2011b) analyzed roadside PM<sub>10</sub> enrichment (RE) of trace elements in Barcelona (Spain) reported 698 more than 70% roadside enrichment ratio for key traffic tracers including EC, Fe, Ba, Cu, Sb, Cr 699 700 and Sn. Gietl et al. (2010) estimated that based upon published emission factors brake dust 701 comprises nearly 13% of the total PM at a roadside site in London (UK) and Amato et al. (2011b) 702 reported approximately 60% enrichment of brake wear particles in a roadside location compared to 703 an urban background location in Barcelona (Spain). Pey et al. (2010) reported that the concentration of EC was found to be 3-8 times higher for the urban site in comparison with the 704 background site in Barcelona (Spain) and Perrone et al. (2012) found EC to be contributing 11-15% 705 706 of PM<sub>2.5</sub> in urban areas compared to 3-5% at a rural site and 1-2% at the background site in Milan 707 (Italy). Vertical concentration gradients have also been reported for trace elements with highest 708 concentrations at kerbside as compared with rooftop (Johansson et al., 2009). Analysis of size-709 resolved PM<sub>2.5</sub> at a roadside site in Beijing revealed that re-suspended dust and construction dust

contribute 32.7-50.4% of total  $PM_{2.5}$  (Song et al., 2012) and a reduction was observed in concentrations of species associated with traffic during the 2008 Olympic Games).

712

Comparatively higher concentrations of hopanes, steranes, n-alkanes and PAHs have been reported 713 714 for highway/near-roadway sites compared to other sites (Yan et al., 2009; Aldabe et al., 2011; 715 Peltier et al., 2011; Perrone et al., 2012). Yan et al. (2009) analysed PM<sub>2.5</sub> concentrations at urban 716 and rural sites and observed a higher concentration for EC and primary organic compounds released 717 by motor vehicle exhaust such as hopanes and steranes. They found organic matter to be 718 contributing between 51-72% to PM2.5 mass at the urban site compared with 40-54% at the rural 719 site. Aldabe et al. (2011) reported a concentration gradient for total carbon with the highest 720 concentration at the roadside site and lowest concentration at a rural site, and a decreasing trend of 721 OC concentration was observed in Prague with a decrease in traffic intensity (Ondracek et al., 2011). 722 Gao et al. (2011) also reported higher concentrations for EC and hopanes at the roadside site compared to a rooftop site, highlighting the contribution of traffic emissions. Jones and Harrison 723 724 (2006) used ratios of concentration to NO<sub>x</sub> for determination of EFs for road traffic or particle 725 number and mass metrics with the assumption that particle dispersion is similar to dispersion of NO<sub>x</sub> in the atmosphere. The EFs that they obtained from this study for particle mass and number 726 727 (Table 6) were comparable with emission factors calculated using other methods such as 728 dynamometer and tunnel/roadway studies. Liacos et al. (2012) observed higher concentrations of 729 trace metals using on-road measurements and postulated that studies using sampling from near 730 roadside conditions might not always capture the pollutant concentrations that drivers are exposed 731 to (Liacos et al., 2012).

732

#### 733 **6.3 NO**<sub>x</sub> as Tracer

With the assumption that road traffic is the primary urban source for  $NO_x$  in the urban atmosphere, Harrison et al. (1997) used regression analysis between particle mass fractions and  $NO_x$  to obtain 736 contributions from the traffic source for Birmingham (UK) with a total of 32% of the fine PM concentration attributed to vehicular exhaust emissions for the winter season. The method is very 737 simple, using the assumption that the regression intercept in PM at zero NO<sub>x</sub> is due to non-traffic 738 739 sources, with the traffic contribution estimated by difference from the measured mean concentration. 740 The same approach was also used by Fuller et al. (2002) who estimated  $PM_{10}$  and  $PM_{2.5}$ concentrations using NO<sub>x</sub> and PM emissions for London (UK) and derived a typical PM<sub>10</sub>: NO<sub>x</sub> 741 742 ratio of  $0.045 \pm 0.003$ . Lim et al. (1999) also used this approach to determine the contribution of 743 PAHs to traffic in Birmingham and concluded that traffic contributed 80-82% and 61-67% of PAH 744 concentrations in the city centre and an urban background site (university campus) respectively. 745 Thorpe et al. (2007) used this method to calculate re-suspension EFs. The  $PM_{10}$  emission factor was calculated using NO<sub>x</sub> as a tracer and coarse particle source strength and abrasion emissions were 746 subtracted from total  $PM_{10}$  emission to a obtain a re-suspension emission factor. However, it is 747 748 important to note that this method is only useful if the predominant source for local ground-level NO<sub>x</sub> is traffic, unless used in a twin-site approach. 749

750

#### 751 **6.4** Other Methods

752 Several studies have reported lower levels of vehicular emissions on weekends (Rose et al., 2006; Barmpadimos et al., 2011). Analysis of the weekly cycle of coarse PM concentrations (2.5-10 µm) 753 754 with the weekly PM coarse traffic emission was carried out in Zurich (Switzerland) and it was estimated that nearly 70% and 53% of the ambient coarse PM mass is contributed by traffic on 755 weekdays and weekends respectively (Barmapadimos et al., 2011). With the assumption that the 756 difference in PM concentrations between weekdays and weekends at an urban background site is 757 758 due to traffic movement, it is possible to calculate the traffic increment to urban PM concentrations, 759 although the results show large uncertainties (Jones et al., 2008).

Vehicle-chase experiments have also been used where vehicles are chased for a specific period of time to collect pollutant concentrations allowing data to be collected for real-world conditions of dilution (Canagaratna et al. 2010; Wang et al., 2011). However, the EFs are only valid for the specific sampling conditions and there is no control over driving conditions (Wang et al., 2011).

764

6.

## **RECEPTOR MODELLING OF TRAFFIC PM**

765 Watson and Chow (2007) describe receptor models as models that "interpret measurements of physical and chemical properties taken at different times and places to infer the possible sources of 766 excessive concentrations and to quantify the contributions from those sources". A number of 767 768 receptor models are used for source apportionment including the Chemical Mass Balance (CMB) model, statistical models such as PCA and PMF, Multilinear Engine (ME), Constrained Physical 769 Receptor Model (COPREM) and UNMIX. With the assumption that the relative concentrations of 770 771 chemical species are preserved between sources and receptors, receptor models use the principle of mass conservation for apportionment of PM mass to different air pollution sources. Thus, the 772 773 concentration of a species measured in a particular sample can be described as (Hopke, 2003):

$$X_{ij} = \sum_{p=1}^{p} g_{ip} f_{pj} + e_{ij}$$
(6)

where  $X_{ij}$  is the measured concentration of the *j*th species in the *i*th sample,  $f_{pj}$  is the concentration of the *j*th species in material emitted by the source *p*,  $g_{ip}$  is the contribution of the *p*th source to the *i*th sample and  $e_{ij}$  is the portion of the measurement that cannot be fitted by the model. Different receptor models use different approaches to solve this equation, for e.g., the CMB model uses the effective-variance least squares method whereas UNMIX uses eigenvector analysis. Concentrations of trace elements (e.g. Si, Fe, Cu), ions (e.g.  $SO_4^{2-}$ ), EC/OC and organic compounds (e.g. PAHs, hopanes, alkanoic acids) are used as inputs for the models.

In recent years, there have been many studies for source apportionment of road traffic-generated
PM, some of which appear in Table 7. There have also been studies that have used both CMB and
factor analysis models (Larsen and Baker, 2003; Bullock et al., 2008).

- 785
- 786 **6.1**

## .1 Multivariate Statistical Methods

There are a number of different methods based on factor analysis including PCA, PMF, UNMIX and ME. Such methods do not require a priori information about source emission characteristics and are useful in cases where relevant source profiles are not available (Hopke, 2003; Viana et al., 2008). In the simplest matrix form, the equation 6 can be represented as (Hopke, 2003)

$$X = GF'$$

A number of studies have been conducted using the different factor analysis methods and studies in 791 792 different locations have often reported different magnitudes of exhaust and non-exhaust emissions. 793 In Taiwan, gasoline, diesel and fuel oil emissions together with dust and brake wear were found to 794 be key contributors to traffic-generated PM using PCA (Lin et al., 2005). Using COPREM, Wahlin 795 et al. (2006) apportioned the local traffic emissions to exhaust, road dust, brake wear and winter 796 salting in Copenhagen (Denmark). Amato et al. (2009) analysed road dust in Barcelona (Spain) 797 using ME-2, and re-suspension was found to contribute 37%, 15% and 3% of total traffic emissions respectively in PM<sub>10</sub>, PM<sub>2.5</sub> and PM<sub>1</sub> respectively. Other identified sources included secondary 798 799 aerosols and a soil source. Sjodin et al. (2010) found road wear to be the most important source of PM<sub>10</sub> emissions in streets as well as background locations in Sweden, which is probably a reflection 800 801 of road sanding and use of studded tyres. Using PMF, three distinct traffic sources including vehicle 802 abrasion (36%), re-suspension (43%) and fuel combustion (21%) were resolved for traffic sources 803 using trace metals in Nice (France) from data collected in a road tunnel (Fabretti et al., 2009). Based on PMF analysis in Switzerland, Bukowiecki et al. (2010) reported 21%, 38% and 41% of the 804 traffic emissions to be contributed by brake wear, re-suspended road dust and exhaust emissions 805

806 respectively for street canyons. For freeways, while the contribution of exhaust emissions remained 807 the same, the contribution of brake wear was much less at 3% whereas re-suspended road dust 808 contributed as much as 56%. Movement of traffic also contributes to re-suspension of road dust and 809 re-suspended road dust has been estimated to contribute as much as 38% of the total PM in a street 810 canyon and 56% on a highway (Bukowiecki et al., 2010; Laidlaw et al., 2012). Analysis of PM in 811 Madrid (Spain) using PMF estimated that nearly 29% of the PM<sub>10</sub> in the city is contributed by road 812 dust while another 31% is contributed by direct vehicular emission (Karanasiou et al., 2011). In 813 Oporto (Portugal), exhaust emissions and re-suspension due to traffic were found to contribute 12-814 55% of the coarse PM and 14-66% of the fine PM based on PCA- multiple regression analysis and 815 inter-site mass balance analysis (Oliviera et al., 2010). They used inter-site mass balance for separating the traffic increment in the road dust and concluded that nearly 30% of the fine fraction 816 and 50-70% of the coarse fraction PM is contributed by road dust. They also used a roadside impact 817 818 factor (defined as the fractional difference between the concentrations at the roadside and urban background) and the highest factors were observed for Cu, Zr, Fe, Cr and Sn indicating a traffic 819 820 source for these metals. Thorpe et al. (2007) reported re-suspended road dust, abrasion source emissions and exhaust and fine-range abrasion emissions to contribute 20-22%, 19% and 60% to 821 roadside incremental PM<sub>10</sub> concentration respectively in a major highway in London. Four major 822 823 sources for road surface dust particles of diameter less than 10 µm including motor exhaust, brake 824 and tyre wear and road surface wear together with background urban dust were reported for European cities with an increase in the concentrations of trace metals and OC/EC with an increase 825 826 in traffic volume (Amato et al., 2011a). Song and Gao (2011) identified three distinct sources of trace metals at a highway site in New Jersey (USA) using factor analysis: brake wear and fuel 827 828 combustion, primary fuel combustion and tyre wear and fuel combustion. PCA analysis was used 829 for determining PM<sub>2.5</sub> sources in Incheon (Korea) and motor vehicles/sea salt and soil emissions were found to contribute 39% and 7% respectively (Choi et al., 2012). Similar results were obtained 830

by Cheng et al. (2011) in Hong Kong where traffic emissions and re-suspended dust emissions werefound to contribute 38% and 5% using PMF.

833 Dreyfus et al. (2009) conducted PMF analysis with time-resolved organic molecular marker data and identified diesel and road dust/car emissions factors which accounted for 2/3<sup>rd</sup> of the total OC 834 concentration. Receptor modelling analysis for PM2.5 was conducted using PMF and UNMIX, and 835 PMF distinguished between diesel and gasoline engine emissions using temperature resolved 836 carbon fractions (Sahu et al., 2011). Pey et al. (2009) performed source apportionment using PCA 837 and regression with a large dataset including information on particle number concentration; size 838 839 distribution and chemical composition for an urban background site in Barcelona, and road traffic 840 was identified as the most significant source (52-86%) for aerosols in the sub-micrometer range.

Liu et al. (2007) attributed the PAHs in road dust in Shanghai (China) to traffic emissions and coal 841 combustion using principal component analysis (PCA) and compositional analysis. Using the same 842 843 technique, Han et al. (2009) attributed PAHs in road dust in Anshan (China) to emissions from 844 traffic, coal combustion, steel industry and cooking. Using MLR, Boonyatumanond et al. (2007) reported tyres, diesel vehicle soot and used engine oil to be the key sources for PAHs found in street 845 846 dust in Bangkok (Thailand). Hassanien and Abdel-Latif (2008) attributed PAHs in road dust in 847 Cairo (Egypt) to emissions from traffic, industry and incomplete combustion of waste based on 848 correlation analysis. Teixeira et al. (2012) reported vehicular emissions and coal and wood combustion to be the key sources for PAHs in Porte Alegre (Brazil) using PCA. 849

Other approaches have also been used for estimation of source contributions. Recently, Harrison et al. (2012b) used field data for estimating the contribution of brake wear, tyre wear and road dust using size distributions of Ba, Zn and Si as the tracers for the sources respectively. By merging size distributions measured with a Scanning Mobility Particle Sizer (SMPS) and an Aerodynamic Particle Sizer (APS), Harrison et al. (2011) were able to generate continuous particle size distributions for a roadside site from 15 nm – 10  $\mu$ m diameter. By applying PMF, they were able to

separately identify nucleation mode and accumulation mode exhaust particles as well as brake dustand re-suspension particles, and several particle classes arising from the urban background.

858

## 859 7.2 Chemical Mass Balance (CMB) Model

The CMB model uses the ambient measurement data for chemical species together with the associated uncertainty and source profiles for different sources as inputs and the output consists of estimates of contribution of each source to the total mass. The CMB model has several assumptions including (Watson et al., 2002):

chemical species do not react with one another and the species composition does not change
between the source(s) and the receptor(s)

- number of species are more than the number of sources
- all of the potential sources are included the model

• source profiles are not collinear

• random and non-correlated uncertainties

870

871 It is important to note that while the multivariate models do not require any information on source emission composition, the CMB model requires the species abundances (in the form of source 872 profiles) in each source as a model input. Source profiles are "the mass abundances, i.e. fraction of 873 874 total mass of chemical species in source emissions, and such profiles are generally representative of 875 source categories rather than individual emitters" (Watson et al., 2002). There are several ways of preparing source profiles for road vehicles including tests with chassis dynamometers, 876 measurements in tunnels/ highways and near roadside measurements (Phuleria et al., 2007; Handler 877 et al., 2008; Yan et al., 2009) as discussed in section 6. Source profiles are used for quantification 878 879 of source contributions to particulate matter using the chemical mass balance (CMB) model as well 880 as to compare and validate results obtained from factor analysis models. The CMB model relies to a 881 large extent on the accuracy of the source profiles used as an input. The typical components of any

882 source profile are found to be more-or-less similar although the mass abundances vary depending on location and emitter characteristics. Thus, different combinations of source profiles can provide 883 884 statistically valid yet completely different solutions and selection of a source profile comparable to 885 ambient data is one of the key prerequisites for a good CMB solution (Robinson et al., 2006). 886 Absence of locally-representative source profiles and inter-region variability can result in errors in estimation of source contributions from the CMB model (Yan et al., 2009; Rutter et al., 2011). 887 888 Significant differences have also been reported between real-world emissions and laboratory-tested 889 emissions (Yan et al., 2009; Ancelet et al., 2011). Analysis of vehicular source profiles in Bangkok, 890 Thailand led to the conclusion that while source profiles for different vehicle categories vary 891 significantly, ranges of the light duty and heavy duty average source profiles overlap and can be represented by a single composite profile (Oanh et al., 2010). Fraser et al. (2003) and Fujita et al. 892 (2007) used the CMB model to analyse the apportionment of PM mass between diesel and gasoline 893 894 vehicles. While high molecular weight (4-7 ring) PAHs (coronene, benzo(ghi)perylene), hopanes and steranes were found to be influential in case of gasoline engine emissions, 2-4 ring methylated 895 896 PAHs and EC were reported to be key species in the case of diesel engine emissions. Hanedar et al. (2008) used the CMB model to resolve the sources of PAHs in Istanbul (Turkey) and identified 897 vehicle emissions (gasoline and diesel), natural gas combustion, coal and wood burning as major 898 899 sources with vehicular emissions contributing more than 50% at both urban and rural location. 900 Perrone et al. (2012) used CMB analysis to resolve PM sources in Milan (Italy) and reported traffic emissions to contribute between 17-24% to PM<sub>2.5</sub> mass in an urban location compared to 7-9% at a 901 902 rural site and El Haddad et al. (2011) reported vehicular emissions to contribute 17% to the PM mass in Marseille (France). In Detroit (USA), traffic emissions and road dust emissions were 903 904 reported to contribute 24-36% and 3-5% respectively of mass using CMB analysis of PM<sub>2.5</sub> data.

905

906 In recent years, there have been several studies focusing on characterization of trace metals and/or907 organic marker compounds in PM contributed by road traffic emissions and preparation of mixed

908 traffic source profiles (El Haddad et al., 2009; Fabretti et al., 2009; Yan et al., 2009; Ancelet et al., 909 2011). El Haddad et al. (2009) prepared a source profile for the vehicle fleet in France using tunnel 910 measurements and nearly 70 organic markers, whereas Fabretti et al. (2009) used trace element 911 marker data from tunnel-based measurements to derive chemical fingerprints for different trafficrelated sources including combustion, re-suspended dust and abrasion-related particles. Tunnel 912 studies were also used in USA for preparation of source profiles for gasoline and diesel engines 913 using inorganic and organic markers (Phuleria et al., 2006; Landis et al., 2007). Yin et al. (2010) 914 915 used organic molecular source tracers to provide separate quantification of gasoline, diesel and 916 smoker vehicle emissions at two UK sites. Recently, an attempt has also been made to characterize 917 the contribution of lubricating oil and fuel to the PM emissions (Sonntag et al., 2012).

918

#### 919 7. CONCLUSIONS

920 Road traffic frequently makes a very significant contribution (5-80% depending on site and location) to airborne concentrations of PM, whether expressed in terms of particle mass or number 921 concentration (Thorpe and Harrison, 2008; Belis et al., 2013). Particles arise not only from the 922 923 engine exhaust but also from abrasion of tyres, road surface and brake components as well as from resuspension from the road surface. The sizes of particles emitted from these sources differ 924 considerably and size association may be a useful guide to particle source, as is chemical 925 926 composition. Fleet mix, type of road, traffic characteristic and fuel type are the factors that impact the emission type and the relative contributions from exhaust and non-exhaust emissions. The 927 determination of particle mass and number increments due to traffic emissions can be based on a 928 number of methods including measurements in well controlled environments such as a road tunnel, 929 930 or twin site studies such as a highway and urban background (allowing for subtraction of the 931 background contribution), as well as using vehicle-specific tracers and the more sophisticated receptor modelling methods based on multi-variate statistics or Chemical Mass Balance which have 932 933 been used in many studies to estimate the contributions to particulate matter concentrations, not

only from road vehicles but also from other sources. However, significant gaps still exist in ourunderstanding of traffic emissions, particularly:

936 1. Characterization and quantification of different non-exhaust sources.

- 937 2. Impacts of non-exhaust emissions upon human health: While there are many studies reporting
- links between exposure to air pollution and adverse health impacts, detailed information on
- the components that contribute to toxicity is missing (Kelly and Fussell, 2012).
- 940 3. Aerosol chemistry in high-traffic environments, and the evolution of particles emitted from941 vehicles
- 942 Further, most of the reported analyses have been carried out in the USA or Europe, and there is a
- 943 lack of reliable information on traffic emissions in areas with high population density in Africa,
- 944 Asia and South America. The field offers much scope for future research , including the
- 945 development of enhanced methods for quantification of non-exhaust contributions to airborne
- 946 concentrations.

## 948 **REFERENCES**

949

953

956

960

964

967

979

983

- Aatmeeyata, K. D. S. and Sharma, M. (2009). Traffic generated non-exhaust particulate emissions
  from concrete pavement: A mass and particle size study for two-wheelers and small cars.
  Atmospheric Environment, 43(35), 5691-5697.
- Aatmeeyata, K. D. S., and Sharma, M. (2010). Polycyclic aromatic hydrocarbons, elemental and
   organic carbon emissions from tire-wear. Science of the Total Environment, 408(20), 4563-4568.
- Abu-Allaban, M., Coulomb, W., Gertler, A.W., Gillies, J., Pierson, W.R., Rogers, C.F., Sagebiel,
  J.C. and Tarnay, L. (2002). Exhaust particle size distribution measurements at the Tuscarora
  Mountain Tunnel. Aerosol Science and Technology, 36(6), 771-789.
- Abu-Allaban, M., Gillies, J.A., Gertler, A.W., Clayton, R. and Proffitt, D. (2003). Tailpipe,
  resuspended road dust, and brake wear emission factors from on-road vehicles. Atmospheric
  Environment, 37, 5283-5293.
- Adachi, K., and Tainosho, Y. (2004). Characterization of heavy metal particles embedded in tire
  dust. Environment International, 30(8), 1009-1017.
- Aldabe, J., Elustondo, D., Santamaría, C., Lasheras, E., Pandolfi, M., Alastuey, A. and Querol, X.
  (2011). Chemical characterisation and source apportionment of PM2.5 and PM10 at rural, urban and traffic sites in Navarra (North of Spain). Atmospheric Research, 102(1-2), 191-205.
  971
- Allen, A. G., Nemitz, E., Shi, J. P., Harrison, R. M., and Greenwood, J. C. (2001). Size distributions
  of trace metals in atmospheric aerosols in the United Kingdom. Atmospheric Environment, 35(27),
  4581-4591.
- Allen, J. O., Alexandrova, O. and Kaloush, K.E. (2006). Tire wear emissions for asphalt rubber and
  portland cement concrete pavement surfaces: Arizona State University. Report submitted to Arizona
  Department of Transportation (Contract KR-04-0720-TRN).
- Almeida, S. M., Pio, C. A., Freitas, M. C., Reis, M. A., and Trancoso, M. A. (2006). Source
  apportionment of atmospheric urban aerosol based on weekdays/weekend variability: evaluation of
  road re-suspended dust contribution. Atmospheric Environment, 40(11), 2058-2067.
- Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A., and Moreno, T. (2009). Spatial and
  chemical patterns of PM10 in road dust deposited in urban environment. Atmospheric Environment,
  43(9), 1650-1659.
- Amato, F., Pandolfi, M., Moreno, T., Furger, M., Pey, J., Alastuey, A., Bukowiecki, N., Prevot,
  A.S.H., Baltensperger, U. and Querol, X. (2011a). Sources and variability of inhalable road dust
  particles in three European cities. Atmospheric Environment, 45(37), 6777-6787.
- 991
- Amato, F., Viana, M., Richard, A., Furger, M., Prévôt, A. S. H., Nava, S., Lucarelli, F., Bukowiecki,
  N., Alastuey, A., Reche, C., Moreno, T., Pandolfi, M., Pey, J. and Querol, X. (2011b). Size and
  time-resolved roadside enrichment of atmospheric particulate pollutants. Atmospheric Chemistry
  and Physics, 11(6), 2917-2931.
- Amato, F., Karanasiou, A., Moreno, T., Alastuey, A., Orza, J.A.G., Lumbreras, J., Borge, R., Boldo,
  E., Linares. C. and Querol, X. (2012). Emission factors from road dust resuspension in a
  Mediterranean freeway. Atmospheric Environment, 61, 580-587.

1000 Ancelet, T., Davy, P. K., Trompetter, W. J., Markwitz, A., and Weatherburn, D. C. (2011). 1001 Carbonaceous aerosols in an urban tunnel. Atmospheric Environment, 45(26), 4463-4469. 1002 1003 Apeagyei, E., Bank, M. S. and Spengler, J. D. (2011). Distribution of heavy metals in road dust 1004 along an urban-rural gradient in Massachusetts. Atmospheric Environment, 45(13), 2310-2323. 1005 1006 Ban-Weiss, G.A., Lunden, M.M, Kirchstetter, T.W. and Harley, R.A. (2010). Size-resolved particle 1007 number and volume emission factors for gasoline and diesel motor vehicles. Aerosol Science, 41, 5-1008 12. 1009 1010 Barmpadimos, I., Nufer, M., Oderbolz, D. C., Keller, J., Aksoyoglu, S., Hueglin, C., Baltensperger, 1011 U. and Prevot, A.S.H. (2011). The weekly cycle of ambient concentrations and traffic emissions of 1012 coarse (PM10-PM2.5) atmospheric particles. Atmospheric Environment, 45(27), 4580-4590. 1013 1014 Beddows, D. C. S. and Harrison, R. M. (2008). Comparison of average particle number emission 1015 factors for heavy and light duty vehicles derived from rolling chassis dynamometer and field studies. 1016 Atmospheric Environment, 42, 7954-7966. 1017 1018 Beddows, D. C. S., Dall'Osto, M. and Harrison, R. M. (2009). Cluster analysis of rural, urban and 1019 curbside atmospheric particle size data. Environmental Science and Technology, 43, 4694-4700. 1020 Belis, C.A., Karagulian, F., Larsen, B.R. and Hopke, P.K. (2013). Critical review and meta-analysis 1021 1022 of ambient particulate matter source apportionment using receptor models in Europe. Atmospheric 1023 Environment, 69, 94-108. 1024 1025 Berger, J. and Denby, B. (2011). A generalized model for traffic induced road dust emissions. 1026 Model description and evaluation. Atmospheric Environment, 45, 3692-3703. 1027 1028 Bhaskar, V.S. and Sharma, M. (2008). Assessment of fugitive road dust emissions in Kanpur, India. 1029 Transportation Research Part D, 13, 400-403. 1030 1031 Birmili, W., Allen, A. G., Bary, F., and Harrison, R. M. (2006). Trace Metal Concentrations and Water Solubility in Size-Fractionated Atmospheric Particles and Influence of Road Traffic. 1032 1033 Environmental Science and Technology, 40(4), 1144-1153. 1034 1035 Blau, P.J. and Meyer, H. M. III (2003). Characteristics of wear particles produced during friction 1036 tests of conventional and unconventional disc brake materials. Wear, 255, 1261-1269. 1037 1038 Boonyatumanond, R., Murakami, M., Wattayakorn, G. Togo, A. and Takeda, H. (2007). Sources of 1039 polycyclic aromatic hydrocarbons (PAHs) in street dust in a tropical Asian mega-city, Bangkok, 1040 Thailand. Science of the Total Environment, 384, 420-432. 1041 1042 Boulter, P. G., Thorpe, A. J., Harrison, R. M. and Allen, A. G. (2005). Road vehicle non-exhaust 1043 particulate matter: Final report on emission modelling. Published Project Report PPR110. Available 1044 at http://uk-air.defra.gov.uk/reports/cat15/0706061624 Report2 Emission modelling.PDF. 1045 1046 Brook, J. R., Graham, L., Charland, J. P., Cheng, Y., Fan, X., Lu, G., Li, S.M., Lillyman, C., MacDonald, P., Caravaggio, G. and MacPhee, J.A. (2007). Investigation of the motor vehicle 1047 1048 exhaust contribution to primary fine particle organic carbon in urban air. Atmospheric Environment, 1049 41(1), 119-135. 1050

- Buckeridge, D.L., Glazier, R., Harvey, B.J., Escobar, M., Amrhein, C. and Frank, J. (2002). Effect
  of motor vehicle emissions on respiratory health in an urban area. Environmental Health
  Perspectives, 110(3), 293-300.
- Bukowiecki, N., Lienemann, P., Hill, M., Furger, M., Richard, A., Amato, F., Prevot, A.S.H.,
  Baltensperger, U., Buchmann, B. and Gehrig, R. (2010). PM10 emission factors for non-exhaust
  particles generated by road traffic in an urban street canyon and along a freeway in Switzerland.
  Atmospheric Environment, 44(19), 2330-2340.
- Bullock, K.R., Duvall, R.M., Norris, G.A., McDow, S.R. and Hays, M.D. (2008). Evaluation of the
  CMB and PMF models using organic molecular markers in fine particulate matter collected during
  the Pittsburgh Air Quality Study. Atmospheric Environmet, 42, 6897-6904.

1063

1071

1075

1079

1086

1090

1093

1096

- Caceres, D., Adonis, M., Gil, L., and Miguel, A. H. (1998). Exhaust PAH in Chilean vehicles:
  Levels, source signatures and receptor modeling. Journal of Aerosol Science, 29, Supplement 2(0),
  S741-S742.
- 1068 Camatani, M., Crosta, G. F., Dolukhanyan, T., Sung, C., Giuliani, G., Corbetta, G. M., Cencetti, S.
  1069 and Regazzoni, C. (2001). Microcharacterization and identification of tire debris in heterogenous
  1070 laboratory and environmental specimens. Materials Characterization, 46, 271-283.
- 1072 Canagaratna, M. R, Onasch, T. B., Wood, E. C., Herndon, S. C., Jayne, J. T., Cross, E. S., Miake1073 Lye, R. C., Kolb, C. E. and Worsnop, D. R. (2010). Evolution of vehicle exhaust particles in the
  1074 Atmosphere. Journal of Air and Waste Management Association, 60, 1192-1203.
- 1076 Canepari, S., Perrino, C., Olivieri, F., and Astolfi, M. L. (2008). Characterisation of the traffic
  1077 sources of PM through size-segregated sampling, sequential leaching and ICP analysis.
  1078 Atmospheric Environment, 42(35), 8161-8175.
- 1080 Cass, G. R. (1998). Organic molecular tracers for particulate air pollution sources. Trends in
  1081 Analytical Chemistry, 17(6), 356-366.
  1082
- 1083 Chan, D. and Stachowiak, G. W. (2004). Review of automotive brake friction materials.
  1084 Proceedings of the Institution of Mechanical Engineers, Part D: Journal of Automobile Engineering,
  1085 218 (9), 953-966.
- 1087 Chan, T. W. and Mozurkewich, M. (2007). Application of absolute principal component analysis to
  1088 size distribution data: identification of particle origins. Atmospheric Chemistry and Physics, 7, 8871089 897.
- 1091 Chang, T. Y., Modzelewski, S. W., Norbeck, J. M., and Pierson, W. R. (1981). Tunnel air quality
  1092 and vehicle emissions. Atmospheric Environment, 15(6), 1011-1016.
- 1094 Charron, A. and Harrison, R. M. (2003). Primary particle formation from vehicle emissions during
  1095 exhaust dilution in the roadside atmosphere. Atmospheric Environment, 37(29), 4109-4119.
- 1097 Charron, A. and Harrison, R. M. (2009). Atmospheric Nanoparticles. Environmental and Human1098 Health Impacts of Nanotechnology, John Wiley and Sons, Ltd, 63-209.
- Chellam, S., Kulkarni, P. and Fraser, M. P. (2005). Emissions of organic compounds and trace
  metals in fine particulate matter from motor vehicles: a tunnel study in Houston, Texas. Journal of
  Air and Waste Management Association, 55, 60-72.

- Chen, J., Zheng, H., Wang, W., Liu, H., Lu, L., Bao, L. and Ren, L. (2006). Resuspension method
  for road surface dust collection and aerodynamic size distribution characterization. China
  Particuology, 4(6), 300-303.
- 1107 Chen, F., Hu, W. and Zhong, Q. (2013). Emissions of particle-phase polycyclic aromatic
  1108 hydrocarbons (PAHs) in the Fu Gui-Shan Tunnel of Nanjing, China. Atmospheric Research, 124,
  1109 53-60.
- 1111 Cheng, Y., Lee, S. C., Ho, K. F., Chow, J. C., Watson, J. G., Louie, P. K. K., Cao, J. J. and Hai, X.
  (2010). Chemically-speciated on-road PM2.5 motor vehicle emission factors in Hong Kong.
  1113 Science of the Total Environment, 408(7), 1621-1627.
- 1114

1124

1129

1132

1135

1139

1143

1147

- Cheng, Y., Zou, S. C., Lee, S. C., Chow, J. C., Ho, K. F., Watson, J. G., Han, Y. M., Zhang, R. J.,
  Zhang, F., Yau, P.S., Huang, Y., Bai, Y. and Wu, W. J. (2011). Characteristics and source
  apportionment of PM1 emissions at a roadside station. Journal of Hazardous Materials, 195, 82-91.
- Chiang, H.-L. and Huang, Y.-S. (2009). Particulate matter emissions from on-road vehicles in afreeway tunnel study. Atmospheric Environment, 43, 4014-4022.
- Chiang, H.-L., Lai, Y.-M. and Chang, S.-Y. (2012). Pollutant constituents of exhaust emitted from
  light-duty diesel vehicles. Atmospheric Environment, 47, 399-406.
- Chirico, R., Prevot A. S. H., DeCarlo, P. F., Heringa, M. F., Richter, R., Weingartner, E. and
  Baltensperger, U. (2011). Aerosol and trace vehicle emission factors measured in a tunnel using an
  Aerosol Mass Spectrometer and other on-line instrumentation. Atmospheric Environment, 45,
  2182-2192.
- Choi, J.-K., Heo, J.-B., Ban, S.-J., Yi, S.-M. and Zoh, K.-D. (2012). Chemical characteristics of
  PM2.5 aerosol in Incheon, Korea. Atmospheric Environment, 60, 583-592.
- 1133 Councell, T. B., Duckenfield, K. U., Landa, E. R. and Callender, E. (2004). Tire wear particles as a
  1134 source of Zn to the environment. Environmental Science and Technology, 38(15), 4206-4214.
- 1136 Crawford, J., Chambers, S., Cohen, D. D., Dyer, L., Wang, T. and Zahorowski, W. (2007).
  1137 Receptor modelling using Positive Matrix Factorisation, back trajectories and Radon-222.
  1138 Atmospheric Environment, 41(32), 6823-6837.
- Cyrys, J., Pitz, M., Heinrich, J., Wichmann, H. E. and Peters, A. (2008). Spatial and temporal
  variation of particle number concentration in Augsburg, Germany. Science of the Total
  Environment, 401(1–3), 168-175.
- Dahl, A., Gharibi, A., Swietlicki, E., Gudmundsson, A., Bohgard, M., Ljungman, A., Blomqvist, G.
  and Gustafsson, M. (2006). Traffic-generated emissions of ultrafine particles from pavement–tire
  interface. Atmospheric Environment, 40(7), 1314-1323.
- Dall'Osto, M., Thorpe, A., Beddows, D. C. S., Harrison, R. M., Barlow, J. F., Dunbar, T., Williams,
  P. I. and Coe, H. (2011). Remarkable dynamics of nanoparticles in the urban atmosphere.
  Atmospheric Physics and Chemistry, 11, 6623-6637.
- 1151
- 1152
- 1153

- Denier van der Gon, H., Gerlofs-Nijland, M.E., Gehrig, R., Gustafsson, M., Janssen, N., Harrison,
  R.M., Hulskotte, J., Johansson, C., Jozwicka, M., Keuken, M., Krijgsheld, K., Ntziachristos, L.,
  Riediker, M. and Cassee, F.R. (2013). The Policy Relevance of Wear Emissions from Road
  Transport, Now and in the Future- An International Workshop Report and Consensus Statement.
- 1158 Journal of the Air and Waste Management Association, 63(2), 136-149.
- 1159

1166

1170

1175

1179

1184

1188

1193

1197

- Dongarra, G., Manno, E. and Varrica, D. (2009). Possible markers of traffic-related emissions.
  Environmental Monitoring and Assessment, 154, 117-125.
- Dreyfus, M. A., Adou, K., Zucker, S. M. and Johnston, M. V. (2009). Organic aerosol source
  apportionment from highly time-resolved molecular composition measurements. Atmospheric
  Environment, 43(18), 2901-2910.
- Duong, T. T. and Lee, B. K. (2011). Determining contamination level of heavy metals in road dust
  from busy traffic areas with different characteristics. Journal of Environmental Management, 92(3),
  554-562.
- El Haddad, I., Marchand, N., Dron, J., Temime-Roussel, B., Quivet, E., Wortham, H., Jaffrezo, J.L.,
  Baduel, C., Voisin, D., Besombes, J.L. and Gille, G. (2009). Comprehensive primary particulate
  organic characterization of vehicular exhaust emissions in France. Atmospheric Environment,
  43(39), 6190-6198.
- El Haddad, I., Marchand, N., Wortham, H., Piot, C., Besombes, J.-L., Cozic, J., Chauvel, C.,
  Armengaud, A., Robin, D. and Jaffrezo, J.-L. (2011). Primary sources of PM2.5 organic aerosol in
  an industrial Mediterranean city, Marseille. Atmospheric Chemistry and Physics, 11, 2039- 2058.
- Etyemezian, V., Kuhns, H., Gillies, J., Chow, J., Hendrickson, K., McGown, M. and Pitchford, M.
  (2003). Vehicle-based road dust emissions measurement (III): effect of speed, traffic volume,
  location, and season on PM10 road dust emissions in the Treasure Valley, ID. Atmospheric
  Environment, 37, 4583-4593.
- Fabretti, J.-F., Sauret, N., Gal, J.-F., Maria, P.-C. and Schärer, U. (2009). Elemental characterization
  and source identification of PM2.5 using Positive Matrix Factorization: The Malraux road tunnel,
  Nice, France. Atmospheric Research, 94(2), 320-329.
- Fan, Y., Meng, Q., Weisel, C., Shalat, S.,Laumbach, R., Ohman-Strickland, P., Black, K.,
  Rodriguez, M. and Bonanno, L. (2006). Acute Short-Term Exposures to PM2.5 Generated by
  Vehicular Emissions and Cardiopulmonary Effects in Older Adults. Epidemiology, 17(6), S213S214.
- Fang, G.-C., Chang, C.-N., Wu, Y.-S., Fu, P. P.-C., Yan, I.-L. and Chen, M.-H. (2004).
  Characterization, identification of ambient air and road dust polycyclic aromatic hydrocarbons in
  central Taiwan, Taichung. Science of the Total Environment, 327, 135-146.
- Franco, V., Kousoulidou, M., Muntean, M., Ntziachristos, L., Hausberger, S. and Dilara, P. (2013).
  Road vehicle emission factors development: a review. Atmospheric Environment, 70, 84-97.
- Fraser, M.P., Cass, G.R. and Simoneit, B.R.T. (1998). Gas-Phase and Particle-Phase Organic
  Compounds Emitted From Motor Vehicle Traffic in a Los Angeles Roadway Tunnel.
  Environmental Science and Technology, 32, 2051-2060.
- 1204

- Fraser, M.P., Buzcu, B., Yue, Z.W., Mcgaughey, G.R., Desai, N.R., Allen, D.T., Seila, R.L.,
  Lonneman, W.A. and Harley, R.A.(2003). Separation of Fine Particulate Matter Emitted from
  Gasoline and Diesel Vehicles Using Chemical Mass Balancing Techniques. Environmental Science
  and Technology, 37, 3904-3909.
- Fujita, E. M., Campbell, D. E. and Zielinska, B. (2006). Chemical analysis of lubrication oil
  samples from a study to characterize exhaust emissions from light-duty gasoline vehicles in the
  Kansas City metropolitan area (Final Report ).
- Fujita, E. M., Campbell, D. E., Arnott, W. P., Chow, J. C. and Zielinska, B. (2007). Evaluations of
  the Chemical Mass Balance Method for Determining Contributions of Gasoline and Diesel Exhaust
  to Ambient Carbonaceous Aerosols. Journal of Air and Waste Management Association, 57, 721740.
- 1218

1230

1233

1238

1243

1246

1251

- Fuller, G. W., Carslaw, D. C. and Lodge, , H. W. (2002). An empirical approach for the prediction
  of daily mean PM10 concentrations. Atmospheric Environment, 36, 1431-1441.
- Fulper, C.R., Kishan, S., Baldauf, R.W., Sabisch, R., Warila, J., Fujita, E.M., Scarbro, C., Crews,
  W.S., Snow, R., Gabele, P., Sanots, R., Tierney, E. and Cantrell, B. (2010). Methods of
  Characterizing the Distribution of Exhaust Emissions from Light-Duty, Gasoline-Powered Motor
  Vehicles in the U.S. Fleet. Journal of Air and Waste Management Association, 60, 1376-1387.
- Funasaka, K., Miyazaki, T., Kawaraya, T., Tsuruho, K. and Mizuno, T. (1998). Characteristics of
  particulates and gaseous pollutants in a highway tunnel. Environmental Pollution, 102(2–3), 171176.
- Furusjo, E., Sternbeck, J. and Cousins, A.P. (2007). PM10 source characterization at urban and
  highway roadside locations. Science of the Total Environment, 387, 206-219.
- Fushimi, A., Hasegawa, S., Takahashi, K., Fujitani, Y., Tanabe, K. and Kobayashi, S. (2008).
  Atmospheric fate of nuclei-mode particles estimated from the number concentrations and chemical
  composition of particles measured at roadside and background sites. Atmospheric Environment,
  42(5), 949-959.
- Gao, B., Yu, J.-Z., Li, S.-X., Ding, X., He, Q.-F. and Wang, X.-M. (2011). Roadside and rooftop
  measurements of polycyclic aromatic hydrocarbons in PM2.5 in urban Guangzhou: Evaluation of
  vehicular and regional combustion source contributions. Atmospheric Environment, 45(39), 71847191.
- Garg, B. D., Cadle, S. H., Mulawa, P. A., Groblicki, P. J., Laroo, C. and Parr, G. A. (2000). Brake
  Wear Particulate Matter Emissions. Environmental Science and Technology, 34(21), 4463-4469.
- Gehrig, R., Hill, M., Buchmann, B., Imhof, D., Weingartner, E. and Baltensperger, U. (2004).
  Separate determination of PM10 emission factors of road traffic for tailpipe emissions and
  emissions from abrasion and resuspension processes. International Journal of Environment and
  Pollution, 22(3): 312-325.
- Gehrig, M., Zeyer, K., Bukowiecki, N., Lienemann, P., Poulikakos, L.D., Furger, M. and
  Buchmann, B. (2010). Mobile load simulators- A tool to distinguish between the emissions due to
  abrasion and resuspension of PM10 from road surfaces. Atmospheric Environment, 44, 4937-4943.

- Gertler, A.W. and Pierson, W.R. (1996). Recent measurements of mobile source emission factors in
  North American tunnels. The Science of the Total Environment, 189/190, 107-113.
- Gertler, A.W., Gillies, J.A., Pierson, W.R., Rogers, C.F., Sagebiel, J.C., Abu-Allaban, M.,
  Coulombe, W., Tarnay, L. and Cahill, T.A. (2002). Real-World Particulate Matter and Gaseous
  Emissions from Motor Vehicles in a Highway Tunnel. Research Report 107, Health Effects
  Institute.
- 1263

1279

1283

1288

1292

1296

1258

- Gertler, A., Kuhns, H., Abu-Allaban, M., Damm, C., Gilies, J., Etyemezian, V., Clayton, R. and
  Proffitt, D. (2006). A case study of the impact of winter road sand/salt and street sweeping on road
  dust re-entrainment. Atmospheric Environment, 40, 5976-5985.
- Gietl, J. K., Lawrence, R., Thorpe, A. J. and Harrison, R. M. (2010). Identification of brake wear
  particles and derivation of a quantitative tracer for brake dust at a major road. Atmospheric
  Environment, 44(2), 141-146.
- Gillies, J.A., Gertler, A.W., Sagebiel, J.C. and Dippel, W.A. (2001). On-Road Particulate Matter
  (PM2.5 and PM10) Emissions in the Sepulveda Tunnel, Los Angeles, California. Environmental
  Science and Technology, 35, 1054-1063.
- Grieshop, A. P., Lipsky, E. M., Pekney, N. J., Takahama, S. and Robinson, A. L. (2006). Fine
  particle emission factors from vehicles in a highway tunnel: Effects of fleet composition and season.
  Atmospheric Environment, 40, 287-298.
- 1280 Gunawardana, C., Goonetilleke, A., Egodawatta, P., Dawes, L. and Kokot, S. (2011). Source
  1281 characterisation of road dust based on chemical and mineralogical composition. Chemosphere,
  1282 87(2), 163-170.
- Gustafsson, M., Blomqvist, G., Gudmundsson, A., Dahl, A., Swietlicki, E., Bohgard, M., Lindbom,
  J. and Ljungman, A. (2008). Properties and toxicological effects of particles from the interaction
  between tyres, road pavement and winter traction material. The Science of the Total Environment,
  393(2-3), 226-240.
- Gustafsson, M., Blomqvist, A., Gudmundsson, A., Dahl, A., Jonsson, P. and Swietlicki, E. (2009).
  Factors influencing PM10 emissions from road pavement wear. Atmospheric Environment, 49, 226-240.
- Han, L., Zhuang, G., Cheng, S., Wang, Y. and Li, J. (2007). Characteristics of re-suspended road
  dust and its impact on the atmospheric environment in Beijing. Atmospheric Environment, 41(35),
  7485-7499.
- Han, B., Bai, Z., Guo, G., Wang, F., Li, F., Liu, Q., Ji, Y., Li, X. and Hu, Y. (2009).
  Characterization of PM(10) fraction of road dust for polycyclic aromatic hydrocarbons (PAHs)
  from Anshan, China. Journal of hazardous materials, 170(2-3), 934-940.
- Handler, M., Puls, C., Zbiral, J., Marr, I., Puxbaum, H. and Limbeck, A. (2008). Size and
  composition of particulate emissions from motor vehicles in the Kaisermühlen-Tunnel, Vienna.
  Atmospheric Environment, 42(9), 2173-2186.
- Hanedar, A., Alp, K., Kaynak, B., Baek, J., Avsar, E. and Odman, M.T. (2008). Concentrations and
  sources of PAHs at three stations in Istanbul, Turkey. Atmospheric Research, 99, 391-399.
- 1307

- Harris, S. J. and Maricq, M. M. (2001). Signature size distributions for diesel and gasoline engine
  exhaust particulate matter. Journal of Aerosol Science, 32(6), 749-764.
- Harrison, R. M., Deacon A. R. and Jones, M. R. (1997). Sources and processes affecting
  concentrations of PM10 and PM2.5 Particulate Matter in Birmingham (U.K.). Atmospheric
  Environment, 31, 4103-4117.
- 1314
- 1315 Harrison, R.M., Shi, J.P., Xi, S., Khan, A., Mark, D., Kinnersley, R. and Yin, J. (2000).
- Measurement of number, mass and size distribution of particles in the atmosphere, Phil. Trans. R.Soc. Lond., A, 358, 2567-2580.
- 1318

1326

1330

- Harrison, R. M., Tilling, R., Romero, M.S.C., Harrad, S. and Jarvis, K. (2003). A study of trace
  metals and polycyclic aromatic hydrocarbons in the roadside environment. Atmospheric
  Environment, 37, 2391-2402.
- Harrison, R. M., Jones, A. M. and Barrowcliffe, R. (2004). Field study of the influence of
  meteorological factors and traffic volumes upon suspended particle mass at urban roadside sites of
  different geometry. Atmospheric Environment, 38, 6361–6369.
- Harrison, R. M. (2009). Airborne Particulate Matter from Road Traffic: Current Status of
  Knowledge and Research Challenges Paper presented at the ETTAO09. In 17th Transport and Air
  Pollution Symposium 3rd Environment and Transport Symposium.
- Harrison, R. M., Giorio, C., Beddows, D. C. and Dall'Osto, M. (2010). Size distribution of airborne
  particles controls outcomes of epidemiological studies. Science of the Total Environment, 409,
  289-293.
- Harrison, R.M., Beddows, D.C.S. and Dall'Osto, M., (2011). PMF analysis of wide-range particle
  size spectra collected on a major highway. Environmental Science and Technology, 45, 5522-5528.
- Harrison, R. M., Dall'Osto, M., Beddows, D. C. S., Thorpe, A. J., Bloss, W. J., Allan, J. D., Coe, H.,
  Dorsey, J. R., Gallagher, M., Martin, C., Whitehead, J., Williams, P. I., Jones, R. L., Langridge, J. M.,
  Benton, A. K., Ball, S. M., Langford, B., Hewitt, C. N., Davison, B., Martin, D., Petersson, K.,
  Henshaw, S. J., White, I. R., Shallcross, D. E., Barlow, J. F., Dunbar, T., Davies, F., Nemitz, E.,
  Phillips, G. J., Helfter, C., Di Marco, C. F. and Smith, S. (2012a). Atmospheric chemistry and
  physics in the atmosphere of a developed megacity (London): An overview of the REPARTEE
  experiment and its conclusions. Atmospheric Chemistry and Physics, 12, 3065-3114.
- Harrison, R. M., Jones, A. M., Gietl, J., Yin, J. and Green, D. C. (2012b). Estimation of the
  contributions of brake dust, tire wear, and resuspension to nonexhaust traffic particles derived from
- 1348 atmospheric measurements. Environmental Science and Technology, 46, 6523-6529.
- Hassanien, M. A. and Abdel-Latif, N. M. (2008). Polycyclic aromatic hydrocarbons in road dust
  over Greater Cairo, Egypt. Journal of Hazardous Materials, 151, 247-254.
- 1351
- Hays, M. D., Cho, S.-H., Baldauf, R., Schauer, J. J. and Shafer, M. (2011). Particle size
  distributions of metal and non-metal elements in an urban near-highway environment. Atmospheric
  Environment, 45(4), 925-934.
- He, L.-Y., Hu, M., Huang, X.-F., Zhang, Y.-H., Yu, B.-D. and Liu, D.-Q. (2006). Chemical
  characterization of fine particles from on-road vehicles in the Wutong tunnel in Shenzhen, China.
  Chemosphere, 62(12), 1565-1573.
- 1359

- He, L.-Y., Hu, M., Zhang, Y.-H., Huang, X.-F. and Yao, T.-T. (2008). Fine Particle Emissions from
  On-Road Vehicles in the Zhujiang Tunnel, China. Environmental Science and Technology, 42(12),
  4461-4466.
- 1363 HEI Panel on the Health Effects of Traffic-Related Air Pollution (2010). Traffic-Related Air
- 1364 Pollution: A Critical Review of the Literature on Emissions, Exposure, and Health Effects. HEI
- 1365 Special Report, 17, Health Effects Institute, Boston, MA.
- 1366
- Hien, T. T., Thanh, L. T., Kameda, T., Takenaka, N. and Bandow, H. (2007). Distribution
  characteristics of polycyclic aromatic hydrocarbons with particle size in urban aerosols at the
  roadside in Ho Chi Minh City, Vietnam. Atmospheric Environment, 41(8), 1575-1586.
- 1370

1381

1385

1389

1393

1398

Hildemann, L. M., Markowski, G. R., Jones, M. C. and Cass, G. R. (1991). Submicrometer aerosol
mass distributions of emissions from boilers, fireplaces, automobiles, diesel trucks and meat
cooking operatios. Aerosol Science and Technology, 14(1), 138-152.

- Hjortenkrans, D. S. T., Bergbäck, B. G. and Häggerud, A. V. (2007). Metal Emissions from Brake
  Linings and Tires: Case Studies of Stockholm, Sweden 1995/1998 and 2005. Environmental
  Science and Technology, 41(15), 5224-5230.
- Hopke P. K. (2003). Recent developments in receptor modelling. Journal of Chemometrics, 17, 255-265.
- Huang, X.-F., Chen, D.-L., Lan, Z.-J., Feng, N., He, L.-Y., Yu, G.-H. and Luan, S.-J. (2012).
  Characterization of organic aerosol in fine particles in a mega-city of South China: Molecular composition, seasonal variation, and size distribution. Atmospheric Research 114-115, 28-37.
- Hussein, T., Johansson, C., Karlsson, H. and Hansson, H.-C. (2008). Factors affecting non-tailpipe
  aerosol particle emissions from paved roads: On-road measurements in Stockholm, Sweden.
  Atmospheric Environment, 42(4), 688-702.
- Iijima, A., Sato, K., Yano, K., Tago, H., Kato, M., Kimura, H. and Furuta, N. (2007). Particle size
  and composition distribution analysis of automotive brake abrasion dusts for the evaluation of
  antimony sources of airborne particulate matter. Atmospheric Environment, 41(23), 4908-4919.
- Imhof, D., Weingartner, E., Prevot, A.S.H., Ordonez, C., Kurtenbach, R., Weisen, P., Rodler, J.,
  Sturm, P., McCrae, I., Sjodin, A. and Baltensperger, U. (2005a). Aerosol and NOx emissions
  factors and submicron particle size distributions in two road tunnels with different traffic regimes.
  Atmospheric Chemistry and Physics Discussions, 5, 5127-5166.
- Imhof, D., Weingartner, E., Ordonez, C., Gehrig, R., Hill, M., Buchmann, B. and Baltensperger, U.
  (2005b). Real-world emission factors of fine and ultrafine aerosol particles for different traffic
  situations in Switzerland. Environmental Science and Technology, 39(21), 5127-5166.
- 1402
  1403 Ingo, G. M., D'Uffizi, M., Falso, G., Bultrini, G. and Padeletti, G. (2004). Thermal and
  1404 microchemical investigation of automotive brake pad wear residues. Thermochimica Acta, 418, 611405 68.
  - Jamriska, M., Morawska, L., Thomas, S. and He, C. (2004). Diesel Bus Emissions Measured in A
    Tunnel Study. Environmental Science and Technology, 38, 6701-6709.
  - 1409 1410

- 1411 Jayaratne, E. R., Wang, L., Heuff, D., Morawska, L. and Ferreiera, L. (2009). Increase in particle
- 1412 number emissions from motor vehicles due to interruption of steady traffic flow. Transportation
- 1413 Research Part D, 14, 521-526.
- 1414

1421

1425

1429

1445

1450

1455

- Johansson, C., Norman, M. and Burman, L. (2009). Road traffic emission factors for heavy metals.
  Atmospheric Environment, 43(31), 4681-4688.
- Jones, A. M. and Harrison, R. M. (2006). Estimation of emisison factors of particle number and
  mass fractions from traffic at a site where mean vehicle speeds vary over short distances.
  Atmospheric Environment, 35, 7125-7137.
- Jones, A. M., Yin, J. and Harrison, R. M. (2008). The weekday-weekend difference and the
  estimation of the non-vehicle contributions to the urban increment of airborne particulate matter.
  Atmospheric Environment, 42, 4467-4479.
- Kam, W., Liacos, J. W., Schauer, J. J., Delfino, R. J. and Sioutas, C. (2012). Size-segregated
  composition of particulate matter (PM) in major roadways and surface streets. Atmospheric
  Environment, 55, 90-97.
- Karanasiou, A., Moreno, T., Amato, F., Lumbreras, J., Narros, A., Borge, R., Tobias, A., Boldo, E.,
  Linares, C., Pey, J., Reche, C., Alastuey, A. and Querol, X. (2011). Road dust contribution to PM
  levels Evaluation of the effectiveness of street washing activities by means of Positive Matrix
  Factorization. Atmospheric Environment, 45(13), 2193-2201.
- Karanasiou, A., Moreno, T., Amato, F., Tobias, A., Boldo, E., Linares, C., Lumbreras, J., Borge, R.,
  Alastuey, A. and Querol, X. (2012). Variation of PM2.5 concentrations in relation to street washing
  activities. Atmospheric Environment, 54, 465- 469.
- Kaunhaniemi, M., Kukkonen, J., Harkonen, J., Nikmo, J., Kangas, L., Omstedt, G., Ketzel, M.,
  Kousa, A., Haakana, M. and Karppinen, A. (2011). Evaluation of a road dust suspension model for
  predicting the concentrations of PM10 in a street canyon. Atmospheric Environment, 45, 3646-3654.
- Kelly, F.J. and Fussell, J.C. (2012). Size, source and chemical composition as determinats of
  toxicity attributable to ambient particulate matter. Atmospheric Environment, 60, 504- 526.
- Kerminen, V.-M., Pakkanen, T. A., Mäkelä, T., Hillamo, R. E., Sillanpää, M., Rönkkö, T., Virtanen,
  A., Keskinen, J., Pirjola, L., Hussein, T. and Hameri, K. (2007). Development of particle number
  size distribution near a major road in Helsinki during an episodic inversion situation. Atmospheric
  Environment, 41(8), 1759-1767.
- Ketzel, M.G., Omstedt, G., Johansson, C., During, I., Pohjola, M., Oettl, D., Gidhagen, L., Wahlin,
  P., Lohmeyer, A. and Berkowicz (2007). Estimation and validation of PM2.5/PM10 exhaust and
  non-exhaust emission factors for practical street pollution modelling. Atmospheric Environment, 41,
  9370-9385.
- Keuken, M., Denier van der Gon, H. and van der Valk, K. (2010). Non-exhaust emissions of PM
  and the efficiency of emission reduction by road sweeping and washing in the Netherlands. The
  Science of the Total Environment, 408(20), 4591-4599.
- Keuken, M., Henzing, B., Zandveld, P., van den Elshout, S. and Karl, M. (2012). Dispersion of
  particle numbers and elemental carbon from road traffic, a harbour and an airstrip in the
  Netherlands. Atmospheric Environment, 54, 320-327.

1469

1473

1476

1480

1483

1487

1491

1495

1503

- 1464 Kim, J. Y., Lee, J. Y., Kim, Y. P., Lee, S. B., Jin, H. C. and Bae, G. N. (2012). Seasonal
  1465 characteristics of the gaseous and particulate PAHs at a roadside station in Seoul, Korea.
  1466 Atmospheric Research, 116, 142-150.
- 1467 Kittleson, D. B., Watts, W. F. and Johnson, J.P. (2004). Nanoparticle emisisons on Minnesota1468 highways. Atmospheric Environment, 38, 9-19.
- 1470 Klems, J. P., Pennington, M. R., Zordan, C. A. and Johnston, M. V. (2010). Ultrafine particles near
  1471 a roadway intersection: Origin and apportionment of fast changes in concentration. Environmental
  1472 Science and Technology, 44(20), 7903-7907.
- Kuhlbusch, T.A.J., John, A.C. and Quass, U. (2009). Sources and source contributions to fineparticles. Biomarkers, 14(1), 23-28.
- 1477 Kumar, P., Fennell, P., Langley, D. and Britter, R. (2008). Pseudo-simultaneous measurements for
  1478 the vertical variation of coarse, fine and ultrafine particles in an urban street canyon. Atmospheric
  1479 Environment, 42(18), 4304-4319.
- Kumar, P., Pirjola, L., Ketzel, M. and Harrison, R.M. (2013). Nanoparticle emissions from 11 nonvehicle exhaust sources- A review. Atmospheric Environment, 67, 252-277.
- Kumata, H., Yamada, J., Masuda, K., Takada, H., Sato, Y., Sakural, T. and Fujiwara, K. (2002).
  Benzothiazolamines as tire-derived molecular markers: sorptive behavious in street runoff and application to source apportioning. Environmental Science and Technology, 36(4), 702-708.
- Kupiainen, K. J., Tervahattu, H., Räisänen, M., Mäkelä, T., Aurela, M. and Hillamo, R. (2005). Size
  and Composition of Airborne Particles from Pavement Wear, Tires, and Traction Sanding.
  Environmental Science and Technology, 39(3), 699-706.
- 1492 Kwon, E.E. and Castaldi, M.J. (2012). Mechanistic understanding of polycyclic aromatic
  1493 hydrocarbons (PAHs) from the therman degradation of tires under various oxygen concentration
  1494 atmospheres. Environmental Science and Technology, 46(23), 12921-6.
- Laidlaw, M. A. S., Zahran, S., Mielke, H. W., Taylor, M. P. and Filippelli, G. M. (2012). Resuspension of lead contaminated urban soil as a dominant source of atmospheric lead in
  Birmingham, Chicago, Detroit and Pittsburgh, USA. Atmospheric Environment, 49, 302-310.
- Landis, M. S., Lewis, C. W., Stevens, R. K., Keeler, G. J., Dvonch, J. T. and Tremblay, R. T.
  (2007). Ft. McHenry tunnel study: Source profiles and mercury emissions from diesel and gasoline
  powered vehicles. Atmospheric Environment, 41(38), 8711-8724.
- Larsen, R.K. and Baker, J.E. (2003). Source apportionment of polycyclic aromatic hydrocarbons in
  the urban atmosphere: A comparison of three methods. Environmental Science and Technology,
  37(9), 1873-1881.
- Lee, W.-J., Wang, Y.-F., Lin, T.-C., Chen, Y.-Y., Lin, W.-C., Ku, C.-C. and Cheng, J.-T. (1995).
  PAH characteristics in the ambient air of traffic-source. The Science of the Total Environment, 159, 185-200.
- 1511
- Liacos, J. W., Kam, W., Delfino, R. J., Schauer, J. J. and Sioutas, C. (2012). Characterization of organic, metal and trace element PM2.5 species and derivation of freeway-based emission rates in Los Angeles, CA. Science of the total Environment, 435-436, 5283-5293.

1516 Lim, L.H., Harrison, R.M. and Harrad, S. (1999). The contribution of traffic to atmospheric concentrations of polyaromatic hydrocarbons. Environmental Science and Technology, 33, 3538-1517 1518 3542. Lim, J., Yu, L.E., Kostetski, Yu. Yu., Lim, C., Ryu, J. and Kim, J. (2008). Effects of driving 1519 1520 conditions on diesel exhaust particles. Journal of Air and Waste Management Association, 58, 1521 1077-1085. 1522 Lin, C.-C., Chen, S.-J., Huang, K.-L., Hwang, W.-I., Chang-Chien, G.-P. and Lin, W.-Y. (2005). 1523 1524 Characteristics of Metals in Nano/Ultrafine/Fine/Coarse Particles Collected Beside a Heavily Trafficked Road. Environmental Science and Technology, 39(21), 8113-8122. 1525 1526 1527 Lin, H.-Y., Hung, C.-H., Yuan, C.-S., Chen, C.-J., Chen, W.-C., Chiang, S.-W. and Wang, C., J.-L. 1528 (2008). Characteristics and source identification of roadside suspended particles. Transportation 1529 Research Part D: Transport and Environment, 13(8), 531-538. 1530 1531 Lingard, J. J. N., Agus, E. L., Young, D. T., Andrews, G. E. and Tomlin, A. S. (2006). Observations 1532 of urban airborne particle number concentrations during rush-hour conditions: analysis of the 1533 number based size distributions and modal parameters. Journal of Environmental Monitoring, 8(12), 1534 1203-1218. 1535 Liu, M., Cheng, S.B., Ou, D.N., Hou, L.J., Gao, L., Wang, L.L., Xie, Y.S., Yang, Y., Xu, S.Y. 1536 1537 (2007). Characterization, identification of road dust PAHs in central Shanghai areas, China. 1538 Atmospheric Environment, 41, 8785-8795. 1539 1540 Lough, G. C., Schauer, J. J., Park, J.-S., Shafer, M. M., DeMinter, J. T. and Weinstein, J. P. (2005). 1541 Emissions of Metals Associated with Motor Vehicle Roadways. Environmental Science and 1542 Technology, 39(3), 826-836. 1543 1544 Lough, G. C., Christensen, C. G., Schauer, J. J., Tortorelli, J., Mani, E., Lawson, D. R., Clark, N. N. 1545 and Gabela, P. A. (2007). Development of molecular marker source profiles for emissions from on-1546 road gasoline and diesel vehicle fleets. Journal of the Air and Waste Management Association, 57(10), 1190-1199. 1547 1548 1549 Luo, X. S., Yu, S., Zhu, Y. G. and Li, X. D. (2011). Trace metal contamination in urban soils of 1550 China. The Science of the Total Environment, 421-422, 17-30. 1551 1552 Majumdar, D., Rajaram, B., Meshram, S. and Chalapati Rao, C.V. (2012). PAHs in Road Dust: Ubiquity, Fate, and Summary of Available Data. Critical Reviews in Environmental Science and 1553 Technology, 42(12), 1191-1232. 1554 1555 Mancilla, Y. and Mendoza, A. (2012). A tunnel study to characterize PM2.5 emissions from 1556 1557 Gasoline-powered vehicles in Monterrey, Mexico. Atmospheric Environment, 59, 449-460. 1558 1559 Manno, E., Varrica, D. and Dongarrà, G. (2006). Metal distribution in road dust samples collected 1560 in an urban area close to a petrochemical plant at Gela, Sicily, Atmospheric Environment, 40(30), 1561 5929-5941. 1562 1563 Mari, M., Harrison, R.M., Schuhmacher, M., Domingo, J.L. and Pongpiachan S. (2010). Inferences 1564 over the sources and processes affecting polycyclic aromatic hydrocarbons in the atmosphere derived 1565 from measured data. Science of the Total Environment, 408, 2387-2393. 1566 51

- Martuzevicius, D., Kliucininkas, L., Prasauskas, T., Krugly, E., Kauneliene, V. and Strandberg, B.
  (2011). Resuspension of particulate matter and PAHs from street dust. Atmospheric Environment,
  45(2), 310-317.
- Mathissen, M., Scheer, V., Vogt, R. and Benter, T. (2011). Investigation on the potential generation
  of ultrafine particles from the tire–road interface. Atmospheric Environment, 45(34), 6172-6179.
- Maricq, M. M. (2007). Chemical characterization of particulate emissions from diesel engines: A
   review. Journal of Aerosol Science, 38(11), 1079-1118.
- Masiol, M., Hofer, A., Squizzato, S., Piazza, R., Rampazzo, G. and Pavoni, B. (2012). Carcinogenic
  and mutagenic risk associated to airborne particle-phase polycyclic aromatic hydrocarbons: A
  source apportionment. Atmospheric Environment, 60, 375- 382.
- Mauderly, J.L. (1994). Toxicological and epidemiological evidence for health risks from inhaled
  engine emissions. Environmental Health Perspectives, 102, 165-171.
- Mclaren, R., Gertler, A.W., Wittorff, D.N., Belzer, W., Dann, T. and Singleton, D.L. (1996). RealWorld Measurements of Exhaust and Evaporative Emissions in the Cassiar Tunnel Predicted by
  Chemical Mass Balance Modeling. Environmental Science and Technology, 30, 3001-3009.
- Miguel, A. H., Kirchstetter, T. W., Harley, R. A. and Hering, S. V. (1998). On-Road Emissions of
  Particulate Polycyclic Aromatic Hydrocarbons and Black Carbon from Gasoline and Diesel
  Vehicles. Environmental Science and Technology, 32(4), 450-455.
- Milani, M., Pucillo, F.P., Ballerini, M., Camatani, M., Gualtieri, M. and Martino, S. (2004). First
  evidence of tyre debris characterization at the nanoscale by focused ion beam. Materials
  Characterization, 52, 283-288.
- Morawska, L., Jayaratne, E.R., Mengersen, K., Jamriska, M. and Thomas, S. (2002). Differences in
  airborne particle and gaseous concentrations in urban air between weekdays and weekends.
  Atmospheric Environment, 36, 4375-4383.
- 1600 Nanzetta, K. M. and Holmen, B. A. (2004). Roadside Particle Number Distributions and
  1601 Relationships between Number Concentrations, Meteorology, and Traffic along a Northern
  1602 California Freeway. Journal of Air and Waste Management Association, 54, 540-554.
- Narvaez, R. F., Hoepner, L., Chillrud, S. N., Yan, B., Garfinkel, R., Whyatt, R., Camann, D., Perera,
  F. P., Kinney, P. L. and Miller, R. L. (2008). Spatial and Temporal Trends of Polycyclic Aromatic
  Hydrocarbons and Other Traffic-Related Airborne Pollutants in New York City. Environmental
  Science and Technology, 42, 7330-7335.
- 1609 Nicholson, K.W. (1988). A review of particle resuspension. Atmospheric Environment (1967),
  1610 22(12), 2639-2651.
- 1612 Nielsen, T. (1996). Traffic contribution of polycyclic aromatic hydrocarbons in the centre of a large1613 city. Atmospheric Environment, 30(20), 3481-3490.
- 1615 Ntziachristos, L., Ning, Z., Geller, M.D., Sheesley, R.J., Schauer, J.J. and Sioutas, C. (2007). Fine,
  1616 ultrafine and nanoparticle trace element compositions near a major freeway with a high heavy-duty
  1617 diesel fraction. Atmospheric Environment, 41, 5864-5696.
- 1618

1576

1580

1583

1587

1591

1595

1599

1603

1608

1611

- Oliveira, C., Pio, C., Caseiro, A., Santos, P., Nunes, T., Mao, H., Luahana, L. and Sokhi, R.S.
  (2010). Road traffic impact on urban atmospheric aerosol loading at Oporto, Portugal. Atmospheric
  Environment, 44(26), 3147-3158.
- 1622

1630

1633

1637

1641

1645

1648

1653

1658

1663

- Oliveira, C., Martins, N., Tavares, J., Pio, C., Cerqueira, M., Matos, M., Silva, H., Oliviera, S. and
  Camoes, F. (2011). Size distribution of polycyclic aromatic hydrocarbons in a roadway tunnel in
  Lisbon, Portugal. Chemosphere, 83(11), 1588-1596.
- 1627 Omar, N. Y. M. J., Abas, M. R. B., Rahman, N. A., Tahir, N. M., Rushdi, A. I. and Simoneit, B. R.
  1628 T. (2007). Levels and distributions of organic source tracers in air and roadside dust particles of
  1629 Kuala Lumpur, Malaysia. Environmental Geology, 52, 1485-1500.
- 1631 Omstedt, G., Bringfelt, B., Johansson, C. (2005). A model for vehicle-induced non-tailpipe
  1632 emissions of particles along Swedish roads. Atmospheric Environment, 39, 6088-6097.
- Oanh, N. T. K., Thiansathit, W., Bond, T. C., Subramanian, R., Winijkul, E. and Paw-armart, I.
  (2010). Compositional characterization of PM2.5 emitted from in-use diesel vehicles. Atmospheric
  Environment, 44(1), 15-22.
- 1638 Ondráček, J., Schwarz, J., Ždímal, V., Andělová, L., Vodička, P., Bízek, V., Tsai, C.-J., Chen, S.-C.
  1639 and Smolik, J. (2011). Contribution of the road traffic to air pollution in the Prague city (busy
  1640 speedway and suburban crossroads). Atmospheric Environment, 45(29), 5090-5100.
- Padro-Martinez, L. T., Patton, A. P., Trull, J. B., Zamore, W., Brugge, D. and Durant, J. L. (2012).
  Mobile monitoring of particle number concentration and other traffic-related air pollutants in a near-highway neighbourhood over the course of a year. Atmospheric Environment, 61, 253-264.
- Pant, P. and Harrison, R. M. (2012). Critical review of receptor modelling for particulate matter: Acase study of India, Atmospheric Environment, 49, 1-12.
- Peltier, R. E., Cromar, K. R., Ma, Y., Fan, Z.-H. and Lippmann, M. (2011). Spatial and seasonal
  distribution of aerosol chemical components in New York City: (2) Road dust and other tracers of
  traffic-generated air pollution. Journal of Exposure Science and Environmental Epidemiology,
  21(5), 484-494.
- Perez, N., Pey, J., Cusack, M., Reche, C., Querol, X., Alastuey, A. and Viana, M. (2010).
  Variability of Particle Number, Black Carbon and PM10, PM2.5 and PM1 levels and speciation:
  Influence of road traffic emissions on urban air quality. Aerosol Science and Technology, 44(7), 487-499.
- Perrone, M. G., Larsen, B. R., Ferrero, L., Sangiorgi, G., De Gennaro, G., Udisti, R., Zangranado,
  R., Gambaro, A. and Bolzacchini, E. (2012). Sources of high PM2.5 concentrations in Milan,
  Northern Italy: Molecular marker data and CMB modelling. Science of the Total Environment, 414,
  343-355.
- Pey, J., Querol, X., Alastuey, A., Rodriguez, R., Putuad, J.-P. and Van Dingenen, R. (2009).
  Source apportionemnt of urban fine and ultra-fine particle number concentration in a Western
  Mediterranean city. Atmospheric Environment, 43, 4407-4415.
- Pey, J., Querol, X. and Alastuey, A. (2010). Discriminating the regional and urban contributions in
  the North-Western Mediterranean: PM levels and composition. Atmospheric Environment, 44(13),
  1587-1596.

- Phuleria, H. C., Geller, M. D., Fine, P. M. and Sioutas, C. (2006). Size-Resolved Emissions of
  Organic Tracers from Light- and Heavy-Duty Vehicles Measured in a California Roadway Tunnel.
  Environmental Science and Technology, 40(13), 4109-4118.
- Phuleria, H. C., Sheesley, R. J., Schauer, J. J., Fine, P. M. and Sioutas, C. (2007). Roadside
  measurements of size-segregated particulate organic compounds near gasoline and dieseldominated freeways in Los Angeles, CA. Atmospheric Environment, 41(22), 4653-4671.
- 1677
  1678 Prichard, H.M. and Fisher, P.C. (2012). Identification of Platinum and Palladium Particles Emitted
  1679 from Vehicles and Dispersed into the Surface Environment. Environmental Science and
  1680 Technology, 46(6), 3149-3154.
- Pulles, T., Denier van der Gon, H., Appelman, W. and Verheul, M. (2012). Emission factors for
  heavy metals from diesel and petrol used in European vehicles. Atmospheric Environment, 61, 641651.
- 1684 1685 Puustinen
- Puustinen, A., Hämeri, K., Pekkanen, J., Kulmala, M., de Hartog, J., Meliefste, K., ten Brink, H.,
  Kos, G., Katsouyanni, K., Karakatsani, A., Kotronarou, A., Kavouras, I., Meddings, C., Thomas, S.,
  Harrison, R., Ayres, J.G., van der Zee, S. and Hok, G. (2007). Spatial variation of particle number
  and mass over four European cities. Atmospheric Environment, 41(31), 6622-6636.
- Reche, C., Querol, X., Alastuey, A., Viana, M., Pey, J., Moreno, T., Rodriguez, S., Gonzalez, Y.,
  Fernandez-Camacho, R., Sanchez de la Campa, A. M., de la Rosa, J., Dall'Osto, M., Prevot, A. S.
  H., Hueglin, C., Harrison, R.M. and Quincey, P. (2011). New considerations for PM, black carbon
  and particle number concentration for air quality monitoring across different European cities.
  Atmospheric Chemistry and Physics, 11, 6207-6227.
- 1696 Reddy, C. M. and Quinn, J. G. (1997). Environmental Chemistry of Benzothiazoles Derived from
  1697 Rubber. Environmental Science and Technology, 31, 2847-2853.
  1698
- 1699 Rexeis, M. and Hausberger, S. (2009). Tend of vehicle emission levels until 2020 Prognosis
  1700 based on current vehicle measurements and future emission legislation. Atmospheric Environment,
  1701 43, 4689-4698.
  1702
- 1703 Riddle, S. G., Robert, M. A., Jakober, C. A., Hannigan, M. P. and Kleeman, M. J. (2007). Size
  1704 Distribution of Trace Organic Species Emitted from Light-Duty Gasoline Vehicles. Environmental
  1705 Science and Technology, 41(21), 7464-7471.
- 1707 Rissler, J., Swietlicki, E., Bengtsson, A., Boman, C., Pagels, J.,Sandstrom, T., Blomberg, A. and
  1708 Londahl, J. (2012). Experimental determination of deposition of diesel exhaust particles in the
  1709 human respiratory tract. Journal of Aerosol Science, 48, 18-33.
- 1711 Robert, M.A., VanBergen, S., Kleeman, M.J. and Jakober, C.A. (2007a). Size and composition
  1712 distributions of particulate matter emisisons: Part 1- Light-duty diesel vehicles. Journal of Air and
  1713 Waste Management Association, 57, 1414-1428.
- 1714

1706

- 1715 Robert, M. A., Kleeman, M. J. and Jakober, C. A. (2007b). Size and composition distributions of
  1716 particulate matter emisisons: Part 2- Heavy-duty diesel vehicles. Journal of Air and Waste
  1717 Management Association, 57, 1429-1438.
  1718
- 1719 Robinson, A. L., Subramanian, R., Donahue, N. M. and Rogge, W. F. (2006). Source
  1720 Apportionment of Molecular Markers and Organic Aerosol1. Polycyclic Aromatic Hydrocarbons
  1721 and Methodology for Data Visualization. Environmental Science and Technology, 40(24), 78031722 7810.

- 1723 Rodriguez, S., Van Dingenen, R., Putuad, J.-P., Dell'Acqua, A., Pey, J., Querol, X., Alastuey, A.,
- 1724 Chenery, S.,Ho, K.-F., Harrison, R., Tardivo, R., Scarnato, B. and Gemelli, V. (2007). A study on
- the relationship between mass concentrations, chemistry and number size distribution of urban fine
- aerosols in Milan, Barcelona and London. Atmospheric Chemistry and Physics, 35, 2217-2232.
- 1727

1739

1743

1754

1758

- Rogak, S. N., Green, S. I. and Pott, U. (1998). Use of Tracer Gas for Direct Calibration of
  Emission-Factor Measurements in a Traffic Tunnel. Journal of the Air and Waste Management
  Association, 48(6), 545-552.
- 1732 Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R. and Simoneit, B. R. T. (1993).
  1733 Sources of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust:
  1734 roads as sources and sinks. Environmental Science and Technology, 27(9), 1892-1904.
- 1735
  1736 Rose, D., Wehner, B., Ketzel, M., Engler, C., Voigtlander, J., Tuch, T. and Wiedensohler, A. (2006).
  1737 Atmospheric number size distributions of soot particles and estimation of emission factors.
  1738 Atmospheric Chemistry and Physics, 6(4), 1021-1031.
- Rosenbohm, E., Vogt, R., Scheer, V., Nielsen, O. J., Dreiseidler, A., Baumbach, G., Imhof, D.,
  Baltensperger, U., Fuchs, J. and Jaeschke, W. (2006). Particulate size distributions and mass
  measured at a motorway during the BAB II campaign. Atmospheric Environment, 37, 5283-5293.
- 1744 Ruellan, S. and Cachier, H. (2001). Characterisation of fresh particulate vehicular exhausts near a
  1745 Paris high flow road. Atmospheric Environment, 35(2), 453-468.
  1746
- 1747 Rutter, A. P., Snyder, D. C., Schauer, J. J., Sheesley, R. J., Olson, M. R. and DeMinter, J. (2011).
  1748 Contributions of resuspended soil and road dust to organic carbon in fine particulate matter in the
  1749 Midwestern US. Atmospheric Environment, 45(2), 514-518.
  1750
- Sadiktsis, I., Bergvall, C., Johansson, C. and Westerholm, R. (2012). Automobile Tires—A
  Potential Source of Highly Carcinogenic Dibenzopyrenes to the Environment. Environmental
  Science and Technology, 46(6), 3326-3334.
- Sahu, M., Hu, S., Ryan, P. H., Le Masters, G., Grinshpun, S. A., Chow, J. C. and Biswas, P. (2011).
  Chemical compositions and source identification of PM2.5 aerosols for estimation of a diesel source surrogate. Science of the Total Environment, 409(13), 2642-2651.
- Samara, C. and Voutsa, D. (2005). Size distribution of airborne particulate matter and associatedheavy metals in the roadside environment. Chemosphere, 59, 1197-1206.
- Sanchez-Ccoyllo, O. R., Ynoue, R. Y., Martins, L. D., Astolfo, R., Miranda, R. M., Freitas, E. D.,
  Borges, A.S., Fornaro, A., Freitas, H., Moreira, A. and Andrade, M. F. (2009). Vehicular particulate
  matter emissions in road tunnels in Sao Paulo, Brazil. Environmental Monitoring and Assessment,
  149(1-4), 241-249.
- Sanders, P. G., Xu, N., Dalka, T. M. and Maricq, M. M. (2003). Airborne brake wear debirs: Size
  distributions, composition, and a comparison of dynamometer and vehicle tests. Environmental
  Science and Technology, 37(4), 4060-4069.
- 1770
- Schaap, M., Manders, A.M.M., Hendriks, E.C.J., Cnossen, J.M., Segers, A.J.S., Denier van der Gon,
  H.A.C., Jozwicka, M., Sauter, F., Velders, G., Matthijsen, J. and Builtjes, P.J.H. (2009). Regional
  modelling of PM10 over the Netherlands. Technical Report 500099008. Available at
  http://www.rivm.nl/bibliotheek/rapporten/500099008.pdf [Accessed September 9, 2012].

- Schauer, J. J., Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R. and Simoneit, B. R. T.
  (1996). Source apportionment of airborne particulate matter using organic compounds as tracers.
  Atmospheric Environment, 30, 3837-3855.
- 1778
- Schauer, J. J., Lough, G. C., Shafer, M. M., Christensen, W. C., Arndt, M. F., DeMinter, J. T. and
  Park, J.-S. (2006). Characterization of Emissions of Metals Emitted from Motor Vehicles: Health
  Effects Institute.
- 1782

1798

1802

- Shen, Z., Han, Y., Cao, J., Tian, J., Zhu, C., Liu, S., Liu, P. and Wang, Y. (2010). Characteristics of
  traffic-related emissions: A case study in roadside ambient air over Xi'an, China. Aerosol and Air
- 1785 Quality Research, 10, 292-300.1786
- Shi, J. P. and Harrison, R. M., (1999). Investigation of ultrafine particle formation during diesel
  exhaust dilution. Environmental Science and Technology, 33, 3730-3736.
- Shi, J. P., Khan, A. A. and Harrison, R. M., (1999). Measurements of ultrafine particle
  concentration and size distribution in the urban atmosphere. Science of the Total Environment, 235,
  51-64.
- Sjodin, A., Ferm, M., Bjork, A., Rahmberg, M., Gudmundsson, A., Swietlickli, E., Johansson, C.,
  Gustafsson, M. and Blomqvist, G. (2010). Wear particles from road traffic: a field, laboratory and
  modelling study. IVL Report B1830. Availabe at: http://www.ivl.se/webdav/files/Brapporter/B1830.pdf.
- Smith, D. J. T. and Harrison, R. M. (1996). Concentrations, trends and vehicle source profile of
  polynuclear aromatic hydrocarbons in the U.K. atmosphere. Atmospheric Environment, 30(14),
  2513-2525.
- Sodeman, D. A., Toner, S. M. and Prather, K. A. (2005). Determination of single particle mass
  spectral signatures from light-duty vehicle emissions. Environmental Science and Technology,
  39(12), 4569-4580.
- Song, F. and Gao, Y. (2011). Size distributions of trace elements associated with ambient particular
  matter in the affinity of a major highway in the New Jersey–New York metropolitan area.
  Atmospheric Environment, 45(37), 6714-6723.
- 1811 Song, S., Wu, Y., Jiang, J., Yang, L., Cheng, Y. and Hao, J. (2012). Chemical characteristics of
  1812 size-resolved PM(2.5) at a roadside environment in Beijing, China. Environmental Pollution, 161,
  1813 215-221.
  1814
- 1815 Sonntag, D. B., Bailey, C. R., Fulper, C. R. and Baldauf, R. W. (2012). Contribution of Lubricating
- 1816 Oil to Particulate Matter Emissions from Light-Duty Gasoline Vehicles in Kansas City.
- 1817 Environmental Science and Technology, 46(7), 4191–4199.
- 1818 Stechmann, H. and Dannecker, W. (1990). Characterization and source analysis of vehicle1819 generated aerosols. Journal of Aerosol Science, 21, Supplement 1, S287-S290.
  1820
- Staehelin, J., Schlapfer, K., Burgin, T., Steinemann, U., Schneider, S., Brunner, D., Baumle, M.,
  Meier, M., Zahner, C., Keiser, S., Stahel, W. and Keller, C. (1995). Emission factors from road
  traffic from a tunnel study (Gubrist tunnel, Switzerland). Part I: concept and first results. Science of
  the Total Environment, 169, 141-147.
- 1825

- 1826 Sternbeck, J., Sjödin, Å. and Andréasson, K. (2002). Metal emissions from road traffic and the
  1827 influence of resuspension—results from two tunnel studies. Atmospheric Environment, 36(30),
  1828 4735-4744.
  1829
- Sun, Y.L., Zhang, Q., Schwab, J.J., Chen, W.-N., Bae, M.-S., Hung, H.-M., Lin, Y.-C., Ng, N.L.,
  Jayne, J., Massoli, P., Williams, L.R. and Demerjian, K.L. (2012). Characterization of nearhighway submicron aerosols in New York City with a high-resolution aerosol mass spectrometer.
  Atmospheric Chemistry and Physics, 12, 2215- 2227.
- Sysalova, J., Sykorova, I., Havelcova, M., Szakova, J., Trejtnarova, H. and Kotlik, B. (2012).
  Toxicologically important trace elements and organic compounds investigated in size –fractionated
  urban particulate matter collected near the Prague highway. Science of the total Environment, 437,
  127-136.
- Tanner, P.A., Hoi-Ling, M. and Yu, P.K.N. (2008). Fingerprinting metals in street dust in Beijing,
  Shanghai and Hong Kong. Environmental Science and Technology, 42, 7111-7117.
- Teixiera, E.C., Agudelo-Castaneda, D.M., Fachel, J.M.G., Leal, K.L., Garcia, K.O. and Wiegand, F.
  (2012). Source identification and seasonal variation of polycyclic aromatic hydrocarbons associated
  with atmospheric fine and coarse particles in the Metropolitan Area of Porto Alegre, RS, Brazil.
  Atmospheric Research, 118: 390-403.
- Tervahattu, H., Kupiainen, K.J., Raisanen, M., Makela, T. and Hillamo, R. (2006). Generation of
  urban road dust from anti-skid and asphalt concrete aggregates. Journal of Harazdous Materials,
  132, 39-46.
- Thorpe, A. J., Harrison, R. M., Boulter, P. G. and McCrae, I. S. (2007). Estimation of particle
  resuspension source strength on a major London Road. Atmospheric Environment, 41, 8007-8020.
- Thorpe, A. and Harrison, R. M. (2008). Sources and properties of non-exhaust particulate matter
  from road traffic: a review. Science of the Total Environment, 400(1-3), 270-282.
- 1858 Van Poppel, M., Panis, L.I., Govarts, E., Van Houtte, J. and Maenhaut, W. (2012). A comparative
  1859 study of traffic related air pollution next to a motorway and a motorway flyover. Atmospheric
  1860 Environment, 60, 132-141
- Varrica, D., Bardelli, F., Dongarra, G. and Tamburro, E. (2012). Speciation of Sb in airborne
  particulate matter, vehicle brake linings and brake pad wear residues. Atmospheric Environment, 64,
  1864 18-24.
- Viana, M., Kuhlbusch, T.A.J., Querol, X., Alastuey, A., Harrison, R.M., Hopke, P.K., Winiwarter,
  W., Vallius, M., Szidat, S., Prevot, A.S.H., Hueglin, C., Bloemen, H., Wahlin, P., Vecchi, R.,
  Miranda, A.I., Kasper-Giebl, A., Maenhaut, W. And Hitzenberger, R. (2008). Source apportionment
  of particulate matter in Europe: A review of methods and results. Aerosol Science, 39, 827-849.
- 1871 Wahlin, P., Berkowicz, R. and Palmgren, F. (2006). Characterisation of traffic-generated particulate
  1872 matter in Copenhagen. Atmospheric Environment, 40(12), 2151-2159.
- 1874 Wang, G., Kawamura, K., Xie, M., Hu, S., Gao, S., Cao, J., An, Z. and Wang, Z. (2009). Size
  1875 distribution on n-alkanes, PAHs and hopanes and their sources in the urban, mountain and marine
  1876 atmospheres over East Asia. Atmospheric Chemistry and Physics, 14: 8869-8882.
- 1877

1839

1842

1847

1851

1854

1861

1865

1870

1880 Copenhagen. Atmospheric Chemistry and Physics, 10, 2745-2764. 1881 1882 Wang, X., Westerdahl, D., Wu, Y., Pan, X. and Zhang, K. M. (2011). On-road emission factor 1883 distributions of individual diesel vehicles in and around Beijing, China. Atmospheric Environment, 1884 45(2), 503-513. 1885 Watson, J.G., Chow, J.C., Lowenthal, D.H., Pritchett, L.C., Frazier, C.A., Neuroth, G.R. and 1886 1887 Robbins, R. (1994). Differences in the carbon composition of source profiles for diesel- and 1888 gasoline- powered vehicles. Atmospheric Environment, 28(15), 2493-2505. 1889 1890 Watson, J. G., Zhu, T., Chow, J. C., Engelbrecht, J., Fujita, E. M. and Wilson, W. E. (2002). 1891 Receptor modeling application framework for particle source apportionment. Chemosphere, 49(9), 1892 1093-1136. 1893 1894 Watson, J.G. and Chow, J.C. (2007). Receptor models for source apportionment of suspended 1895 particles.In Introduction to Environmental Forensics, 2nd Edition, 2, B. Murphy and R. Morrison 1896 (Eds.), Academic Press, New York, NY, 279-316. 1897 1898 Wehner, B., Birmili, W., Gnauk, T. and Wiedensohler, A. (2002). Particle number size distributions 1899 in a street canyon and their transformation into the urban-air background: measurements and a 1900 simple model study. Atmospheric Environment, 36, 2215-2223. 1901 1902 Wei, B. and Yang, L. (2010). A review of heavy metal contaminations in urban soils, urban road 1903 dusts and agricultural soils from China. Microchemical Journal, 94(2), 99-107. 1904 1905 Weingartner, E., Keller, C., Stahel, W. A., Burtscher, H. and Baltensperger, U. (1997). Aerosol 1906 emission in a road tunnel. Atmospheric Environment, 31(3), 451-462. 1907 Weijers, E. P., Khlystov, A. Y., Kos, G. P. A. and Erisman, J. W. (2004). Variability of particulate 1908 1909 matter concentrations along roads and motorways determined by a moving measurement unit. 1910 Atmospheric Environment, 38, 2993-3002. 1911 1912 Westerdahl, D., Wang, X., Pan, X. and Zhang, M.K. (2009). Characterization of on-road vehicle emission factors and microenvironmental air quality in Beijing, China. Atmospheric Environment, 1913 1914 43, 697-705. 1915 1916 Wik, A. and Dave, G. (2009). Occurrence and effects of tire wear particles in the environment--a 1917 critical review and an initial risk assessment. Environmental pPllution, 157(1), 1-11. 1918 1919 Wu, S.-P., Tao, S., Zhang, Z.-H., Lan, T. and Zuo, Q. (2007). Characterization of TSP-bound n-1920 alkanes and polycyclic aromatic hydrocarbons at rural and urban sites of Tianjin, China. 1921 Environmental Pollution, 147(1), 203-210. 1922 Yan, B., Zheng, M., Hu, Y., Ding, X., Sullivan, A. P., Weber, R. J., Baek, J., Edgerton, E. S. and 1923 1924 Russell, A.G. (2009). Roadside, Urban, and Rural Comparison of Primary and Secondary Organic Molecular Markers in Ambient PM2.5. Environmental Science and Technology, 43(12), 4287-4293. 1925 1926 1927 Yin, J., Harrison, R. M., Chen, Q., Rutter, A. and Schauer, J. J. (2010). Source apportionment of 1928 fine particles at urban background and rural sites in the UK atmosphere. Atmospheric Environment,

Wang, F., Ketzel, M., Ellermann, T., Wahlin, P., Jensen, S.S., Fang, D. and Massling, A. (2010).

Particle number, particle mass and NOx emission factors at a highway and an urban street in

1929 44, 841-851.

1878

- 1930 Zhang, Y., Stedman, D. H., Bishop, G. A., Guenther, P.L. and Beaton, S. P. (1995). Worldwide On1931 Road Vehicle Exhaust Emissions Study by Remote Sensing. Environmental Science
  1932 andTechnology, 29, 2286-2294.
- 1934 Zhu, Y., Hinds, W. C., Kim, S. and Sioutas, C. (2002a). Concentration and Size Distribution of
  1935 Ultrafine Particles Near a Major Highway. Journal and Air and Waste Management Association, 52,
  1936 1032-1042.
- 1937

1944

1933

- Zhu, Y., Hinds, W. C., Kim, S., Shen, S. and Sioutas, C. (2002b). Study of ultrafine particles near a
  major highway with heavy-duty diesel traffic. Atmospheric Environment, 36(27), 4323-4335.
- 1941 Zhu, Y., Kuhn, T., Mayo, P. and Hinds, W.C. (2006). Comparison of daytime and nighttime
  1942 concentration profiles and size distributions of ultrafine particles near a major highway.
  1943 Environmental Science and Technology, 40, 2531-2536.
- 1945 Zhu, C.-S., Chen, C.-C., Cao, J.-J, Tsai, C.-J., Chou, C. C.-K., Liu, S.-C. and Roam, G.-D. (2010).
  1946 Characterization of carbon fractions for atmospheric fine particles and nanoparticles in a highway
- 1947 tunnel. Atmospheric Environment, 44, 2668-2673.
- 1948

1949

1951	TABLE L	EGENDS
1952		
1953	Table 1:	Mechanisms for non-exhaust particle emission (Rogge et al., 1993;
1954		Boulter et al., 2005; Wahlin et al., 2006; Thorpe et al. 2007; Amato et al.,
1955		2009; Harrison, 2009; Gietl et al., 2010; Barmapadimos et al., 2011;
1956		Hays et al., 2011; Denier van der Gon, 2013).
1957	<b>T-11-0</b>	
1958	Table 2:	Key tracers used for non-exhaust PM.
1959	Table 2.	Datio of Cy to Sh characteristic of bushes wear resticles in ambient DM
1960	Table 5:	Ratio of Cu to Sb characteristic of brake wear particles in ambient PM
1961		reported in the interature.
1962	Tabla 1.	(a) Results from mass size distribution analysis for traffic related PM as
1903	1 abic 4.	(a) Results from mass size distribution analysis for traffic-related 1 W as reported in the literature (in um): (b) Results from mass size distribution
1065		analysis for traffic-related elements as reported in the literature (in $\mu$ m)
1966		analysis for traffic felded elements as reported in the incrature (in µin).
1967	Table 5:	Comparison between dynamometer and tunnel/roadway measurements
1968		(Allen et al., 2001: Imhof et al., 2005a: Phuleria et al., 2006: Phuleria et
1969		al., 2007; Handler et al., 2008; He et al., 2008; El Haddad et al., 2009;
1970		Sanchez-Ccoyllo et al., 2009; Franco et al., 2013).
1971		
1972	Table 6:	Emission factors as reported in the literature (selected studies).
1973		
1974	Table 7:	Source apportionment analyses for road traffic-generated PM (selected
1975		studies).
1976		
1977		
1978	FIGURE I	LEGEND
1979		
1980	Figure 1:	Methods for direct road traffic emissions analyses.
1981		
1982		
1983		
1984		

**Table 1:** Mechanisms for non-exhaust particle emission (Rogge et al., 1993; Boulter et al., 2005;

Wahlin et al., 2006; Thorpe et al. 2007; Amato et al., 2009; Harrison, 2009; Gietl et al., 2010;
Barmapadimos et al., 2011; Hays et al., 2011; Denier van der Gon, 2013)

Emission type	Mechanism	Includes
Direct	Abrasion and wear and	Tyre, brake, clutch,
	tear	road surface
	Corrosion	Vehicle, street furniture
Indirect	Re-suspension (due to	
	tyre shear, wind and	
	vehicle turbulence)	

1988

**Table 2:** Key tracers used for non-exhaust PM

Reference	Brake wear	Tyre wear	Re-suspension
Adachi and Tainosho		Zn	-
(2004)			
	Fe, Ba, Cu, Sb, Zr		
Schauer et al. (2006)	Fe, Cu, Ba		-
Grieshop et al. (2006)	Cu, Sb, Ba and Ga		
Wahlin et al. (2006)	Cr, Fe, Cu, Zn, Zr, Mo,	Al, Si, K, Ca, Ti, Mn, Fe,	Zn and Sr (together with
	Sn, Sb, Ba and Pb	road dust)	
Tanner et al. (2008)	Cu, Cd	Zn	
Canepari et al. (2008)	Ba, Fe, Sb, Sr	-	-
Harrison (2009)	Ba, Cu	-	Al
Dongarra et al. (2009)	Cu, Mo, Sb	-	
Fabretti et al. (2009)	Cu, Zn, Sb, Sn (vehicular	-	Rb, Sr, Mn, Fe, As
	abrasion)		
Keuken et al. (2010)	Cu	Zn	-
Bukowiecki et al. (2010)	Fe, Cu, Zn, Zr, Mo, Sn,		
	Sb and Ba		
Pey et al. (2010)	-	-	Fe, Ca, Sb, Sn, Cu, Zn
Perez et al. (2010)	Sb, Cu, Ni, Sn (wear of bra	ake, tire and other parts)	Fe
Amato et al. (2011a)	Fe, Cu, Zn, Cr, Sn, Sb	OC, S, Zn	Al, Ca, Fe, V
Apeagyei et al. (2011)	Fe, Ti, Cu, Ba	Zn, Ca, W, K, Fe, Ti, Cr,	-
		Мо	
Duong and Lee (2011)	Ni, Cu	Zn	
Ondracek et al. (2011)	Cu, Ba, Fe, Zn	-	
Song and Gao (2011)	Sb, Cu, Fe, Pb	Zn, Co	
Sahu et al. (2011)	Zn (brake and tyre wear)	-	
Peltier et al. (2011)	-	-	Al, Si, Ti, Fe
Harrison et al. (2012b)	Ba, Cu, Fe, Sb	Zn	Si, Al

1990

**Table 3:** Ratio of Cu to Sb characteristic of brake wear particles in ambient PM reported in theliterature

Reference	Location	Cu:Sb ratio
Stechmann and Dannecker (1990)	Hamburg, Germany	5.6
Adachi and Tainosho (2004)	Kobe, Japan	1.3
Lin et al. (2005)	Taiwan	9.1±1.79 (coarse fraction)
		2.17±0.83 (fine fraction)
Wahlin et al. (2006)	Copenhagen, Denmark	4.4±0.3
Hjortenkrans et al. (2007)	Stockholm, Sweden	5.3
Fabretti et al. (2009)	Nice, France	7.0±1.3
Dongarra et al. (2009)	Palermo, Italy	4.9 (coarse fraction)
		2.5 (fine fraction)
Amato et al. (2011b)	Barcelona, Spain	8.0 ±1.5 (roadside)
		$6.8 \pm 1.9$ (urban)
Bukowiecki et al. (2010)	Zürich-Weststrasse, Switzerland	6.2 (coarse fraction)
Pey et al. (2010)	Barcelona, Spain	9 (coarse fraction)
		10 (fine fraction)
Gietl et al. (2010)	London, UK	9.1

**Table 4:** (a) Results from mass size distribution analysis for traffic-related PM as reported in the
 literature (in μm)

Reference	Location	Site type	Component	Distribution type	Primary peak
Allen et al. (2001)	USA	Tunnel	PM <sub>10</sub>	Unimodal	0.1-0.18
Hildemann et al.(1991)	USA	Freeway	PM	Trimodal	Ultrafine (<0.2), fine (0.2-1) and coarse (1-10)
Robert et al.	USA	Dynamometer	PM <sub>1.8</sub> (HDGV)	Unimodal	0.1-0.18
(2007)			$PM_{1.8}$ (LDGV)	Unimodal	0.1-0.18

1998

(b) Results from mass size distribution analysis for traffic-related elements as reported in theliterature (in μm)

Reference	Location	Metal(s)	Distribution type	Primary peak	Secondary
Stechmann and Dannecker (1990)	Germany	Cu, Ba	Unimodal	2.5	peak
Funasaka et al.	Japan (tunnel)	EC	Unimodal		
(1998)		OC	Bimodal	1	2-10
Lin et al., 2005	Taiwan	Si, Al, Fe, Ca	Bimodal	3.2-5.6	Nano size range
		Ba	Bimodal	1.0-1.8	0.18-0.32
		Pb	Bimodal	3.2-5.6	0.32- 0.56
		Zn	Bimodal	1.0-1.8	0.32-0.56
		Mn, Cu, Cd, Sr	Bimodal		
		Cr, V	Trimodal		
Lough et al. (2005)	USA	Cu	Bimodal	3.0	0.1
Wahlin et al.	Denmark	Ba, Sb	Unimodal	3	
(2006)		Cu	Bimodal	3	0.1
Hien et al. (2007)	Vietnam	PAHs in PM <sub>2.5</sub>	Bimodal	Coarse and fine	
Riddle et al. (2007)	USA	Hopanes and steranes in PM <sub>1.8</sub>	Unimodal	0.10-0.18	0.32 (vehicles without catalysts
Wang et al. (2009)	China	n-Alkanes, PAHs and Hopanes	Unimodal	0.7-1.1	
Gietl et al.	UK	Ba, Fe	Unimodal	1.2-7.2	
(2010)		Cu, Sb	Bimodal	1.2-7.2	0.2-0.4
Huang et al. (2012)	China	n-Alkanes	Unimodal	0.32-0.56	
		PAHs	Unimodal	0.56-1.0	
		Hopanes	Unimodal	0.56-1.0	3.2 (in some cases)
Ondráček et al.	Czech Republic	Fe, Cu, Mn, Zn	Unimodal	2.5	
(2011)	(freeway)	Si, Al, Ca and coarse mode K	Unimodal	5	
Song and Gao ( <u>20122011</u> )	USA	Al, Fe, Cu, Sb, Sc, Mn	Bimodal	3.2- 5.6	1.0-1.8
		Cd	Bimodal	0.18- 0.56	3.2-5.6
		Pb, Zn	Trimodal	0.32-0.56	1.0-1.8, 3.2- 5.6
		Ni, V	Bimodal	0.18-0.32	3.2-5.6

Table 5: Comparison between dynamometer and tunnel/roadway measurements (Allen et al., 2001;
Imhof et al., 2005a; Phuleria et al., 2006; Phuleria et al., 2007; Handler et al., 2008; He et al., 2008;
El Haddad et al., 2009; Sanchez-Ccoyllo et al., 2009; Franco et al., 2013)

	Dynamometer measurements	Tunnel/Roadway measurements
Testing conditions	Precise and controlled; different test cycles and driving conditions can be analysed; can include cold- start emissions	Ambient conditions- cannot be physically controlled; cannot include cold-start emissions
		Better known boundary conditions and dilution effects
		In some cases, atmospheric conditions in tunnel may not be the same as ambient environment
Representativeness for the vehicle fleet	Low, since the tests are conducted on a sub-set of the in-use fleet and variations in engine type, vehicle age and maintenance and mixing of emissions from different vehicles etc are not accounted for. Such studies do not account for particle aging effects	High, since the measurements are made in ambient environment and for in-use mixed fleet
Emission type accounted for	Exhaust	Exhaust and non-exhaust

**Table 6:** Emission factors as reported in the literature (selected studies)

• Particle Number

Reference	City, Country	Study type	Particle number EF
Gertler et al. (2002)	USA	Tunnel (mixed LDV and	5x 10 <sup>12</sup> particles/veh/km (low HDV)
		HDV)	$2-3 \times 10^{13}$ particles/veh/km (high HDV)
Abu-Allaban et al. (2002)	USA	Tunnel	$5.2-5.4 \times 10^{13}$ particles/veh/km (low HD fraction)
			2.1-3.1x 10 <sup>14</sup> particles/veh/km (high HD
			fraction)
Jamriska et al. (2004)	Australia	Tunnel (diesel powered buses)	$3.11 \pm 2.41 \ge 10^{14}$ particles/km
Kittleson et al. (2004)	USA	Roadside (particles $> 3$ nm)	1.9-9.9 x 10 <sup>14</sup> particles/km (gasoline-dominant
			fleet)
Jones and Harrison (2006)	UK	Roadside	$5.84 \times 10^{13}$ particles/veh/km
			$6.36 \times 10^{14}$ particles/veh/km (LDV)
Westerdahl et al. (2009) China Roadside- LDV (prevalent		1.8 x 10 <sup>15</sup> particles/kg fuel	
		during daytime)	
		Roadside- HDV (prevalent	1.1 x 10 <sup>16</sup> particles/kg fuel
		during nighttime)	
Wang et al. (2009)	Denmark	Highway	$(215\pm15)$ 10 <sup>12</sup> particles/ veh/km
Ban-Weiss et al. (2010)	USA	Tunnel- LDV	$(3.9\pm1.4)$ 10 <sup>14</sup> particles/kg fuel
		Tunnel-HDV	$(3.3\pm1.3)$ 10 <sup>15</sup> particles/kg fuel
Wang et al. (2011)	China	Vehicle chase	$7.1 \times 10^{16}$ particles/kg fuel

## • Particle Mass

Reference	City, Country	Study type	PM size range	EF
Weingartner et al. (1997)	Switzerland	Tunnel (mixed LDV and	PM <sub>3</sub>	310 mg/km (63% vehicles running on
		HDV with more LDV)		diesel)
Gillies et al. (2001)	USA	Tunnel (mixed LDV and	PM <sub>2.5</sub>	$0.052 \pm 0.027$ g/km
		HDV)	$PM_{10}$	$0.069 \pm 0.030$ g/km
Gertler et al. (2002)	USA	Tunnel (mixed LDV and	PM <sub>2.5</sub>	$0.062 \pm 0.042$ g/km

		HDV)	PM <sub>10</sub>	$0.087 \pm 0.054$ g/km
Jamriska et al. (2004) Australia Tunnel (diesel p		Tunnel (diesel powered	PM <sub>2.5</sub>	583± 451 mg/km
		buses)		
Greishop et al. (2006)	USA	Tunnel	PM <sub>2.5</sub> (rush hour)	189±23 mg/kg fuel
			PM <sub>2.5</sub> (mid-day)	158±29 mg/kg fuel
			$PM_{2.5}$ (early morning-HDV)	437±76 mg/kg fuel
Handler et al. (2008)	Austria	Tunnel	TSP	129±45 mg/veh/km
			$PM_{10}$	62±18 mg/veh/km
			PM <sub>2.5</sub>	26±10 mg/veh/km
Sanchez-Ccoyllo et al.,	Brazil		Coarse	127 mg/km
(2009)			Fine	92 mg/km
Wang et al. (2010)	Denmark	Highway	PM <sub>2.5</sub>	29 mg/veh/km
Chiang and Huang (2009) Taiwan Freeway tunnel F		PM <sub>2.5-10</sub>	18±6.5 mg/veh/km	
			PM <sub>2.5</sub>	39±11 mg/veh/km
Cheng et al. (2010)	Hong Kong	Tunnel, roadside sites	PM <sub>2.5</sub>	257±31 mg/veh/km
		and urban site		
Bukowiecki et al. (2010)	Switzerland	Roadside	PM <sub>10</sub> - LDV	24±8 mg/veh/km
			PM <sub>10</sub> -HDV	498±86 mg/veh/km
Wang et al. (2011)	China	Vehicle chase	PM <sub>0.5</sub>	2.35 g/kg fuel
Mancilla and Mendoza	Mexico	Tunnel	$PM_{10}$	22.8±7.4 mg/veh/km
(2012)				
Chen et al. (2013)	China	Tunnel (mixed with higher proportion of gasoline		687 mg/veh/km (summer)
vehicles)			714 mg/veh/km (winter)	

## • LDV and HDV vehicles (mass)

Reference	Country	Study type	PM size range	LDV EF	HDV EF
Weingartner et al. (1997)	Switzerland	Tunnel	PM <sub>3</sub>	$8.53 \pm 0.47 \text{ mg/km}$	$383.5 \pm 10.7$ mg/km
Gertler et al. (2002)	USA	Tunnel (mixed LDV and HDV)	PM <sub>2.5</sub>	$0.014 \pm 0.013 \text{ g/km}$	$0.135 \pm 0.018$ g/km 0.181 ± 0.013 g/km
Abu-Allaban et al.	USA	Roadside (LD-Spark	PM10	10-70 mg/km/vehicle	60-570 mg/km

(2003)		ignition and HD-	PM <sub>2.5</sub>	10-50 mg/km/vehicle	60-480 mg/km
		diesel)			
Chiang et al. (2012)	Taiwan	Dynamometer	PM <sub>2.5</sub>	0.172 g/km (diesel)	
Oanh et al. (2010)	Bangkok	Dynamometer	PM <sub>2.5</sub>	230 mg/km	176 mg/km
Jones and Harrison	UK	Roadside	$PM_{10}$	0.033±0.006 g/veh/km	0.370±0.032 g/veh/km
(2006)			PM <sub>2.5</sub>	0.010±0.004 g/veh/km	0.179±0.022 g/veh/km
Liacos et al. (2012)	USA	Freeway	PM <sub>2.5</sub>	151189.3 mg/km/h (low HDV)	169629.5 mg/km/h
					(high HDV)

• Non-exhaust emissions

Reference	City, Country	Study type	PM size range	Road dust	Brake wear	Tyre wear
Abu-Allaban et al.	USA	Roadside (LD-	PM <sub>10</sub> LDV	40-780 mg/km/vehicle	0-80	-
(2003)		Spark ignition			mg/km/vehicle	
		and HD-diesel)	PM <sub>10</sub> HDV	230-7800 mg/km/vehicle	0-610	
					mg/km/vehicle	
			PM <sub>2.5</sub> LDV	2-25 mg/km/vehicle	0-5 mg/km/vehicle	-
			PM <sub>2.5</sub> HDV	15-300 mg/km/vehicle	0-15	
					mg/km/vehicle	
Bukowiecki et al.	Switzerland	Roadside	PM10	27 mg/vehicle/km	15 mg/ vehicle /km	
(2010)						
Amato et al. (2012)	Spain	Freeway	PM <sub>10</sub>	$22.7 \pm 14.2$ mg/vehicle	-	-
				kilometer travelled		

 Table 7: Source apportionment analyses for road traffic-generated PM (selected studies)

Reference	Country	Method	Particle type	Sources
Lin et al.	Taiwan	PCA	Coarse particles	Fugitive dust and brake lining, diesel, fuel oil,
(2005)				gasoline
			Fine particles	Gasoline, diesel, brake lining/tyre wear, fuel oil
			Ultrafine and	Gasoline, diesel, industry, fuel oil
			nano particles	
Furujso et al.	Stockholm,	PMF	$PM_{10}$	Resuspension
(2007)	Sweden			Vehicle derived (brake wear)
				Road salt
				Regional combustion
Char and	Ontonia	A la a 1	Doutial a normality	Long range transport
Chan and Morrielianiah	Ontario,	Adsolute	Particle number	Common sources across 5 sites: Nucleation
(2007)	Canada	PCA	uistribution	particles (photochemically produced), regional
(2007)			with trace gas	pollution factor, boundary layer dynamics factor
			meteorological	Specific sources at the three sites: Local
			data	industrial emissions (urban)
			uata	Processed nucleation mode narticles (polluted
				rural)
				Transported fine particles (downwind Toronto)
Amato et al.	Barcelona.	PMF2	$PM_{10}$ and $PM_{25}$	Road dust factor accounted for 17% PM10, 8%
(2009)	Spain	with ME	10	$PM_{25}$ and less than 2% of $PM_{1}$ .
× ,	1			2.5
				All traffic sources contributed to 46% PM10,
				51% PM <sub>2.5</sub> and 48% of PM <sub>1</sub> .
Fabretti et al.	Nice, France	PMF	PM <sub>2.5</sub>	Vehicle abrasion: Cu, Zn, Sb (36%)
(2009)				
				Re-suspension: Mn, Fe, As, Rb, Sr (43%)
				Fuel combustion: V, Ni, Co (21%)
Dreyfus et al.	Wilmington,	PMF	Organic carbon	Diesel factor: Alkyl fragment, benzothiazole,
(2009)	USA			benzenedicarboxylic acids/3,4-
				dimethoxybenzaldenyde, 3,4-
				dimethoxybenzoic acid, hopanes
				Road dust/aar amissions: honoras honzothiozola
				and triacontanoic acid
Mancilla and	Monterrey	FΔ	PM <sub>e</sub> -	Brake emissions
Mendoza	Mexico	17	1 1/12.5	Exhaust emissions
(2012)	MEXICO			L'Aldust emissions
Ondracek et	Prague,			Abrasion of vehicle parts: Fe, Cu, Mn, Zn
al. (2011)	Czech			Re-suspension of road dust: Si, Al, Ca
· · /	Republic			Long-range transport/regional background: Ca,
	1			K
Song and	New Jersey,	FA	РМ	Brake wear and fuel combustion: Fe, Sb, Pb, Cd
Gao (2011)	USA			(~35%)
				Primary fuel combustion: Cr, Ni, V, Cu
				(~28.3%)
				Tyre wear and fuel combustion: Zn, Co
				(~23.7%)
Yin et al.	United	CMB	PM <sub>2.5</sub>	Vegetative detritus; woodsmoke; natural gas;
(2010)	Kingdom			diesel engines (11.6%); gasoline engines
				(1.4%); smoking engines (9.7%); coal;
				dust/soil; secondary organic matter; sea salt;
			1	ammonium sulphate; ammonium nitrate



Figure 2: Methods for direct road traffic emissions analyses