

A discusión

DIFFUSION IN COMPLEX SOCIAL NETWORKS*

Dunia López-Pintado

WP-AD 2004-33

Corresponding author: Dunia López-Pintado, Universidad de Alicante, Apto. Correos 99, E. 03080 Alicante.
Tel: +34 96 590 34 81, Fax: +34 96 590 34 64. E-mail: dunia@merlin.fae.ua.es

Editor: Instituto Valenciano de Investigaciones Económicas, S.A.

Primera Edición Septiembre 2004.

Depósito Legal: V-3982-2004

IVIE working papers offer in advance the results of economic research under way in order to encourage a discussion process before sending them to scientific journals for their final publication.

* I wish to thank my advisor Fernando Vega-Redondo for his continuous encouragement and valuable suggestion. I am also grateful to Juan D. Moreno-Ternero, Ricardo Martínez, Giovanni Ponti and Leon Danon for their helpful comments. Financial support was provided by the Instituto Valenciano de Investigaciones Económicas (IVIE).

DIFFUSION IN COMPLEX SOCIAL NETWORKS

Dunia López-Pintado

ABSTRACT

This paper studies the problem of spreading a product (an idea or a technology) among agents in a social network. An agent obtains the product with a probability that depends on the spreading rate (or degree of contagion) of the product as well as on the behaviour of the agent's neighbours. This paper shows, using *a mean field* approach, that there exists a threshold for the spreading rate that shapes the pattern of the product's diffusion. This threshold, that depends on the network structure and the mechanism of contagion, determines whether the product spreads and becomes persistent or it does not spread and vanishes.

Key words: social networks, diffusion, contagion.

JEL Classification: C73

1 Introduction

Introducing a new product, technology or idea, in the market is an issue of major social-economic relevance. Innovations do not necessarily spread at once, but often spread gradually through social and geographic networks. In fact, many products promote rather easily in a social system through a domino effect. In a first stage a few innovators adopt the product, and this makes more likely that their neighbors do the same, then their neighbors' neighbors and so forth. Indeed, these products or ideas can spread more efficiently from “consumer-to-consumer dialogue”, rather than from sellers to consumers. Indeed, the opinion on these products among the agents in the social system heavily depends on their interpersonal ties. These communication channels are more trusted and have greater effectiveness than mass media advertisements. Thus, traditional marketing is being replaced by new strategies in which the product is turned into “epidemics” where consumers do the marketing themselves.¹ A recent example is that of the mobile phones. They became popular in the mid 90's and, at present, almost every individual possesses a phone, which is considered as an essential commodity in developed countries. Apart from the intrinsic advantages that the new product might provide to its users, the fast spreading of it in the population is reinforced by more subtle aspects, such as fashion and benefits from coordinating in the decision with your contacts. The spreading of these products share common features with the contagion of an infectious disease in a population. The aim of this paper is to bring these issues to a common setting to describe how a new technology or an idea propagates in a population where agents only interact with their neighbors. In particular, we address the following questions: How many initial adopters are needed to spread a product? and How does the spreading pattern depend on the interaction structure among individuals and on the contagion mechanism?

We consider a large population with a complex pattern of interaction among agents. Moreover, the social system is described through a network structure. Traditionally, the study of networks has been a topic of graph theory. Graph theory, however, concentrated in small networks with a high degree of regularity. This paper focuses on the large-scale statistical properties of the network instead of on the properties of single vertices. We assume that the precise topology of the network is unknown and thus it is considered as a “random ensemble”. The number of edges a node has -the connectivity of the node- is characterized by a distribution function $P(k)$, which gives the probability that a randomly selected node has exactly k edges. Throughout this paper, the network is exogenously given and it is characterized by its connectivity distribution $P(k)$.

Random graphs have been widely studied in the literature of complex networks. The seminal paper by Erdos and Renyi (1959) defines a *random graph* by a group of N nodes such that every pair of nodes is connected with a certain

¹In a recent book, Seith (2002) describes how an “idea” can spread in a population the same way a “virus” does.

probability p . The graphs generated in this manner have a connectivity distribution which is a *Poisson distribution* with its peak at the average connectivity, denoted by $\langle k \rangle$. In this case, the majority of nodes have similar connectivity. Recent empirical studies show that most large complex networks are characterized by a connectivity distribution different to a Poisson distribution (e.g., Barabási et al., 2000; Faloutsos et al., 1999; Lijeros et al., 2001; Yook et al., 2001, etc.). For instance, WWW, Internet and human sexual contacts, among others, have a power-law connectivity distribution, i.e. $P(k) \sim k^{-\gamma}$ where γ ranges between 2 and 3. This implies that each node has a statistically significant probability of having a very large number of connections compared to the average connectivity $\langle k \rangle$ which generates an extreme heterogeneity in the connectivity of agents. Such random networks are called *scale-free*. This class of networks can be easily simulated by imposing that every period new nodes are introduced in the network and these are linked preferentially to the most highly connected existing nodes. Therefore, two principles underlay scale-free networks: preferential attachment and growth (see Barabási and Albert, 1999).

This work attempts to be of general applicability, i.e. the results are formulated for any given connectivity distribution ($P(k)$). We pay special attention, however, to the differential properties of Poisson and scale-free networks. We have considered a simple diffusion model. Each agent, classified as either an “active” or a “potential” consumer, is represented by a node in the complex network of social contacts. The transition from a potential to an active consumer depends on the intrinsic properties of the product as well as on the number and behavior of neighbors. Conversely, an active consumer becomes potential at an exogenously given rate. This reflects the idea that, with a certain probability, independent of the behavior of neighbors, an agent may need to replace the product because it is lost or deteriorated. The framework considered in this work is closely related to the so-called “susceptible-infected-susceptible” (SIS) model, commonly used in epidemiology. Some paradigmatic examples that are described using the SIS model are the diffusion of AIDS in a sexual contact network or the spreading of a computer virus via Internet (e.g. Pastor-Satorras and Vespignani, 2000; Lloyd and May, 2001). Each agent is represented by a node and can be either “healthy” or “infected”. In each time step a healthy node is infected at a rate ν if it is directly connected to at least one infected agent. Conversely, an infected agent is cured at a rate $\delta > 0$.

This paper extends the SIS model in several ways. For instance, the SIS model considers the contagion of a disease as a linear function of the absolute number of infected neighbors, whereas the present model allows for non-linear mechanisms. For example, adding one active consumer in the neighborhood of an agent with only one active neighbor could have a different effect than doing the same in a situation where many neighbors are already active. Furthermore, a richer framework is introduced in which the intensity of each interaction can depend on the total number of interactions. This possibility has been ignored in the epidemiology literature although it is a natural assumption in most economic

contexts. Thus, in our model, the contagion could depend not only on the absolute number of neighboring active consumers but also on the size of the neighborhood, i.e. the connectivity of the agent.

The theoretical results of our model are derived using the so-called *mean field theory*. Heuristically, this theory simplifies the description of the exact contagion process by substituting some local variables of the dynamics by their global mean values. Although our model is an attempt to predict events in a social-economic environment, this approach is commonly used in physics and biology since it provides a reasonable guide of the qualitative behavior of complex systems. We show that there exists a threshold for the degree of contagion (or spreading rate) of the product, such that, above the threshold the technology spreads and becomes persistent. This threshold depends crucially on two features: the *mechanism of diffusion* and the *connectivity distribution* of agents in the population. Indeed, when the mechanism of diffusion is such that the intensity of each interaction is independent of the total number of neighbors, i.e. the contagion of the product only depends on the *absolute* number of active consumers among neighbors, the diffusion of the product is easier and greater in scale-free networks than in Poisson networks. In contrast, if the intensity of each interaction decreases in parallel with the number of neighbors, or specifically, the contagion of the product depends on the *relative* proportion of active consumers among neighbors, all networks exhibit the same spreading behavior. Finally, for concave diffusion functions there always exists a continuous transition from the absence to the existence of diffusion. Nevertheless, for some particular non-concave diffusion functions, this transition is non-continuous, i.e. a *phase transition* and hysteresis phenomenon occurs.

In summary, this paper builds on two literatures. The formal framework considered is close to the literature on epidemiology and complex systems, as mentioned above, mean field theory is often used. However, the inspiration for this work comes mainly from the fast expanding pure game theory literature on social and economic networks. Recent instances of this literature show that the pattern of interaction between individuals is crucial in determining the nature of outcomes. A wide number of papers have focused on the analysis of lattices; that is, regular networks in which all players have the same number of direct connections (e.g., Anderlini and Ianni, 1996; Ellison, 1993; Goyal, 1996; Young, 2002; Blume, 1995). One step beyond comes from Morris (2000) who has developed techniques to study coordination games in general networks. Although, the present paper shares the flavour of these previous works it introduces important novelties. First, we study very general contagion mechanisms characterized by the fact that the transition from one individual state to the other (active to potential consumer and vice versa) is typically stochastic and asymmetric. Second, we consider complex random networks rather than networks with a deterministic geometric form. The paper is organized as follows. The model is contained in Section 2. Section 3 provides a game theoretical context where the model can be applied. Section 4 returns to the general model and introduces the mean

field theory. Section 5 presents the main results. In Section 6 we run some simulations of the original dynamics in order to test the validity of the theoretical results. Finally, Section 7 concludes. Some proofs have been relegated to the Appendix.

2 The model

Let $N = \{1, 2, \dots, i, \dots, n\}$ be a finite but large set of agents. Assume agents are communicated one with another through certain channels which determine the social system. More precisely, each agent interacts only with her fixed group of neighbors, i.e. direct connections. These interactions represent personal and professional contacts. To describe the social system formally, consider an undirected network $\Gamma \equiv (V, L)$ where V is the set of nodes and L is the set of undirected links. Each node represents one agent in the population. A link $\{i, j\}$ belongs to the set L if and only if agents i and j are directly connected. Let $K_i \subseteq N$ be the set of neighbors of player i and let k_i be its cardinality which is referred as her *connectivity* from here onwards.

Assume that the population is large and the pattern of interactions between agents is complex. Moreover, the network structure has a high degree of randomness and thus can only be described by its large-scale statistical properties. Denote by $P(k)$ to the connectivity distribution of the network, i.e. the fraction of agents in the population that have exactly k direct neighbors. Equivalently, $P(k)$ is the probability that an agent chosen uniformly at random has connectivity k . Throughout this paper, the network is characterized by being “random” and having a connectivity distribution $P(k)$ which is exogenously given. These networks have been referred in the literature as *generalized random networks* since they extend the Erdos-Renyi random graphs by incorporating the property of non-Poisson connectivity distributions.² Roughly speaking, a random graph is characterized by the fact that when generating the network, the link-formation operates in a stochastically independent way across nodes. This process rules out correlations among connectivity of connected nodes and generates a graph with a low clustering coefficient. One of the aims of this work is to explicitly account for the influence of $P(k)$ in the spreading behavior of the product.

Assume there is a new product in the market. We focus on its spreading among the population N . To do so, consider that an agent $i \in N$ can only exist in two discrete states $s_i \in \{0, 1\}$, where $s_i = 0$ if i is a “potential” consumer and $s_i = 1$ if i is an “active” consumer. A potential consumer is an agent that does not have the product but is susceptible of obtaining it if exposed to someone who does. An active consumer is an agent that has already adopted the product and so can influence her neighbors in favor of obtaining it.

²For further details on generalized random networks the reader is referred to M.E.J. Newman (2003)

Consider a stochastic continuous time dynamics process as follows. At time t , the state of the system is a vector

$$s_t = (s_{1t}, s_{2t}, \dots, s_{it}, \dots, s_{nt}) \in S^n \equiv \{0, 1\}^n$$

where $s_{it} = 0$ if i is a potential consumer at time t whereas $s_{it} = 1$ if i is an active consumer at time t . Assume i is a potential consumer at time t . She becomes an active consumer at a rate that depends crucially on: her connectivity k_i , the number of neighbors that are active consumers at time t (a_i hereafter) and the spreading rate (or degree of contagion) of the product, denoted by $\nu \geq 0$. More precisely, the transition rate from potential to active consumer is given by a function $F(\nu, k_i, a_i)$ that determines the properties of the mechanism of diffusion. We assume independence of the spreading rate effect and the effect that the behavior of neighbors has over the agent's decision. Thus,

$$F(\nu, k_i, a_i) = \nu \cdot f(k_i, a_i)$$

where $f(k_i, a_i)$, named as the *diffusion function* from here onwards, is a non-negative function only defined for $(k_i, a_i) \in N \times N$ such that $0 \leq a_i \leq k_i$.³ It is worth noting that, the connectivity of an agent is fixed throughout the dynamics. Instead, the number of active consumers among neighbors " a_i " might change over time. We suppose that a necessary condition for the adoption of the product is that at least one neighbor has already adopted it. More precisely,

$$f(k, 0) = 0 \text{ for all } k \geq 1 \tag{A-1}$$

Roughly speaking, the transition from a potential to an active consumer can be interpreted as follows. At a rate ν any given agent becomes aware of the existence of the product -e.g. through mass media advertisement- and considers the possibility of adopting it. The agent's final decision, however, depends crucially on her neighbors' behavior. More precisely, the agent responds to her neighbors current configuration by choosing an action according to some choice rule. The particular choice rule considered is characterized by $f(k_i, a_i)$.

Conversely, consider agent $i \in N$ is an active consumer at time t . Then, i becomes a potential consumer at some stochastically constant rate $\delta > 0$ which indicates the rate at which the agent may need to replace the product because it is lost or deteriorated. Notice that, this transition is independent of her neighbors' behavior. It is implicit in this formulation that the cost of "maintaining" the product is approximately zero and thus agents never have incentives for getting rid of it. Finally, let us define the *effective spreading rate* of the product by $\lambda = \frac{\nu}{\delta}$.

For concreteness, we will now define formally what we mean by the mechanism of diffusion.

³The SIS model can be obtained as a particular case of this model by simply assuming $f(k, a) = a$.

Definition 1 A mechanism of diffusion is a pair $m = (\lambda, f(\cdot))$ where λ denotes the effective spreading of the product and $f(\cdot)$ denotes the diffusion function.

Notice that, since the transition rates only depend on the properties of the present state, the dynamics induced by the connectivity distribution $P(k)$ and the mechanism of diffusion m determines a continuous *Markov chain* over the space of possible states S^n .

The aim of this work is to analyze *whether* and *how* the product spreads in the population. Several questions raise as natural:

- Is there prevalence of the product in the long-run of the dynamics?
- Are small perturbations of the initial state in which there are no active consumers enough to converge to states with a positive fraction of active consumers?
- Is there a discontinuity (or phase transition) in the proportion of active consumers as we increase λ ?

The next section describes briefly a particular context where this model could be applied.

3 An example

Consider a population of agents $N = \{0, 1, \dots, n\}$. As before, agents interact only with their fixed group of neighbors. The pattern of interaction among them is described through a social network where each node represents one agent and the connections among them are represented by links.

Let x be a new technology. Assume that the cost incurred by an individual i in case of adopting x is randomly determined by \tilde{c}_i . For the sake of concreteness, suppose $\tilde{c}_i \sim U[0, C]$ where C is the highest possible cost. Also assume that $(\tilde{c}_j)_{j \in N}$ are i.i.d. Therefore (ex-post) the cost can be different across agents. For simplicity, assume that, once adopting the product, the cost of maintaining it is zero.

Suppose that, if two players are neighbors, there is a pairwise interaction that can generate mutual payoffs. The common set of strategies is $S = \{0, 1\}$ where $s_i = 1$ means agents i is an active consumer whereas $s_i = 0$ otherwise. For each pair of strategies $s, s' \in S$, the payoff earned by a player i choosing s when interacting with her partner j choosing s' is $b > 0$ if both players are active consumers and zero otherwise.

At a constant rate, $\nu > 0$ a potential consumer considers the possibility of adopting the new technology. If this were the case, the player uses a myopic best response to update her strategy. Thus, the player compares the benefits obtained next period in the case of adopting with those obtained in case of remaining as a potential consumer. We can think of two different settings:

- *Case 1 (absolute dependence)*

In this formulation, players interact with all their neighbors every period. Heuristically, this implies agents are continuously observing at all neighbors and thus benefits are computed as the sum of the benefits obtained from each bilateral interaction. Hence, a potential consumer i with connectivity k_i and with a_i active consumers among her neighbors becomes an active consumer iff,

$$a_i b - c_i \geq 0$$

Consequently, i 's rate of transition from potential to active consumer is the probability that agent i 's cost is below her benefits, i.e.

$$P(\tilde{c}_i \leq a_i b) = f(a_i) = \begin{cases} \frac{b}{C} a_i & \text{if } a_i \leq \frac{C}{b} \\ 1 & \text{if } a_i > \frac{C}{b} \end{cases}$$

Note that, the reverse transition, i.e. from active to potential consumer, is never a best response of the player. Nevertheless, we assume that at a rate $\delta > 0$ the product deteriorates and needs to be replaced. If this were the case, agents have to re-consider the possibility of adopting it or not.

Observe that, here, the diffusion function only depends on the absolute number of active consumers among neighbors. In consequence, two agents with the same number of neighboring active consumers have the same probability of becoming active consumers and this is independent of their respective connectivities. This feature depends crucially on the specific context considered as illustrated through the alternative setting presented below.

- *Case 2 (frequency dependence)*

Assume agents only interact with one of their neighbors in each time period. Moreover, the individual with whom to interact is selected uniformly at random across neighbors every period. If we take any potential consumer i with connectivity k_i and with a_i active consumers among her neighbors. Then, this agent will become an active consumer iff,

$$\frac{a_i}{k_i} b - c_i \geq 0$$

In other words, she computes her expected benefits in case of adopting the product and compares this with her cost.

Consequently, i 's rate of transition from potential to active consumer is the probability that agent i 's cost is below her benefits, i.e.

$$P(\tilde{c}_i \leq \frac{a_i}{k_i} b) = f(a_i, k_i) = \begin{cases} \frac{b}{C} \frac{a_i}{k_i} & \text{if } \frac{a_i}{k_i} \leq \frac{C}{b} \\ 1 & \text{if } \frac{a_i}{k_i} > \frac{C}{b} \end{cases}$$

As before, assume that at a rate $\delta > 0$ an agent needs to replace the product because it is lost or deteriorated.

Note that, in contrast with the previous case, the diffusion function depends both on the absolute number of active consumers among neighbors and on the total number of neighbors. More precisely, it depends on the relative density of active consumers among neighbors. We will see later in the text that the results on the diffusion pattern of the product depend crucially on the setting considered.

In the next section, we return to the general model to study when and how the product spreads in the population. The analytical results of the exact model are extremely complicated and thus will not be tackled in this paper. Nevertheless, to proceed, two complementary approaches can be considered. On the one hand, the analysis of the model can be simplified using the so-called mean field theory. This approach is described and studied in detail in the next section. On the other hand, we can simulate the dynamics in order to obtain numerical approximations of the results for the exact model. This second alternative will be tackled in Section 6 below.

4 Mean field theory

The analytical study of this model can be undertaken in terms of a dynamical *mean-field theory*. Other reports show that mean-field approximations can be expected to give a reasonable guide to the qualitative behavior of complex dynamics.

Before describing the theoretical framework, we will present some additional notation. Let $\rho_k(t)$ be the relative density of active consumers at time t with connectivity k . Consequently, $\rho(t) = \sum_k P(k)\rho_k(t)$ is the relative density of active consumers at time t . From here onwards, the state of the system at any given time t , will be characterized by the profile $(\rho_k(t))_{k \geq 1}$.

Denote by θ to the probability that any given link points to an active consumer. Therefore, the probability that a potential agent with k links has exactly a neighboring active consumers is $\binom{k}{a} \theta^a (1 - \theta)^{(k-a)}$ since this event follows a binomial distribution with parameters k and θ . Obviously, there is an approximation inherent in this formulation because we have assumed that θ is the same for all vertices, when in general it too will be dependent on vertex connectivity. This is precisely the nature of a mean-field approximation.

Consider a potential consumer with k neighbors and a active consumers among them. She becomes an active consumer at a rate $\nu f(k, a)$. Thus, the transition rate from potential to active consumer for an agent with connectivity k is given by

$$\tilde{g}_{\nu,k}(\theta) = \sum_{a=0}^k \nu f(k, a) \binom{k}{a} \theta^a (1 - \theta)^{(k-a)}$$

The dynamical mean-field equation can thus be written as,

$$\frac{d\rho_k(t)}{dt} = -\rho_k(t)\delta + (1 - \rho_k(t))\tilde{g}_{\nu,k}(\theta) \quad (1)$$

Roughly speaking, equation (1) says the following: the variation of the relative density of active consumers with k links at time t equals the proportion of potential consumers with k neighbors at time t that become active consumers (i.e. $(1 - \rho_k(t))\tilde{g}_{\nu,k}(\theta)$) minus the proportion of active consumers with k neighbors at time t that become potential consumers (i.e. $\rho_k(t)\delta$).

Assume that the time scale of the dynamics is much smaller than the life-span of the agents in the population; therefore terms reflecting birth or death of individuals are not included. Moreover, several assumptions are implicit in equation (1). First, we assume the size of the population is large, i.e. $n \rightarrow +\infty$. Second, we consider the so-called *homogeneous mixing hypothesis*. This implies, on the one hand, no correlation between the connectivity of connected agents and, on the other hand, an homogeneous distribution of initial adopters in the population. In consequence, the only source of heterogeneity in the population is the connectivity of agents.

After imposing the stationary condition $\frac{d\rho_k(t)}{dt} = 0$ in equation (1) for all $k \geq 1$, the equation, valid for the behavior of the system at large times is,

$$\rho_k = \frac{g_{\lambda,k}(\theta)}{1 + g_{\lambda,k}(\theta)} \quad (2)$$

where

$$g_{\lambda,k}(\theta) = \frac{1}{\delta}\tilde{g}_{\nu,k}(\theta) = \sum_{a=0}^k \lambda f(k,a) \binom{k}{a} \theta^a (1-\theta)^{(k-a)}$$

The exact calculation of θ for general networks is a difficult task. However, we can calculate its value for the case of a random network, in which there are no correlations among the connectivities of different nodes.⁴ For this case, it is straightforward to see that,

$$\theta = \frac{1}{\langle k \rangle} \sum_k k P(k) \rho_k \quad (3)$$

where $\langle k \rangle$ is the average connectivity of the network, i.e. $\langle k \rangle = \sum_k k P(k)$.

⁴It is left for further research to allow for correlations in the connectivity of neighbors in this model. Pointing in this direction, there are some papers providing answers to this issue for the SIS model. See Beguña et al. (2002) and Eguíluz et al.(2002). Their main result is that, allowing for first order correlations among connectivity of nodes does not change the conclusions obtained in Pastor-Sarorrás and Vespignani (2000) and thus the epidemic threshold vanishes to zero for scale-free networks.

The system formed by the equations (2) and (3) determine the stationary values for θ and $(\rho_k)_k$. To solve this system, we should simply replace equation (2) in equation (3) and obtain,

$$\theta = H_\lambda(\theta) \tag{4}$$

where

$$H_\lambda(\theta) \equiv \frac{1}{\langle k \rangle} \sum_k k P(k) \frac{g_{\lambda,k}(\theta)}{1 + g_{\lambda,k}(\theta)} \tag{5}$$

The solutions of equation (4) are the stationary values of θ . Note that, these values correspond to the set of fixed points of the function $H_\lambda(\theta)$. Although the exact stationary values for θ are generally difficult to obtain, the main questions raised at the introduction of the paper can be answered by simply analyzing the shape of all the functions in the family $\{H_\lambda(\theta)\}_{\lambda \geq 0}$. Upon replacing θ in equation (2) we also determine the stationary values $(\rho_k)_k$.

5 Results

In what follows we will present the main results of the paper. For concreteness, we will define first the concepts of *sustainable diffusion*, *positive diffusion* and *unique diffusion* of the product.

Definition 2 *Given $P(k)$ and m , we say that there is sustainable diffusion of the product if there exists a locally stable state of the dynamics with a positive fraction of active consumers.*

The concept of stability required in this definition is the standard one. Roughly speaking, a state is stable if it is a stationary state of the dynamics resistant to small perturbations. Notice that, sustainable diffusion implies that, under certain initial conditions, the dynamics converges to a state with a positive fraction of active consumers. Next, we will define the concept of positive diffusion.

Definition 3 *Given $P(k)$ and m , we say that there is positive diffusion of the product if, starting at any initial state $\theta_0 \neq 0$, the dynamics converges to a stable state with a positive fraction of active consumers.*

Notice that, positive diffusion does not imply uniqueness of the non-null stable state. Thus, the long-run behavior of the dynamics can depend on the initial conditions. However, it implies that, if we slightly perturb the initial state with no active consumers, i.e. we introduce a “small” number of initial adopters, the dynamics leads towards a non-null stable state. Finally, the following definition addresses the global behavior of the dynamics.

Definition 4 *Given $P(k)$ and m , we say that there is unique diffusion of the product if there exists a unique stable state of the dynamics with a positive fraction of active consumers.*

In other words, in the case of unique diffusion, the long-run behavior of the dynamics does not depend on the initial conditions.

It is straightforward to show that the following implications hold;

$$\text{unique diffusion} \Rightarrow \text{positive diffusion} \Rightarrow \text{sustainable diffusion}$$

Notice that, the existence of a non-null solution of equation (4) implies the existence of a non-null stable state θ^* of the dynamics, which also implies sustainable diffusion of the product.

Let $\rho_k(\lambda)$ be a function that provides for every given value of the effective spreading rate $\lambda \geq 0$, the relative density of active consumers with connectivity k predicted in the long-run of the dynamics, when the initial state is taken to be infinitesimally close to the one with no active consumers. Moreover, let $\rho(\lambda) = \sum_k P(k)\rho_k(\lambda)$ be the *degree of diffusion function*.

The aim of this section is to describe in some detail the relationship between the connectivity distribution of the network $P(k)$ and the mechanism of diffusion m with the spreading behavior of the product. It is straightforward to show that, given (A-1) the state with no active consumers ($\theta = 0$) is stationary. Thus, to spread the product in the population there must be an initial shock of active consumers. This section analyzes a situation where the initial state of the dynamics is such that there is a “small” proportion of initial adopters, i.e. $\theta_0 \sim 0$. One interpretation for this is that the firm interested in the diffusion of the product initially gives it “for free”. It is reasonable to assume that the firm is going to choose a small number of initial adopters and then rely on the contagion process for the diffusion of it to a larger fraction of agents. Given the nature of the question, we will first focus on the concept of *positive diffusion* defined above. *Unique* and *sustainable diffusion*, will be studied later in the paper.

Theorem 5 *Given a network with connectivity distribution $P(k)$, and a diffusion function $f(k, a)$ satisfying (A-1), there exists a threshold for the effective spreading rate $\lambda_p = \frac{\langle k \rangle}{\sum_k k^2 P(k) f(k, 1)}$ such that, there is positive diffusion of the product if and only if $\lambda > \lambda_p$.*

A detailed proof of the Theorem is presented in the Appendix. The sketch of the proof, however, is the following. For every value of $\lambda \geq 0$, the stationary states of the dynamics are given by the fix points of $H_\lambda(\theta)$. Notice that, assumption (A-1) implies that $H_\lambda(0) = 0$ and therefore, as mentioned above, the state $\theta = 0$ is stationary. Since $g_{\lambda, k}(\theta) \geq 0$ then $0 \leq H_\lambda(\theta) < 1$. In particular, this implies that, positive diffusion occurs if and only if the state $\theta = 0$ is unstable (i.e. there exist an $\epsilon > 0$ such that $H_\lambda(\theta) > \theta$ for all $\theta \in (0, \epsilon)$) or equivalently $\frac{dH_\lambda(\theta)}{d\theta} \Big|_{\theta=0} > 1$. The threshold is obtained simply by solving for λ in the previous equation with the equality condition.

Several interesting points follow from this result. The threshold that determines the diffusion of the product, depends both on the connectivity distribution of the network (i.e. $P(k)$) and on the particular diffusion function considered (i.e. $f(k, 1)$). Specifically, in order to assess the existence or not of some positive prevalence, it is enough to consider what happens in a neighborhood with only one active agent. As highlighted above, this is merely a consequence of the fact that, for positive diffusion to occur, the state with no active consumers has to be unstable. Notice that, if $\lambda > \lambda_p$ then, in the long-run, the product spreads and becomes persistent in a fraction of the population. The degree of the diffusion, however, might depend on the initial conditions. If, on the contrary, we assume $\lambda \leq \lambda_p$ then, if there is only a small fraction of initial adopters, in the long-run, the product will disappear from the market. In other words, we either never reach a state with a positive fraction of active consumers or, if we do, it must be because there is a sufficiently high “stock” of initial adopters.

The following corollary is obtained directly from the above result.

Corollary 6 *If the transition rate from potential to active consumer is independent of the connectivity of the agent (i.e. $f(k, a) = f(k', a) \equiv f(a) \forall k, k' \geq 0$) then the threshold is $\lambda_p = \frac{1}{f(1)} \frac{\langle k \rangle}{\langle k^2 \rangle}$.*

One of the main conclusions obtained from Corollary 1 is that the threshold depends on the connectivity distribution $P(k)$. In particular, it depends on the ratio between its first and second order moments. The examples below illustrate the main insights of this result. Consider three type of networks - scale-free, homogeneous and Poisson- and assume they have the same average connectivity.

1 Scale-free networks

Scale-free networks are characterized by having a power-law connectivity distribution. In particular,

$$P(k) \propto k^{-\gamma}$$

where γ ranges between 2 and 3. This property implies that there exists a significant proportion of agents with large connectivity with respect to the average ($\langle k \rangle$). Hence, the variance of the connectivity of agents tends to infinity in parallel with the size of the population (i.e. $\langle k^2 \rangle \rightarrow \infty$ when $n \rightarrow +\infty$). These agents behave as “hubs” and are capable of spreading the product quickly. Consequently, the threshold for the spreading rate tends to 0. In other words, no matter how small the spreading rate is, positive diffusion of the product in the population will always occur.⁵

2 Homogeneous networks

⁵This result is an extension of the one obtained by Pastor-Satorras and Vespignani (2000) for linear diffusion functions with the absence of neighborhood effects.

Homogeneous networks are such that all nodes have approximately the same connectivity. In particular,

$$P(k) \sim \begin{cases} 0 & \text{if } k \neq \langle k \rangle \\ 1 & \text{if } k = \langle k \rangle \end{cases}$$

Since the variance in the connectivity of nodes is approximately zero, the threshold for the spreading rate is roughly $\frac{1}{f(1)} \frac{1}{\langle k \rangle}$. In homogeneous networks there exists a positive threshold that separates the values of the spreading rate for which the product spreads from those for which it does not. This threshold is inversely proportional to the average connectivity of the network. Moreover, it depends on the transition rate from potential to active consumer when only one neighbor is consuming the product. Consequently, there are different diffusion functions that provide the same threshold. For example, consider $f_1(a) = a^{1/2}$, $f_2(a) = a$ and $f_3(a) = a^2$. All of them have the same threshold. However, it is worthwhile mentioning that, whenever the spreading rate of the product is higher than the corresponding threshold, the degree of the diffusion, is higher for f_2 than for f_1 and higher for f_3 than for f_2 .⁶

3 Poisson networks

Poisson networks are characterized by having a Poisson connectivity distribution. In particular,

$$P(k) = \frac{1}{k!} e^{-\langle k \rangle} \langle k \rangle^k$$

It is straightforward to show that the threshold for Poisson networks is in between the threshold for scale-free and homogeneous networks. Thus, the following holds:

$$\lambda_p^{SF} < \lambda_p^P < \lambda_p^H$$

In general, any other type of network with the same average connectivity, has a threshold that lies in between these two. The reason for this is the following: the variance of the connectivity distribution $P(k)$ is given by $var(k) = \langle k^2 \rangle - \langle k \rangle^2$, thus $\langle k^2 \rangle = var(k) + \langle k \rangle^2$. Since λ_p is inversely proportional to the second order moment, if we compare two networks with the same average connectivity, the one with the highest variance has the lowest threshold.

The next corollary is also a consequence of Theorem 5. Consider now the case where the neighborhood considerations is affected by the neighborhood size. Despite its significance, this issue has been ignored in the epidemiology literature. In the present framework the network represents the system of social contacts. Therefore, it is plausible to assume that agents have a limited amount of time to spend in social acquaintances. Consequently, the time spent with each

⁶This is because $f_1(a) \leq f_2(a) \leq f_3(a)$ for all $a \geq 0$.

of her neighbors (or intensity of each of her interactions) decreases in parallel with the total number of neighbors. In this respect, for example, the effect of having one active neighbor for a very well connected agent is not the same than for an agent with few neighbors. A natural, although extreme, candidate to consider as a diffusion function would be the relative number $\frac{a}{k}$ of active neighbors. This corollary shows that, in this case, the critical spreading rate is equal to unity, independently of what might be the underlying network (scale-free, Poisson, homogeneous, etc.).

Corollary 7 *If $f(k, a) = \frac{a}{k}$ then $\lambda_p = 1$.*

Hence, for scale-free networks, in this case, there exists a positive threshold determining under which conditions there is prevalence of the product in the population. The intuition behind this result is that, although scale-free network have a significant fraction of hubs that facilitate the contagion of the product, this effect is cancelled out by the fact that a hub agent is very difficult to “convince” given that what matters here is the relative density of active neighbors.

Up to now we have analyzed whether there is or not prevalence of the product in the population when the initial state is “close” to the state with no active consumers. We now want to go one step beyond and study more general properties of the dynamics.

5.1 Concave diffusion functions

In this section we find conditions over the diffusion mechanisms that guarantee a unique long-run behavior of the dynamics. In other words, we analyze the convergence of the dynamics independently of initial conditions. Consider a diffusion function satisfying an additional assumption. For all $k \geq 1$, $f(k, a)$ as a function of a is (weakly) *concave*. In other words, the following must hold:

$$f(k, a) - f(k, a - 1) \geq f(k, a + 1) - f(k, a) \text{ for all } 0 < a < k \quad (\text{A-2})$$

Hence, for any given agent, adding one more active consumer among her neighbors has an impact over her probability of obtaining the product, which is (weakly) decreasing with respect to the existing number of active consumers among her neighbors.

The following proposition determines the threshold for unique diffusion of the product.

Proposition 8 *Given a network with connectivity distribution $P(k)$, and a diffusion function $f(k, a)$ satisfying (A-1) and (A-2) there exists a threshold for the effective spreading rate $\lambda_u = \frac{\langle k \rangle}{\sum_k k^2 P(k) f(k, 1)}$ such that, there is unique diffusion of the product if and only if $\lambda > \lambda_u$. Moreover, if $\lambda \leq \lambda_u$ the dynamics converge to the state with no active consumer.*

A detailed proof of Proposition 8 is presented in the Appendix. The sketch of the proof is the following. Assumption (A-2) implies $H_\lambda(\theta)$ is concave for all $\lambda \geq 0$ (this is proved in the Appendix). Thus, as illustrated by Figure ??, if $\frac{dH_\lambda(\theta)}{d\theta}\big|_{\theta=0} > 1$ there is a non-null stationary state which is globally stable.⁷ However, if $\frac{dH_\lambda(\theta)}{d\theta}\big|_{\theta=0} \leq 1$ then $\theta = 0$ is the unique stable state. Let λ_s denote the threshold for sustainable diffusion. Note that, we obtain:

$$\lambda_s = \lambda_p = \lambda_u$$

As a consequence of this result we have the following corollaries.

Corollary 9 *Given a network with connectivity distribution $P(k)$, and a diffusion function $f(k, a)$ satisfying (A-1) and (A-2), then the degree of diffusion function $\rho(\lambda)$ is continuous.*

The proof of this result is straightforward. Notice that, as aforementioned, for all $\lambda \geq 0$, $H_\lambda(\theta)$ is concave. Moreover, for all $\theta \in [0, 1]$, $H_\lambda(\theta)$ as a function of λ is increasing and continuous. This implies that, the solution of equation (4), i.e. $\theta(\lambda)$, as a function of λ is also continuous, and by the same token the degree of diffusion function $\rho(\lambda)$ is continuous as well.

As an example, consider a diffusion function satisfying (A-1) and (A-2) and such that it is independent of the connectivity of agents (i.e. $f(k, a) = f(k', a) \equiv f(a) \forall k, k' \geq 0$). Then, the degree of diffusion function for the three types of networks aforementioned -scale-free, homogeneous and Poisson- is continuous (see the graphs represented in Figure 1). Notice that, for low values of λ the degree of diffusion is higher for scale-free networks than for Poisson networks and higher for Poisson than for homogeneous networks.

Corollary 10 *If $f(k, a) = \frac{a}{k}$ then $\rho(\lambda) = \frac{\lambda}{1-\lambda}$ if $\lambda > \lambda_p$ and $\rho(\lambda) = 0$ otherwise.*

The proof of this result is attained by simply substituting $f(k, a) = \frac{a}{k}$ in the expression for $g_{\lambda,k}(\theta)$. That is,

$$g_{\lambda,k}(\theta) = \frac{1}{k} \sum_{a=0}^k \lambda a \binom{k}{a} \theta^a (1-\theta)^{(k-a)} = \frac{1}{k} \lambda \theta k = \lambda \theta$$

Notice that, $g_{\lambda,k}(\theta)$ does not depend on k . Then, replacing $g_{\lambda,k}(\theta)$ in equation (5) the following holds:

$$H_\lambda(\theta) = \frac{\lambda \theta}{1 + \lambda \theta} \tag{6}$$

⁷A state θ is globally stable if for any initial state $\theta_0 \in (0, 1)$, the dynamics converges to this state.




Figure 1: Degree of diffusion function for scale-free, homogeneous and Poisson networks when $f(k, a) = f(k', a) \equiv f(a) \forall k, k' \geq 0$.

It can be easily shown that, equation (6) has a unique non-null solution when $\lambda > 1$ which is $\theta^* = \frac{\lambda-1}{\lambda}$. Thus, after simple algebraic operations, the degree of diffusion is:

$$\rho(\lambda) = \begin{cases} 0 & \text{if } \lambda \leq 1 \\ \frac{\lambda}{1-\lambda} & \text{if } \lambda > 1 \end{cases}$$

It is worthwhile mentioning that, in this case, the degree of diffusion function does not depend on the connectivity distribution of the network (see Figure 2).

5.2 Other diffusion functions: phase transition

Assume that for all $k \geq 1$, $f(k, a)$ as a function of a is (weakly) *convex*. The following must hold:

$$f(k, a+1) - f(k, a) \geq f(k, a) - f(k, a-1) \text{ for all } 0 < a < k$$

Hence, for any given agent, adding one more active consumer among her neighbors has an impact over her probability of obtaining the product, which is (weakly) increasing with respect to the existing number of active consumers among her neighbors.

Due to its operational complexity, general results for convex diffusion functions are not easy to obtain. In what follows, we analyze an illustrative example to highlight the difference with the results obtained for concave diffusion functions. The diffusion function considered is,

$$f(k, a) = \left(\frac{a}{k}\right)^2 \quad (7)$$




Figure 2: Degree of diffusion function for scale-free, homogeneous and Poisson networks when $f(k, a) = \frac{a}{k}$

This contagion mechanism takes into account the relative density of active consumers among an agent's neighbors in a convex way. The threshold for positive diffusion in this case is equal to the average connectivity, $\lambda_p = \langle k \rangle$ (see Theorem 5). To study the threshold for unique and sustainable diffusion we will analyze the shape of the family of functions $\{H_\lambda(\theta)\}_{\lambda \geq 0}$. Note that, in this case, function $g_{\lambda, k}(\theta)$ is equal to the expression,

$$g_{\lambda, k}(\theta) = \lambda \sum_{a=0}^k \left(\frac{a}{k}\right)^2 \binom{k}{a} \theta^a (1-\theta)^{(k-a)} = \frac{\lambda}{k^2} E[\chi^2]$$

where χ is a random variable that follows a binomial distribution with parameters θ, k . That is, $\chi \sim \text{Bin}(k, \theta)$. Therefore, the following holds,

$$E[\chi^2] = \text{Var}[\chi] + E[\chi]^2 = k\theta(1-\theta) + (\theta k)^2 = (k^2 - k)\theta^2 + k\theta$$

and thus,

$$g_{\lambda, k}(\theta) = \frac{\lambda}{k^2} ((k^2 - k)\theta^2 + k\theta) = \frac{\lambda}{k} ((k-1)\theta^2 + \theta)$$

The function $H_\lambda(\theta)$ in this case has the form,

$$H_\lambda(\theta) \equiv \frac{1}{\langle k \rangle} \sum_k k P(k) \frac{\frac{\lambda}{k} ((k-1)\theta^2 + \theta)}{1 + \frac{\lambda}{k} ((k-1)\theta^2 + \theta)}$$

The shape of $H_\lambda(\theta)$ depends crucially on $P(k)$. Therefore, for the sake of simplicity, we will focus on two specific types of networks: (i) a scale-free network with $\gamma = 3$, i.e. $P(k) \propto k^{-3}$ and (ii) an homogeneous network.

It is straightforward to show that, if $\langle k \rangle$ is sufficiently high (higher than 5) for both cases (i) and (ii) the family of functions $\{H_\lambda(\theta)\}_{\lambda \geq 0}$ exhibits the following

pattern. For low values of λ the function is convex. As λ increases the function has an *S-shape*, i.e. convex for low values of θ and concave for high values of θ . Finally, if λ is sufficiently high $H_\lambda(\theta)$ is concave. For simplicity, a family of functions $\{H_\lambda(\theta)\}_{\lambda \geq 0}$ satisfying this property is referred as an *S-shape family*.

Note that, if $\{H_\lambda(\theta)\}_{\lambda \geq 0}$ is an *S-shape family* of functions, the following holds:

$$\text{positive diffusion} \Leftrightarrow \text{unique diffusion}$$

and thus $\lambda_p = \lambda_u$.

Moreover, there exists a threshold $\tilde{\lambda}$ (*concavity threshold*) for the spreading rate such that, if $\lambda > \tilde{\lambda}$ then $H_\lambda(\theta)$ is concave. This value is implicitly obtained by the expression $H_\lambda''(0) = 0$. After some simple algebraic operations we obtain the following expression,

$$H_\lambda''(0) = \frac{\lambda}{\langle k \rangle} \sum_k P(k) \frac{2k(k-1) - 2\lambda k^2}{k}$$

It is straightforward to show that, the two types of networks considered satisfy,

$$H_{\lambda_p}''(0) > 0$$

This implies, in particular, that the threshold for positive diffusion is below the threshold for concavity, i.e. $\lambda_p < \tilde{\lambda}$. In other words, the function associated with the threshold for positive (or unique) diffusion, has an *S-shape*. Therefore,

$$\lambda_s < \lambda_p$$

The main consequences of this result are the following:

- If $\lambda_s < \lambda < \lambda_p$, there are two different (non-null) stationary states of the dynamics; an unstable state, denoted by θ_1^* and a stable state denoted by θ_2^* . The convergence to the stable state depends crucially on the initial conditions since initial conditions above θ_1^* would lead the dynamics towards θ_2^* , whereas initial conditions below θ_1^* would lead the dynamics towards $\theta^* = 0$. Two effects take place when the spreading rate becomes higher. On the one hand, the proportion of active consumers in the non-null stable state (θ_2^*) increases. On the other hand, its basin of attraction also becomes larger.
- There is a *phase transition* or discontinuity in the degree of diffusion function. In other words, when the spreading rate λ is “slightly” above the threshold λ_p , the degree of diffusion $\rho(\lambda)$ is significantly positive.
- The effect of varying the value of the spreading rate λ can be analyzed using a different approach. Assume that the contagion dynamics has already reached a certain stable state. Where would the dynamics converge if there was an increase or decrease of the effective spreading rate? In other




Figure 3: Hysteresis phenomenon for homogeneous and scale-free networks (with $\gamma = 3$) and $f(k, a) = (\frac{a}{k})^2$

words, taking as initial condition the previously established stable state, what would be the new long-run prediction of the dynamics? As illustrated by the graph represented in Figure 3 if the spreading rate increases (upward arrows in the figure) then the long-run behavior of the dynamics would coincide with the one exhibited by function $\rho(\lambda)$, thus, having a discontinuity at $\lambda = \lambda_p$. However, if the spreading rate decreases (downward arrows in the figure), the degree of diffusion will continue to be positive until λ reaches the threshold for sustainable diffusion λ_s . The existence of two different thresholds depending on the direction of the spreading rate is a well-known occurrence, present in many other phenomena referred as *hysteresis*.

6 Simulations

In this section we have developed some simulations to test the validity of the mean-field approximations used throughout the paper. We have generated two different random networks in terms of their connectivity distributions: (i) a scale-free (specifically, $P(k) \propto k^{-3}$) network (ii) a Poisson network.⁸ Both networks have a total of 1000 nodes and an average connectivity of approximately 9 neighbors per node. We have considered the discrete version of the continuous dynamics used to derive the theoretical results. In this respect, we have assumed that, in every period one (and only one) agent is chosen to revise her “strategy”. For the sake of concreteness, we have focused on testing the contents of Corollaries 6 and 7.

All figures presented below have in common the following characteristics. We

⁸Both of these networks were generated using the program *Pajek*, software package for Large Network Analysis.




Figure 4: Number of active consumers $n(t)$ for the scale-free network when $f(k, a) = a$, $\lambda = 0.05, 0.5$ and 5 , and $t \in [1, 10^5]$

represent how the number of active consumers ($n(t)$, ordinate) changes as a function of time (t periods, abscissa) at different values of the spreading rate λ . The data are the average of 40 simulations. For each simulation, the initial condition is randomly chosen such that individuals are active in round $t = 1$ with probability 0.01.

For Corollary 6, we considered a diffusion function that depends linearly on the absolute number of active consumers in the neighborhood of an agent. Specifically, $f(a) = a$. We wanted to test if the threshold for the scale-free network tends to zero and if it is lower than the threshold for the Poisson network.

The graph in Figure 4 represents how the number of active consumers changes over a total of 10^5 periods for the scale-free network at three different values of the spreading rate ($\lambda = 0.05, 0.5$ and 5 ; green, blue and red line, respectively). Observe that, as expected, the degree of diffusion is higher the higher the spreading rate is. Moreover, as the period increase, the number of active consumers increases as well. Also note that, for $\lambda = 0.05$ there is prevalence of the product in the long-run, thus this could indicate that, in this case, the threshold for positive diffusion tends to zero.

The graph in Figure 5 represents how the number of active consumers changes over a total of 5×10^4 periods for the Poisson network at three different values of the spreading rate ($\lambda = 0.05, 0.5$ and 5 ; green, blue and red line, respectively). In contrast with the previous case, for $\lambda = 0.05$ there is no prevalence of the product in the long-run. This indicates that the threshold for positive diffusion in the Poisson network is above 0.05 and thus higher than for the scale-free network.




Figure 5: Number of active consumers $n(t)$ for the Poisson network when $f(k, a) = a$, $\lambda = 0.05, 0.5$ and 5 , and $t \in [1, 5 \times 10^4]$

These simulations also provide relevant information concerning the rate of convergence to the stationary state, an aspect of the dynamics that has not been addressed in the theoretical analysis. Observe that, there is a significant evidence reflecting a higher rate of convergence in the Poisson network than in the scale-free network.

For Corollary 7, we considered the diffusion function $f(k, a) = \frac{a}{k}$. We wanted to test if the diffusion threshold for the scale-free and Poisson networks is equal to 1.

The graph in Figure 6 represents how the number of active consumers changes over a total of 10^5 periods for the scale-free network at four different values of the spreading rate ($\lambda = 0.8, 1, 1.2$ and 1.4 ; yellow, green, blue and red line, respectively). Notice that, the threshold for positive diffusion is close to 1 (between $\lambda = 1$ and $\lambda = 1.2$) and thus significantly higher than the threshold obtained for the diffusion function considered previously as predicted by the theoretical results.

The last set of simulations, presented in Figure 7 represent how the number of active consumers changes over a total of 10^5 periods for the Poisson network at four different values of the spreading rate ($\lambda = 0.8, 1, 1.2$ and 1.4 ; yellow, green, blue and red line, respectively). The threshold for positive diffusion is approximately at $\lambda = 1$, thus, in this case, “close” to the threshold obtained for the scale-free network.




Figure 6: Number of active consumers $n(t)$ for the scale-free network when $f(k, a) = \frac{a}{k}$, $\lambda = 0.8, 1, 1.2$ and 1.4 , and $t \in [1, 10^5]$




Figure 7: Number of active consumers $n(t)$ for the Poisson network when $f(k, a) = \frac{a}{k}$, $\lambda = 0.8, 1, 1.2$ and 1.4 , and $t \in [1, 10^5]$

7 Conclusion

The objective of this paper is to analyze how the diffusion of a new product or technology takes place on a social complex network. The network is characterized by one of its large-scale statistical properties -the connectivity distribution- rather than by a specific geometric form (such as lines, circles, lattices and so forth). A wide class of diffusion dynamics (or mechanisms) has been considered. In all of them, the probability of agents adopting the product depends on the product's spreading rate and the behavior of the agents' closest neighbors.

The main contribution of this paper is to *characterize* the contagion (diffusion) threshold in terms of the properties of the network and the diffusion mechanism. One of the principal findings is that, the threshold depends crucially on the network considered when the intensity of each interaction is assumed to be independent of the neighborhoods size. More specifically, the higher the variance in the connectivity distribution of the network, the lower the threshold. This implies, in particular, that scale-free networks are optimal for spreading the product. In contrast with this result, if the diffusion mechanism considered is such that the intensity of each interaction is inversely proportional to the neighborhoods size, the variance of the connectivity distribution of the network has no effect over the threshold. In other words, all networks (with the same average connectivity) have the same positive threshold. Finally, we also show that, for some particular diffusion mechanisms, there is a *phase transition* in the degree of the diffusion function. In other words, there is a discontinuity in the fraction of active consumers in the contagion threshold.

The simulations presented in the last section of the paper show that, the theoretical results, obtained using mean field approximations provide a reasonable guide of the qualitative properties and long-run predictions of the diffusion dynamics.

8 Appendix

Proof of Theorem 5:

Assumption (A-1) implies that $\theta = 0$ is a stationary state of the dynamics (i.e. $H_\lambda(0) = 0$). In addition, $0 \leq H_\lambda(\theta) \leq 1$ for all $\theta \in [0, 1]$ since $g_{\lambda,k}(\theta) \geq 0$. Therefore, given $\lambda \geq 0$, there exists a non-null stable state of the diffusion dynamics if and only if $\theta = 0$ is *unstable*. Formally, this means that there must exist an $\varepsilon > 0$ such that, $\theta < H_\lambda(\theta)$ for all $\theta \in (0, \varepsilon)$. Equivalently,

$$\left. \frac{dH_\lambda(\theta)}{d\theta} \right|_{\theta=0} \equiv H'_\lambda(0) > 1 \quad (8)$$

Let us now calculate the exact value for the threshold. Recall that,

$$H_\lambda(\theta) \equiv \frac{1}{\langle k \rangle} \sum_k k P(k) \frac{g_{\lambda,k}(\theta)}{1 + g_{\lambda,k}(\theta)} \quad (9)$$

where

$$g_{\lambda,k}(\theta) = \sum_{a=0}^k \lambda f(k, a) \binom{k}{a} \theta^a (1-\theta)^{(k-a)}.$$

Then, using equation (9) we express $H'_\lambda(\theta)$ as follows,

$$H'_\lambda(\theta) = \frac{1}{\langle k \rangle} \sum_k k P(k) \frac{g'_{\lambda,k}(\theta)}{(1 + g_{\lambda,k}(\theta))^2} \quad (10)$$

where

$$g'_{\lambda,k}(\theta) = \sum_{a=0}^k \lambda f(k, a) \binom{k}{a} (a\theta^{a-1}(1-\theta)^{(k-a)} - \theta^a(k-a)(1-\theta)^{(k-a-1)}). \quad (11)$$

If $\theta = 0$ is substituted in equation (10) we obtain that,

$$H'_\lambda(0) = \frac{\lambda \sum_k k^2 P(k) f(k, 1)}{\langle k \rangle} > 1 \Leftrightarrow \lambda > \frac{\langle k \rangle}{\sum_k k^2 P(k) f(k, 1)}.$$

□

Proof of Proposition 8:

It is straightforward to show that, for every given $0 \leq \theta \leq 1$, $H_\lambda(\theta)$ as a function of λ , is increasing. That is,

$$H_\lambda(\theta) \leq H_{\lambda'}(\theta) \Leftrightarrow \lambda \leq \lambda'. \quad (12)$$

Let us show that, given $\lambda \geq 0$, assumption (A-2) implies $H_\lambda(\theta)$ is concave for all $\theta \in [0, 1]$. Notice that,

$$H''_\lambda(\theta) = \frac{1}{\langle k \rangle} \sum_k k P(k) \frac{g''_{\lambda,k}(\theta)(1 + g_{\lambda,k}(\theta)) - 2(g'_{\lambda,k}(\theta))^2}{(1 + g_{\lambda,k}(\theta))^3}.$$

Thus, it is enough to prove that $g''_{\lambda,k}(\theta) \leq 0$, since this would imply that $H''_\lambda(\theta) \leq 0$ as well. If we group the coefficients of the same polynomial on θ in equation (11) we obtain,

$$g'_{\lambda,k}(\theta) = \sum_{a=0}^{k-1} \lambda (-f(k, a) \binom{k}{a} (k-a) + f(k, a+1) \binom{k}{a+1} (a+1)) \theta^a (1-\theta)^{(k-a-1)}. \quad (13)$$

Note that, the coefficients of $f(k, a)$ and $f(k, a+1)$ are equal but with opposite sign since

$$\binom{k}{a} (k-a) = \binom{k}{a+1} (a+1) = \frac{k!}{a!(k-a-1)!}.$$

Therefore, we can simplify equation (13) as follows:

$$g'_{\lambda,k}(\theta) = \sum_{a=0}^{k-1} \frac{k!}{a!(k-a-1)!} \lambda(f(k, a+1) - f(k, a)) \theta^a (1-\theta)^{(k-a-1)}$$

whose second derivative is,

$$g''_{\lambda,k}(\theta) = \sum_{a=0}^{k-1} \frac{k!}{a!(k-a-1)!} \lambda(f(k, a+1) - f(k, a)) \\ (a\theta^{a-1}(1-\theta)^{(k-a-1)} - \theta^a(k-a-1)(1-\theta)^{(k-a-2)})$$

Again, grouping the coefficients of the same polynomials on θ we obtain,

$$g''_{\lambda,k}(\theta) = \sum_{a=0}^{k-1} \lambda \binom{k}{a+1} (k-a-1)(a+1)(f(k, a+2) - f(k, a+1)) \\ - \binom{k}{a} (k-a)(k-a-1)(f(k, a+1) - f(k, a)) \theta^a (1-\theta)^{(k-a-2)}.$$

Since,

$$\binom{k}{a+1} (k-a-1)(a+1) = \binom{k}{a} (k-a)(k-a-1) = \frac{k!}{a!(k-a-2)!}$$

we thus can simplify $g''_{\lambda,k}(\theta)$ as follows:

$$g''_{\lambda,k}(\theta) = \sum_{a=0}^{k-2} \frac{k!}{a!(k-a-2)!} \lambda((f(k, a+2) - f(k, a+1)) \\ - (f(k, a+1) - f(k, a)) \theta^a (1-\theta)^{(k-a-2)}).$$

To conclude, observe that assumption (A-2) implies that $g''_{\lambda,k}(\theta) \leq 0$.

It is straightforward to show that, the concavity of $H_\lambda(\theta)$ for all $\lambda \geq 0$, assumption (A-1) and condition (12) completes the proof. \square

References

- [1] Anderlini, L. and A. Ianni (1996): "Path Dependence and Learning from Neighbors", *Games and Economic Behavior*, 13, 141-177.
- [2] Barabási, A. and R. Albert (1999): "Emergence of Scaling in Random Networks", *Science* Vol. 286, 509-512.
- [3] Barabási, A., R. Albert and H. Jeong (2000): "Scale-free Characteristics of Random Networks: the Topology of the World-Wide Web", *Physica A*, 281, 2115.

- [4] Blume, L. (1995): “The Statistical Mechanics of Best-Response Strategy Revision”, *Games and Economic Behavior* 11, 111-145.
- [5] Boguña, M. and R. Pastor-Satorras (2002): “Epidemic Spreading in Correlated Complex Networks”, e-print cond-mat/0205621.
- [6] Ellison, G. (1993): “Learning, Local Interaction, and Coordination”, *Econometrica* 61, 1047-1071.
- [7] Erdos P., A. Reny (1959): “On Random Graphs”, *Publ. Math. Debrecen*, 6, 290-297.
- [8] Equiluz, V.M. and Konstantin, K. (2002): “Epidemic Threshold in Structured Scale-Free Networks”, *Physical Review Letters*, Vol. 89, No. 10.
- [9] Faloutsos, M., P. Faloutsos and C. Faloutsos (1999): “On Power-Law Relationships of the Internet Topology”, *Computer Communications Rev.*, 29, 251-262.
- [10] Goyal, S. (1996): “Interaction Structure and Social Change”, *Journal of Institutional and Theoretical Economics* 152, 472-495.
- [11] Lijeros, F., C.R. Edling, L.A.N. Amaral, H.E. Stanley and Y. Aberg (2001): “The Web of Human Sexual Contacts”, *Nature* 411, 907-908.
- [12] Lloyd, A. and R. May (2001): “How Viruses Spread Among Computers and People”, *Science* Vol. 292, 1316-1317.
- [13] Morris, S. (2000): “Contagion”, *Review of Economic Studies* 67, 57-78.
- [14] Newman, M. E. J. (2003): “The Structure and Function of Complex Networks”, *SIAM Review*, Vol. 45, 2, 167-256.
- [15] Pastor-Satorras, R. and A. Vespignani (2000): “Epidemic Spreading in Scale-Free Networks”, *Physical Review Letters* 86, No. 14, 3200-3203.
- [16] Seth, G. (2002): “Unleashing the Ideavirus”, The Free Press.
- [17] Yook, S.H., H. Jeong and A.L. Barabási (2001): “Modeling the Internet’s Large-Scale Topology” e-print cond-mat/0107417.
- [18] Young, P. (2002): “The Diffusion of Innovation in Social Networks”, in L. Blume and S. Durlauf, (ed), *The Economy as a Complex Evolving System*, vol. III, Oxford University Press, forthcoming.