

Manuscript Number:

Title: Oral health, dental care and mouthwash associated with upper aerodigestive tract cancer

Article Type: Original Research article

Keywords: Head and neck neoplasms; oral health; oral hygiene; genetic polymorphisms.

Corresponding Author: Prof. Dr. Wolfgang Ahrens,

Corresponding Author's Institution: Leibniz-Institute for Prevention Research and Epidemiology – BIPS GmbH

First Author: Wolfgang Ahrens

Order of Authors: Wolfgang Ahrens; Hermann Pohlabein; Ronja Foraita; Mari Nelis; Pagona Lagiou; Areti Lagiou; Christine Bouchardy; Alena Slamova; Miriam Schejbalova; Franco Merletti; Lorenzo Richiardi; Kristina Kjaerheim; Antonio Agudo; Xavier Castellsague; Tatiana V Macfarlane; Gary J Macfarlane; Yuan-Chin Amy Lee; Renato Talamini; Luigi Barzan; Cristina Canova; Lorenzo Simonato; Peter Thomson; Patricia A McKinney; Alex D McMahon; Ariana Znaor; Claire M Healy; Bernard E McCartan; Andres Metspalu; Manuela Marron; Mia Hashibe; David I Conway; Paul Brennan

Abstract: Objective: To evaluate the association of oral health (OH), dental care (DC) and mouthwash with tumors of the upper-aerodigestive tract (UADT), and to which extent enzymes involved in the metabolism of alcohol modify the effect of alcohol-based mouthwashes.'

Materials and Methods: Subjects were interviewed about OH and DC behaviors. Mouthwash use was analyzed in relation to alcohol dehydrogenase genes (ADH1B,ADH1C,ADH7) and one aldehyde dehydrogenase gene (ALDH2) known to be associated with head and neck cancers. Adjusted odds ratios (ORs) and 95%-confidence intervals [CI] were estimated with multiple logistic regression models.

Results: ORs of low versus high scores of DC and OH were 2.36[CI=1.51-3.67] and 2.22[CI=1.45-3.41], respectively, for all UADT sites combined. The OR for use of mouthwash 3 or more times/day was 3.23[CI=1.68-6.19]. The OR for the rare variant ADH7 coding for fast ethanol metabolism was lower in mouthwash-users (OR=0.53[CI=0.35-0.81]) as compared to never-users (OR=0.97[CI=0.73-1.29]) indicating effect modification (p=0.005) while no relevant differences were observed between users and non-users for the variant alleles of ADH1B, ADH1C or ALDH2.

Conclusions: Poor OH and DC are independent risk factors for UADT. Whether mouthwash use may entail some risk through the alcohol content in most formulations on the market remains to be clarified.

## **Oral health, dental care and mouthwash associated with upper aerodigestive tract cancer**

Wolfgang Ahrens<sup>1,2</sup>, Hermann Pohlabein<sup>3</sup>, Ronja Foraita<sup>3</sup>, Mari Nelis<sup>4,5,6</sup>, Pagona Lagiou<sup>7</sup>, Areti Lagiou<sup>8</sup>, Christine Bouchardy<sup>9</sup>, Alena Slamova<sup>10</sup>, Miriam Schejbalova<sup>10</sup>, Franco Merletti<sup>11</sup>, Lorenzo Richiardi<sup>11</sup>, Kristina Kjaerheim<sup>12</sup>, Antonio Agudo<sup>13</sup>, Xavier Castellsague<sup>13</sup>, Tatiana V. Macfarlane<sup>14</sup>, Gary J. Macfarlane<sup>14</sup>, Yuan-Chin Amy Lee<sup>15</sup>, Renato Talamini<sup>16</sup>, Luigi Barzan<sup>17</sup>, Cristina Canova<sup>18</sup>, Lorenzo Simonato<sup>19</sup>, Peter Thomson<sup>20</sup>, Patricia A. McKinney<sup>21,22</sup>, Alex D. McMahon<sup>23</sup>, Ariana Znaor<sup>24</sup>, Claire M. Healy<sup>25</sup>, Bernard E. McCartan<sup>25,26</sup>, Andres Metspalu<sup>4,5</sup>, Manuela Marron<sup>27,28</sup>, Mia Hashibe<sup>27</sup>, David I. Conway<sup>22</sup>, Paul Brennan<sup>27</sup>

<sup>1</sup> Department of Epidemiological Methods and Etiologic Research, Leibniz-Institute for Prevention Research and Epidemiology - BIPS, Bremen, Germany

<sup>2</sup> Institute for Statistics, University Bremen, Bremen, Germany

<sup>3</sup> Department of Biometry and Data Management, Leibniz-Institute for Prevention Research and Epidemiology - BIPS, Bremen, Germany

<sup>4</sup> Institute of Molecular and Cell Biology, University of Tartu, Estonia

<sup>5</sup> Estonian Genome Center and Center of Translational Genomics, University of Tartu, Estonia

<sup>6</sup> Department of Genetic Medicine and Development, University of Geneva Medical School, Geneva, Switzerland

<sup>7</sup> Department of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Athens, Greece

<sup>8</sup> Faculty of Health Professions, Athens Technological Educational Institute, Athens, Greece

<sup>9</sup> Institute of Social and Preventive Medicine, Geneva Cancer Registry, IMSP, University of Geneva, Switzerland

<sup>10</sup> Institute of Hygiene and Epidemiology, First Faculty of Medicine, Charles University of Prague, Prague, Czech Republic

<sup>11</sup> Department of Medical Sciences, Cancer Epidemiology Unit, CeRMS and CPO Piemonte University of Turin, Turin, Italy

<sup>12</sup> Cancer Registry of Norway, Oslo, Norway

- <sup>13</sup> Institut Catala`d'Oncologia (ICO), CIBERESP, IDIBELL Hospitalet de Llobregat, Barcelona, Spain
- <sup>14</sup> School of Medicine and Dentistry, University of Aberdeen, Aberdeen, United Kingdom
- <sup>15</sup> Department of Family and Preventive Medicine, Division of Public Health, School of Medicine, University of Utah, Utah, USA
- <sup>16</sup> Aviano Cancer Centre, Aviano, Italy
- <sup>17</sup> General Hospital of Pordenone, Pordenone, Italy
- <sup>18</sup> Centre for Environment and Health, National Heart and Lung Institute, MRC-HPA, Imperial College London, London, United Kingdom
- <sup>19</sup> Universita degli Studi di Padova, Padova, Italy
- <sup>20</sup> University of Newcastle Dental School, Newcastle, United Kingdom
- <sup>21</sup> Centre for Epidemiology and Biostatistics, University of Leeds, Leeds, United Kingdom
- <sup>22</sup> NHS NSS ISD, Edinburgh, Scotland, United Kingdom
- <sup>23</sup> Veterinary and Life Sciences, College of Medicine, University of Glasgow, Glasgow, United Kingdom
- <sup>24</sup> Croatian National Cancer Registry, Zagreb, Croatia
- <sup>25</sup> Trinity College School of Dental Science, Dublin, Ireland
- <sup>26</sup> School of Medicine, Royal College of Surgeons in Ireland, Dublin, Ireland
- <sup>27</sup> International Agency for Research on Cancer (IARC), Lyon, France
- <sup>28</sup> Institute of Medical Biostatistics, Epidemiology and Informatics (IMBEI), University of Mainz, Mainz, Germany

### **Corresponding authors:**

Wolfgang Ahrens

E-Mail: [ahrens@bips.uni-bremen.de](mailto:ahrens@bips.uni-bremen.de)

Address: Leibniz-Institute for Prevention Research and Epidemiology – BIPS GmbH

Achterstrasse 30,

28359 Bremen, Germany

Tel.: +49 (0)421 218 56820

Fax: +49 (0)421 218 56821

Paul Brennan

International Agency for Research on Cancer (IARC)

150 cours Albert Thomas,  
69008 Lyon, France  
Tel.: +33 (0)4 7273 8391  
Fax: +33 (0)4 7273 8342  
E-mail: brennan@iarc.fr

**Keywords:** head and neck neoplasms; oral health; oral hygiene; genetic polymorphisms.

### **Funding**

This work was supported by the European Community (5th Framework Programme) [grant number QLK1-CT-2001-00182], University of Athens Medical School and Bureau of Epidemiologic Research Academy of Athens, for the Athens center; Padova University (Contract No. CPDA057222) for the Padova center; Compagnia San Paolo, AIRC, and Piedmont Region for the Turin center. The Estonian study was supported by Targeted Financing from Estonian Government: SF0180142, European Union through the European Regional Development Fund in the frame of Centre of Excellence in Genomics and 7FP Project ECOGENE.

**Running head:** Oral health and risk of UADT

**Word count - content:** 3.590 (3.807 including funding, acknowledgements and competing interest)

**Word count - abstract:** 200

**Number of tables:** 5

**Number of figures:** 1

## **ABSTRACT**

### **Objective**

To evaluate the association of oral health (OH), dental care (DC) and mouthwash with tumors of the upper-aerodigestive tract (UADT), and to which extent enzymes involved in the metabolism of alcohol modify the effect of alcohol-based mouthwashes.'

### **Materials and Methods**

Subjects were interviewed about OH and DC behaviors. Mouthwash use was analyzed in relation to alcohol dehydrogenase genes (ADH1B,ADH1C,ADH7) and one aldehyde dehydrogenase gene (ALDH2) known to be associated with head and neck cancers. Adjusted odds ratios (ORs) and 95%-confidence intervals [CI] were estimated with multiple logistic regression models.

### **Results**

ORs of low versus high scores of DC and OH were 2.36[CI=1.51-3.67] and 2.22[CI=1.45-3.41], respectively, for all UADT sites combined. The OR for use of mouthwash 3 or more times/day was 3.23[CI=1.68-6.19]. The OR for the rare variant ADH7 coding for fast ethanol metabolism was lower in mouthwash-users (OR=0.53[CI=0.35-0.81]) as compared to never-users (OR=0.97[CI=0.73-1.29]) indicating effect modification ( $p_{\text{heterogeneity}}=0.65$ ) while no relevant differences were observed between users and non-users for the variant alleles of ADH1B, ADH1C or ALDH2.

### **Conclusions**

Poor OH and DC are independent risk factors for UADT. Whether mouthwash use may entail some risk through the alcohol content in most formulations on the market remains to be clarified.

## Introduction

Tumors of the oral cavity, larynx, oropharynx, hypopharynx and esophagus, i.e. upper aerodigestive tract tumors (UADT), account for approximately 129,000 new cancer cases annually in the European Union(1), making them together the 4<sup>th</sup> and 10<sup>th</sup> most common cancer site in men and women, respectively.

The most important risk factors are consumption of alcohol and tobacco with a greater than multiplicative joint effect(2). Among never smokers it is likely that alcohol has little or no effect(1;3). There is also increasing evidence that high consumption of fruits and vegetables are protective(4-8), and emerging evidence for the role of HPV infection in some subsites(9). Occupational exposures are known risk factors for UADT(10-14). Low socioeconomic status (SES) and a downward trajectory of social position over the life-course are also both associated with an increased risk after adjustment for confounding factors(15;16).

Both a lack of oral hygiene and poor oral health seem to be risk factors for cancers of the UADT, independently from smoking and alcohol consumption (17-20). Frequent use of mouthwash may be a risk factor due to the alcohol contained in many formulations(21;22). Ethanol is metabolized to acetaldehyde by alcohol dehydrogenase (ADH) and is further metabolized to acetic acid by aldehyde dehydrogenase (ALDH). Polymorphic variants of the genes *ADH1B*, *ADH1C* and *ADH7* encoding for rapid metabolism of ethanol have been shown to decrease the risk of UADT(23-27) indicating that fast metabolism of ethanol reduces its carcinogenic effect. In a genome-wide association study comprising over 8,000 cases and 11,000 controls (including the ARCAGE study participants), variants in all 4 genes were robustly associated with UADT, with the association for the *ADH1B* gene variant (rs1229984) being particularly prominent among heavy drinkers(24). Similarly, we hypothesize that an effect of mouthwash may be modified by *ADH* genes, in particular by the *ADH1B* variant rs1229984.

The evidence for a risk entailed by alcohol-containing mouthwashes is limited. Although a recent meta-analysis showed no statistically significant association between mouthwash-use and oral cancer(28), this issue may still be considered as controversial(29). A new case-control study also found a positive association between daily mouthwash use and head and neck carcinoma with no difference between the effects of alcohol-containing and non-alcoholic mouthwashes(20). The objective of this analysis is to disentangle the effects of mouthwash from major potential confounding factors such as smoking and alcohol and to investigate the role of mouthwash and oral health/dental care more comprehensively(30).

Using a novel approach, two composite weighted scores were constructed using indicators of oral health and dental care that were associated with UADT in previous studies. In addition, we investigate the potential risk due to frequent use of mouthwash and its potential effect modification by genetic variants that have been shown to modify the risk of UADT in heavy drinkers.

## **Materials and methods**

Briefly, the ARCAGE multicenter case-control study for 13 centers (cf. Tab.1) in 9 European countries collected extensive lifestyle data including oral health (OH) and dental care (DC)(31). The study was approved by the ethical review board of the coordinating center, IARC, and the respective local boards in all centers. All subjects signed an informed consent form. With a common protocol, cases were defined as those patients newly diagnosed with primary squamous cell tumors of the UADT between 2002 and 2005. Diagnoses included malignant cancers of the oral cavity (ICD-O-3 topography: C00-06), oropharynx (C09, 10), hypopharynx (C12, 13), pharynx (C14), larynx (C32), or esophagus (C15). Controls were frequency-matched to cases by sex and age (5-year groups). In the UK centers, population controls were randomly selected from the same medical practice list as the corresponding cases. In all other countries patients admitted to the same hospitals as the cases for a wide

spectrum of medical conditions not associated with alcohol consumption and smoking were selected as controls.

Subjects were interviewed face-to-face by trained interviewers about socio-demographic characteristics, anthropometric measures, smoking and alcohol consumption, frequency of intake of selected foods, a detailed occupational history and a medical and dental history including oral hygiene habits.

We used the number of years of fulltime education (school, high school, university) as a measure of SES. Lifetime smoking history of cigarettes, cigars, and pipes was used to calculate 'pack-years'. Lifetime duration of drinking and average daily alcohol consumption over the lifetime (drinks/day) was assessed for all beverage categories. Daily consumption was converted into standard units with one drink corresponding to 8g of ethanol. Weekly consumption of fruits and vegetables recorded by a food frequency questionnaire was categorized by country-specific tertiles(31).

In the absence of any previous established scoring system we *a-priori* constructed a composite weighted score of oral health (OH) based on the following three variables that repeatedly were reported as being associated with one or more UADT sites. The categories were chosen to ensure equal distribution of study subjects while the weights (in brackets) were assigned according to the strength of associations in previous studies without knowing the corresponding risk estimates in our data:

1. Wearing of dentures (none=0; partial denture in upper or lower jaw=1; partial denture in both jaws=2; complete denture in one jaw=3; complete denture in both jaws=4)
2. Age at starting to wear dentures (no denture=0; denture at age 55 years or older=1; denture at age 35-54 years=2; denture at age below 35 years=3)
3. Frequency of gum bleeding from brushing teeth (sometimes or never=0; always or almost always=1; zero in subjects wearing complete dentures in both jaws).

Weights were added up over these variables to give a maximum score of seven for the poorest OH level. History of oral, pharyngeal or laryngeal biopsies, while included in the questionnaire (and analyzed separately), were not included in the OH score as we could not rule out that part of the biopsies reported by cases were disease-related.

The composite weighted score of dental care (DC) was constructed *a-priori* in a similar way as the OH score based on the following three variables:

1. Frequency of tooth cleaning (at least twice/day=0; once/day=1; 1-4 times/week=2; less often or never=3).
2. Use of toothbrush, toothpaste or dental floss (two or three of these=0; only one of these three =1; none of these=2).
3. Frequency of visiting a dentist (at least once/year=0; every 2-5 years=1; less than every 5 years=2; never=3).

The maximum DC score of eight indicated poor DC. Mouthwash use was not included in the DC score as previous studies suggested that frequent mouthwash use carries a potential risk rather than a benefit for UADT. Those with complete upper and lower dentures were not scored and excluded.

Composite weighted scores were grouped into four categories such that OH and DC scores of zero served as the reference. The other categories were defined such that each included a meaningful number of subjects while keeping the range between low and high scores as wide as possible.

The sample size was chosen to allow identification of a twofold relative risk for rare exposures with 1% prevalence, or a 50% risk increase for more common exposures of at least 5% prevalence. Regarding an assessment of gene-environment interaction, the sample size was chosen to allow identification of a twofold relative risk for polymorphisms with a population prevalence of 5% or more. For inter-country comparisons, the sample size was chosen to allow detection of effect modification between countries for common

risk factors (e.g. 10% or more), which have a moderate increased risk of between 2- and 2.5-fold. The study was designed to provide a sufficient number of cases to enable subgroup analyses.

Odds ratios (OR) and corresponding 95% confidence intervals (CI) were estimated by unconditional logistic regression using the PROC LOGISTIC function of the SAS software package, Version 9.2.

The strength and robustness of the association between indicators of the OH and DC scores and UADT were explored by consecutive inclusion of potential confounders in the statistical model to examine attenuation of the ORs. The basic model 1 (OR<sub>1</sub>) includes sex, age (9 categories) and study center (using dummy variables). As additional variables, model 2 (OR<sub>2</sub>) includes smoking status (never, former, current smoking), tobacco consumption (pack-years), duration of alcohol drinking (6 dummies for duration: 1-9 years, 10-19 years, 20-29 years, 30-39 years, 40+ years, unknown) and frequency of alcohol drinking. Frequency of drinking was weighted by multiplying the number of drinks/week by years of alcohol drinking. We imputed 44 missing values for frequency by the median (of cases and controls) within each drinking duration category. Further adjustment for education was included in model 3 (OR<sub>3</sub>) using two dummy variables (intermediate/upper secondary school [corresponding to 8-13 years of school], university degree). Model 4 (OR<sub>4</sub>) extended model 3 by adding two tertiles of frequency of fruit and vegetable intake, respectively (two dummy variables each). In sensitivity analyses, we stratified by sex and by subsite of UADT (esophagus; hypopharynx and larynx; oral cavity and oropharynx).

We analyzed 4 SNPs from the genes *ADH1B* (rs1229984), *ADH1C* (rs698), *ADH7* (rs1573496) and *ALDH2* (rs4767364) assuming a dominant model. All four SNPs have been conclusively shown to be associated with a risk of UADT, with the most comprehensive evidence from a recent genome-wide study of UADT that involved over 8,000 cases and 11,000 controls including the ARCAGE study. Of particular interest was the greater effect

among alcohol users that was observed with the *ADH1B* variant rs1229984, implying a gene-environment interaction(24). We hypothesized that the association with these SNPs may be more pronounced in mouthwash users, in particular for *ADH1B* (rs1229984).

## **Results**

The overall response proportion was 80% in cases and 66% in controls. The study group comprised 1,963 cases and 1,993 controls after exclusion of prevalent cases, subjects with tumors of the salivary glands, carcinoma in situ and non UADT as well as controls with non-eligible diseases (Tab.1).

The majority of the cases (48%) had tumors of the mouth and oropharynx, followed by hypopharynx and larynx (36%) and esophagus (12%). The site of origin was not assigned in 5% because the tumor was with overlapping sites.

Table 2 shows the distribution of known risk factors and their adjusted ORs. Our data corroborated the dose-effect relationship of smoking and alcohol consumption with UADT risk. SES and consumption of fruits and vegetables were inversely associated with risk of these tumors after adjustment for smoking and alcohol.

Table 3 displays the risk estimates for each indicator of OH and DC and the impact of adjustment on these estimates.

### *Indicators of oral health*

Wearing of dentures was associated with an elevated risk, with complete dentures in both jaws carrying the highest risk ( $OR_1=2.60; CI=2.15-3.15$ ). The risk estimate was substantially reduced after adjustment for smoking and alcohol ( $OR_2$ ) while further adjustment for education and diet had little impact ( $OR_4=1.73; CI=1.39-2.15$ ). Similarly, age at starting to wear dentures below 55 years was associated with elevated risk by about 40%. Frequent gum bleeding when cleaning teeth was associated with no noticeable risk differences.

### *Indicators of dental care*

Compared to subjects cleaning teeth at least twice per day the risk estimates increased with decreasing frequency of tooth cleaning. Subjects cleaning their teeth less than once per week showed the highest risk estimate ( $OR_1=3.51; CI=2.55-4.83$ ) but adjustment attenuated this effect substantially ( $OR_4=1.37; CI=0.95-1.99$ ). Similarly, the elevated risk for not using a toothbrush ( $OR_1=1.96; CI=0.94-4.06$ ) came close to unity after adjustment ( $OR_4=1.15; CI=0.48-2.80$ ). Neither non-use of toothpaste nor non-use of dental floss showed a marked risk after adjustment for confounding. Risk estimates decreased with increasing frequency of visiting a dentist during the last 20 years. Compared to participants who were seeing the dentist at least once a year, participants who never visited the dentist had an estimated  $OR_4$  of 1.93 ( $CI=1.48-2.51$ ).

Frequent use of mouthwash (three and more times/day) was positively associated with UADT as compared to never use with adjustment for confounders increasing the risk estimate further ( $OR_4=3.23; CI=1.68-6.19$ ). While this effect was strong, only 1.8% of cases and 0.8% of controls reported such frequent use. The risk was most pronounced for oral cavity ( $OR_4=3.53; CI=1.65-7.57$ ) and pharyngeal cancers ( $OR_4=3.50; CI=1.55-7.89$ ).

When restricting the analysis to non-smokers or to non-drinkers, the risk persisted. Only four (3.1%) of 128 non-drinking cases and three (1.2%) of 254 non-drinking controls reported frequent use of mouthwash, resulting in an  $OR_4$  of 4.42 ( $CI=0.79-24.75$ ). Frequent mouthwash users did not differ from less frequent users in terms of amount of drinking and smoking, frequency of toothbrushing or use of toothpaste or dental floss. Although frequent mouthwash users are more likely to wear complete dentures (28.9% vs. 17.5%), risk estimates for frequent mouthwash use were elevated in both groups with  $OR_4=7.27$  ( $CI=1.32-40.0$ ) among subjects wearing complete dentures and  $OR_4=2.57$  ( $CI=1.24-5.33$ ) among subjects not having complete dentures.

### *Weighted scores of oral health and dental care*

ORs for OH/DC scores are reported in Table 4. OH scores 1-4 did not show noticeable risk elevations while scores 5, 6 and 7 did. Poor OH (score=7) was associated with a more than two-fold risk of developing UADT after adjustment for confounding (OR<sub>4</sub>=2.22;CI=1.45-3.41). Even a moderate lack of DC (score=1) was associated with an elevated risk of developing UADT after adjustment (OR<sub>4</sub>=1.36;CI=1.07-1.72). ORs increased substantially with increasing DC score reaching their maximum for scores 5-8 (OR<sub>1</sub>=6.01;CI=4.07-8.87). Smoking and alcohol were the main confounders (OR<sub>2</sub>=2.77;CI=1.79-4.27) while further adjustment for SES and diet had little impact (OR<sub>4</sub>=2.36;CI=1.51-3.67).

Subgroup analyses showed consistent risk elevations for OH/DC across all subsites of UADT. It is noteworthy, however, that the highest risk estimates for the DC score were observed for hypopharyngeal/laryngeal tumors, and not oral/oropharyngeal tumors (Tab.5). We observed a more than two-fold though statistically non-significant increased risk for frequent mouthwash use even in the subgroup of cases and controls with the best OH score (score=0)(OR<sub>4</sub>=2.33;CI=0.72-7.47).

### *Genetic markers*

As expected, and in line with results from the larger GWA study(24), strongly significant inverse associations were observed for the rare variants of *ADH1B* (OR=0.55(0.41-0.74)) and *ADH7* (OR=0.79(0.64-0.98))(Fig.1). Both of these rare variants code for fast alcohol metabolism. Correspondingly, a positive association was observed for the rare variant of *ADH1C* that codes for slow ethanol metabolism (OR=1.23(1.04-1.46)). A moderate non-significantly increased risk was seen for the *ALDH2* variant (OR=1.07(0.91-1.27))(not included in Fig.1). No strong effect modification by any of these variants was observed after stratifying by level of alcohol consumption. Similarly, the strong effect modification seen in

the larger GWA analysis for *ADH1B* was not replicated here, indicating that the sample size with ARCADE alone is likely to be insufficient to robustly detect effect modifications. Further, the only indication for effect modification observed after stratifying for mouthwash use was for the *ADH7* variant ( $p_{\text{heterogeneity}}=0.065$ ). When this analysis was restricted to subjects who consumed less than 5 drinks/month the reduced risk for the *ADH7* variant persisted among mouthwash users as compared to non-users but the risk estimates became rather unstable, because this restriction reduced the sample size by 74%. Given that no clear effect modification was observed for alcohol use in this analysis or the larger GWA analysis for the *ADH7* variant, it is unclear to what extent this could represent real effect modification and not just random fluctuation.

## **Discussion**

Our study showed that both poor oral health and poor dental care were associated with the risk of UADT independently of other known risk factors. Semi-quantitative scores of OH and DC revealed a dose-effect relationship with the risk of UADT tumors. Frequent use of mouthwashes (3+ times/day) entailed an elevated risk of developing UADT.

The strengths of our study include: (i) the statistical power—this is among the largest case-control studies to-date which have examined the etiology of UADT associated with oral hygiene; (ii) the restriction to histologically confirmed incident cases; (iii) the control for multiple confounding variables including smoking, alcohol, consumption of fresh fruits and vegetables as well as SES(8;15;16;32;33).

Some methodological issues require attention. ARCADE relies on hospital controls in all but the UK centers. Hospital controls were chosen to achieve better participation because blood samples had to be taken, but this choice may be more vulnerable to selection bias. To minimize such bias, only conditions not related to tobacco smoking, alcohol intake or diet were eligible as controls(8). However, lower socioeconomic groups have a higher risk of

hospital admission(34). Our study may thus underestimate the association between UADT and OH/DC since both are associated with low SES.

Residual confounding by tobacco smoking, alcohol drinking, and possibly other factors such as poor diet cannot be ruled out as adjustment had an important effect on most risk estimates. It cannot be excluded that some misclassification of these confounders has taken place, leading to incomplete adjustment. However, residual confounding should be small since associations persisted in stratified analyses and since we were able to carefully adjust for alcohol and tobacco for which histories detailed every change in their consumption. Alcohol and tobacco consumption of our controls was comparable to other case-control studies(6;19;35-38).

The association between poor OH/DC and UADT tumors was attenuated but not eliminated after full adjustment. Furthermore, despite methodological differences, our findings are corroborated by many previous studies(19;39-41)(17;18;42-49). As these studies used retrospective designs they are all vulnerable to reporting bias which might explain some of the apparent consistency across studies. However, as such reporting bias, e.g. due to underreporting of smoking by cases, would primarily affect factors known to be associated with UADT like smoking and alcohol but not DC or OH, strong associations of the latter with UADT may not be explained by such bias.

Some studies assessed OH by clinical examination, but they did not describe precisely how the various indicators were combined into a single variable. Nevertheless, aggregate variables of OH were consistently associated with elevated risks of head and neck cancers. Inspection of the mouth was not done in the ARCAGE study. Instead, we developed a composite weighted score for OH and for DC, respectively, in order to obtain robust semi-quantitative variables that capture the various dimensions of oral health and dental care based on the self-reported data. This approach has never been used before. The score

allows for subgroup analyses and for the assessment of dose-effect relationships. The two scores both showed consistent associations with the tumor sites investigated in our study and may thus serve as a useful tool in future assessments.

In agreement with previous IARC studies(19) we found the strongest association with frequent mouthwash use (3+ times/day) for oral cavity and pharyngeal cancers. However, this is based on 36 (17/17+2) cases and 16 controls only. In their meta-analysis Gandini et al. observed a dose-effect relationship with increasing frequency of mouthwash use, but the corresponding risk estimates were unstable and not statistically significant(28). The Carolina Head and Neck Cancer Study did not observe such association(49). In a previous study, Winn et al. observed an elevated risk only for the use of mouthwash containing alcohol(43). A recent study by Eliot et al.(20) reported an elevated risk of head and neck carcinomas for frequent use of both, alcohol-containing and alcohol-free mouthwashes. However, the validity of this distinction based on self-reports may be questionable. Unfortunately our questionnaire did not allow to distinguish between both types of mouthwashes. An in-depth analysis of a Puerto Rican case-control study of oral and pharyngeal cancer(50) showed no overall increased risk but revealed an effect of alcohol-containing mouthwash in women and in subjects who neither drank alcohol nor smoked cigarettes. Our study supports these findings as well as the results of subgroup analyses among alcohol abstainers from the recent study by Guha et al.(19) suggesting mouthwash to be a risk factor for the development of UADT that may act through the alcohol contained in most formulations on the market(21). Given alcohol consumption among never smokers appears to have little or no risk for head and neck cancers(2;3) we may expect that similarly, any risk associated with alcohol from mouthwash use may be restricted to smokers. The sample size of our current study was however too limited to test such a hypothesis.

Further, similar to our previous studies that have identified effect modification between alcohol metabolizing genes and alcohol consumption, we investigated whether any effect modification existed between alcohol and acetaldehyde genes known to be associated with head and neck cancers and mouthwash use. On the one hand, the apparent effect modification by the variant ADH7 would support a carcinogenic effect of mouthwash but it is surprising that the effect is stronger than in heavy drinkers. In retrospect, it is likely that the sample size for this study (up to 1,353 cases and 1,488 controls with genetic and mouthwash data available) was too limited to robustly detect effect modification. On the other hand, the results on the small group of users of mouthwash 3+ times/day might indicate some underlying oral disease as people with oral problems are more likely to use mouthwash(51). It should be noted that study subjects who used mouthwash 3+ times/day were more likely to wear dentures. Moreover, reverse causation cannot be excluded since UADT patients are often prescribed mouthwashes like chlorhexidine for prevention of mouth infection. Also the lack of association in the more informative group of regular users (1-2 times/day) does not support the hypothesis of an association. Neither does the similar magnitude of an effect on oral cavity/oropharynx and hypopharynx/larynx.

In conclusion our study provides further evidence for an elevated risk of UADT due to poor OH and poor DC that is not explained by smoking, alcohol or other confounding factors. Excessive use (3+ times/day) of mouthwash also increased risk for UADT but the question whether this effect is mediated by alcohol content or just due to bias or reverse causation cannot be decided with our data.

UADT prevention strategies may want to consider promoting oral health and dental care, regularly attending the dentist, good personal oral hygiene practices, and avoiding excessive mouthwash use.

## **Acknowledgements**

The authors thank the patients and their families for their participation and gratefully acknowledge the study interviewers and our clinical colleagues in hospitals and primary care who supported this study.

GJM and TVM partly worked on this study while at the University of Manchester. We acknowledge the help of Dr. Ann-Marie Biggs, Professor Martin Tickle and Professor Phil Sloan in study conduct in the Manchester center.

We are grateful for the editorial support by Ina Alvarez in revising previous versions of this manuscript.

## **Competing interests**

The authors report no conflict of interest and declare no financial relationships with any organization that might have an interest in the submitted work in the previous three years, as well as no other relationships or activities that could appear to have influenced the submitted work.

## Reference List

- (1) Ferlay J, Parkin DM, Steliarova-Foucher E. Estimates of cancer incidence and mortality in Europe in 2008. *Eur J Cancer* 2010; 46(4):765-781.
- (2) Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C et al. Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol Biomarkers Prev* 2009; 18(2):541-550.
- (3) Hashibe M, Brennan P, Benhamou S, Castellsague X, Chen C, Curado MP et al. Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *J Natl Cancer Inst* 2007; 99(10):777-789.
- (4) Kjaerheim K, Gaard M, Andersen A. The role of alcohol, tobacco, and dietary factors in upper aerogastric tract cancers: a prospective study of 10,900 Norwegian men. *Cancer Causes Control* 1998; 9(1):99-108.
- (5) Freedman ND, Park Y, Subar AF, Hollenbeck AR, Leitzmann MF, Schatzkin A et al. Fruit and vegetable intake and head and neck cancer risk in a large United States prospective cohort study. *Int J Cancer* 2008; 122(10):2330-2336.
- (6) Sapkota A, Hsu CC, Zaridze D, Shangina O, Szeszenia-Dabrowska N, Mates D et al. Dietary risk factors for squamous cell carcinoma of the upper aerodigestive tract in central and eastern Europe. *Cancer Causes Control* 2008; 19(10):1161-1170.
- (7) Boeing H, Dietrich T, Hoffmann K, Pischon T, Ferrari P, Lahmann PH et al. Intake of fruits and vegetables and risk of cancer of the upper aero-digestive tract: the prospective EPIC-study. *Cancer Causes Control* 2006; 17(7):957-969.
- (8) Lagiou P, Talamini R, Samoli E, Lagiou A, Ahrens W, Pohlabein H et al. Diet and upper-aerodigestive tract cancer in Europe: the ARCAGE study. *Int J Cancer* 2009; 124(11):2671-2676.
- (9) D'Souza G, Kreimer AR, Viscidi R, Pawlita M, Fakhry C, Koch WM et al. Case-control study of human papillomavirus and oropharyngeal cancer. *N Engl J Med* 2007; 356(19):1944-1956.
- (10) Purdue MP, Jarvholm B, Bergdahl IA, Hayes RB, Baris D. Occupational exposures and head and neck cancers among Swedish construction workers. *Scand J Work Environ Health* 2006; 32(4):270-275.
- (11) Tarvainen L, Kyyronen P, Kauppinen T, Pukkala E. Cancer of the mouth and pharynx, occupation and exposure to chemical agents in Finland [in 1971-95]. *Int J Cancer* 2008; 123(3):653-659.
- (12) Dietz A, Ramroth H, Urban T, Ahrens W, Becher H. Exposure to cement dust, related occupational groups and laryngeal cancer risk: results of a population based case-control study. *Int J Cancer* 2004; 108(6):907-911.

- (13) Ramroth H, Dietz A, Ahrens W, Becher H. Occupational wood dust exposure and the risk of laryngeal cancer: a population based case-control study in Germany. *Am J Ind Med* 2008; 51(9):648-655.
- (14) Richiardi L, Corbin M, Marron M, Ahrens W, Pohlabeln H, Lagiou P et al. Occupation and risk of upper aerodigestive tract cancer: The ARCAGE study. *Int J Cancer* 2011.
- (15) Conway DI, Petticrew M, Marlborough H, Berthiller J, Hashibe M, Macpherson LM. Socioeconomic inequalities and oral cancer risk: a systematic review and meta-analysis of case-control studies. *Int J Cancer* 2008; 122(12):2811-2819.
- (16) Schmeisser N, Conway DI, McKinney PA, McMahon AD, Pohlabeln H, Marron M et al. Life course social mobility and risk of upper aerodigestive tract cancer in men. *Eur J Epidemiol* 2010; 25(3):173-182.
- (17) Bundgaard T, Wildt J, Frydenberg M, Elbrond O, Nielsen JE. Case-control study of squamous cell cancer of the oral cavity in Denmark. *Cancer Causes Control* 1995; 6(1):57-67.
- (18) Rosenquist K, Wennerberg J, Schildt EB, Bladstrom A, Goran HB, Andersson G. Oral status, oral infections and some lifestyle factors as risk factors for oral and oropharyngeal squamous cell carcinoma. A population-based case-control study in southern Sweden. *Acta Otolaryngol* 2005; 125(12):1327-1336.
- (19) Guha N, Boffetta P, Wunsch F, V, Eluf NJ, Shangina O, Zaridze D et al. Oral health and risk of squamous cell carcinoma of the head and neck and esophagus: results of two multicentric case-control studies. *Am J Epidemiol* 2007; 166(10):1159-1173.
- (20) Eliot MN, Michaud DS, Langevin SM, McClean MD, Kelsey KT. Periodontal disease and mouthwash use are risk factors for head and neck squamous cell carcinoma. *Cancer Causes Control* 2013; 24(7):1315-1322.
- (21) McCullough MJ, Farah CS. The role of alcohol in oral carcinogenesis with particular reference to alcohol-containing mouthwashes. *Aust Dent J* 2008; 53(4):302-305.
- (22) Lachenmeier DW, Gumbel-Mako S, Sohnius EM, Keck-Wilhelm A, Kratz E, Mildau G. Salivary acetaldehyde increase due to alcohol-containing mouthwash use: a risk factor for oral cancer. *Int J Cancer* 2009; 125(3):730-735.
- (23) Hashibe M, McKay JD, Curado MP, Oliveira JC, Koifman S, Koifman R et al. Multiple ADH genes are associated with upper aerodigestive cancers. *Nat Genet* 2008; 40(6):707-709.
- (24) McKay JD, Truong T, Gaborieau V, Chabrier A, Chuang SC, Byrnes G et al. A genome-wide association study of upper aerodigestive tract cancers conducted within the INHANCE consortium. *PLoS Genet* 2011; 7(3):e1001333.
- (25) Asakage T, Yokoyama A, Haneda T, Yamazaki M, Muto M, Yokoyama T et al. Genetic polymorphisms of alcohol and aldehyde dehydrogenases, and drinking, smoking and diet in Japanese men with oral and pharyngeal squamous cell carcinoma. *Carcinogenesis* 2007; 28(4):865-874.

- (26) Oze I, Matsuo K, Suzuki T, Kawase T, Watanabe M, Hiraki A et al. Impact of multiple alcohol dehydrogenase gene polymorphisms on risk of upper aerodigestive tract cancers in a Japanese population. *Cancer Epidemiol Biomarkers Prev* 2009; 18(11):3097-3102.
- (27) Tanaka F, Yamamoto K, Suzuki S, Inoue H, Tsurumaru M, Kajiyama Y et al. Strong interaction between the effects of alcohol consumption and smoking on oesophageal squamous cell carcinoma among individuals with ADH1B and/or ALDH2 risk alleles. *Gut* 2010; 59(11):1457-1464.
- (28) Gandini S, Negri E, Boffetta P, La VC, Boyle P. Mouthwash and oral cancer risk quantitative meta-analysis of epidemiologic studies. *Ann Agric Environ Med* 2012; 19(2):173-180.
- (29) Warnakulasuriya S. Causes of oral cancer--an appraisal of controversies. *Br Dent J* 2009; 207(10):471-475.
- (30) Conway D. Oral health, mouthwashes and cancer--what is the story? *Evid Based Dent* 2009; 10(1):6-7.
- (31) Lagiou P, Georgila C, Minaki P, Ahrens W, Pohlabein H, Benhamou S et al. Alcohol-related cancers and genetic susceptibility in Europe: the ARCAGE project: study samples and data collection. *Eur J Cancer Prev* 2009; 18(1):76-84.
- (32) Lee YC, Marron M, Benhamou S, Bouchardy C, Ahrens W, Pohlabein H et al. Active and involuntary tobacco smoking and upper aerodigestive tract cancer risks in a multicenter case-control study. *Cancer Epidemiol Biomarkers Prev* 2009; 18(12):3353-3361.
- (33) Macfarlane TV, Macfarlane GJ, Oliver RJ, Benhamou S, Bouchardy C, Ahrens W et al. The aetiology of upper aerodigestive tract cancers among young adults in Europe: the ARCAGE study. *Cancer Causes Control* 2010; 21(12):2213-2221.
- (34) Saxena S, George J, Barber J, Fitzpatrick J, Majeed A. Association of population and practice factors with potentially avoidable admission rates for chronic diseases in London: cross sectional analysis. *J R Soc Med* 2006; 99(2):81-89.
- (35) Jansson C, Johansson AL, Nyren O, Lagergren J. Socioeconomic factors and risk of esophageal adenocarcinoma: a nationwide Swedish case-control study. *Cancer Epidemiol Biomarkers Prev* 2005; 14(7):1754-1761.
- (36) Altieri A, Bosetti C, Gallus S, Franceschi S, Dal ML, Talamini R et al. Wine, beer and spirits and risk of oral and pharyngeal cancer: a case-control study from Italy and Switzerland. *Oral Oncol* 2004; 40(9):904-909.
- (37) Ferraroni M, Negri E, La VC, D'Avanzo B, Franceschi S. Socioeconomic indicators, tobacco and alcohol in the aetiology of digestive tract neoplasms. *Int J Epidemiol* 1989; 18(3):556-562.
- (38) Morse DE, Psoter WJ, Cleveland D, Cohen D, Mohit-Tabatabai M, Kosis DL et al. Smoking and drinking in relation to oral cancer and oral epithelial dysplasia. *Cancer Causes Control* 2007; 18(9):919-929.
- (39) Abnet CC, Kamangar F, Islami F, Nasrollahzadeh D, Brennan P, Aghcheli K et al. Tooth loss and lack of regular oral hygiene are associated with higher risk of esophageal squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 2008; 17(11):3062-3068.

- (40) Maier H, Zoller J, Herrmann A, Kreiss M, Heller WD. Dental status and oral hygiene in patients with head and neck cancer. *Otolaryngol Head Neck Surg* 1993; 108(6):655-661.
- (41) Velly AM, Franco EL, Schlecht N, Pintos J, Kowalski LP, Oliveira BV et al. Relationship between dental factors and risk of upper aerodigestive tract cancer. *Oral Oncol* 1998; 34(4):284-291.
- (42) Zheng TZ, Boyle P, Hu HF, Duan J, Jian PJ, Ma DQ et al. Dentition, oral hygiene, and risk of oral cancer: a case-control study in Beijing, People's Republic of China. *Cancer Causes Control* 1990; 1(3):235-241.
- (43) Winn DM, Blot WJ, McLaughlin JK, Austin DF, Greenberg RS, Preston-Martin S et al. Mouthwash use and oral conditions in the risk of oral and pharyngeal cancer. *Cancer Res* 1991; 51(11):3044-3047.
- (44) Moreno-Lopez LA, Esparza-Gomez GC, Gonzalez-Navarro A, Cerero-Lapiedra R, Gonzalez-Hernandez MJ, Dominguez-Rojas V. Risk of oral cancer associated with tobacco smoking, alcohol consumption and oral hygiene: a case-control study in Madrid, Spain. *Oral Oncol* 2000; 36(2):170-174.
- (45) Fernandez Garrote L, Herrero R, Ortiz Reyes RM, Vaccarella S, Lence Anta J, Ferbeyre L et al. Risk factors for cancer of the oral cavity and oro-pharynx in Cuba. *British Journal of Cancer* 85[1], 46-56. 2001. Cancer Research Campaign.

Ref Type: Magazine Article

- (46) Lissowska J, Pilarska A, Pilarski P, Samolczyk-Wanyura D, Piekarczyk J, Bardin-Mikollajczak A et al. Smoking, alcohol, diet, dentition and sexual practices in the epidemiology of oral cancer in Poland. *Eur J Cancer Prev* 2003; 12(1):25-33.
- (47) Balaram P, Sridhar H, Rajkumar T, Vaccarella S, Herrero R, Nandakumar A et al. Oral cancer in southern India: the influence of smoking, drinking, paan-chewing and oral hygiene. *Int J Cancer* 2002; 98(3):440-445.
- (48) Subapriya R, Thangavelu A, Mathavan B, Ramachandran CR, Nagini S. Assessment of risk factors for oral squamous cell carcinoma in Chidambaram, Southern India: a case-control study. *Eur J Cancer Prev* 2007; 16(3):251-256.
- (49) Divaris K, Olshan AF, Smith J, Bell ME, Weissler MC, Funkhouser WK et al. Oral health and risk for head and neck squamous cell carcinoma: the Carolina Head and Neck Cancer Study. *Cancer Causes Control* 2010; 21(4):567-575.
- (50) Winn DM, Diehl SR, Brown LM, Harty LC, Bravo-Otero E, Fraumeni JF, Jr. et al. Mouthwash in the etiology of oral cancer in Puerto Rico. *Cancer Causes Control* 2001; 12(5):419-429.
- (51) Macfarlane TV, Kawecki M, Cunningham C, Bovaird I, Morgan R, Rhodes K et al. Mouthwash use in general population: results from adult dental health survey in Grampian, Scotland. *Journal of Oral & Maxillofacial Research* 2010; 1(4).

Table 1. Characteristics of Study Subjects of the ARCAGE Study Included in the Analysis of Oral Health and Dental Care

	Men				Women				All			
	Cases		Controls		Cases		Controls		Cases		Controls	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
<b>Age (years)</b>												
<40	16	(1.0)	45	(3.1)	23	(5.5)	49	(8.9)	39	(2.0)	94	(4.7)
40-44	55	(3.6)	62	(4.3)	23	(5.5)	24	(4.4)	78	(4.0)	86	(4.3)
45-49	152	(9.9)	122	(8.5)	25	(5.9)	41	(7.4)	177	(9.0)	163	(8.2)
50-54	254	(16.5)	212	(14.7)	39	(9.2)	61	(11.1)	293	(14.9)	273	(13.7)
55-59	322	(20.9)	259	(18.0)	82	(19.4)	88	(16.0)	404	(20.6)	347	(17.4)
60-64	264	(17.1)	218	(15.1)	60	(14.2)	66	(12.0)	324	(16.5)	284	(14.2)
65-69	219	(14.2)	234	(16.2)	67	(15.9)	81	(14.7)	286	(14.6)	315	(15.8)
70-74	151	(9.8)	173	(12.0)	55	(13.0)	68	(12.3)	206	(10.5)	241	(12.1)
75+	108	(7.0)	117	(8.1)	48	(11.4)	73	(13.2)	156	(7.9)	190	(9.5)
<b>Center</b>												
Prague	159	(10.3)	154	(10.7)	31	(7.3)	33	(6.0)	190	(9.7)	187	(9.4)
Bremen	236	(15.3)	264	(18.3)	51	(12.1)	64	(11.6)	287	(14.6)	328	(16.5)
Athens	192	(12.5)	143	(9.9)	47	(11.1)	51	(9.3)	239	(12.2)	194	(9.7)
Aviano	119	(7.7)	118	(8.2)	32	(7.6)	33	(6.0)	151	(7.7)	151	(7.6)
Padova	109	(7.1)	97	(6.7)	26	(6.2)	33	(6.0)	135	(6.9)	130	(6.5)
Turin	123	(8.0)	144	(10.0)	43	(10.2)	54	(9.8)	166	(8.5)	198	(9.9)
Dublin	33	(2.1)	6	(0.4)	11	(2.6)	13	(2.4)	44	(2.2)	19	(1.0)
Oslo	121	(7.9)	109	(7.6)	51	(12.1)	75	(13.6)	172	(8.8)	184	(9.2)
Glasgow	61	(4.0)	52	(3.6)	32	(7.6)	39	(7.1)	93	(4.7)	91	(4.6)
Manchester	104	(6.7)	122	(8.5)	45	(10.7)	64	(11.6)	149	(7.6)	186	(9.3)
Newcastle	71	(4.6)	95	(6.6)	17	(4.0)	18	(3.3)	88	(4.5)	113	(5.7)
Barcelona	168	(10.9)	101	(7.0)	27	(6.4)	65	(11.8)	195	(9.9)	166	(8.3)
Zagreb	45	(2.9)	37	(2.6)	9	(2.1)	9	(1.6)	54	(2.8)	46	(2.3)
<b>Subsite</b>												
Mouth/												
oropharynx	680	(44.1)			254	(60.2)			934	(47.6)		

	Men				Women				All			
	Cases		Controls		Cases		Controls		Cases		Controls	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
Hypopharynx/ larynx	619	(40.2)			79	(18.7)			698	(35.6)		
Esophagus	173	(11.2)			61	(14.5)			234	(11.9)		
Overlapping multiple sites	69	(4.5)			28	(6.6)			97	(4.9)		
<b>All</b>	1541	(100.0)	1442	(100.0)	422	(100.0)	551	(100.0)	1963	(100.0)	1993	(100.0)

Table 2. Distribution of Known Risk Factors of UADT and Corresponding ORs with 95%-Confidence Intervals.

	Cases		Controls				OR <sub>2</sub>	95%-CI
	N	(%)	N	(%)	OR <sub>1</sub>	95%-CI		
<b>Smoking status</b> <sup>*1</sup>								
Never †	217	(11.1)	733	(36.8)	1.00	(ref.)	1.00	(ref.)
Former	472	(24.0)	688	(34.5)	2.25	(1.84-2.75)	2.02	(1.64-2.49)
Current	1274	(64.9)	572	(28.7)	8.45	(6.96-10.26)	6.66	(5.45-8.14)
<b>Pack-years</b> <sup>**1</sup>								
Never †	217	(11.1)	733	(36.8)	1.00	(ref.)	1.00	(ref.)
(0-20)	341	(17.4)	570	(28.6)	2.14	(1.73-2.63)	2.04	(1.65-2.53)
[20-40)	650	(33.1)	416	(20.9)	5.98	(4.86-7.35)	4.98	(4.02-6.16)
[40-60)	427	(21.8)	180	(9.0)	9.72	(7.61-12.42)	7.34	(5.69-9.46)
[60-max]	316	(16.1)	91	(4.6)	14.69	(10.91-19.77)	10.41	(7.65-14.17)
Missing	12	(0.6)	3	(0.2)	15.73	(4.33-57.13)	14.60	(3.97-53.67)
<b>Alcohol drinking (frequency)</b> <sup>**2</sup>								
Never †	128	(6.5)	254	(12.7)	1.00	(ref.)	1.00	(ref.)
<1 drink/day	516	(26.3)	798	(40.0)	1.35	(1.04-1.74)	1.02	(0.77-1.36)
1-2 drinks/day	500	(25.5)	586	(29.4)	2.03	(1.54-2.66)	1.25	(0.93-1.68)

	Cases		Controls						
	N	(%)	N	(%)	OR <sub>1</sub>	95%-CI	OR <sub>2</sub>	95%-CI	
3-4 drinks/day	325	(16.6)	199	(10.0)	4.13	(3.04-5.61)	1.93	(1.38-2.71)	
5 drinks/day	463	(23.6)	122	(6.1)	9.81	(7.12-13.52)	3.95	(2.79-5.60)	
Missing	31	(1.6)	34	(1.7)	1.85	(1.08-3.20)	1.28	(0.71-2.29)	
<b>Alcohol drinking (duration)<sup>†‡</sup></b>									
Never †	128	(6.5)	254	(12.7)	1.00	(ref.)	1.00	(ref.)	
1-9 years	39	(2.0)	62	(3.1)	1.34	(0.84-2.15)	1.02	(0.62-1.70)	
10-19 years	92	(4.7)	117	(5.9)	1.76	(1.22-2.54)	1.22	(0.82-1.81)	
20-29 years	292	(14.9)	250	(12.5)	2.58	(1.92-3.47)	1.59	(1.15-2.20)	
30-39 years	587	(29.9)	506	(25.4)	2.52	(1.93-3.30)	1.43	(1.06-1.92)	
40+ years	810	(41.3)	775	(38.9)	2.23	(1.71-2.90)	1.28	(0.96-1.72)	
Missing †	15	(0.8)	29	(1.5)	1.23	(0.63-2.41)	0.79	(0.38-1.66)	
<b>Education</b>									
General secondary school	719	(36.6)	517	(25.9)	1.00	(ref.)	1.00	(ref.)	
Intermediate or advanced secondary school	1147	(58.4)	1274	(63.9)	0.63	(0.53-0.75)	0.77	(0.64-0.93)	
University degree	97	(4.9)	202	(10.1)	0.32	(0.24-0.43)	0.60	(0.44-0.82)	
<b>Fruit intake (tertiles)</b>									
Low †	963	(49.1)	596	(29.9)	1.00	(ref.)	1.00	(ref.)	

	Cases		Controls						
	N	(%)	N	(%)	OR <sub>1</sub>	95%-CI	OR <sub>2</sub>	95%-CI	
Medium	707	(36.0)	899	(45.1)	0.49	(0.42-0.56)	0.65	(0.55-0.76)	
High	293	(14.9)	498	(25.0)	0.34	(0.29-0.41)	0.53	(0.43-0.64)	
<b>Vegetable intake (tertiles)</b>									
Low †	932	(47.5)	703	(35.3)	1.00	(ref.)	1.00	(ref.)	
Medium	858	(43.7)	1042	(52.3)	0.64	(0.56-0.73)	0.75	(0.64-0.87)	
High	173	(8.8)	248	(12.4)	0.49	(0.39-0.62)	0.60	(0.47-0.78)	
<b>All</b>	1963	(100.0)	1993	(100.0)					

OR<sub>1</sub>: adjusted for age (9 categories), sex and study center

OR<sub>2</sub>: adjusted as OR<sub>1</sub> plus smoking status (never, former, current), plus cumulative tobacco consumption (packyears),

plus alcohol drinking duration (1-9 ys, 10-19 ys, 20-29 ys, 30-39 ys, 40+ ys, missing),

plus a cumulative alcohol consumption (alcohol drinking frequency weighted by duration of the drinking period)

\*1: OR<sub>2</sub> not adjusted for cumulative tobacco consumption

\*2: OR<sub>2</sub> not adjusted for cumulative alcohol consumption

†: Reference category (ref.)

Table 3. Distribution of Indicators of Oral Hygiene, Dental Care and Oral Health With Corresponding ORs and 95%-Confidence Intervals.

	Cases (%)	Contr. (%)	(%)	OR <sub>1</sub>	95%-CI	OR <sub>2</sub>	95%-CI	OR <sub>3</sub>	95%-CI	OR <sub>4</sub>	95%-CI
<b>Wearing dentures</b>											
None †	974 (49.6)	1226 (61.5)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	(ref.)
Partial (one jaw)	136 (6.9)	163 (8.2)	1.13	(0.88-1.45)	1.03	(0.78-1.36)	1.02	(0.77-1.34)	1.03	(0.78-1.36)	
Partial (lower and upper jaw)	152 (7.7)	154 (7.7)	1.35	(1.05-1.72)	1.29	(0.98-1.69)	1.27	(0.97-1.67)	1.24	(0.94-1.66)	
Complete (one jaw)	246 (12.5)	184 (9.2)	1.84	(1.48-2.28)	1.29	(1.01-1.64)	1.25	(0.98-1.59)	1.22	(0.96-1.56)	
Complete (lower and upper jaw)	445 (22.7)	251 (12.6)	2.60	(2.15-3.15)	1.81	(1.46-2.25)	1.76	(1.42-2.19)	1.73	(1.39-2.15)	
Missing	10 (0.5)	15 (0.8)									
<b>Age at starting to wear dentures</b>											
No denture †	974 (49.6)	1226 (61.5)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	(ref.)
55 years or older	259 (13.2)	269 (13.5)	1.28	(1.03-1.58)	1.09	(0.86-1.39)	1.07	(0.84-1.36)	1.05	(0.83-1.34)	
35-54 yrs	477 (24.3)	321 (16.1)	1.97	(1.66-2.34)	1.50	(1.24-1.82)	1.46	(1.20-1.77)	1.44	(1.19-1.75)	
Below 35 years	203 (10.3)	146 (7.3)	1.95	(1.53-2.47)	1.41	(1.08-1.84)	1.38	(1.06-1.80)	1.35	(1.03-1.77)	
Missing	50 (2.5)	31 (1.6)									
<b>Gum bleeding</b>											
[complete dentures excluded]											
Never †	719 (47.4)	824 (47.3)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	(ref.)
Sometimes	517 (34.1)	726 (41.7)	0.79	(0.68-0.92)	0.95	(0.80-1.13)	0.95	(0.80-1.13)	0.94	(0.79-1.12)	
Almost always	85 (5.6)	94 (5.4)	1.03	(0.75-1.41)	1.15	(0.81-1.63)	1.16	(0.81-1.65)	1.07	(0.75-1.53)	
Missing	197 (13.0)	98 (5.6)									

	Cases	(%)	Contr.	(%)	OR <sub>1</sub>	95%-CI	OR <sub>2</sub>	95%-CI	OR <sub>3</sub>	95%-CI	OR <sub>4</sub>	95%-CI
<b>Teeth cleaning</b>												
[complete dentures excluded]												
Twice / day †	622	(41.0)	1043	(59.9)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
Once / day	464	(30.6)	442	(25.4)	1.69	(1.42-2.01)	1.36	(1.12-1.65)	1.35	(1.11-1.64)	1.25	(1.03-1.53)
1-4x/week	172	(11.3)	128	(7.3)	2.18	(1.68-2.83)	1.53	(1.14-2.05)	1.49	(1.11-2.00)	1.39	(1.03-1.87)
Less than once/ week or never	153	(10.1)	69	(4.0)	3.51	(2.55-4.83)	1.61	(1.12-2.31)	1.56	(1.08-2.25)	1.37	(0.95-1.99)
Missing	107	(7.0)	60	(3.4)								
<b>Use of toothbrush</b>												
[complete dentures excluded]												
Yes †	1308	(86.2)	1639	(94.1)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
No	20	(1.3)	12	(0.7)	1.96	(0.94-4.06)	1.27	(0.54-3.01)	1.26	(0.53-3.00)	1.15	(0.48-2.80)
Missing	190	(12.5)	91	(5.2)								
<b>Use of toothpaste</b>												
[complete dentures excluded]												
Yes †	1279	(84.3)	1626	(93.3)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
No	47	(3.1)	24	(1.4)	2.26	(1.36-3.74)	1.40	(0.79-2.48)	1.38	(0.78-2.45)	1.33	(0.75-2.37)
Missing	192	(12.6)	92	(5.3)								
<b>Use of dental floss</b>												
[complete dentures excluded]												
Yes †	180	(11.9)	349	(20.0)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
No	1136	(74.8)	1296	(74.4)	1.57	(1.28-1.93)	1.26	(1.00-1.57)	1.24	(0.99-1.55)	1.20	(0.95-1.50)
Missing	202	(13.3)	97	(5.6)								

	Cases	(%)	Contr.	(%)	OR <sub>1</sub>	95%-CI	OR <sub>2</sub>	95%-CI	OR <sub>3</sub>	95%-CI	OR <sub>4</sub>	95%-CI
<b>Visiting a dentist</b>												
Every year †	583	(29.7)	965	(48.4)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
Every 2 - 5 years	401	(20.4)	423	(21.2)	1.68	(1.40-2.01)	1.32	(1.08-1.61)	1.30	(1.07-1.59)	1.25	(1.02-1.53)
Less than every 5 years	577	(29.4)	405	(20.3)	2.67	(2.22-3.20)	1.64	(1.34-2.01)	1.59	(1.29-1.95)	1.52	(1.23-1.87)
Never	345	(17.6)	176	(8.8)	3.87	(3.08-4.88)	2.13	(1.65-2.76)	2.02	(1.56-2.63)	1.93	(1.48-2.51)
Missing	57	(2.9)	24	(1.2)								
<b>Frequency of mouthwash</b>												
Never †	1199	(61.1)	1218	(61.1)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
Less than daily	407	(20.7)	447	(22.4)	0.94	(0.80-1.10)	1.04	(0.87-1.25)	1.05	(0.88-1.26)	1.06	(0.89-1.27)
1-2 times / day	290	(14.8)	300	(15.1)	1.02	(0.84-1.23)	1.01	(0.81-1.25)	1.03	(0.83-1.27)	1.06	(0.86-1.32)
3 and more times / day	36	(1.8)	16	(0.8)	2.31	(1.27-4.21)	2.91	(1.52-5.57)	3.01	(1.56-5.79)	3.23	(1.68-6.19)
Missing	31	(1.6)	12	(0.6)								

OR<sub>1</sub>: adjusted for age (9 categories), sex and study center

OR<sub>2</sub>: adjusted as OR<sub>1</sub> plus smoking status (never, former, current)

plus cumulative tobacco consumption (pack-years)

plus alcohol drinking duration (1-9 yrs, 10-19 yrs, 20-29 yrs, 30-39 yrs, 40+ yrs, missing)

plus a cumulative alcohol consumption (alcohol drinking frequency weighted by duration of the drinking period)

OR<sub>3</sub>: adjusted as OR<sub>2</sub> plus SES / professional education (primary school / worker, further school / clerks, University degree / manager)

OR<sub>4</sub>: adjusted as OR<sub>3</sub> plus consumption of fruits and vegetables (country-specific tertiles each)

† Reference category (ref.)

Table 4. Distribution of Indices for Oral Health and Dental Care With Corresponding ORs and 95%-Confidence Intervals.

	Cases (%)	Contr. (%)	OR <sub>1</sub>	95%-CI	OR <sub>2</sub>	95%-CI	OR <sub>3</sub>	95%-CI	OR <sub>4</sub>	95%-CI
<b>Oral Health Score *</b>										
OH Score=0 †	829 (42.2)	1122 (56.3)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
OH-Score =1-4	293 (14.9)	370 (18.6)	1.15	(0.96-1.38)	1.00	(0.82-1.23)	0.99	(0.81-1.21)	0.97	(0.79-1.19)
OH-Score =5,6	508 (25.9)	334 (16.8)	2.32	(1.94-2.77)	1.62	(1.33-1.97)	1.57	(1.29-1.92)	1.53	(1.25-1.87)
OH-Score =7	97 (4.9)	41 (2.1)	3.95	(2.68-5.83)	2.34	(1.54-3.57)	2.27	(1.49-3.47)	2.22	(1.45-3.41)
Missing **	236 (12.0)	126 (6.3)								
<b>Dental Care Score §</b>										
[complete dentures excluded]										
DC Score=0 †	298 (19.2)	675 (38.7)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)	1.00	(ref.)
DC Score =1	269 (17.7)	390 (22.4)	1.69	(1.36-2.09)	1.45	(1.15-1.83)	1.45	(1.15-1.83)	1.36	(1.07-1.72)
DC Score =2-4	619 (40.8)	521 (29.9)	3.17	(2.58-3.90)	2.04	(1.63-2.56)	2.01	(1.60-2.53)	1.88	(1.49-2.37)
DC Score =5-8	109 (7.2)	49 (2.8)	6.01	(4.07-8.87)	2.77	(1.79-4.27)	2.68	(1.73-4.15)	2.36	(1.51-3.67)
Missing §§	223 (14.7)	107 (6.1)								

OR<sub>1</sub>: adjusted for age (9 categories), sex and study center

OR<sub>2</sub>: adjusted as OR<sub>1</sub> plus smoking status (never, former, current)

plus cumulative tobacco consumption (pack-years)

plus alcohol drinking duration (1-9 yrs, 10-19 yrs, 20-29 yrs, 30-39 yrs, 40+ yrs, missing)

plus a cumulative alcohol consumption (alcohol drinking frequency weighted by duration of the drinking period)

OR<sub>3</sub>: adjusted as OR<sub>2</sub> plus SES / professional education (primary school, further school, University degree)

OR<sub>4</sub>: adjusted as OR<sub>3</sub> plus consumption of fruits and vegetables (country-specific tertiles each)

† Reference category (ref.)

\*Oral Health Score is the sum of the three following items:

**Denture:** No denture = 0, Partial denture (one jaw) = 1, Partial denture (upper and lower jaw) = 2, Complete denture (one jaw) = 3,  
Complete denture (upper and lower jaw)=4

**Age at starting to wear dentures:** No denture = 0, 55 years or older = 1, 35-54 years = 2, below 35 years = 3

**Gum bleeding:** never / sometimes and complete denture in the upper and lower jaw = 0, almost always = 1

\*\*Missing value for denture or age at start

§Dental Care Score is the sum of the three following items:

**Frequency of tooth cleaning:** At least twice/day = 0, once/day = 1, 1-4 times/week = 2, less or never = 3

**Use of toothbrush, toothpaste and dental floss:** Two or three of these = 0, only one of these three = 1, none of these three = 2

**Frequency of visiting a dentist:** At least once/year = 0, every 2-5 years=1, less than every 5 years=2, never=3

<sup>ss</sup>Missing value for at least one item

Table 5. Odds Ratios for Indices of Oral Health and Dental Care With Corresponding 95%-Confidence Intervals.

Frequency of	Mouth/ oropharynx					Hypopharynx/ larynx					Esophagus					Multiple sites						
	Contr.	(%)	Cases	(%)	OR <sub>4</sub>	95%-CI	Cases	(%)	OR <sub>4</sub>	95%-CI	Cases	(%)	OR <sub>4</sub>	95%-CI	Cases	(%)	OR <sub>4</sub>	95%-CI	Cases	(%)	OR <sub>4</sub>	95%-CI
<b>mouthwash</b>																						
Never †	1218	(61.1)	547	(58.6)	<b>1.00</b>	(ref.)	431	(61.7)	<b>1.00</b>	(ref.)	159	(67.9)	<b>1.00</b>	(ref.)	62	(63.9)	<b>1.00</b>	(ref.)	62	(63.9)	<b>1.00</b>	(ref.)
Less than daily	447	(22.4)	209	(22.4)	<b>1.10</b>	(0.89-1.38)	133	(19.1)	<b>0.99</b>	(0.76-1.29)	45	(19.2)	<b>1.14</b>	(0.77-1.70)	20	(20.6)	<b>0.94</b>	(0.51-1.72)	20	(20.6)	<b>0.94</b>	(0.51-1.72)
1-2 times / day	300	(15.1)	152	(16.3)	<b>1.11</b>	(0.86-1.44)	104	(14.9)	<b>1.08</b>	(0.80-1.47)	21	(9.0)	<b>0.76</b>	(0.44-1.33)	13	(13.4)	<b>0.99</b>	(0.48-2.03)	13	(13.4)	<b>0.99</b>	(0.48-2.03)
3 and more times / day	16	(0.8)	17	(1.8)	<b>3.53</b>	(1.65-7.57)	17	(2.4)	<b>3.50</b>	(1.55-7.89)	0	(0.0)	-	( - )	2	(2.1)	<b>3.89</b>	(0.59-25.52)	2	(2.1)	<b>3.89</b>	(0.59-25.52)
Missing	12	(0.6)	9	(1.0)			13	(1.9)			9	(3.8)			0	(0.0)			0	(0.0)		
<b>Oral Health</b>																						
<b>Score</b>																						
OH Score=0 †	1122	(56.3)	443	(47.4)	<b>1.00</b>	(ref.)	253	(36.2)	<b>1.00</b>	(ref.)	99	(42.3)	<b>1.00</b>	(ref.)	34	(35.1)	<b>1.00</b>	(ref.)	34	(35.1)	<b>1.00</b>	(ref.)
OH-Score =1-4	370	(18.6)	141	(15.1)	<b>0.94</b>	(0.73-1.21)	102	(14.6)	<b>1.00</b>	(0.74-1.36)	39	(16.7)	<b>1.05</b>	(0.68-1.64)	11	(11.3)	<b>0.63</b>	(0.30-1.36)	11	(11.3)	<b>0.63</b>	(0.30-1.36)
OH-Score =5,6	334	(16.8)	216	(23.1)	<b>1.31</b>	(1.03-1.68)	217	(31.1)	<b>1.94</b>	(1.48-2.55)	47	(20.1)	<b>1.08</b>	(0.70-1.67)	28	(28.9)	<b>1.98</b>	(1.04-3.75)	28	(28.9)	<b>1.98</b>	(1.04-3.75)
OH-Score =7	41	(2.1)	47	(5.0)	<b>2.00</b>	(1.21-3.31)	28	(4.0)	<b>2.14</b>	(1.18-3.88)	12	(5.1)	<b>2.43</b>	(0.98-5.99)	10	(10.3)	<b>2.56</b>	(0.96-6.84)	10	(10.3)	<b>2.56</b>	(0.96-6.84)
Missing *	126	(6.3)	87	(9.3)			98	(14.0)			37	(15.8)			14	(14.4)			14	(14.4)		
<b>Dental Care</b>																						
<b>Score</b>																						
DC Score=0 †	675	(38.7)	163	(22.1)	<b>1.00</b>	(ref.)	81	(15.5)	<b>1.00</b>	(ref.)	46	(24.1)	<b>1.00</b>	(ref.)	8	(11.8)	<b>1.00</b>	(ref.)	8	(11.8)	<b>1.00</b>	(ref.)
DC Score =1	390	(22.4)	138	(18.7)	<b>1.37</b>	(1.02-1.83)	90	(17.2)	<b>1.49</b>	(1.04-2.16)	28	(14.7)	<b>0.72</b>	(0.41-1.26)	13	(19.1)	<b>2.33</b>	(0.89-6.11)	13	(19.1)	<b>2.33</b>	(0.89-6.11)
DC Score =2-4	521	(29.9)	308	(41.8)	<b>1.95</b>	(1.47-2.59)	219	(42.0)	<b>2.00</b>	(1.40-2.86)	62	(32.5)	<b>1.21</b>	(0.72-2.03)	30	(44.1)	<b>3.60</b>	(1.43-9.06)	30	(44.1)	<b>3.60</b>	(1.43-9.06)
DC Score =5-8	49	(2.8)	46	(6.2)	<b>2.23</b>	(1.29-3.84)	45	(8.6)	<b>2.89</b>	(1.57-5.32)	13	(6.8)	<b>1.95</b>	(0.81-4.69)	5	(7.4)	<b>3.54</b>	(0.84-14.84)	5	(7.4)	<b>3.54</b>	(0.84-14.84)
Missing §	107	(6.1)	82	(11.1)			87	(16.7)			42	(22.0)			12	(17.6)			12	(17.6)		

OR<sub>4</sub>: adjusted for age (9 categories), sex and study center; plus smoking status (never, former, current)

plus cumulative tobacco consumption (pack-years)

plus alcohol drinking duration (1-9 yrs, 10-19 yrs, 20-29 yrs, 30-39 yrs, 40+ yrs, missing)

plus a cumulative alcohol consumption (alcohol drinking frequency weighted by duration of the drinking period)

plus SES / professional education (primary school, further school, University degree)

plus consumption of fruits and vegetables (country-specific tertiles each)

† Reference category (ref.)

\*Missing value for denture or age at start

§Missing value for at least one item




Figure 1 Odds ratio (OR) of upper aero-digestive cancer by rs1229984 (ADH1B), rs1573496 (ADH7) and rs1693482 (ADH1C) genotype. Rare allele carriers (dominant model) versus common allele homozygous genotype. ORs are standardized by age, sex, centre, cumulative alcohol consumption and, when relevant, smoking. P values for heterogeneity are derived from a Chi Square statistic with 2df. P values for trend test are derived from a Chi Square statistic with 1df.