

# NIH Public Access

**Author Manuscript** 

JAm Chem Soc. Author manuscript; available in PMC 2011 April 14.

#### Published in final edited form as:

*JAm Chem Soc.* 2010 April 14; 132(14): 5010–5011. doi:10.1021/ja1017046.

## Enantioselective Alkenylation via Nickel-Catalyzed Cross-Coupling with Organozirconium Reagents

#### Sha Lou and Gregory C. Fu\*

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

### Abstract



A new family of organometallic compounds, organozirconium reagents, are shown to serve as suitable partners in cross-coupling reactions of (activated) secondary alkyl electrophiles. Thus, the first catalytic method for coupling secondary  $\alpha$ -bromoketones with alkenylmetal reagents has been developed, specifically a mild, versatile, and stereoconvergent carbon–carbon bond-forming process that generates potentially labile  $\beta$ , $\gamma$ -unsaturated ketones with good enantioselectivity.

Because olefin-containing molecules are ubiquitous targets,<sup>1</sup> the development of effective methods for incorporating this functional group is an important goal. If absolute stereochemistry can also be defined during the bond construction, the utility of a new process is further enhanced. The reaction of secondary alkyl electrophiles with alkenylmetal reagents is an attractive approach to the introduction of olefins, but there are relatively few reports of such cross-couplings.<sup>2,3</sup> Furthermore, there has been only one investigation of an enantioselective variant of this type of process (three examples of cross-couplings of  $\alpha$ -haloesters with alkenyltrimethoxysilanes).<sup>3b</sup>

Alkenylzirconium compounds are appealing alkenylmetals for cross-couplings,<sup>4</sup> in part because they are readily accessible by reacting commercially available Schwartz's reagent (Cp<sub>2</sub>ZrHCl) with alkynes. Although secondary alkyl electrophiles have now been coupled with many different families of organometallic partners (e.g., boron, zinc, magnesium, silicon, tin, and indium compounds),<sup>5</sup> to the best of our knowledge, they have not been cross-coupled with organozirconium reagents.<sup>6</sup> In this report, we describe a mild and versatile method for coupling organozirconium compounds with secondary alkyl halides, specifically alkenylzirconium reagents with racemic  $\alpha$ -bromoketones; in addition, we establish that this stereoconvergent carbon–carbon bond-forming process can be accomplished with good enantioselectivity to generate potentially labile  $\beta$ , $\gamma$ -unsaturated ketones (eq 1).<sup>7,8,9</sup>

gcf@mit.edu.

Supporting Information Available: Experimental procedures and compound characterization data (PDF). This material is available free of charge via the Internet at http://pubs.acs.org.

Lou and Fu



In initial studies, we attempted to apply previously described methods for nickel-catalyzed asymmetric cross-couplings<sup>8,9</sup> to the alkenylation of 2-bromo-1-phenylpropan-1-one; unfortunately, we did not obtain the desired product in satisfactory yield or ee. We therefore decided to explore the potential utility of a new family of coupling partners, alkenylzirconium reagents. In addition to their ready availability, we anticipated that they could provide another crucial attribute: carbon–carbon bond formation might be achieved under sufficiently mild, non-basic conditions that racemization of the  $\alpha$  stereocenter and isomerization to the  $\alpha$ , $\beta$ -unsaturated isomer would be avoided. Such complications have thus far precluded a Buchwald-Hartwig-type approach to the catalytic asymmetric alkenylation of carbonyl compounds to generate  $\alpha$  tertiary stereocenters.<sup>10</sup>

We have determined that, under the appropriate conditions, a nickel/bis(oxazoline) catalyst can achieve the stereoconvergent cross-coupling of racemic  $\alpha$ -bromoketones with alkenylzirconium compounds to generate  $\beta$ , $\gamma$ -unsaturated ketones in good yield and ee (Table 1).<sup>11</sup> Carbon–carbon bond formation occurs below ambient temperature (10 °C) and without the need for any additives (e.g., Brønsted bases) that might erode the enantioselectivity. As illustrated in Table 1, a broad array of alkenylzirconium reagents are suitable cross-coupling partners. Thus, the R<sup>2</sup> group can range in steric demand from hydrogen to *t*-butyl (entries 1–5). Furthermore, functional groups such as aromatic rings, oxygen substituents, and alkenes are compatible with the reaction conditions (entries 3 and 6–9).

We have also examined the scope of this new cross-coupling process with respect to the  $\alpha$ bromoketone. As illustrated in Table 2, a diverse set of aryl alkyl ketones are suitable electrophiles. A variety of different types of substituents can be present on the aromatic ring (electron-donating or electron-withdrawing: entries 4–7; ortho, meta, or para: entries 4–9), and the aromatic group can be a heterocycle (e.g., a thiophene in entry 10). In addition, an array of alkyl groups on the ketone and R<sup>2</sup> substituents on the alkenylzirconium reagent are tolerated.

The same method can be applied directly to enantioselective cross-couplings of dialkyl ketones with alkenylzirconium reagents (Table 3). This contrasts with our study of asymmetric Kumada reactions of ketones with aryl Grignard reagents, wherein different

JAm Chem Soc. Author manuscript; available in PMC 2011 April 14.

Lou and Fu

coupling conditions (ligand and temperature) were necessary for aryl alkyl ketones versus dialkyl ketones.9c

The chiral nickel/bis(oxazoline) catalyst can dictate the stereochemical outcome of a crosscoupling of an  $\alpha$ -bromoketone that bears another stereocenter (i.e., catalyst-controlled stereoselectivity: eq 2 and eq 3). Furthermore, stereoselective functionalizations of the crosscoupling product can be achieved (eq 4).





(4)

In summary, we have demonstrated that a new family of organometallic compounds, organozirconium reagents, can serve as suitable partners in cross-coupling reactions of (activated) secondary alkyl electrophiles. Thus, we have developed the first catalytic method for coupling secondary  $\alpha$ -haloketones with alkenylmetal reagents, specifically a mild, versatile, and stereoconvergent carbon-carbon bond-forming process that generates potentially labile  $\beta$ , $\gamma$ -unsaturated ketones with good enantioselectivity. Additional efforts to expand the scope of cross-couplings of alkyl electrophiles are underway.

#### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

#### Acknowledgments

Support has been provided by the National Institutes of Health (National Institute of General Medical Sciences, grant R01-GM62871), Merck Research Laboratories, and Novartis. We thank Shaun D. Fontaine, Dr. Peter Mueller, and Dr. Michael R. Reithofer for assistance.

J Am Chem Soc. Author manuscript; available in PMC 2011 April 14.

#### References

- 1. For example, see quinine, epothilone A, and palytoxin.
- 2. For leading references to reactions of aryl electrophiles with alkenylmetal reagents, see: Denmark SE, Butler CR. Chem Commun 2009:20–33.
- For some examples, see: Unactivated electrophiles: (a) Zhou J-R, Fu GC. J Am Chem Soc 2004;126:1340–1341. [PubMed: 14759182] Powell DA, Maki T, Fu GC. J Am Chem Soc 2005;127:510–511. [PubMed: 15643860] Guérinot A, Reymond S, Cossy J. Angew Chem, Int Ed 2007;46:6521–6524.Czaplik WM, Mayer M, von Wangelin AJ. Angew Chem, Int Ed 2009;48:607– 610.Activated electrophiles (asymmetric): (b) Dai X, Strotman NA, Fu GC. J Am Chem Soc 2008;130:3302–3303. [PubMed: 18302392]
- For an overview, see: Negishi, E-i; Zeng, X.; Tan, Z.; Qian, M.; Hu, Q.; Huang, ZIn. Metal-Catalyzed Cross-Coupling Reactions. de Meijere, A.; Diederich, F., editors. Wiley–VCH; New York: 2004. p. 815-889.
- 5. For leading references, see: Rudolph A, Lautens M. Angew Chem, Int Ed 2009;48:2656–2670.
- We are aware of one report of the cross-coupling of a primary alkyl electrophile with an organozirconium reagent: Wiskur SL, Korte A, Fu GC. J Am Chem Soc 2004;126:82–83. [PubMed: 14709070]
- 7. For a catalytic asymmetric method for the  $\alpha$ -alkenylation of aldehydes to generate tertiary stereocenters, see: Kim H, MacMillan DWC. J Am Chem Soc 2008;130:398–399. [PubMed: 18095690]
- For an overview and leading references to asymmetric cross-couplings of secondary alkyl halides, see: Glorius F. Angew Chem, Int Ed 2008;47:8347–8349.
- 9. For recent work on asymmetric cross-couplings of secondary alkyl halides, see: (a) Smith SW, Fu GC. J Am Chem Soc 2008;130:12645–12647. [PubMed: 18763769] (b) Lundin PM, Esquivias J, Fu GC. Angew Chem, Int Ed 2009;48:154–156. (c) Lou S, Fu GC. J Am Chem Soc 2010;132:1264–1266. [PubMed: 20050651]
- For a discussion and leading references to catalytic asymmetric alkenylations of enolates to generate quaternary stereocenters, see: Taylor AM, Altman RA, Buchwald SL. J Am Chem Soc 2009;131:9900–9901. [PubMed: 19580273]
- 11. Notes: (a) In the absence of NiCl<sub>2</sub>•glyme or ligand **1**, the desired product is generated in <5% yield. NiBr<sub>2</sub>•glyme or Ni(cod)<sub>2</sub> may be used in place of NiCl<sub>2</sub>•glyme. (b) For the cross-coupling illustrated in entry 3 of Table 1: a reaction on a gram-scale proceeded at room temperature in 90% yield and 92% ee; use of 1.1, rather than 2, equivalents of the organozirconium reagent led to a slower reaction (after 3 days at room temperature: 85% yield, 91% ee). (c) During the cross-coupling, no kinetic resolution of the starting material (the racemic secondary alkyl bromide) is detected, and the ee of the product is constant. (d) After a competition experiment with an  $\alpha$ -bromoketone, an unactivated secondary alkyl bromide was recovered in essentially quantitative yield. (e) In preliminary studies under our standard conditions, an  $\alpha$ -chloroketone, a cyclic  $\alpha$ -bromoketone, and an alkenylzirconium reagent derived from hydrozirconation of an internal alkyne were not suitable cross-coupling partners. (f) The rate law for the cross-coupling is first-order in the catalyst and in the nucleophile, and it is zero-order in the electrophile (NMR studies in d<sub>8</sub>-THF).

#### Table 1

Catalytic Asymmetric Alkenylations with Organozirconium Reagents: Scope with Respect to the Nucleophile



All data are the average of two experiments.

<sup>a</sup>Yield of purified product.

<sup>b</sup>Catalyst loading: 10% NiCl<sub>2</sub> • glyme/12% (–)-1.

	ee (%)	92	80	8
In respect to the call of the	yield (%) <sup>a</sup>	83	95	∞ ∞
	$\mathbb{R}^2$	(CH <sub>2</sub> ) <sub>4</sub> Cl	<i>m</i> -tolyl	CH <sub>2</sub> Ph
		$\mathbf{R}^{1} = \mathbf{E} \mathbf{t}$	CH <sub>2</sub> CH <sub>2</sub> CI	<i>i-</i> Bu
				بع م
	entry	-	2	5 JAm Chem Soc. Author manuscript; available in PMC 2011 April 14.

NIH-PA Author Manuscript

Table 2

**NIH-PA** Author Manuscript

.

**NIH-PA** Author Manuscript



Page 7

**NIH-PA** Author Manuscript

**NIH-PA** Author Manuscript

#### Table 3

Catalytic Asymmetric Alkenylations with Organozirconium Reagents: Scope with Respect to the Dialkyl Ketone (for the reaction conditions, see eq 1)



Lou and Fu



All data are the average of two experiments.

<sup>a</sup>Yield of purified product.

 $^b$  The reaction was run at 40 °C.