



Published in final edited form as:

J Am Chem Soc. 2015 March 11; 137(9): 3157–3160. doi:10.1021/ja512015w.

Alkyne Hydroacylation: Switching Regioselectivity by Tandem Ruthenium Catalysis

Qing-An Chen, Faben A. Cruz, and Vy M. Dong*

Department of Chemistry, University of California, Irvine, 4403 Natural Sciences 1, Irvine, CA 92697, United States

Abstract

By using tandem ruthenium-catalysis, internal alkynes can be coupled with aldehydes for the synthesis of β,γ -unsaturated ketones. The catalyst promotes alkyne transformations with high regioselectivity, with examples that include the differentiation of a methyl versus ethyl substituent on the alkyne. Mechanistic studies suggest that the regioselectivity results from a selective allene formation that is governed by allylic strain.

In contrast to an enzyme, which is typically specific, a single transition-metal complex can catalyze diverse transformations. This generality can be exploited in *tandem catalysis*, a strategy where one catalyst promotes two or more transformations in a cascade.¹ Ru-complexes are especially versatile in developing efficient cascade reactions, including examples that feature olefin metathesis,² reductive C–C bond coupling,³ and amide synthesis.⁴ It occurred to us that tandem catalysis⁵ could be used to overcome challenges in hydroacylation, the addition of an aldehyde C–H bond across an unsaturated π -bond.⁶ Towards expanding ketone hydroacylation, we reported a cascade involving alcohol oxidation, enantioselective ketone reduction, and lactol oxidation to generate lactones, a process in which all three transformations were promoted by Noyori's Ru-catalyst.^{5a} Herein, we apply tandem Ru-catalysis to achieve the hydroacylation of alkynes with unprecedented regiocontrol to afford β,γ -unsaturated ketones.

The coupling of an aldehyde to an alkyne is a promising route to ketones, which has been studied using both transition metal catalysts (*e.g.*, Ni,⁷ Rh,^{8–9} Ru,¹⁰ Ir,¹¹ Au,¹² Cu¹³) and organocatalysts (*e.g.*, *N*-heterocyclic carbenes) (Figure 1a). In alkyne hydroacylation, α,β -unsaturated ketones are afforded as the major constitutional isomer for both intramolecular and intermolecular variants. Most of these methods require terminal or symmetric alkynes to achieve high regioselectivity. Others require aldehydes bearing directing groups to promote C–H bond functionalization in preference to decarbonylation.

*Corresponding Author: dongv@uci.edu.

Supporting Information

Experimental procedures and spectral data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

The authors declare no competing financial interests.

In light of these challenges, we proposed using a metal-hydride catalyst to perform a cascade involving (1) alkyne-allene isomerization and (2) allene-aldehyde coupling (Figure 1b). This protocol would tolerate a wider scope of coupling partners and provide access to an alternative ketone isomer. We judged this cascade to be feasible on the basis of literature precedence for metal-hydride catalysis of the independent transformations. Yamamoto demonstrated alkynes as convenient surrogates for allenes in Pd-catalyzed hydroamination.¹⁵ Ishii¹⁶ and Breit¹⁷ achieved alkyne isomerizations by Ir and Rh metal-hydride pathways, respectively. Krische proposed that Ru(0) isomerized alkynes to allenes, albeit by a C–H propargyl activation mechanism.¹⁸ While Willis has reported allene hydroacylations with chelating aldehydes,¹⁹ we envisioned the resulting allene would undergo metal-hydride addition to generate a metal-allyl species that could couple to non-chelating aldehydes (Figure 1b). This mechanistic step has been shown in the coupling of allenes and aldehydes to generate homoallylic alcohols.²⁰ In our proposal, subsequent β -hydride elimination would generate the desired β,γ -unsaturated ketone and regenerate the metal-hydride catalyst. Thus, we set out to identify a single metal catalyst that could promote both transformations in sequence to afford a novel alkyne hydroacylation.


To test our hypothesis, we chose the coupling of benzaldehyde **1a** and 2-pentyne **2a** as the model. The unsymmetrical alkyne **2a** is a challenging substrate for regioselective transformations due to the similar steric bulk of methyl and ethyl substituents (A-values of 1.74 versus 1.79 kcal/mol, respectively).²¹ An initial survey of commercially available metal hydrides complexes (Rh, Ru, Ir) revealed that RuHCl(CO)(PPh₃)₃ was a promising catalyst precursor (Table 1). Regioselectivity for β,γ -unsaturated ketone **3aa** over **4aa** and **5aa** depended on the structure of the ligand chosen (entries 1–7). Only traditional hydroacylation products (α,β -unsaturated ketones) were obtained (**4aa** and **5aa**) using non ferrocene-based ligands, BINAP and Tangphos (entries 1–2). Switching to ferrocene-based ligands gave the desired β,γ -unsaturated product **3aa** (entries 3–6).²² Josiphos **L1** provided the highest reactivity and regioselectivity (entry 5). Removing the backbone methyl group in Josiphos **L1** resulted in a decrease in regioselectivity (entry 7 vs 5). Increasing the catalyst loading to 4 mol% improved the yield (entry 8). We used racemic **L1** in these initial studies. The stereocenter of Josiphos **L1** has little impact on enantioselectivity²³ but appears essential for both regioselectivity and reactivity.²⁴

With this protocol, we explored the hydroacylation of 2-pentyne **2a** with various aldehydes **1** (Table 2). Moderate to good yields (66–85%) and high regioselectivities (>20:1) were obtained with aldehydes **1** possessing varying electronic properties (entries 1–6). The coupling of **2a** tolerated aldehydes **1g–h** with increased steric bulk (72–78% yield, entries 7–8). Heteroaromatic β,γ -unsaturated ketones were obtained through the hydroacylation of 2-pentyne **2a** with aldehydes bearing furan (**1i–j**), thiophene (**1k**), and indole (**1l**) rings (entries 9–12).

We next examined the scope of alkynes (Table 3) and found that secondary (Cy) and tertiary (^tBu) alkyl substituted alkynes were suitable coupling partners (entries 1–2). High isolated yields (83–95%) and regioselectivities (>20:1) were observed in the hydroacylation of 1-phenyl-1-propyne with various aldehydes (entries 3–9). Alkynes bearing aryl groups of varying electronic and steric properties could be coupled with aldehyde **1b** with 61–83%

yields (entries 10–15). Switching the methyl group to an ethyl group by using in 1-phenyl-1-butyne resulted in no desired β,γ -unsaturated ketone product (entry 16). This lack of reactivity may be ascribed to increased allylic strain disfavoring β -hydride elimination to form the allene (Scheme 1).


To support the proposed allene intermediate, we prepared phenylallene **6** independently and then subjected this allene to benzaldehyde **1a** under otherwise standard conditions (eq. 1). This control experiment provides the same product **3ad** as that obtained in the hydroacylation 1-phenyl-1-propyne. While allene **6** was completely consumed, multiple unidentified side products were observed. We propose that tandem catalysis overcomes possible decomposition pathways by forming allenes in low concentration. A similar phenomenon was observed by Krische in his recent study on the synthesis of homoallylic alcohols.¹⁸




(1)

On the basis of literature and our own observations, we propose the mechanism featuring tandem Ru-catalysis (Scheme 1). The reversible insertion of 2-pentyne **2a** into ruthenium hydride (Ru–H) can yield Ru-vinyl intermediate **A** or **B**. The interception of intermediate **A** with aldehyde **1** generates isomer **5**, whereas interception with **B** generates isomer **4**. Indeed, we observed formation of both **5aa** and **4aa** (~3:1 ratio) when using BINAP or TangPhos (Table 1, entries 1–2). While formation of both **A** and **B** are feasible, we propose that β -hydride elimination from **B** to generate the 1,3-disubstituted allene **D** will be disfavored due to higher allylic strain. In contrast, the β -hydride elimination of metal-vinyl complex **A** will form 1-substituted allene **C** *in situ* and regenerate a Ru–H species.²⁶ The reinsertion of Ru–H into the terminal allene **C** will occur regioselectively to yield Ru-allyl **E**. Trapping of aldehyde **1** through a six-membered intermediate **F** yields ruthenium alkoxide **G**.²⁰ A second β -hydride elimination from **G** delivers the observed ketone **3** and regenerates the Ru–H.

When the hydroacylation was performed with deuterio-alkyne **2d-d₃**, the deuterium label was scrambled into the α -, β - and γ -positions of ketone **3ad-d₃** (eq. 2). The incorporation of deuterium into both α - and β - positions of ketone **3ad-d₃** suggests that the Ru–H species generated from both β -H elimination steps (**A** to **C** and **G** to **3**) are indistinguishable (Scheme 1). The observed incorporation of hydrogen at the γ -position of product **3ad-d₃** indicates the reversibility of β -hydride elimination in allene formation (Scheme 1). This hypothesis is supported by deuterium scrambling in the isotope-labeling study with aldehyde **1a-d₁** (eq. 3).



(2)



(3)

This tandem strategy can be used to access other useful motifs. For example, we investigated the coupling of aldehyde **1b** with 1-(trimethylsilyl)propyne **7** for the formation of silyl dienol ether **8** (Scheme 2). A 1,3-silyl migration²⁷ of intermediate **9** generates a nucleophile that can be used in asymmetric vinylogous aldol reactions.²⁸

In addition, through a procedure involving tandem Ru-catalysis followed by base-catalyzed isomerization, we can generate α,β -unsaturated ketones **4** with 70–87% yields (Scheme 3). This one-pot protocol allows us to differentiate between the methyl and ethyl groups on 2-pentyne **2a** with high regiocontrol and generate α,β -unsaturated ketones that would be difficult to access using traditional hydroacylation methods.

By identifying the Ru-L1 catalyst, we have been able to expand the power of aldehyde-alkyne cross couplings through a cascade mechanism that enables unique regioselectivity. Due to the formation of a 1-substituted allene intermediate, our protocol differentiates a methyl from an ethyl substituent on 2-pentyne with high regioselectivity. Future studies will focus on stereoselective variants. More importantly, insights from these studies will contribute to the emerging use of alkynes as allene surrogates and guide future developments in tandem catalysis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding provided by UC Irvine and the National Institutes of Health (GM105938). We are grateful to Eli Lilly for a Grantee Award. We thank Michael Yamano for preparing phenylallene **6**.


References

1. For a definition and review on tandem catalysis, see a) Fogg DE, dos Santos EN. *Coord Chem Rev.* 2004; 248:2365. b) Chapman CJ, Frost CG. *Synthesis.* 2007; 1c) Shindoh N, Takemoto Y, Takasu K. *Chem Eur J.* 2009; 15:12168. [PubMed: 19847824]

2. a) Alcaide B, Almendros P, Luna A. *Chem Rev.* 2009; 109:3817. [PubMed: 19572654] b) Vougioukalakis GC, Grubbs RH. *Chem Rev.* 2010; 110:1746. [PubMed: 20000700]
3. a) Zbieg JR, Yamaguchi E, McInturff EL, Krische MJ. *Science.* 2012; 336:324. [PubMed: 22442385] b) Geary LM, Glasspoole BW, Kim MM, Krische MJ. *J Am Chem Soc.* 2013; 135:3796. [PubMed: 23448269] c) Park BY, Montgomery TP, Garza VJ, Krische MJ. *J Am Chem Soc.* 2013; 135:16320. [PubMed: 24156560]
4. a) Gunanathan C, Ben-David Y, Milstein D. *Science.* 2007; 317:790. [PubMed: 17690291] b) Gunanathan C, Milstein D. *Chem Rev.* 2014; 10.1021/cr5002782
5. For examples of tandem catalysis from our laboratory, see: a) Murphy SK, Dong VM. *J Am Chem Soc.* 2013; 135:5553. [PubMed: 23566215] b) Oldenhuis NJ, Dong VM, Guan Z. *J Am Chem Soc.* 2014; 136:12548. [PubMed: 25170560] c) Dornan PK, Kou KGM, Houk KN, Dong VM. *J Am Chem Soc.* 2014; 136:291. [PubMed: 24350903]
6. For selected reviews on hydroacylation, see: a) Bosnich B. *Acc Chem Res.* 1998; 31:667. b) Fu, GC. *Modern Rhodium-Catalyzed Reactions.* Evan, PA., editor. Wiley-VCH; New York: 2005. p. 79-91. c) Jun CH, Jo EA, Park JW. *Eur J Org Chem.* 2007; 1869d) Willis MC. *Chem Rev.* 2010; 110:725. [PubMed: 19873977] e) Leung JC, Krische MJ. *Chem Sci.* 2012; 3:2202.
7. a) Tsuda T, Kiyoi T, Saegusa T. *J Org Chem.* 1990; 55:2554. b) Yang F, Jin T, Yamamoto Y. *Tetrahedron.* 2012; 68:5223.
8. a) Kokubo K, Matsumasa K, Miura M, Nomura M. *J Org Chem.* 1997; 62:4564. b) Jun CH, Lee H, Hong JB, Kwon BI. *Angew Chem Int Ed.* 2002; 41:2146. c) Willis MC, Randell-Sly HE, Woodward RL, McNally SJ, Currie GS. *J Org Chem.* 2006; 71:5291. [PubMed: 16808518] d) González-Rodríguez C, Parsons SR, Thompson AL, Willis MC. *Chem Eur J.* 2010; 16:10950. [PubMed: 20715218] e) González-Rodríguez C, Pawley RJ, Chaplin AB, Thompson AL, Weller AS, Willis MC. *Angew Chem Int Ed.* 2011; 50:5134. f) Lenden P, Entwistle DA, Willis MC. *Angew Chem Int Ed.* 2011; 50:10657. g) Poingdestre S-J, Goodacre JD, Weller AS, Willis MC. *Chem Commun.* 2012; 48:6354. h) Chaplin AB, Hooper JF, Weller AS, Willis MC. *J Am Chem Soc.* 2012; 134:4885. [PubMed: 22324763] i) Pawley RJ, Huertos MA, Lloyd-Jones GC, Weller AS, Willis MC. *Organometallics.* 2012; 31:5650. j) Castaing M, Wason SL, Estepa B, Hooper JF, Willis MC. *Angew Chem Int Ed.* 2013; 52:13280.
9. a) Tanaka K, Fu GC. *J Am Chem Soc.* 2001; 123:11492. [PubMed: 11707133] b) Tanaka K, Fu GC. *J Am Chem Soc.* 2002; 124:10296. [PubMed: 12197729] c) Tanaka K, Fu GC. *J Am Chem Soc.* 2003; 125:8078. [PubMed: 12837058] d) Takeishi K, Sugishima K, Sasaki K, Tanaka K. *Chem Eur J.* 2004; 10:5681. [PubMed: 15470696] e) Chung LW, Wiest O, Wu Y-D. *J Org Chem.* 2008; 73:2649. [PubMed: 18324834]
10. a) Williams VM, Leung JC, Patman RL, Krische MJ. *Tetrahedron.* 2009; 65:5024. [PubMed: 20613891] b) Miura H, Wada K, Hosokawa S, Inoue M. *Chem Eur J.* 2013; 19:861. [PubMed: 23233450]
11. Hatanaka S, Obora Y, Ishii Y. *Chem Eur J.* 2010; 16:1883. [PubMed: 20029924]
12. Shi S, Wang T, Weingand V, Rudolph M, Hashmi ASK. *Angew Chem Int Ed.* 2014; 53:1148.
13. Chen S, Li X, Zhao H, Li B. *J Org Chem.* 2014; 79:4137. [PubMed: 24724722]
14. a) Biju AT, Wurz NE, Glorius F. *J Am Chem Soc.* 2010; 132:5970. [PubMed: 20384303] b) Wang Z, Yu Z, Wang Y, Shi D. *Synthesis.* 2012; 44:1559.
15. a) Kadota I, Shibuya A, Gyoung YS, Yamamoto Y. *J Am Chem Soc.* 1998; 120:10262. b) Kadota I, Shibuya A, Lutete LM, Yamamoto Y. *J Org Chem.* 1999; 64:4570. [PubMed: 11674520] c) Lutete LM, Kadota I, Yamamoto Y. *J Am Chem Soc.* 2004; 126:1622. [PubMed: 14871079]
16. Obora Y, Hatanaka S, Ishii Y. *Org Lett.* 2009; 11:3510. [PubMed: 19719194]
17. a) Lumbroso A, Koschker P, Vautravers NR, Breit B. *J Am Chem Soc.* 2011; 133:2386. [PubMed: 21291216] b) Lumbroso A, Abermil N, Breit B. *Chem Sci.* 2012; 3:789. c) Xu K, Khakyzadeh V, Bury T, Breit B. *J Am Chem Soc.* 2014; 136:16124. [PubMed: 25365339]
18. Park BY, Nguyen KD, Chaulagain MR, Komanduri V, Krische MJ. *J Am Chem Soc.* 2014; 136:11902. [PubMed: 25075434]
19. a) Kokubo K, Matsumasa K, Nishinaka Y, Miura M, Nomura M. *Bull Chem Soc Jpn.* 1999; 72:303. b) Osborne JD, Randell-Sly HE, Currie GS, Cowley AR, Willis MC. *J Am Chem Soc.*

- 2008; 130:17232. [PubMed: 19053453] c) Randell-Sly HE, Osborne JD, Woodward RL, Currie GS, Willis MC. *Tetrahedron*. 2009; 65:5110.
20. a) Bower JF, Skucas E, Patman RL, Krische MJ. *J Am Chem Soc*. 2007; 129:15134. [PubMed: 18020342] b) Skucas E, Bower JF, Krische MJ. *J Am Chem Soc*. 2007; 129:12678. [PubMed: 17900123] c) Ngai MY, Rucas E, Krische MJ. *Org Lett*. 2008; 10:2705. [PubMed: 18533665] d) Han SB, Kim IS, Han H, Krische MJ. *J Am Chem Soc*. 2009; 131:6916. [PubMed: 19453190] e) Skucas E, Zbieg JR, Krische MJ. *J Am Chem Soc*. 2009; 131:5054. [PubMed: 19317402] f) Zbieg JR, McInturff EL, Krische MJ. *Org Lett*. 2010; 12:2514. [PubMed: 20459077] g) Moran J, Preetz A, Mesch RA, Krische MJ. *Nature Chem*. 2011; 3:287. [PubMed: 21430686] h) Zbieg JR, McInturff EL, Leung JC, Krische MJ. *J Am Chem Soc*. 2011; 133:1141. [PubMed: 21175178] i) Sam B, Montgomery TP, Krische MJ. *Org Lett*. 2013; 15:3790. [PubMed: 23841678]
21. a) Booth H, Everett JR. *J Chem Soc Perk Trans*. 2(1980):255. b) Eliel, EL.; Wilen, SH.; Mander, LN. *Stereochemistry of Organic Compounds*. Wiley-VCH; New York: 1994.
22. For examples of synthesis β,γ -unsaturated ketones by hydroacclation of dienes, see: a) Omura S, Fukuyama T, Horiguchi J, Murakami Y, Ryu I. *J Am Chem Soc*. 2008; 130:14094. [PubMed: 18841894] b) Shibahara F, Bower JF, Krische MJ. *J Am Chem Soc*. 2008; 130:14120. [PubMed: 18841895] c) Chen Q-A, Kim DK, Dong VM. *J Am Chem Soc*. 2014; 136:3772. [PubMed: 24588202]
23. Although we could not separate the two enantiomers of **3aa** by chiral SFC analysis. We observed an ee of 10% for the synthesis of an analogous **3ad** using (*R,S_p*)-**L1**. Efforts to develop more efficient chiral ligands are warranted.
24. a) Shen QL, Shekhar S, Stambuli JP, Hartwig JF. *Angew Chem Int Ed*. 2005; 44:1371. b) Shen Q, Hartwig JF. *J Am Chem Soc*. 2006; 128:10028. [PubMed: 16881628] c) Shen Q, Ogata T, Hartwig JF. *J Am Chem Soc*. 2008; 130:6586. [PubMed: 18444639]
25. Hoffmann RW. *Chem Rev*. 1989; 89:1841.
26. Hill AF, Ho CT, WiltonEly J. *Chem Commun*. 1997; 2207
27. Takahashi M, Kira M. *J Am Chem Soc*. 1999; 121:8597.
28. Denmark SE, Heemstra JR Jr. *J Org Chem*. 2007; 72:5668. [PubMed: 17583959]

a) Previous alkyne hydroacylations by various catalysts



b) Proposed alkyne hydroacylation by tandem Ru-catalysis







Figure 1.
Intermolecular Hydroacylation of Alkynes

**Scheme 1.**

Tandem Ru-Catalysis: Mechanism for Hydroacylation of Alkynes




Scheme 2.
Silyl Dienol Ether Prepared by Tandem Catalysis




Scheme 3.
Regioselective Hydroacylation of 2-Pentyne.

Table 1

Ligand Effects on Hydroacylation of 2-Pentyne^a

Entry	Ligand	Yield (%) ^b		
		3aa	4aa	5aa
1	BINAP	0	1	3
2	Tangphos	0	5	18
3	dppf	11	43	40
4	dcypf	30	0	25
5	L1	54	0	2
6	L2	32	3	58
7	L3	41	0	8
8 ^c	L1	81	1	2




^a **1a** (0.20 mmol), **2a** (0.60 mmol), RuHCl(CO)(PPh₃)₃ (2 mol%), Ligand (2 mol%), toluene (0.5 mL), 80 °C, 18 h.

^b Determined by ¹H NMR or GC-FID with 1,3,5-trimethoxybenzene (0.05 mmol) as the internal standard.

^c RuHCl(CO)(PPh₃)₃ (4 mol%), **L1** (4 mol%), 15 h.

Table 2


Hydroacylation with Various Aldehydes^a

Entry	R in 1	Yield (%) ^b
1	Ph (1a)	81 (3aa)
2	4-MeOC ₆ H ₄ (1b)	69 (3ba)
3	4-MeC ₆ H ₄ (1c)	85 (3ca)
4	4-FC ₆ H ₄ (1d)	79 (3da)
5	4-ClC ₆ H ₄ (1e)	74 (3ea)
6	4-BrC ₆ H ₄ (1f)	66 (3fa)
7	3-MeOC ₆ H ₄ (1g)	78 (3ga)
8	2-MeOC ₆ H ₄ (1h)	72 (3ha)
9	2-Furyl (1i)	71 (3ia)
10	3-Furyl (1j)	76 (3ja)
11	2-Thienyl (1k)	87 (3ka)
12	3- <i>N</i> -Ts-indolyl (1l)	83 (3la)

^a **1** (0.20 mmol), **2a** (0.60 mmol), RuHCl(CO)(PPh₃)₃ (4 mol%), **L1** (4 mol%), toluene (0.5 mL), 80 °C, 15–18 h.

^b Isolated yields of **3**, >20:1 regioselectivity based on ¹H NMR or GC-FID analyses of the reaction mixtures.

Table 3

Hydroacylation with Various Alkynes^a

Entry	3	R or R ¹ in 3	Yield (%) ^b
1		Cy	68 (3bb)
2		^t Bu	93 (3bc)
3		Ph	95 (3ad)
4		4-MeOC ₆ H ₄	94 (3bd)
5		3-MeOC ₆ H ₄	90 (3gd)
6		2-Furyl	87 (3id)
7		3-Furyl	94 (3jd)
8		2-Thienyl	88 (3kd)
9		3- <i>N</i> -Ts-indolyl	83 (3ld)
10		4-ClC ₆ H ₄	77 (3be)
11		4-BrC ₆ H ₄	61 (3bf)
12		4-CF ₃ C ₆ H ₄	82 (3bg)
13		4-MeOC ₆ H ₄	82 (3bh)
14		3-MeC ₆ H ₄	83 (3bi)
15		2-MeC ₆ H ₄	67 (3bj)
16		Ph	0 (3ak)

^a **1** (0.20 mmol), **2a** (0.24–0.40 mmol), RuHCl(CO)(PPh₃)₃ (4 mol%), **L1** (4 mol%), toluene (0.5–1.0 mL), 80–90 °C, 15–18 h.

^b Isolated yields of **3**, >20:1 regioselectivity based on ¹H NMR or GC-FID analyses of the reaction mixtures.