

HHS Public Access

Author manuscript *J Am Chem Soc.* Author manuscript; available in PMC 2016 March 11.

Published in final edited form as: *J Am Chem Soc.* 2015 March 11; 137(9): 3161–3164. doi:10.1021/jacs.5b00747.

Enantioselective Ruthenium Catalyzed Carbonyl Allylation *via* Alkyne-Alcohol C-C Bond Forming Transfer Hydrogenation: Allene Hydrometallation *vs.* Oxidative Coupling

Tao Liang, **Khoa D. Nguyen**, **Wandi Zhang**, and **Michael J. Krische**^{*} University of Texas at Austin, Department of Chemistry, Austin, TX 78712, USA

Abstract

Chiral ruthenium(II) complexes modified by Josiphos ligands catalyze the reaction of alkynes with primary alcohols to form homoallylic alcohols with excellent control of regio-, diastereo- and enantioselectivity. These processes represent the first examples of enantioselective carbonyl allylation using alkynes as allylmetal equivalents.

Enantioselective carbonyl allylation is of broad use in chemical synthesis.¹ As exemplified in the seminal work of Hoffmann (1978),^{2a} preformed allylmetal reagents modified by chiral auxiliaries are effective in enantioselective carbonyl allylation (Scheme 1, eq. 1).² Alternatively, as first demonstrated by Yamamoto (1991), achiral allylmetal reagents can used in combination with chiral catalysts (Scheme 1, eq. 1).³ Finally, enantioselective carbonyl allylation can be achieved through the reductive coupling of allylic halides, as in Nozaki-Hiyama-Kishi type allylations,⁴ and umpoled reactions of allylic carboxylates under the conditions of metal catalysis.⁵ The latter reaction type typically requires stoichiometric metallic or metal-based reductants. Exploiting hydrogen embedded in alcohol reactants, we have developed redox-neutral enantioselective carbonyl allylations,⁶ wherein alcohol oxidation is coupled to reductive cleavage of allylic carboxylates⁷ or the hydrometallation of dienes,⁸ allenes⁹ or enynes¹⁰ to generate aldehyde-organometal pairs (Scheme 1, eq. 2).

In the course of our studies, Obora and Ishii reported the iridium catalyzed reaction of 1aryl-1-propynes with alcohols to form racemic products of carbonyl allylation.¹¹ Having encountered such products in allene-alcohol C-C couplings catalyzed by iridium^{9,12} and ruthenium,^{12b} alkyne-to-allene isomerization is likely operative in this process. The possibility of promoting such transformations using a *ruthenium* catalyst appeared tenuous, as established alkyne-alcohol C-C couplings are known to form allylic alcohols.^{13,14} Despite this precedent, we recently found that ruthenium catalysts generated upon the acid-base reaction of H₂Ru(CO)(PPh₃)₃ and 2,4,6-(2-Pr)₃PhSO₃H convert 2-alkynes and primary alcohols to (*Z*)-homoallylic alcohols through pathways involving alkyne-to-allene isomerization with subsequent allene-carbonyl oxidative coupling (Scheme 1, eq. 3).¹⁵ In

Corresponding Author. mkrische@mail.utexas.edu.

Supporting Information Available: Experimental procedures and spectral data. HPLC traces corresponding to racemic and enantiomerically enriched samples. Single crystal X-ray diffraction data for compound **3m**. This material is available free of charge *via* the internet at http://pubs.acs.org.

this communication, we disclose key alterations to the reaction conditions that suppress allene-carbonyl oxidative coupling of the transient allene to favor allene hydrometallation,¹⁶ resulting in the first examples of catalytic enantioselective carbonyl allylation using alkynes as chiral allylmetal equivalents (Scheme 1, eq. 4).¹⁷

Under established conditions for the ruthenium catalyzed coupling of 4-methyl-2-pentyne 1a with p-bromobenzyl alcohol 2a, the linear (Z)-homoallylic alcohol 4a is generated in 70% isolated yield as a single alkene stereoisomer (Table 1, entry 1). It was found that introduction of Bu_4NI (10 mol%) inhibited conversion to the (Z)-homoallylic alcohol 4a (Table 1, entry 2). Using the chelating phosphine ligand dppf in the absence of Bu_4NI , roughly equimolar quantities of the branched allylation product 3a and the (Z)-homoallylic alcohol 4a were obtained in modest yield (Table 1, entry 3). To our delight, the combination of Bu_4NI and dppf led to an 83% yield of the branched allylation product **3a** as a 15:1 (anti:syn) diastereomeric mixture (Table 1, entry 4). Using dippf as ligand, the branched allylation product **3a** was formed as a single *anti*-diastereomer in 57% yield (Table 1, entry 5). In general, P-alkyl substituted chelating ligands provided superior levels of diastereoselectivity compared to their P-phenyl congeners (Table 1, entry 4 vs. 5; entry 6 vs. 7). At this stage, diverse chiral chelating phosphine ligands were evaluated. Promising levels of asymmetric induction were observed using the Josiphos ligands SL-J009-1 and SL-J002-1 (Table 1, entries 8 and 9). Lowering the loading of 2,4,6-tri(2-propyl)phenylsulfonic acid (5 mol%),^{18,19} the branched allylation product **3a** could be obtained in excellent yield and 94% enantiomeric excess using either SL-J009-1 and SL-J002-1 as ligand (Table 1, entries 10 and 11). Omission of 2,4,6-tri(2-propyl)phenylsulfonic acid led to over-oxidation of 3a to form ketones, that is, products of allene hydroacylation.¹⁴ Omission of 2-PrOH led to a roughly 20% reduction in yield of 3a and accumulation of unreacted aldehyde, dehydro-2a.

With these optimized conditions in hand, diverse alcohols **2a–21** were surveyed for their ability to participate in this new protocol for asymmetric allylation (Table 2). Benzylic alcohols **2a–2f** were converted to adducts **3a–3f**, respectively, in good yield with complete levels of *anti*-diastereoselectivity and uniformly high levels of enantiomeric enrichment (94–96% ee). Allylic alcohols **2g–2i** could potentially engage in internal redox isomerization to form the corresponding aldehydes,²⁰ but such products are not observed. Rather, adducts **3g–3i** are formed in a highly regio- and stereoselective manner. Similarly high levels of selectivity were observed in the conversion of aliphatic alcohols **2j–2l** to adducts **3j–3l**. Under these first-generation conditions, branched primary alcohols such as isobutyl alcohol, provided adducts in the range of 30–40% yield. Under these conditions, secondary alcohols oxidize to form ketone products and do not engage in C-C coupling.

To further explore the scope of this method for catalytic enantioselective allylation, a series of substituted propynes **1b–1d** possessing *tert*-butyl (**1b**), 4-(*N*-Boc-piperidinyl) (**1c**) and phenyl (**1d**) moieties were explored in couplings to benzylic alcohol **2a** and aliphatic alcohol **2j** (Table 3). In the case of the *tert*-butyl substituted alkyne **1b**, adducts **3m** and **3p** were generated in good yield with complete levels of *anti*-diastereoselectivity and excellent levels of enantioselectivity, 94% ee and 90% ee, respectively. Similarly, for 4-(*N*-Boc-piperidinyl) substituted alkyne **1c**, adducts **3n** and **3q** were formed with good control of relative and

J Am Chem Soc. Author manuscript; available in PMC 2016 March 11.

absolute stereochemistry, despite potential cleavage of the acid-sensitive Boc-protecting group. Finally, the phenyl substituted alkyne **1d** coupled to alcohols **2a** and **2j** to provide adducts **3o** and **3r**, respectively, in good yield as single diastereomers and good to excellent control of enantioselectivity, 86% ee and 90% ee, respectively. Attempted use of 2-pentyne resulted in a complex mixture of products. The relative and absolute stereochemical assignment of adducts **3a–3r** is made in analogy adduct **3m**, which was determined by single crystal X-ray diffraction analysis.

A general catalytic mechanism was proposed (Scheme 2). Allene hydrometallation forms a nucleophilic allylruthenium complex. The stoichiometric reaction of $HXRu(CO)(PR_3)_3$ (X = Cl, Br) with allenes to form π -allylruthenium complexes is known.²¹ We have found that such π -allylruthenium species are highly fluxional in nature, undergoing rapid geometrical isomerization by way of the σ -bound haptomers.^{12b} Coordination of aldehyde to the pentacoordinate (E)- σ -allylruthenium species leads to stereospecific addition through a closed transition structure to form the homoallylic ruthenium alkoxide. Protonolytic cleavage of the alkoxide by the arylsulfonic acid releases the product and provides the indicated ruthenium sulfonate, which is displaced by a primary alcohol. Subsequent β hydride elimination generates aldehyde along with the ruthenium hydride to close the catalytic cycle. Notably, in the absence of acid, β -hydride elimination occurs at the stage of the homoallylic ruthenium alkoxide to furnish the β , γ -enone.¹⁴ Hence, substoichiometric arylsulfonic acid appears to catalyze alkoxide exchange at the metal center. Using HIRu(CO)(PR₃)₃ as precatalyst in the absence of arylsulfonic acid under otherwise standard conditions, alcohol **2a** is converted to the branched allylation product **3a** in 68% yield, >20:1 dr and 81% ee, corroborating intervention of iodide as a counterion for ruthenium. The strong σ -donicity of iodide may destabilize ruthenium(0), suppressing competitive oxidative coupling pathways. The indicated deuterium labelling experiments are consistent with this interpretation of the mechanism, and further reveal that the reaction products are inert with respect to alcohol dehydrogenation, which prevents any erosion in kinetic stereoselectivity under the reaction conditions. It should be noted that protons from the reactant alcohol or adventitious water may contribute to diminished levels of deuterium incorporation.²²

In summary, we report a ruthenium catalyzed enantioselective carbonyl allylation wherein alkynes and primary alcohols serve as redox pairs. A notable feature of these transformations is the use of alkynes as a reservoir for allenes, which are kinetically more reactive and which form *in situ* through a second, parallel ruthenium catalyzed isomerization process. Of further note are the subtle changes in reaction conditions that enable partitioning of two competing catalytic pathways, allene-carbonyl oxidative coupling *vs* allene hydrometallation, while the parallel ruthenium catalyzed alkyne isomerization pathway is maintained. More broadly, this work contributes to a growing body of catalytic processes developed in our laboratory wherein the native reducing ability of alcohols is used to generate transient organometallics from π -unsaturated precursors, enabling carbonyl addition from the alcohol oxidation level in the absence of stoichiometric organometallic reagents.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The Robert A. Welch Foundation (F-0038) and the NIH-NIGMS (RO1-GM069445) are acknowledged for partial support of this research.

References

- For selected reviews on enantioselective carbonyl allylation, see: Ramachandran PV. Aldrichim Acta. 2002; 35:33.Denmark SE, Fu J. Chem Rev. 2003; 103:2763. [PubMed: 12914480] Kennedy, Jason WJ.; Hall, DG. Angew Chem, Int Ed. 2003; 42:473.Yu CM, Youn J, Jung HK. Bull Korean Chem Soc. 2006; 27:462.Marek I, Sklute G. Chem Commun. 2007:1683.Hall DG. Synlett. 2007:1644.Lachance H, Hall DG. Org React. 2008; 73:1.Yus M, González-Gómez JC, Foubelo F. Chem Rev. 2011; 111:7774. [PubMed: 21923136]
- 2. For selected examples of enantioselective carbonyl allylation using chirally modified allyl metal reagents, see: Herold T, Hoffmann RW. Angew Chem, Int Ed. 1978; 17:768.Hoffmann RW, Herold T. Chem Ber. 1981; 114:375.Hayashi T, Konishi M, Kumada M. J Am Chem Soc. 1982; 104:4963.Brown HC, Jadhav PK. J Am Chem Soc. 1983; 105:2092.Roush WR, Walts AE, Hoong LK. J Am Chem Soc. 1985; 107:8186.Reetz MT. Pure Appl Chem. 1988; 60:1607.Short RP, Masamune S. J Am Chem Soc. 1989; 111:1892.Corey EJ, Yu CM, Kim SS. J Am Chem Soc. 1989; 111:5495.Seebach D, Beck AK, Imwinkelzied R, Roggo S, Wonnacott A. Helv Chim Acta. 1987; 70:954.Riediker M, Duthaler RO. Angew Chem, Int Ed. 1989; 28:494.Panek JS, Yang M. J Am Chem Soc. 2002; 124:7920. [PubMed: 12095334] Hackman BM, Lombardi PJ, Leighton JL. Org Lett. 2004; 6:4375. [PubMed: 15524487] Burgos CH, Canales E, Matos K, Soderquist JA. J Am Chem Soc. 2005; 127:8044. [PubMed: 15926828]
- For selected examples of enantioselective carbonyl allylation using achiral allylmetal reagents in combination with chiral catalysts, see: Furuta K, Mouri M, Yamamoto H. Synlett. 1991:561.Costa AL, Piazza MG, Tagliavini E, Trombini C, Umani-Ronchi A. J Am Chem Soc. 1993; 115:7001.Keck GE, Tarbet KH, Geraci LS. J Am Chem Soc. 1993; 115:8467.Denmark SE, Coe DM, Pratt NE, Griedel BD. J Org Chem. 1994; 59:6161.Denmark SE, Fu J. J Am Chem Soc. 2001; 123:9488. [PubMed: 11562250] Lachance H, Lu X, Gravel M, Hall DG. J Am Chem Soc. 2003; 125:10160. [PubMed: 12926924] Wada R, Oisaki K, Kanai M, Shibasaki M. J Am Chem Soc. 2004; 126:8910. [PubMed: 15264818] Rauniyar V, Hall DG. Angew Chem, Int Ed. 2006; 45:2426.Rauniyar V, Zhai H, Hall DG. J Am Chem Soc. 2008; 130:8481. [PubMed: 18540580] Lou S, Moquist PN, Schaus SE. J Am Chem Soc. 2006; 128:12660. [PubMed: 17002355] Barnett DS, Moquist PN, Schaus SE. Angew Chem, Int Ed. 2009; 48:8679.Jain P, Antilla JC. J Am Chem Soc. 2010; 132:11884. [PubMed: 20690662]
- For selected reviews on enantioselective carbonyl allylation via Nozaki-Hiyama-Kishi coupling, see: Avalos M, Babiano R, Cintas P, Jiménez JL, Palacios JC. Chem Soc Rev. 1999; 28:169.Bandini M, Cozzi PG, Umani-Ronchi A. Chem Commun. 2002:919.Hargaden GC, Guiry PJ. Adv Synth Catal. 2007; 349:2407.
- 5. For selected reviews covering carbonyl allylation via umpolung of π-allyls, see: Masuyama Y. Liebeskind LS. Advances in Metal-Organic Chemistry. JAI PressGreenwich1994; 3:255.Tamaru Y. Negishi, E-i; de Meijere, A.In Handbook of Organopalladium Chemistry for Organic Synthesis. WileyNew York2002; 2:1917.Tamaru Y. Tsuji J. In Perspectives in Organopalladium Chemistry for the XXI Century. ElsevierAmsterdam1999:215.Kondo T, Mitsudo TA. Curr Org Chem. 2002; 6:1163.Tamaru Y. Eur J Org Chem. 2005; 13:2647.Zanoni G, Pontiroli A, Marchetti A, Vidari G. Eur J Org Chem. 2007; 22:3599.
- 6. For a recent review, see: Ketcham JM, Shin I, Montgomery TP, Krische MJ. Angew Chem, Int Ed. 2014; 53:9142.

J Am Chem Soc. Author manuscript; available in PMC 2016 March 11.

- For selected examples of catalytic enantioselective redox-triggered carbonyl allylations employing allylic carboxylates as pronucleophiles, see: Kim IS, Ngai MY, Krische MJ. J Am Chem Soc. 2008; 130:6340. [PubMed: 18444616] Kim IS, Ngai MY, Krische MJ. J Am Chem Soc. 2008; 130:14891. and literature cited in reference 6. [PubMed: 18841896]
- For catalytic enantioselective redox-triggered carbonyl allylations employing dienes as pronucleophiles, see: Zbieg JR, Yamaguchi E, McInturff EL, Krische MJ. Science. 2012; 336:324. [PubMed: 22442385] McInturff EL, Yamaguchi E, Krische MJ. J Am Chem Soc. 2012; 134:20628. literature cited in reference 6. [PubMed: 23234459]
- 9. For catalytic enantioselective redox-triggered carbonyl allylations employing allenes as pronucleophiles, see: Han SB, Kim IS, Han H, Krische MJ. J Am Chem Soc. 2009; 131:6916. and literature cited in reference 6. [PubMed: 19453190]
- For catalytic enantioselective redox-triggered carbonyl propargylations employing 1,3-enynes or propargyl chlorides as pronucleophiles, see: Geary LM, Woo SK, Leung JC, Krische MJ. Angew Chem, Int Ed. 2012; 51:2972. and literature cited in reference 6.
- 11. Obora Y, Hatanaka S, Ishii Y. Org Lett. 2009; 11:3510. [PubMed: 19719194]
- For catalytic non-enantioselective redox-triggered carbonyl allylations employing allenes as pronucleophiles, see: Zbieg JR, McInturff EL, Leung JC, Krische MJ. J Am Chem Soc. 2011; 133:1141. and references cited therein. [PubMed: 21175178]
- (a) Patman RL, Chaulagain MR, Williams VM, Krische MJ. J Am Chem Soc. 2009; 131:2066.
 [PubMed: 19173651] (b) Williams VM, Leung JC, Patman RL, Krische MJ. Tetrahedron. 2009; 65:5024. [PubMed: 20613891] (c) Leung JC, Patman RL, Sam B, Krische MJ. Chem Eur J. 2011; 17:12437. [PubMed: 21953608]
- 14. For related ruthenium catalyzed alkyne hydroacylations to form α, β-unsaturated ketones or β,γketones, which occur by way of alkyne-to-allene isomerization, see: Hatanaka S, Obora Y, Ishii Y. Chem Eur J. 2010; 16:1883. [PubMed: 20029924] Chen Q-A, Cruz FA, Dong VM. J Am Chem Soc. 201510.1021/ja512015w
- Park BY, Nguyen KD, Chaulagain MR, Komanduri V, Krische MJ. J Am Chem Soc. 2014; 136:11902. [PubMed: 25075434]
- 16. For alkyne-to-allene isomerization followed by hydrometallation to form electrophilic allylpalladium or allylrhodium intermediates, see: Trost BM, Brieden Baringhaus KH. Angew Chem, Int Ed. 1992; 31:1335.Gellrich U, Meißner A, Steffani A, Kähny M, Drexler HJ, Heller D, Plattner DA, Breit B. J Am Chem Soc. 2014; 136:1097. [PubMed: 24377792]
- Allylboron reagents prepared from 1-alkynes in one-pot alkyne hydroboration-olefin isomerization participate in chiral Brønsted acid catalyzed enantioselective carbonyl allylation: Miura T, Nishida Y, Murakami M. J Am Chem Soc. 2014; 136:6223. [PubMed: 24735355] Miura T, Nishida Y, Morimoto M, Murakami M. J Am Chem Soc. 2013; 135:11497. [PubMed: 23886015]
- s2,4,6-tri(2-propyl)phenylsulfonic acid hydrate: Jautze S, Peters R. Angew Chem, Int Ed. 2008; 47:9284.
- For the reaction of H₂Ru(CO)(PPh₃)₃ with bronsted acids, see: Dobson A, Robinson SR, Uttley MF. J Chem Soc, Dalton Trans. 1974:370. Also see references 8b, 8c and 12b.
- 20. For a review, see: Larionov E, Li H, Mazet C. Chem Commun. 2014:9816.
- 21. For the stoichiometric reaction of HXRu(CO)(PR₃)₃ (X = Cl, Br) with allenes or dienes to furnish π -allylruthenium complexes, see: Hiraki K, Ochi N, Sasada Y, Hayashida H, Fuchita Y, Yamanaka S. J Chem Soc, Dalton Trans. 1985:873.Hill AF, Ho CT, Wilton-Ely IDET. Chem Commun. 1997:2207.Xue P, Bi S, Sung HHY, Williams ID, Lin Z, Jia G. Organometallics. 2004; 23:4735.
- 22. Tse SKS, Xue P, Lin Z, Jia G. Adv Synth Catal. 2010; 352:1512.

Author Manuscript

Page 5





Prior Work: Alkyne-mediated (Z)-allylation to form racemic linear adducts



This Work: Enantioselective alkyne-mediated allylation to form branched adducts





JAm Chem Soc. Author manuscript; available in PMC 2016 March 11.

Scheme 2.

Deuterium labeling studies and general catalytic mechanism.^a

^aThe extent of ²H incorporation was determined using ¹H and ²H NMR. For the deuterium labelling experiments, reactions were conducted using dippf as ligand. See Supporting Information for further experimental details, including equations accounting for the regioselectivity and extent of deuterium incorporation at position H_a -H_e.

Table 1

Selected experiments illustrating the partitioning of hydrometallative and oxidative coupling pathways in the reaction of 1a and 2a to form isomers 3a and 4a.^{*a*}

$\begin{array}{c} Me \\ Ia \\ 2-PrOH (200 \text{ mol}\%) \\ Ia \\ 2a \\ THF (0.2 \text{ M}), 95 \ ^{\circ}\text{C} \\ 3a, via \text{ allene} \\ hydrometallation \\ Ar = p\text{-BrPh} \\ \end{array} \begin{array}{c} OH \\ C \\ $				
Entry	Bu ₄ NI	ligand	3a (yield, dr)	4a (yield, Z:E)
1^b	-	-	-	70%, >20:1
2	(10 mol%)	-	-	24%, 10:1
3	-	dppf	21%, 7:1	16%, 5:1
4	(10 mol%)	dppf	83%, 15:1	-
5	(10 mol%)	dippf	57%, >20:1	-
6	(10 mol%)	dppb	36%, 13:1	-
7	(10 mol%)	dCypb	49%,>20:1	-
8	(10 mol%)	SL-J009-1	59%, >20:1, 84% ee	-
9	(10 mol%)	SL-J002-1	86%, >20:1, 80% ee	-
10^{C}	(10 mol%)	SL-J002-1	86%, >20:1, 94% ee	-
11 ^{c,d}	(10 mol%)	SL-J009-1	79%, >20:1, 94% ee	-



 a Yields are of material isolated by silica gel chromatography. See Supporting Information for further experimental details.

^b2,4,6-(2-Pr)3PhSO3H (14 mol%).

^с2,4,6-(2-Pr)3PhSO3H (5 mol%).

^dTHF (1 M).

Table 2

Regio-, *anti*-diastereo- and enantioselective C-C coupling of alkyne **1a** and alcohols **2a–2l** to form branched products of carbonyl allylation **3a–3l**.^{*a*}



 a Yields are of material isolated by silica gel chromatography. Diastereoselectivities were determined by ¹H NMR analysis of crude reaction mixtures. See Supporting Information for further experimental details.

^b2-PrOH was omitted.

^с125 °С.

^d48 h.

Author Manuscript

Table 3

Regio-, *anti*-diastereo- and enantioselective C-C coupling of alkynes 1b-1d and alcohols 2a and 2j to form branched products of carbonyl allylation 3m-3r.^{*a*}



^aYields are of material isolated by silica gel chromatography. Diastereoselectivities were determined by ¹H NMR analysis of crude reaction mixtures. See Supporting Information for further experimental details.

^b72 h.

^с75 °С.

^d₂-PrOH was omitted.

 $^e7.5$ mol% of H2Ru(CO)(PPh3)3, ArSO3H, SL-J009-1 and 15 mol% Bu4NI.

 $f_{\rm 48 \ h.}$

J Am Chem Soc. Author manuscript; available in PMC 2016 March 11.