

HHS Public Access

Author manuscript *J Am Chem Soc*. Author manuscript; available in PMC 2016 August 05.

Published in final edited form as:

J Am Chem Soc. 2015 August 5; 137(30): 9523–9526. doi:10.1021/jacs.5b04725.

Stereoconvergent Negishi Arylations of Racemic Secondary Alkyl Electrophiles: Differentiating Between a CF₃ and an Alkyl Group

Yufan Liang and Gregory C. Fu*

Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States

Abstract

In this report, we establish that a readily available nickel/bis(oxazoline) catalyst accomplishes a wide array of enantioconvergent cross-couplings of arylzinc reagents with CF_3 -substituted racemic secondary alkyl halides, a process that necessitates that the chiral catalyst be able to effectively distinguish between a CF_3 and an alkyl group in order to provide good ee. We further demonstrate that this method can be applied without modification to the catalytic asymmetric synthesis of other families of fluorinated organic compounds.

The incorporation of a trifluoromethyl (CF₃) group in an organic molecule can lead to substantially altered properties, including biological activity, as compared to a non-fluorinated compound.¹ Whereas significant advances have been described in the synthesis of molecules that contain an Ar–CF₃ bond,² progress in the development of general methods for the synthesis of compounds that include a C_{sp3} –CF₃ bond has been more limited, particularly in the case wherein the carbon that bears the CF₃ group is a tertiary stereogenic center.^{3,4} Examples of catalytic asymmetric reactions that generate the latter class of compounds include the α -trifluoromethylation of aldehydes,⁵ the conjugate addition to CF₃-containing electron-deficient alkenes;⁶ the trifluoromethylation of allylic electrophiles,⁷ and the hydrogenation of CF₃-substituted alkenes;⁸ these methods often require a suitable additional functional group (e.g., carbonyl, alkenyl, or aryl substituent) to achieve the desired bond formation and good enantioselectivity. The development of a general approach that proceeds in high ee without the need for such a functional group could thus complement existing strategies.

Recently, we have established that chiral nickel catalysts can accomplish enantioconvergent cross-coupling reactions of a number of racemic secondary electrophiles.⁹ These methods have typically employed alkyl halides that are either activated (e.g., benzylic or allylic) or that include a directing group (e.g., a carbonyl or a sulfonamide). In order to achieve the

^{*}Corresponding Author. gcfu@caltech.edu.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures and compound characterization data. This material is available free of charge via the Internet at http://pubs.acs.org.

The authors declare no competing financial interest.

Liang and Fu

Page 2

(1)

(2)

objective outlined above, we decided to pursue the development of a general method for the stereoconvergent cross-coupling of an electrophile of type **A** (eq 1);^{10,11} to furnish good enantioselectivity in such a process, the catalyst must effectively differentiate between a CF_3 and an alkyl group,^{12,13} which was not achieved in the only previous report of such a cross-coupling.¹⁴



In this report, we establish that cross-couplings of CF_3 -substituted racemic alkyl electrophiles can indeed be achieved with good enantioselectivity, without the need for an additional activating/directing group elsewhere in the molecule. Specifically, we establish that a nickel/bis(oxazoline) catalyst can effect asymmetric Negishi arylations of secondary alkyl halides (eq 2).



The data in Table 1 illustrate the impact of various reaction parameters on the efficiency of this stereoconvergent cross-coupling.¹⁵ If either of the catalyst components (NiCl₂•glyme or bis(oxazoline) ligand **1**, each of which is commercially available and can be handled in air) is omitted, little or no carbon–carbon bond formation is observed (entries 2 and 3). Other ligands, including related bis(oxazolines), a pybox, and a 1,2-diamine,¹⁶ furnish lower ee and yield (entries 4–7). The use of less of the organozinc nucleophile, a lower catalyst loading, or THF (without diglyme) as the solvent results in no erosion in enantioselectivity but in a modest loss in yield (entries 8–10). At room temperature, side reactions such as hydrodebromination and debromodefluorination become significant (entry 11). This enantioconvergent Negishi arylation is not highly water-sensitive: the addition of 0.1 equiv of water has no deleterious effect on either ee or yield (entry 12).

Next, we explored the scope of this new method for the catalytic enantioselective synthesis of stereogenic centers that bear a CF_3 substituent (Table 2).¹⁷ Although the use of an orthosubstituted arylzinc reagent leads to a modest yield of the desired cross-coupling product (entry 1), an array of meta- and para-substituted nucleophiles, either electron-rich or

JAm Chem Soc. Author manuscript; available in PMC 2016 August 05.

Liang and Fu

electron-poor, generally furnish very good ee's and yields (entries 2–11). Functional groups such as an aryl ether,¹⁸ an aryl halide (Br, Cl, and F),¹⁹ an aryl thioether,²⁰ and oxygen and nitrogen heterocycles are compatible with the coupling conditions. On a gram scale (1.34 g of product), the stereoconvergent Negishi reaction illustrated in entry 2 of Table 2 proceeds in 96% ee and 91% yield; it is noteworthy that this cross-coupling is not especially air-sensitive–when conducted in a capped vial under air, the product is generated with similar efficiency (96% ee, 87% yield).

With regard to the electrophilic coupling partner, the scope of this method for enantioselective Negishi cross-couplings of fluorinated secondary alkyl halides is also fairly broad (Table 3).²¹ It is worth noting that, in the case of entry 1, the catalyst must effectively differentiate between a CF₃ and a CH₃ group in order to deliver the product in very good ee.¹² A variety of functional groups are compatible with the reaction conditions, including a silyl ether, a primary alkyl chloride,²² a primary alkyl bromide,²³ a primary alkyl tosylate,²⁴ an aryl ether,¹⁸ a ketone, an aryl iodide,¹⁹ a carbamate, an ester, and a furan.

The choice of electrophile is not limited to alkyl bromides. Thus, without modification we can apply to a racemic secondary alkyl iodide the method that we have developed for stereoconvergent cross-couplings of bromides (eq 3).²⁵



Significantly, the method can also be applied to electrophiles that bear an electronwithdrawing group other than CF₃. Thus, excellent enantioselectivity is observed with an array of fluorinated substituents, including perfluoroalkyl, CF₂Cl, CF₂COPh, and CF₂CO₂Et (eq 4–7).²⁶



(3)

Liang and Fu



(7)

In conclusion, we have developed a versatile stereoconvergent Negishi arylation of racemic secondary alkyl halides wherein a chiral nickel catalyst differentiates between a CF_3 and an alkyl substituent in order to deliver high enantiomeric excess. In contrast to many other catalytic asymmetric processes that generate tertiary stereocenters that bear a CF_3 substituent, this method does not require an extraneous activating/directing group; furthermore, this approach is effective at producing a variety of other enantiomerically enriched fluorinated compounds under the same conditions. From a practical perspective, it is noteworthy that both of the catalyst components are commercially available and can be handled in air, and that the method is not particularly air- or water-sensitive. Further studies of nickel-catalyzed cross-couplings of alkyl electrophiles are underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

Support has been provided by the National Institutes of Health (National Institute of General Medical Sciences: R01–GM62871) and the Gordon and Betty Moore Foundation (Caltech Center for Catalysis and Chemical Synthesis). We thank Dr. Allen G. Oliver (University of Notre Dame) and Dr. Nathan D. Schley for assistance.

J Am Chem Soc. Author manuscript; available in PMC 2016 August 05.

REFERENCES

- For leading references, see: Gouverneur V, Müller K. Fluorine in Pharmaceutical and Medicinal Chemistry: From Biophysical Aspects to Clinical Applications. 2012LondonImperial College Press Zhu W, Wang J, Wang S, Gu Z, Aceña JL, Izawa K, Liu H, Soloshonok VA. J. Fluorine Chem. 2014; 167:37–54.
- For discussions and leading references, see: Cho EJ, Senecal TD, Kinzel T, Zhang Y, Watson DA, Buchwald SL. Science. 2010; 328:1679–1681. [PubMed: 20576888] Alonso C, Martínez de Marigorta E, Rubiales G, Palacios F. Chem. Rev. 2015; 115:1847–1935. [PubMed: 25635524]
- For discussions and leading references, see: Yang X, Wu T, Phipps RJ, Toste FD. Chem. Rev. 2015; 115:826–870. [PubMed: 25337896] (b) References 1b and 2b. Liang T, Neumann CN, Ritter T. Angew. Chem. Int. Ed. 2013; 52:8214–8264. Xu J, Liu X, Fu Y. Tetrahedron Lett. 2014; 55:585– 594.
- For a few examples of target compounds, see: (a) Proline derivative: Del Valle JR, Goodman M. Angew. Chem. Int. Ed. 2002; 41:1600–1602. Qiu, X-l; Qing, F-l. J. Org. Chem. 2002; 67:7162– 7164. [PubMed: 12354018] (b) Nucleoside analogue: Jeannot F, Gosselin G, Standring D, Bryant M, Sommadossi J-P, Loi AG, La Colla P, Mathé C. Bioorg. Med. Chem. 2002; 10:3153–3161. [PubMed: 12150860] (c) Estradiol derivative: Blazejewski J-C, Wilmshurst MP, Popkin MD, Wakselman C, Laurent G, Nonclercq D, Cleeren A, Ma Y, Seo H-S, Leclercq G. Bioorg. Med. Chem. 2003; 11:335–345. [PubMed: 12517429]
- For example, see: Nagib DA, Scott ME, MacMillan DWC. J. Am. Chem. Soc. 2009; 131:10875– 10877. [PubMed: 19722670] Allen AE, MacMillan DWC. J. Am. Chem. Soc. 2010; 132:4986– 4987. [PubMed: 20297822]
- For a recent example, see: Massolo E, Benaglia M, Orlandi M, Rossi S, Celentano G. Chem. Eur. J. 2015; 21:3589–3595. [PubMed: 25573847]
- 7. For a recent example, see: Nishimine T, Fukushi K, Shibata N, Taira H, Tokunaga E, Yamano A, Shiro M, Shibata N. Angew. Chem. Int. Ed. 2014; 53:517–520.
- 8. For example, see: Dong K, Li Y, Wang Z, Ding K. Angew. Chem. Int. Ed. 2013; 52:14191-14195.
- 9. (a) For the initial report, see: Fischer C, Fu GC. J. Am. Chem. Soc. 2005; 127:4594–4595.
 [PubMed: 15796523] (b) For the most recent study, see: Choi J, Martín-Gago P, Fu GC. J. Am. Chem. Soc. 2014; 136:12161–12165. [PubMed: 25127186]
- 10. For examples of metal-catalyzed cross-coupling reactions of CF₃CH₂OTs and CF₃CH₂I (*primary* alkyl electrophiles), see: Zhao Y, Hu J. Angew. Chem. Int. Ed. 2012; 51:1033–1036. Liang A, Li X, Liu D, Li J, Zou D, Wu Y, Wu Y. Chem. Commun. 2012; 48:8273–8275. Feng Y-S, Xie C-Q, Qiao W-L, Xu H-J. Org. Lett. 2013; 15:936–939. [PubMed: 23373673] Leng F, Wang Y, Li H, Li J, Zou D, Wu Y, Wu Y. Chem. Commun. 2013; 49:10697–10699.
- 11. To the best of our knowledge, there are no examples of metal-catalyzed cross-couplings of perfluorinated alkyl nucleophiles with unactivated alkyl electrophiles, which could represent a complementary route to the products illustrated in eq 1.
- 12. For an early example of an enantioselective reaction wherein differentiation between a CF₃ and a CH₃ group is achieved, see: Feigl DM, Mosher HS. Chem. Commun. 1965:615–616.
- For a recent review on the influence of fluorine in asymmetric catalysis, see: Cahard D, Bizet V. Chem. Soc. Rev. 2014; 43:135–147. [PubMed: 24162874]
- 14. Liang Y, Fu GC. Angew. Chem. Int. Ed. accepted: DOI: 10.1002/anie.201503297 (alkylation reactions). This method furnishes very modest enantioselectivity (endnotes 17 and 20).
- 15. Notes: (a) The method described in Reference 14 is not effective for the arylation process illustrated in Table 1, furnishing poor ee and yield. (b) PhZnCl is generated from PhLi. If PhZnCl is produced instead from PhMgCl, then the arylation proceeds in 95% ee and 81% yield. Ph2Zn is not effective under these conditions, furnishing 88% ee and 9% yield.
- These families of ligands have previously been employed in stereoconvergent nickel-catalyzed cross-couplings of alkyl electrophiles: (a) Bis(oxazoline): Lou S, Fu GC. J. Am. Chem. Soc. 2010; 132:1264–1266. [PubMed: 20050651] (b) Pybox: Ref 9a. (c) 1,2-Diamine: Saito B, Fu GC. J. Am. Chem. Soc. 2008; 130:6694–6695. [PubMed: 18447357]

JAm Chem Soc. Author manuscript; available in PMC 2016 August 05.

- 17. Notes: (a) During the course of these couplings, we have not observed kinetic resolution of the electrophile (<5% ee), consumption of the cross-coupling product, or erosion in the ee of the product. (b) In a preliminary study under our standard conditions, a thienylzinc, a pyridylzinc, and an alkenylzinc reagent were not suitable cross-coupling partners. (c) Hydrodebromination of the electrophile is the predominant side reaction.
- For examples of the reactivity of aryl ethers in nickel-catalyzed cross-couplings, see: Rosen BM, Quasdorf KW, Wilson DA, Zhang N, Resmerita A-M, Garg NK, Percec V. Chem. Rev. 2011; 111:1346–1416. [PubMed: 21133429]
- For examples of the reactivity of aryl halides (I, Br, Cl, and F) in nickel-catalyzed cross-couplings, see: Han F-S. Chem. Soc. Rev. 2013; 42:5270–5298. [PubMed: 23460083]
- For early examples of nickel-catalyzed cross-couplings of an aryl thioether, see: Wenkert E, Ferreria TW, Michelotti EL. J. Chem. Soc. Chem. Commun. 1979:637–638. Okamura H, Miura M, Takei H. Tetrahedron Lett. 1979; 20:43–46. Takei H, Miura M, Sugimura H, Okamura H. Chem. Lett. 1979:1447–1450.
- 21. In a preliminary study under the standard conditions, an attempt to cross-couple an electrophile wherein alkyl=cyclohexyl was not successful.
- For an early example of a nickel-catalyzed cross-coupling of a primary alkyl chloride, see: Terao J, Watanabe H, Ikumi A, Kuniyasu H, Kambe N. J. Am. Chem. Soc. 2002; 124:4222–4223. [PubMed: 11960446]
- 23. (a) For an early example of a nickel-catalyzed cross-coupling of a primary alkyl bromide, see: Devasagayaraj A, Stüdemann T, Knochel P. Angew. Chem. Int. Ed. 1995; 34:2723–2725. (b) The lower reactivity of the primary alkyl bromide may be due to activation of the secondary alkyl bromide by the CF3 substituent (Ref 14).
- 24. For an early example of a nickel-catalyzed cross-coupling of a primary alkyl tosylate, see Reference 22.
- 25. Under the same conditions, the corresponding alkyl tosylate, alkyl chloride, and alkyl triflate are not useful coupling partners (<5% yield).
- 26. For recent references to the synthesis/utility of chlorodifluoromethyl- and difluorocarbonylcontaining compounds, see: Salomon P, Zard SZ. Org. Lett. 2014; 16:2926–2929. [PubMed: 24828527] Han C, Salyer AE, Kim EH, Jiang X, Jarrard RE, Powers MS, Kirchhoff AM, Salvador TK, Chester JA, Hockerman GH, Colby DA. J. Med. Chem. 2013; 56:2456–2465. [PubMed: 23428109] Ge S, Arlow SI, Mormino MG, Hartwig JF. J. Am. Chem. Soc. 2014; 136:14401– 14404. [PubMed: 25254966]

Table 1

Stereoconvergent Negishi Arylations of Fluorinated Electrophiles: Effect of Reaction Parameters^a

F₃C	Br 6.0% Ni Ph Ph-ZnCl 7.8% racemic 1.5 equiv "standar	$\begin{array}{c} Cl_2 \cdot glyme \\ (S,S)-1 \\ \text{diglyme} \\ 20 \ ^{\circ}C \\ d'' \ \text{conditions} \end{array}$	Ph Ph Ph
entry	variation from the "standard" c	onditions ee (%)	yield (%) ^b
1	none	95	89
2	no NiCl ₂ •glyme	-	<2
3	no (<i>S</i> , <i>S</i>)- 1	-	3
4	(<i>S</i> , <i>S</i>)- 2 , instead of (<i>S</i> , <i>S</i>)- 1	33	26
5	(<i>S</i> , <i>S</i>)- 3 , instead of (<i>S</i> , <i>S</i>)- 1	40	27
6	(<i>R</i> , <i>R</i>)-4, instead of (<i>R</i> , <i>R</i>)-1	-	<2
7	(<i>S</i> , <i>S</i>)- 5 , instead of (<i>S</i> , <i>S</i>)- 1	11	4
8	1.1 equiv Ph–ZnCl	95	78
9	3.0% NiCl ₂ •glyme, 3.9% (<i>S</i> , <i>S</i>)-1	94	72
10	THF only	95	82
11	r.t., instead of -20 °C	86	15
12	0.1 equiv H_2O added	95	88



^aAll data are the average of two experiments.

 b The yields were determined through analysis by 19 F NMR spectroscopy (with the aid of an internal standard).

Table 2

Stereoconvergent Negishi Arylations of Fluorinated Electrophiles: Scope with Respect to the Nucleophile^a

Br	<u></u>	6.0% Ni 7.8%	Cl ₂ •glyme (<i>S,S</i>)– 1	Ar
F ₃ C ∽ racemic	Ph Ar-ZnCl c 1.5 equiv	THF/c	liglyme	F ₃ C ∽ Ph
	-	-20 01	-15 C	
entry	Ar		ee (%)	yield (%) ^b
1		_	87	50
	OMe			
2	х	= OMe	96	90
3	<u> </u>	Me	95	90
4) _ `	CF ₃ F	97	87
5	^	CI	96	79
6			96	86
7		X = H	95	86
8		Me	95	89
9	x-{}-§-	SMe F	94	52
10		Br	96	84
11			95	87
12	o-{``		94	60
13	MeN	}_ફ-	94	68

^aAll data are the average of two experiments.

^bYield of purified product.

J Am Chem Soc. Author manuscript; available in PMC 2016 August 05.

Table 3

Stereoconvergent Negishi Arylations of Fluorinated Electrophiles: Scope with Respect to the Electrophile^a

Br F ₃ C	MeO alkyl ZnCl - ic 1.5 equiv	6.0% NiCl₂•glyme 7.8% (<i>S</i> , <i>S</i>)−1 THF/diglyme –20 °C	F ₃ C alkyl
entry	alkyl	ee (%)	yield (%) ^b
1	Me	91	81
2	<i>n</i> -Oct	95	88
3		96	81
4	OTE	BDPS 94	77
5	, sh	96	89
6	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	∕_ Br 96	80
7	² 2 ² 2 ²	DTs 97	91
8	, pd O	96 Me	89
9) j ² j ²	95	60
10	N Bo	oc 97	82
11	»ť∽∽∽°∦ O	96	81

^aAll data are the average of two experiments.

^bYield of purified product.