

ORIGINAL ARTICLE

Nitrification expanded: discovery, physiology and genomics of a nitrite-oxidizing bacterium from the phylum *Chloroflexi*

Dimitry Y Sorokin^{1,2}, Sebastian Lückner³, Dana Vejmelkova^{2,4}, Nadezhda A Kostrikina¹, Robbert Kleerebezem², W Irene C Rijpstra⁵, Jaap S Sinninghe Damsté⁵, Denis Le Paslier^{6,7,8}, Gerard Muyzer^{2,9}, Michael Wagner³, Mark CM van Loosdrecht² and Holger Daims³

¹Winogradsky Institute of Microbiology, Russian Academy of Sciences, Moscow, Russia; ²Department of Biotechnology, Delft University of Technology, Delft, The Netherlands; ³Department of Microbial Ecology, Ecology Centre, University of Vienna, Vienna, Austria; ⁴Department of Water Technology and Environmental Engineering, ICT Prague, Prague, Czech Republic; ⁵Department of Marine Organic Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, Den Burg, The Netherlands; ⁶Commissariat à l'Energie Atomique, Genoscope, Evry, France; ⁷Centre National de la recherche Scientifique, Evry, France and ⁸Université d'Evry Val d'Essonne, Evry, France

Nitrite-oxidizing bacteria (NOB) catalyze the second step of nitrification, a major process of the biogeochemical nitrogen cycle, but the recognized diversity of this guild is surprisingly low and only two bacterial phyla contain known NOB. Here, we report on the discovery of a chemolithoautotrophic nitrite oxidizer that belongs to the widespread phylum *Chloroflexi* not previously known to contain any nitrifying organism. This organism, named *Nitrolancetus hollandicus*, was isolated from a nitrifying reactor. Its tolerance to a broad temperature range (25–63 °C) and low affinity for nitrite ($K_s = 1 \text{ mM}$), a complex layered cell envelope that stains Gram positive, and uncommon membrane lipids composed of 1,2-diols distinguish *N. hollandicus* from all other known nitrite oxidizers. *N. hollandicus* grows on nitrite and CO₂, and is able to use formate as a source of energy and carbon. Genome sequencing and analysis of *N. hollandicus* revealed the presence of all genes required for CO₂ fixation by the Calvin cycle and a nitrite oxidoreductase (NXR) similar to the NXR forms of the proteobacterial nitrite oxidizers, *Nitrobacter* and *Nitrococcus*. Comparative genomic analysis of the *nxr* loci unexpectedly indicated functionally important lateral gene transfer events between *Nitrolancetus* and other NOB carrying a cytoplasmic NXR, suggesting that horizontal transfer of the NXR module was a major driver for the spread of the capability to gain energy from nitrite oxidation during bacterial evolution. The surprising discovery of *N. hollandicus* significantly extends the known diversity of nitrifying organisms and likely will have implications for future research on nitrification in natural and engineered ecosystems.

The ISME Journal (2012) 6, 2245–2256; doi:10.1038/ismej.2012.70; published online 5 July 2012

Subject Category: microbial engineering

Keywords: *Chloroflexi*; evolution; nitrification; nitrite-oxidizing bacteria; *Nitrolancetus hollandicus*; wastewater treatment

Introduction

Chemolithoautotrophic nitrite-oxidizing bacteria (NOB) gain energy from the oxidation of nitrite to nitrate, which is the second step of nitrification and

thus represents a major biogeochemical process in aquatic and terrestrial ecosystems. Under oxic conditions, nitrite oxidation by NOB is the principal source of nitrate, which constitutes about 88% of the fixed nitrogen in the oceans (Gruber, 2004) and is the predominant inorganic nitrogen form in aerated soils (Dechorgnat *et al.*, 2011). The only other microbial process known to contribute to the global nitrate pool is anaerobic ammonium oxidation ('anammox') that forms nitrate via anaerobic nitrite oxidation (Strous *et al.*, 2006). Nitrate is a major nitrogen source for many organisms and an important electron acceptor for anaerobic respiration. In addition to their crucial function in natural

Correspondence: H Daims, Department of Microbial Ecology, Ecology Centre, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria.

E-mail: daims@microbial-ecology.net

⁹Present address: Department of Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 1090 GE Amsterdam, The Netherlands.

Received 12 March 2012; revised 10 May 2012; accepted 10 May 2012; published online 5 July 2012

ecosystems, NOB are important to mankind as they contribute to nitrogen loss in fertilized soils, but are essential for N-removal from sewage in most biological wastewater treatment plants. Winogradsky (1892) isolated the first NOB from the genus *Nitrobacter* in 1892. Since then the number of newly isolated NOB has been very limited owing to difficulties with growing NOB under laboratory conditions. Consequently, our understanding of the physiological and genomic properties of this important microbial group is restricted to a few representatives. For example, the Genomes Online Database (Kyrpides, 1999) lists only eight genome projects for NOB as opposed to 727 just for *Escherichia coli* (status of February 2012). All currently known NOB belong either to the *Proteobacteria* (Teske *et al.*, 1994) or to the genus *Nitrospira* of the phylum *Nitrospirae* (Ehrich *et al.*, 1995); the phylogenetic affiliation of the marine NOB genus *Nitrospina* with the *Deltaproteobacteria* or a distinct phylum remains uncertain (Teske *et al.*, 1994; Schloss and Handelsman, 2004). Occasionally, new NOB enrichments or isolates are reported (for example, Lebedeva *et al.*, 2011), but these strains usually belong to one of the already established groups. The only exceptions in the past 25 years were the description of *Nitrotoga* (Alawi *et al.*, 2007) as a new nitrite-oxidizing genus in the *Betaproteobacteria* and the discovery of anoxygenic phototrophic NOB from the *Alpha*- and *Gammaproteobacteria* (Griffin *et al.*, 2007; Schott *et al.*, 2010). Thus, the diversity of NOB still appears to be restricted to a few phylogenetic groups, most of which are *Proteobacteria*. In contrast to other N-cycle processes, such as denitrification (Shapleigh, 2006) or N_2 fixation (Zehr *et al.*, 2003), there has been no sign that nitrite oxidation became established in a larger number of microbial phyla in the course of evolution. This was also thought about the first step of nitrification, the oxidation of ammonia to nitrite, as all known ammonia-oxidizing bacteria (AOB) were either *Beta*- or *Gammaproteobacteria* (Purkhold *et al.*, 2000). This picture changed drastically with the discovery of ammonia-oxidizing archaea (Könneke *et al.*, 2005) and their dominance in marine (Wuchter *et al.*, 2006) and terrestrial (Leininger *et al.*, 2006) ecosystems. In analogy, several nitrite oxidizers may have not been identified yet. Recognizing the natural diversity of NOB is essential for assessing the consequences of the increased human discharge of nitrogen, in the form of fertilizers and sewage (Canfield *et al.*, 2010), on N-cycling in natural and engineered ecosystems. A better understanding of NOB is also important for reducing the release of the greenhouse gas N_2O from sewage treatment plants. Nitrite is considered the main factor for N_2O formation (Kampschreur *et al.*, 2008), and the activity of NOB decreases the nitrite concentrations and consequently the N_2O emissions. Moreover, knowing the diversity and genomic makeup of NOB is crucial for reconstructing the evolution of nitrite

oxidation as part of a major biogeochemical cycle on Earth.

Here we report on the discovery and physiological, as well as genomic, characterization of a thermotolerant nitrite-oxidizing bacterium from a bioreactor that showed high nitrite-oxidizing activity, but did not contain functionally significant amounts of known NOB. This nitrite oxidizer, belonging to a deep-branching lineage of the *Chloroflexi*, is the only identified nitrifier from this bacterial phylum and represents the only known non-proteobacterial nitrite oxidizer besides the genus *Nitrospira*. Owing to its distinct phylogenetic affiliation and physiological traits, this bacterium was overlooked in past research on nitrification.

Materials and methods

Cultivation conditions

Nitrolancetus hollandicus was cultured in mineral medium that contained 1.1 mM KH_2PO_4 and 0.3 mM $CaCl_2$ dissolved in distilled water. This solution was dispensed in serum bottles (ratio 1:5 of liquid to air headspace), which were closed with rubber stoppers. Following injection of CO_2 into the gas phase to a final concentration of 10% (v/v), the bottles were sterilized at 120 °C for 20 min. After sterilization, the medium was supplemented with 1 ml l⁻¹ of trace metal solution (Pfennig and Lippert, 1966), 1 mM $MgSO_4$, 5 mM NH_4HCO_3 and 20–50 mM KNO_2 . The medium for mixotrophic cultures was also amended with 40 mM $NaHCOO$. The pH was maintained at 6.9–7.4 by adding either NH_4HCO_3 or CO_2 . The bottles were incubated on a rotary shaker under agitation (100–120 r.p.m) at 37 °C. The solid medium was prepared by 1:1 mixing of double-strength liquid medium and 4% washed Noble agar at 50 °C.

Microscopy and FISH

Phase contrast micrographs of *N. hollandicus* cultures were recorded with a Zeiss Axioplan 2 imaging microscope. Details on electron microscopy are provided in Supplementary Text S1. Fluorescence *in situ* hybridization (FISH) with rRNA-targeted probes was carried out according to the protocol described elsewhere (Daims *et al.*, 2005) using the probes listed in Supplementary Table S2. New 16S rRNA-targeted oligonucleotide probes specific for *N. hollandicus* were designed using the software ARB (Ludwig *et al.*, 2004), and the hybridization conditions were optimized by using the software DAIME (Daims *et al.*, 2006) as described elsewhere (Daims *et al.*, 2005). Confocal FISH images were recorded using a Zeiss LSM 510 Meta confocal laser scanning microscope (Zeiss, Jena, Germany). The biovolume fraction of *N. hollandicus* in activated sludge was measured by digital analysis of confocal FISH

images by using the program DAIME as described previously (Daims and Wagner, 2007).

Chemical analyses

Nitrite was determined colorimetrically by the diazotation reaction-based assay (Griess-Romijn van Eck, 1966) and nitrate by the salicylate method (Bhandari and Simlat, 1986). Formate was measured by HPLC (high performance liquid chromatography) anionic chromatography using an Aminex HPX-87H column (300 × 7.8 mm, $t = 60^\circ\text{C}$, flow rate 0.6 ml min^{-1} , eluent $1.5\text{ mM H}_3\text{PO}_4$). Cell protein concentrations were analyzed by the Lowry method (Lowry *et al.*, 1951) after cell lysis in 1 M NaOH . For details of other physiological and chemical analyses, please refer to Supplementary Text S1.

Genomics and phylogeny

High-molecular-weight genomic DNA was extracted from a *N. hollandicus* culture by the Hexadecyltrimethylammonium bromide (CTAB) method as recommended by the DOE Joint Genome Institute (JGI, <http://my.jgi.doe.gov/general/index.html>), with the only modification that the phenol:chloroform:isoamyl alcohol extraction was performed before the chloroform:isoamyl alcohol purification step in order to avoid carryover of residual phenol. Subsequently, genomic sequences were obtained using GS FLX Titanium (Roche, Basel, Switzerland) and Illumina (Illumina, San Diego, CA, USA) sequencing technologies. One 1/4 picotiter plate Titanium run with a 3-kb paired-end library yielded 199 240 sequence reads that were assembled using the software Newbler (Roche). The 25 258 338 sequence reads (36 bp) obtained from a 1/8 Illumina sequencing run were used to correct Titanium sequencing errors. Contigs were arranged in scaffolds that were ordered by size. The MaGe software system (Vallenet *et al.*, 2006) was used for the automatic prediction and annotation of coding sequences (CDS). The annotation of all CDS in key pathways, including those for nitrite oxidation, respiration and carbon fixation, was manually refined by using the respective tools of MaGe as described in detail elsewhere (Lücker *et al.*, 2010). For details on *nxrA* gene sequencing and phylogenetic analyses, please refer to Supplementary Text S1.

Sequences from *N. hollandicus* have been deposited in the GenBank database (16S rRNA gene: accession no. JQ345500; *nxrA* genes: accession no. JQ279817 to JQ279820; genome sequence: accession no. CAGS01000001 to CAGS01000736).

Results

Isolation and physiological characterization

A laboratory-scale bioreactor was operated at 35°C to achieve partial nitrification of digester effluent containing up to 428 mM of ammonia (Vejmelkova

et al., 2011). The objective was to convert NH_4HCO_3 to NH_4NO_3 and CO_2 during a study on nitrogen recovery from sewage (Vejmelkova *et al.*, 2011). As desired, the reactor showed high ammonia- and also nitrite-oxidizing activities. Fluorescence *in situ* hybridization revealed that this activated sludge was dominated by cell aggregates of AOB, which were related to *Nitrosomonas europaea* and *Nitrosomonas eutropha* (Vejmelkova *et al.*, 2011), but with the exception of a small number of *Nitrobacter* cells no known NOB were detected. Interestingly, however, microscopic observation of the sludge revealed a high density of large, conspicuous cells whose morphology resembled the tip of a lancet. Serial dilution of sludge aliquots and incubation at 30°C in nitrite-containing liquid mineral medium buffered with $\text{NaHCO}_3/\text{CO}_2$ yielded enrichments of *Nitrobacter*. However, a temperature increase to 37°C and replacement of the buffer by $\text{NH}_4\text{HCO}_3/\text{CO}_2$ resulted in a nitrite-oxidizing culture dominated by the lancet-shaped cells (Figure 1). As the AOB were residing in flocs, they could be removed by short centrifugation steps during the enrichment procedure despite the addition of NH_4HCO_3 . The lancet-shaped organism was highly enriched after serial dilutions of this culture, and finally a pure culture was obtained (Supplementary Text S1). The isolated bacterium was designated '*Nitrolance-tus hollandicus* strain Lb'.

N. hollandicus aerobically grew as a chemolithoautotrophic nitrite oxidizer in culture media containing NO_2^- as the sole energy source and $\text{HCO}_3^-/\text{CO}_2$ as the sole carbon source (Figure 2, Supplementary Table S1). However, in contrast to all other known NOB, *N. hollandicus* was not able to assimilate nitrogen from nitrite and required ammonium for growth. Several organic substrates (acetate, pyruvate, lactate, propionate, butyrate, ethanol, glycerol, glucose, fructose and yeast extract) added to the medium either had no detectable effect or inhibited nitrite oxidation and growth. However, when newly incubated in media containing both nitrite and formate, *N. hollandicus* first consumed most of the nitrite and then started to utilize formate (Supplementary Figure S1a). The sequential use of nitrite and formate resembled diauxy, but the ability to use formate depended on preincubation of the culture with nitrite. This might be explained by the nitrite-dependent expression of genes also needed for the respiration of formate. The use of formate is not uncommon among NOB, as *Nitrobacter* is known to grow on formate (Van Gool and Laudelout, 1966) and the genome of '*Ca. Nitrospira defluvii*' encodes a formate dehydrogenase and a formate transporter (Lücker *et al.*, 2010). Nevertheless, the biochemical details of formate use may differ substantially between these species (Supplementary Figure S2 and Supplementary Text S1). No growth of *N. hollandicus* was observed with other inorganic electron donors such as H_2 , thiosulfate and CO at various concentrations and with




Figure 1 Morphology and *in situ* detection of *N. hollandicus*. (a) Phase contrast micrograph of an autotrophically grown pure culture. (b) Transmission electron micrograph of cells from the same culture as in panel a. (c) *In situ* detection of *N. hollandicus* and AOB in activated sludge from the bioreactor that was the source of the isolated *N. hollandicus* strain. The populations were detected by FISH using the probes Ntlc804 (specific for *N. hollandicus*), Cluster6a192 (specific for AOB of the *N. oligotropha* lineage) and the EUB338 probe mix targeting most *Bacteria*. Cells of *N. hollandicus* appear magenta, AOB green-blue and other bacteria dark blue. (d) Thin section electron micrograph of a cell from a culture that was grown mixotrophically in medium containing NO_2^- , $\text{HCO}_3^-/\text{CO}_2$ and formate. The layered cell envelope is clearly visible. N, nucleoid; CW-EL, external layer of the cell wall; CW-PG, peptidoglycane; CM, cytoplasmic membrane.




Figure 2 Growth of *N. hollandicus* in mineral medium containing NO_2^- as the sole energy source. NO_2^- was stoichiometrically oxidized to NO_3^- while biomass protein increased. Results of two replicate experiments are shown. Nitrate was measured only in one experiment.

different O_2 levels (2–20%) in the gas phase. Additional tests showed that CO was also not consumed by nongrowing cultures, but 5% CO in the gas phase did not inhibit growth on nitrite. Under anoxic conditions, small amounts of nitrate were reduced to nitrite in the presence of formate as electron donor, but the culture did not grow.

The optimal NO_2^- concentration range for *N. hollandicus* was 5–20 mM, but the organism displayed a high tolerance to NO_2^- (up to 75 mM) and ammonium (>200 mM) at pH 7–7.5. *N. hollandicus* is a thermotolerant nitrite oxidizer with a growth temperature optimum and maximum at 40 and 46 °C, respectively. Nitrite-oxidizing activity was observed even up to 63 °C with an optimal range from 35 to 55 °C (Supplementary Figure S1b). The organism grew at pH values from 6.2 to 8.3, with the optimal range being pH 6.8–7.5 (Supplementary Figure S1c).

Phylogeny and *in situ* detection

Phylogenetic analyses based on 16S rRNA gene sequences revealed that *N. hollandicus* is affiliated with the thermophilic heterotrophs *Sphaerobacter*

thermophilus and *Thermomicrobium roseum* (Figure 3) in the deep-branching class *Thermomicrobia* of the bacterial phylum *Chloroflexi* (Hugenholtz and Stackebrandt, 2004). To confirm that the isolated organism was identical to the lancet-shaped cells in the source bioreactor, a specific 16S rRNA-targeted oligonucleotide probe (Supplementary Table S2) was designed based on the 16S rRNA gene sequence obtained from the culture. In FISH experiments, this probe hybridized to the *N. hollandicus* culture and to the lancet-shaped bacteria in the reactor biomass (Figure 1c). The same cells were simultaneously detected by probes targeting the phylum *Chloroflexi*, confirming their phylogenetic affiliation with this line of descent. FISH also demonstrated a tight spatial clustering of *N. hollandicus* with AOB in the sludge flocs (Figure 1c) similar to the colocalization frequently reported to occur between AOB and other NOB (Schramm et al., 1999; Maixner et al., 2006), which reflects the mutualistic symbiosis between the two guilds of nitrifiers (Stein and Arp, 1998). *N. hollandicus* was a dominant population representing 34% of the total bacterial biovolume in the bioreactor. Thus, this highly abundant organism most likely caused the strong nitrite-oxidizing activity of the biomass (Vejmelkova et al., 2011) rather than the low numbers of *Nitrobacter* cells, which were sporadically detected by FISH in the same samples.

Cell morphology, ultrastructure and chemical composition

N. hollandicus cells are relatively large (1 to 1.2×2 to $4 \mu\text{m}$), non-motile, rod- (lancet-) shaped and occur

as single cells, pairs or short chains with individual cells attached at their poles (Figure 1). The cells stained Gram-positive and their envelope comprised a thin electron-dense peptidoglycan layer and several outer layers that may consist of carbohydrates or protein (Figure 1d). Complex cell envelope structures with multiple layers are not uncommon among *Chloroflexi*, although they lack an outer membrane as found in Gram-negative bacteria (Sutcliffe, 2011). Meso-diaminopimelic acid, as characteristic for the cell wall of Gram-negative bacteria, was not detected by chemical analysis of peptidoglycan from *N. hollandicus*. Instead, the diamino acids ornithine and lysine were present, suggesting that the peptidoglycan composition of *N. hollandicus* resembles that of Gram-positive bacteria. The predominant membrane lipids were uncommon and were composed of C_{20} alkyl 1,2-diols with a polar head group (sugar or phosphorus-sugar moiety) attached to the first OH group, whereas the second OH group was esterified with a fatty acid (predominantly 12 methyl-octadecanoic acid) (Supplementary Figure S3). Similar glycolipids have only been identified in thermophilic *Chloroflexi* including *Thermomicrobium*, a close relative of *Nitrolancetus* (Figure 3) and *Roseiflexus castenholzii*, as well as in *Thermus* (Pond et al., 1986; Wait et al., 1997; van ver Meer et al., 2002). Alkyl 1,2-diol lipids also occur in hot spring microbial mats (van ver Meer et al., 2002), but a specific role for the adaptation to high temperatures is debatable (Wait et al., 1997). Thin sections revealed the absence of intracytoplasmic membranes (ICM) that are found in *Nitrobacter* and *Nitrococcus* (Figure 1d).




Figure 3 Phylogenetic affiliation of *N. hollandicus*. 16S rRNA-based Bayesian inference tree (s.d. = 0.009202) showing the affiliation of *N. hollandicus* (boldface) with the class *Thermomicrobia* (gray box) of the phylum *Chloroflexi*. Pie charts indicate statistical support of nodes based on bootstrap analysis or Bayesian inference, respectively. Conflicting topologies among the different methods are also indicated. Representatives of the phylum *Actinobacteria* were used as outgroup. The scale bar represents 10% estimated sequence divergence. MB, Bayesian inference; ML, maximum likelihood; MP, maximum parsimony; NJ, neighbor joining.

Genomic analysis

The genome of *N. hollandicus* strain Lb was partially reconstructed by high-throughput sequencing. The assembled sequence comprised 3 999 304 bp in 736 contigs organized on 42 scaffolds (Supplementary Table S3). Thirty-nine of the 40 universally occurring clusters of orthologous groups used by MLTreeMap (Stark *et al.*, 2010) and tRNAs for all 20 amino acids were represented, suggesting that the obtained sequences cover a major part of the *N. hollandicus* genome (Supplementary Table S3).

All known NOB oxidize nitrite by the enzyme NXR, which belongs to the type II group in the dimethyl sulfoxide reductase family of molybdopterin-cofactor-binding enzymes (Meinke *et al.*, 1992; Lucker *et al.*, 2010). NXR is membrane associated and contains an α subunit (NxrA) with the catalytic site and a β subunit (NxrB) that channels electrons derived from nitrite to downstream components of the respiratory chain (Kirstein and Bock, 1993). Most likely, a γ subunit (NxrC) functions as a membrane anchor that may be involved in electron transport (Lucker *et al.*, 2010). The genome sequence of *N. hollandicus* contains four highly similar (90.3–95.8% protein sequence identity) CDS for NxrA as fragments on contig boundaries. We obtained the full sequences of these genes by PCR, cloning and Sanger sequencing. Only one CDS was found for each of the other NXR subunits. Three *nxA* genes form a cluster also containing a two-component response regulator. The fourth *nxA* copy is directly upstream of *nxB*, a NarJ homolog, and *nxC* (Supplementary Figure S4a), suggesting that these genes represent a functional unit. The genomic localization of this *nxA* copy was confirmed by PCR and Sanger sequencing. NarJ is the δ subunit of dissimilatory nitrate reductase (NAR) and inserts the molybdenum cofactor into the α subunit (Blasco *et al.*, 1998). It may have a similar role in NXR biosynthesis (Starkenbourg *et al.*, 2006). Neither NxrA nor NxrB contained predicted signal peptides for protein secretion, indicating a cytoplasmic localization of these subunits.


Electrons derived from NO_2^- most likely are accepted by cytochrome *a* or *c* and then transferred to a terminal cytochrome *c* oxidase, which in *N. hollandicus* is of the aa_3 type (Supplementary Table S4). This short respiratory chain, including a canonical F_1F_0 ATPase, appears to be highly similar to that of *Nitrobacter* (Spieck and Bock, 2005). Genome annotation also revealed complexes I–III (Supplementary Table S4) that are not required for nitrite oxidation, but may function in the reverse flow of electrons from nitrite to NAD^+ or in energy conservation by the oxidation of organic compounds such as formate. The presence of a [NiFe]-hydrogenase, which contains the signature residues of group 1 (H_2 uptake) hydrogenases (Vignais and Billoud, 2007), implies that H_2 might serve as an

electron donor, although the use of H_2 was not observed in our cultivation-based assays.

N. hollandicus possesses an ammonia transporter of the Amt type and a nitrite/nitrate transporter (NarK). The latter is needed for nitrite import into the cell, because NXR is cytoplasmic, and for the export of nitrate. An urea transporter was identified, but no urea catabolic genes were found in the assembled contigs. Further, while an assimilatory nitrate reductase (NasC) is encoded in the genome, no known assimilatory nitrite reductase was found. One CDS similar to ferredoxin-dependent nitrite or sulfite reductase (NirA/Sir) is present, but the gene product, based on the genomic context, most probably functions as Sir (Supplementary Table S4). Cytochrome *c* nitrite reductase (NrfA) is also absent, indicating that *N. hollandicus* is unable to reduce nitrite to ammonia and thus depends on ammonia as N-source. This would explain the observed requirement of the culture for ammonium. Alternatively, ammonia might have gene regulatory functions in *N. hollandicus* and thus be indispensable. *N. hollandicus* appears to lack genes for denitrification (Supplementary Text S1). Solely nitrate reduction to nitrite was observed (see above) and may be explained by NXR working in reverse as in *Nitrobacter* (Sundermeyer-Klinger *et al.*, 1984) or by the activity of the assimilatory nitrate reductase NasC. However, the genome of *N. hollandicus* has not been completely sequenced yet, and the remaining parts may contain the missing genes for nitrite reduction to ammonia and denitrification.

The only autotrophic carbon fixation pathway present in the genome of *N. hollandicus* is the Calvin–Benson–Bassham cycle (Figure 4 and Supplementary Table S4). This pathway is highly unusual within the *Chloroflexi* where only the genus *Oscillochloris* is known to use the Calvin cycle, whereas other autotrophic members of this phylum use the 3-hydroxypropionate cycle for CO_2 fixation (Berg, 2011). The ribulose-1,5-bisphosphate carboxylase (RubisCO) of *N. hollandicus* belongs to a distinct phylogenetic lineage that is related to the red form I RubisCO and also contains the RubisCO of *Oscillochloris trichoides* (Supplementary Figure S4e).

The presence of a formate transporter and a molybdopterin-dependent formate dehydrogenase (Figure 4 and Supplementary Table S4) is consistent with the utilization of formate by the culture (Supplementary Figure S1a). Only few other putative transporters for organic substrates, including amino acids, were identified in the genome (Figure 4). *N. hollandicus* encodes the complete oxidative tricarboxylic acid cycle that provides building blocks for biosynthesis, but may also be involved in the degradation of storage compounds. Indeed, the glycolytic pathway and gluconeogenesis are present, as well as genes for the production and utilization of glycogen deposits (Supplementary Table S4).



The ISME Journal




Figure 5 Phylogeny of NXR and related enzymes, based on the α subunit. Maximum likelihood tree showing the phylogenetic positioning of selected type II enzymes of the dimethyl sulfoxide reductase family, based on the catalytic (α) subunits. For the analysis 1102 amino acid positions were considered. Names of validated enzymes are indicated (Nxr, boldface; Nar, membrane-bound respiratory nitrate reductase; Pcr, perchlorate reductase; Ebd, ethylbenzene dehydrogenase; Ddh, dimethylsulfide dehydrogenase; Clr, chlorate reductase; Ser, selenate reductase). Parentheses contain the number of sequences within a group or the accession number, respectively. Pie charts indicate statistical support of nodes based on bootstrap analysis or Bayesian inference, respectively. More distantly related molybdoenzymes were used as outgroup.

organism and thus may explain the lower growth rates. The NXR forms of *Nitrobacter*, *Nitrococcus* and *Nitrolancetus* are closely related to each other and to the dissimilatory nitrate reductases (NARs) (Figure 5). The α -subunit NxrA, containing the substrate-binding site, is oriented toward the cytoplasmic side of the cytoplasmic membrane (and of the ICM) in *Nitrobacter* and *Nitrococcus* (Spieck and Bock, 2005), and probably also in *Nitrolancetus* because of the lack of any signal peptide for protein secretion. Thus, known NOB with this cytoplasmic form of NXR consistently require high nitrite concentrations to be competitive. In contrast, the NXR of *Nitrospira* forms a distinct phylogenetic lineage together with NXR of the anammox organism 'Ca. Kuenenia stuttgartiensis' (Lücker *et al.*, 2010) (Figure 5) and it faces the periplasmic space (Spieck *et al.*, 1998; Lücker *et al.*, 2010). The phylogeny of NXR from the genus *Nitrospira* remains unknown because of lacking sequences, but this enzyme is known to be periplasmic as well (Spieck and Bock, 2005). Periplasmic forms of NXR are thought to be more efficient, because more proton-motive force should be generated per oxidized nitrite and no nitrite/nitrate transport across the cytoplasmic membrane is needed (Lücker *et al.*, 2010). Accordingly, *Nitrospira* and *Nitrospina* are much better adapted to lower nitrite concentrations than most *Nitrobacter*, *Nitrococcus* and *Nitrolancetus* (Watson and Waterbury, 1971; Bartosch *et al.*, 1999; Schramm *et al.*, 1999) (Supplementary Table S1). On the basis of the properties of characterized NOB, the cellular localization of NXR and its phylogenetic placement within the molybdopterine-cofactor-binding enzymes (Figure 5 and Supplementary

Figure S4b) seem to be useful indicators for predicting the ecological strategies of nitrite oxidizers.

The only previously known thermotolerant NOB belong to the genus *Nitrospira* (Lebedeva *et al.*, 2011). These organisms are inactive below 37 °C, are inhibited already by nitrite concentrations of 6 mM and do not utilize formate or other organic compounds. In contrast, *N. hollandicus* tolerates a broad temperature range (Supplementary Figure S1b), as well as high nitrite concentrations, and can grow mixotrophically on nitrite and formate (Supplementary Figure S1a). These physiological properties distinguish *N. hollandicus* from all other known NOB. An additional unique feature is the apparent dependence of *N. hollandicus* on ammonia as N-source for growth. This may explain why *N. hollandicus* was not enriched or isolated in previous studies, because mineral media usually applied for the cultivation of NOB contain only trace amounts of ammonia, added as ammonium molybdate (Ehrlich *et al.*, 1995). The growth requirements of *N. hollandicus* match the conditions in the source bioreactor operated at 35 °C with high concentrations of ammonium. On the contrary, most conventional nitrifying wastewater treatment plants are operated at lower temperatures and low ammonia and nitrite concentrations in the bulk liquid. These conditions are optimal for K-strategists such as *Nitrospira* (Daims *et al.*, 2001), but select against NOB requiring higher nitrite levels (Schramm *et al.*, 1999). Thus, owing to its high $K_s(\text{NO}_2^-)$ of 1 mM (Supplementary Table S1) and its relatively high temperature optimum, the functional importance of *N. hollandicus* in such conventional wastewater

treatment plants is unlikely. Indeed, by screening several full-scale wastewater treatment plants and laboratory-scale reactors using specific PCR and FISH, we did not detect *Nitrolancetus* except in the reactor that was the source of isolation. From a technical perspective, however, *N. hollandicus* will likely become relevant with the introduction of increased biogas production and with the treatment of the resulting ammonium-rich digestate. Interestingly, a screening of publicly available 16S rRNA gene sequences retrieved the sequence of a close relative to *N. hollandicus*, which had been obtained from a SHARON reactor in Korea (Figure 3). The high temperatures (30–40 °C) and short sludge retention times in such reactors favor the growth of AOB over that of NOB such as *Nitrobacter* and *Nitrospira*, thus enabling partial nitrification of ammonia to nitrite. The effluent is highly suitable for cost-efficient N-removal by anammox bacteria (van Dongen *et al.*, 2001). *Nitrolancetus*-like NOB, which persist under these conditions, would disturb the process by oxidizing nitrite to nitrate. Under this premise, knowledge of *N. hollandicus* and its ecophysiology could be extremely helpful for process design and optimization.

On the basis of the observed properties of *N. hollandicus*, the natural habitat of these NOB is likely to be characterized by elevated temperatures and high ammonia and nitrite concentrations. Such conditions may be found, for instance, in piles of decomposing plant biomass or dung. Consistently, uncultured relatives of *N. hollandicus* and *S. thermophilus* have been detected in municipal compost (Figure 3).

Evolutionary history of NOB

As *Nitrobacter* and *Nitrococcus* form ICM and are related to phototrophic *Proteobacteria* also containing ICM, a phototrophic origin was suggested for these NOB (Teske *et al.*, 1994). This hypothesis gained strong support from the recent isolation of extant phototrophic NOB in the genera *Rhodopseudomonas* and *Thiocapsa*, which are closely affiliated with *Nitrobacter* and *Nitrococcus*, respectively (Schott *et al.*, 2010). The NXR of *Nitrobacter* and *Nitrococcus* is oriented toward the cytoplasm (see also above). In contrast, genome analysis revealed an evolutionary link between *Nitrospira* and anammox organisms (planctomycetes) that likely involved horizontal gene transfer of a periplasmic NXR (not closely related to the NXR of *Nitrobacter* and *Nitrococcus*, Figure 5 and Supplementary Figure S4b) and other proteins involved in nitrite oxidation and electron transport (Lücker *et al.*, 2010). These lines of evidence suggested two independent lines of evolution of nitrite oxidation, exemplified by the different forms of NXR, in *Nitrobacter* and *Nitrococcus* versus *Nitrospira* and anammox (Lücker *et al.*, 2010). The discovery of *Nitrolancetus* adds more complexity to

this picture, because this member of the *Chloroflexi* clearly did not evolve from the same ancestors as any of the other known NOB. The affiliation of *N. hollandicus* with the *Chloroflexi* is supported not only by 16S rRNA phylogeny (Figure 3) but also by its distinct lipid composition, phylogenetic analysis of 39 universally occurring clusters of orthologous groups (Supplementary Figure S4f) and by numerous highly similar sequence matches between the *N. hollandicus* contig sequences and the genomes of *Sphaerobacter* and *Thermomicrobium* (Supplementary Figure S4g). Although several other *Chloroflexi* are anoxygenic phototrophs (Bryant and Frigaard, 2006), *N. hollandicus* is not closely related to these organisms (Figure 3), has no remnants of a photosynthetic apparatus in its genome, and its next known relatives are chemoorganoheterotrophs. Thus, a direct phototrophic origin of nitrite oxidation in *N. hollandicus* appears unlikely. However, the cytoplasmic NXR of *N. hollandicus* is closely related to the NXR of *Nitrobacter* and *Nitrococcus* despite the large phylogenetic distance between the *Chloroflexi* and *Proteobacteria* (Figure 5 and Supplementary Figure S4b). Interestingly, a similar NXR-like enzyme is found in the anaerobic, nitrite-dependent methane oxidizer ‘*Candidatus Methylo- mirabilis oxyfera*’ that belongs to the distinct bacterial phylum NC10 (Ettwig *et al.*, 2010). The high similarity of NXR in these organisms (Figure 5 and Supplementary Figure S4a and S4b) indicates an unexpected evolutionary link. The most parsimonious explanation is that horizontal gene transfer established the cytoplasmically oriented NXR in ancestors of these very distantly related bacteria. This hypothesis gains support from analyses of the genes upstream and downstream of *nrx* (Supplementary Text S1 and Supplementary Figures S4a, S4c and d). Whether this evolutionary scenario supports or contrasts a phototrophic origin of nitrite oxidation in *Nitrobacter* and *Nitrococcus* (Teske *et al.*, 1994) is a fascinating question that might be better illuminated once first sequences of *nrx* loci from the proteobacterial phototrophic NOB (Schott *et al.*, 2010) become available.

As the presence of NXR in *N. hollandicus* may result from horizontal gene transfer, it is tempting to speculate about the lifestyle of its ancestor before the *nrx* gene cluster was acquired. Although utilization of CO or H₂ was not observed, the genome of *N. hollandicus* contains the required genes (see above and Supplementary Text S1). Thus, it appears possible that the ancestor used CO or H₂ as energy source. Extant relatives of *N. hollandicus* might then still be able to utilize these substrates. Indeed, *T. roseum* can oxidize CO and, similar to *N. hollandicus*, it has genes coding for a [NiFe]-hydrogenase (Wu *et al.*, 2009).

Interestingly, the two known NXR forms are not restricted to NOB but occur in functionally different N-cycle bacteria. The periplasmic form of *Nitrospira* is present in anammox organisms, which also

oxidize nitrite (Strous *et al.*, 2006; Lückner *et al.*, 2010). The NXR type of *Nitrobacter*, *Nitrococcus* and *Nitrolanceus* is shared with *Ca. M. oxyfera* where it may function reversibly and reduce nitrate to nitrite (Ettwig *et al.*, 2010). Hence, NXR is a beautiful example for the exchange and functional adaptation of genetic modules as a key motif in the evolution of the N-cycle (Klotz and Stein, 2008).

Acknowledgements

Alexander Galushko, Jasmin Schwarz and Martina Grill are acknowledged for technical support. We thank Galina Streshinskaya for peptidoglycan analyses. This work was supported by the Russian Foundation for Basic Research (RFBR; grant 10-04-00152), the Vienna Science and Technology Fund (WWTF; grant LS09-40), the Austrian Science Fund (FWF; grant P24101-B22) and the Czech Ministry of Education, Youth and Sports (grant MSM 6046137308).

References

- Alawi M, Lipski A, Sanders T, Pfeiffer E-M, Spieck E. (2007). Cultivation of a novel cold-adapted nitrite oxidizing betaproteobacterium from the Siberian Arctic. *The ISME Journal* **1**: 256–264.
- Andrews JH, Harris RF. (1986). r- and K-selection and microbial ecology. *Adv Microb Ecol* **9**: 99–147.
- Bartosch S, Wolgast I, Spieck E, Bock E. (1999). Identification of nitrite-oxidizing bacteria with monoclonal antibodies recognizing the nitrite oxidoreductase. *Appl Environ Microbiol* **65**: 4126–4133.
- Berg IA. (2011). Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl Environ Microbiol* **77**: 1925–1936.
- Bhandari B, Simlat MM. (1986). Rapid micro-method for determination of nitrate in presence of nitrite for biochemical studies. *Ind J Exp Biol* **24**: 323–325.
- Blasco F, Dos Santos JP, Magalon A, Frixon C, Guigliarelli B, Santini CL *et al.* (1998). NarJ is a specific chaperone required for molybdenum cofactor assembly in nitrate reductase A of *Escherichia coli*. *Mol Microbiol* **28**: 435–447.
- Bryant DA, Frigaard NU. (2006). Prokaryotic photosynthesis and phototrophy illuminated. *Trends Microbiol* **14**: 488–496.
- Canfield DE, Glazer AN, Falkowski PG. (2010). The evolution and future of Earth's nitrogen cycle. *Science* **330**: 192–196.
- Daims H, Lückner S, Wagner M. (2006). *daime*, a novel image analysis program for microbial ecology and biofilm research. *Environ Microbiol* **8**: 200–213.
- Daims H, Nielsen JL, Nielsen PH, Schleifer KH, Wagner M. (2001). In situ characterization of *Nitrospira*-like nitrite-oxidizing bacteria active in wastewater treatment plants. *Appl Environ Microbiol* **67**: 5273–5284.
- Daims H, Stoecker K, Wagner M. (2005). Fluorescence in situ hybridisation for the detection of prokaryotes. In: Osborn AM, Smith CJ (eds). *Molecular microbial ecology*. Bios-Garland: Abingdon, UK, pp 213–239.
- Daims H, Wagner M. (2007). Quantification of uncultured microorganisms by fluorescence microscopy and digital image analysis. *Appl Microbiol Biotechnol* **75**: 237–248.
- Dechorgnat J, Nguyen CT, Armengaud P, Jossier M, Diatloff E, Filleur S *et al.* (2011). From the soil to the seeds: the long journey of nitrate in plants. *J Exp Bot* **62**: 1349–1359.
- Ehrich S, Behrens D, Lebedeva E, Ludwig W, Bock E. (1995). A new obligately chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its phylogenetic relationship. *Arch Microbiol* **164**: 16–23.
- Ettwig KF, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MM *et al.* (2010). Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* **464**: 543–548.
- Griess-Romijn van Eck E. (1966). *Physiological and chemical tests for drinking water*. NEN 1056 IV-2 Netherlands Normalisatie Instituut: Rijswijk, The Netherlands.
- Griffin BM, Schott J, Schink B. (2007). Nitrite, an electron donor for anoxygenic photosynthesis. *Science* **316**: 1870.
- Gruber N. (2004). The dynamics of the marine nitrogen cycle and its influence on atmospheric CO₂ variations. In Follows M, Oguz T (eds). *NATO ASI Series*. Kluwer Academic: Dordrecht, pp 97–148.
- Hanada S, Pierson BK. (2006). The family *Chloroflexaceae*. In: Dworkin M, Falkow S, Rosenberg E, Schleifer KH, Stackebrandt E (eds). *The Prokaryotes: A handbook on the biology of Bacteria*. Springer Science + Business Media: New York, pp 815–842.
- Hugenholtz P, Stackebrandt E. (2004). Reclassification of *Sphaerobacter thermophilus* from the subclass *Sphaerobacteridae* in the phylum *Actinobacteria* to the class *Thermomicrobia* (emended description) in the phylum *Chloroflexi* (emended description). *Int J Syst Evol Microbiol* **54**: 2049–2051.
- Juretschko S, Loy A, Lehner A, Wagner M. (2002). The microbial community composition of a nitrifying-denitrifying activated sludge from an industrial sewage treatment plant analyzed by the full-cycle rRNA approach. *System Appl Microbiol* **25**: 84–99.
- Kampschreur MJ, Tan NCG, Kleerebezem R, Picioroanu C, Jetten MSM, Loosdrecht MCM. (2008). Effect of dynamic process conditions on nitrogen oxides emission from a nitrifying culture. *Environ Sci Technol* **42**: 429–435.
- Kirstein K, Bock E. (1993). Close genetic relationship between *Nitrobacter hamburgensis* nitrite oxidoreductase and *Escherichia coli* nitrate reductases. *Arch Microbiol* **160**: 447–453.
- Klotz MG, Stein LY. (2008). Nitrifier genomics and evolution of the nitrogen cycle. *FEMS Microbiol Lett* **278**: 146–156.
- Kyrpides NC. (1999). Genomes OnLine Database (GOLD 1.0): a monitor of complete and ongoing genome projects world-wide. *Bioinformatics* **15**: 773–774.
- Könneke M, Bernhard AE, de la Torre JR, Walker CB, Waterbury JB, Stahl DA. (2005). Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* **437**: 543–546.
- Lebedeva EV, Off S, Zumbragel S, Kruse M, Shagzhina A, Lucker S *et al.* (2011). Isolation and characterization of a moderately thermophilic nitrite-oxidizing bacterium from a geothermal spring. *FEMS Microbiol Ecol* **75**: 195–204.

- Leininger S, Urich T, Schlöter M, Schwark L, Qi J, Nicol GW *et al.* (2006). Archaea predominate among ammonia-oxidizing prokaryotes in soils. *Nature* **442**: 806–809.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. (1951). Protein measurement with the Folin phenol reagent. *J Biol Chem* **193**: 265–275.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhav K *et al.* (2004). ARB: a software environment for sequence data. *Nucleic Acids Res* **32**: 1363–1371.
- Lücker S, Wagner M, Maixner F, Pelletier E, Koch H, Vacherie B *et al.* (2010). A *Nitrospira* metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing bacteria. *Proc Natl Acad Sci USA* **107**: 13479–13484.
- Maixner F, Noguera DR, Anneser B, Stoecker K, Wegl G, Wagner M *et al.* (2006). Nitrite concentration influences the population structure of *Nitrospira*-like bacteria. *Environ Microbiol* **8**: 1487–1495.
- Meincke M, Bock E, Kastrau D, Kroneck PMH. (1992). Nitrite oxidoreductase from *Nitrobacter hamburgensis*: redox centers and their catalytic role. *Arch Microbiol* **158**: 127–131.
- Pfennig N, Lippert KD. (1966). Über das Vitamin B12-Bedürfnis phototropher Schwefelbakterien. *Arch Microbiol* **55**: 245.
- Pond JL, Langworthy TA, Holzer G. (1986). Long-chain diols—a new class of membrane-lipids from a thermophilic bacterium. *Science* **231**: 1134–1136.
- Purkhold U, Pommering-Röser A, Juretschko S, Schmid MC, Koops H-P, Wagner M. (2000). Phylogeny of all recognized species of ammonia oxidizers based on comparative 16S rRNA and amoA sequence analysis: implications for molecular diversity surveys. *Appl Environ Microbiol* **66**: 5368–5382.
- Schloss PD, Handelsman J. (2004). Status of the microbial census. *Microbiol Mol Biol Rev* **68**: 686–691.
- Schott J, Griffin BM, Schink B. (2010). Anaerobic phototrophic nitrite oxidation by *Thiocapsa* sp. strain KS1 and *Rhodopseudomonas* sp. strain LQ17. *Microbiol-ogy* **156**: 2428–2437.
- Schramm A, de Beer D, van den Heuvel JC, Ottengraf S, Amann R. (1999). Microscale distribution of populations and activities of *Nitrosospira* and *Nitrospira* spp. along a macroscale gradient in a nitrifying bioreactor: quantification by in situ hybridization and the use of micro-sensors. *Appl Environ Microbiol* **65**: 3690–3696.
- Shapleigh JP. (2006). The denitrifying prokaryotes. In: Dworkin M, Falkow S, Rosenberg E, Schleifer KH, Stackebrandt E (eds). *The Prokaryotes: a handbook on the biology of Bacteria*, 3 edn. Springer Science + Business Media: New York, pp 769–792.
- Spieck E, Aamand J, Bartosch S, Bock E. (1996). Immunocytochemical detection and location of the membrane-bound nitrite oxidoreductase in cells of *Nitrobacter* and *Nitrospira*. *FEMS Microbiol Lett* **139**: 71–76.
- Spieck E, Bock E. (2005). The lithoautotrophic nitrite-oxidizing bacteria. In: Staley JT, Boone DR, Brenner DJ, de Vos P, Garrity GM, Goodfellow M *et al.* (eds). *Bergey's manual of systematic bacteriology*, 2 edn. Springer Science + Business Media: New York, pp 149–153.
- Spieck E, Ehrlich S, Aamand J, Bock E. (1998). Isolation and immunocytochemical location of the nitrite-oxidizing system in *Nitrospira moscoviensis*. *Arch Microbiol* **169**: 225–230.
- Stark M, Berger SA, Stamatakis A, von Mering C. (2010). MLTreeMap—accurate Maximum Likelihood placement of environmental DNA sequences into taxonomic and functional reference phylogenies. *BMC Genomics* **11**: 461.
- Starkenburger SR, Chain PS, Sayavedra-Soto LA, Hauser L, Land ML, Larimer FW *et al.* (2006). Genome sequence of the chemolithoautotrophic nitrite-oxidizing bacterium *Nitrobacter winogradskyi* Nb-255. *Appl Environ Microbiol* **72**: 2050–2063.
- Stein LY, Arp DJ. (1998). Loss of ammonia monooxygenase activity in *Nitrosomonas europaea* upon exposure to nitrite. *Appl Environ Microbiol* **64**: 4098–4102.
- Strous M, Pelletier E, Mangenot S, Rattei T, Lehner A, Taylor MW *et al.* (2006). Deciphering the evolution and metabolism of an anammox bacterium from a community genome. *Nature* **440**: 790–794.
- Sundermeyer-Klinger H, Meyer W, Warninghoff B, Bock E. (1984). Membrane-bound nitrite oxidoreductase of *Nitrobacter*: evidence for a nitrate reductase system. *Arch Microbiol* **140**: 153–158.
- Sutcliffe IC. (2011). Cell envelope architecture in the *Chloroflexi*: a shifting frontline in a phylogenetic turf war. *Environ Microbiol* **13**: 279–282.
- Taylor MW, Radax R, Steger D, Wagner M. (2007). Sponge-associated microorganisms: evolution, ecology, and biotechnological potential. *Microbiol Mol Biol Rev* **71**: 295–347.
- Teske A, Alm E, Regan JM, Toze S, Rittmann BE, Stahl DA. (1994). Evolutionary relationships among ammonia- and nitrite-oxidizing bacteria. *J Bacteriol* **176**: 6623–6630.
- Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S *et al.* (2006). MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res* **34**: 53–65.
- van Dongen U, Jetten MS, van Loosdrecht MC. (2001). The SHARON-Anammox process for treatment of ammonium rich wastewater. *Water Sci Technol* **44**: 153–160.
- Van Gool A, Laudelout H. (1966). Formate utilization by *Nitrobacter winogradskyi*. *Biochim Biophys Acta* **127**: 295–301.
- van der Meer MT, Schouten S, Hanada S, Hopmans EC, Damste JS, Ward DM. (2002). Alkane-1,2-diol-based glycosides and fatty glycosides and wax esters in *Roseiflexus castenholzii* and hot spring microbial mats. *Arch Microbiol* **178**: 229–237.
- Vejmelkova D, Sorokin DY, Abbas B, Kovaleva OL, Kleerebezem R, Kampschreur MJ *et al.* (2011). Analysis of ammonia-oxidizing bacteria dominating in lab-scale bioreactors with high ammonium bicarbonate loading. *Appl Microbiol Biotechnol* **93**: 401–410.
- Vignais PM, Billoud B. (2007). Occurrence, classification, and biological function of hydrogenases: an overview. *Chem Rev* **107**: 4206–4272.
- Wait R, Carreto L, Nobre MF, Ferreira AM, da Costa MS. (1997). Characterization of novel long-chain 1,2-diols in *Thermus* species and demonstration that *Thermus* strains contain both glycerol-linked and diol-linked glycolipids. *J Bacteriol* **179**: 6154–6162.
- Watson SW, Waterbury JB. (1971). Characteristics of two marine nitrite oxidizing bacteria, *Nitrospina gracilis* nov. gen. nov. sp. and *Nitrococcus mobilis* nov. gen. nov. sp. *Arch Mikrobiol* **77**: 203–230.
- Winogradsky S. (1892). Contributions a la morphologie des organismes de la nitrification. *Arch Sci Biol (St Petersburg)* **1**: 88–137.

- Wu DY, Raymond J, Wu M, Chatterji S, Ren QH, Graham JE *et al.* (2009). Complete genome sequence of the aerobic CO-oxidizing thermophile *Thermomicrobium roseum*. *PLoS One* 4 e4207.
- Wuchter C, Abbas B, Coolen MJ, Herfort L, van Bleijswijk J, Timmers P *et al.* (2006). Archaeal nitrification in the ocean. *Proc Natl Acad Sci USA* **103**: 12317–12322.
- Zehr JP, Jenkins BD, Short SM, Steward GF. (2003). Nitrogenase gene diversity and microbial community

structure: a cross-system comparison. *Environ Microbiol* **5**: 539–554.



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on The ISME Journal website ([/http://www.nature.com/ismej](http://www.nature.com/ismej))