



Published in final edited form as:

Nature. 2008 March 13; 452(7184): 225–229. doi:10.1038/nature06642.

A skin microRNA promotes differentiation by repressing ‘stemness’

Rui Yi^{1,2}, Matthew N. Poy³, Markus Stoffel³, and Elaine Fuchs^{1,2}

¹ Howard Hughes Medical Institute, The Rockefeller University, New York City, New York 10065, USA ² Laboratory of Mammalian Cell Biology and Development, The Rockefeller University, New York City, New York 10065, USA ³ Swiss Federal Institute of Technology ETH Zurich, Institute of Molecular Systems Biology, CH-8093 Zürich, Switzerland

Abstract

In stratified epithelial tissues, homeostasis relies on the self-renewing capacity of stem cells located within the innermost basal layer¹. As basal cells become suprabasal, they lose proliferative potential and embark on a terminal differentiation programme^{2,3}. Here, we show that microRNA-203 is induced in the skin concomitantly with stratification and differentiation. By altering miR-203’s spatiotemporal expression *in vivo*, we show that miR-203 promotes epidermal differentiation by restricting proliferative potential and inducing cell-cycle exit. We identify *p63* as one of the conserved targets of miR-203 across vertebrates. Notably, *p63* is an essential regulator of stem-cell maintenance in stratified epithelial tissues^{4–9}. We show that miR-203 directly represses the expression of *p63*: it fails to switch off suprabasally when either *Dicer1* or miR-203 is absent and it becomes repressed basally when miR-203 is prematurely expressed. Our findings suggest that miR-203 defines a molecular boundary between proliferative basal progenitors and terminally differentiating suprabasal cells, ensuring proper identity of neighbouring layers.

MicroRNAs are small, non-coding RNAs that regulate gene expression post-transcriptionally by directly targeting RNA-induced silencing complex (RISC) to cognate messenger RNA targets¹⁰. When miRNAs are globally ablated in skin epithelium by conditionally targeting the gene that encodes the miRNA-processing enzyme *Dicer1*, hair follicles fail to invaginate. This distorts epidermal morphology, compromising the barrier and underscoring the functional importance of these small RNAs in skin development^{11,12}. To gain further insight into the possible significance of different skin miRNAs, we constructed epidermal miRNA libraries using total RNAs isolated from pure epidermis starting from embryonic day 13.5 (E13.5), when it is still a single-layered epithelium, to postnatal day 4.5 (P4.5), when it is fully stratified. Among more than 100 epidermal

Correspondence and requests for materials should be addressed to E.F. (fuchslb@rockefeller.edu).

Author Contributions R.Y. and E.F. designed the research. R.Y. performed the experiments. M.N.P. and M.S. designed and contributed to the antagonism experiments. R.Y. and E.F. analysed the data and wrote the paper.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare competing financial interests: details accompany the full-text HTML version of the paper at www.nature.com/nature.

miRNAs cloned, miR-203 barely surfaced in the pool of E13.5 clones but emerged as one of the most abundant epidermal miRNAs from E15.5 onwards (Fig. 1a).

The significant upregulation of miR-203 between E13.5 and E15.5 was suggestive that this miRNA may be absent in multipotent progenitors of single-layered epidermis, but is induced upon stratification and differentiation. By *in situ* hybridization¹³, miR-203 was largely confined to suprabasal cells of epithelial tissues, and was especially prominent in skin (Fig. 1b, d, e and Supplementary Fig. 1). The specificity of hybridization was confirmed by analysing conditionally ablated *Dicer1* skin, in which expression of miR-203 and other mature miRNAs was abolished¹¹ (Supplementary Fig. 1a). Quantification of its differential expression by quantitative PCR with reverse transcription (qRT-PCR) revealed ~25-fold more miR-203 in E15.5 suprabasal cells than in their basal counterparts (Fig. 1c). Similarly, miR-203 was rapidly upregulated when primary mouse keratinocytes were induced by calcium to differentiate *in vitro* (Supplementary Fig. 1b).

Epidermal development precedes that of its appendages. However, at early stages of hair follicle development, miR-203 was not detected. By E17.5, faint miR-203 hybridization was detected within the emerging suprabasal cells of developing hair follicles and expression was also seen in stratified layers of developing tongue epithelia (Supplementary Fig. 1f, g). As development advanced, miR-203 expression intensified in differentiating cells of epidermis, hair follicles and sebaceous glands (Fig. 1d, e). Present throughout transcriptionally active, terminally differentiating cells of skin epithelium, miR-203 was conspicuously absent in its proliferating progenitor compartments. Interestingly, miR-203's sequence and expression pattern seemed to be conserved among vertebrates, in which the epidermis is stratified, but not in eukaryotes that have a single-layered epidermis (Supplementary Fig. 2a, b).

If miR-203 functions in the switch between proliferative and terminally differentiating compartments in vertebrate skin, it may be expected to repress its basal targets once basal cells become suprabasal and enter the terminal differentiation programme. To test this hypothesis, we generated transgenic mice expressing miR-203 under the control of the *keratin 14* (*K14*) promoter, active by E15 in basal progenitors of stratified epithelia¹⁴. Most *K14-miR-203* mice died shortly after birth owing to apparent dehydration and/or malnutrition. By E18.5, the level of transgenic basal miR-203 was comparable to endogenous suprabasal miR-203 (Fig. 2a). Moreover, transgenic miR-203 expression did not interfere with endogenous miRNA processing¹⁵ (Supplementary Fig. 3).

At E18.5, *K14-miR-203* back skin epidermis was noticeably thinner than that of its wild-type littermates (Supplementary Fig. 4a). By the time of birth (P0), transgenic epidermis consisted of a layer of flattened basal cells and one layer of suprabasal cells (Fig. 2b). Hence, a thin epidermis can reflect either defective differentiation, as in mice conditionally targeted by *K14-Cre* for Notch effector protein RBPj¹⁶, or impaired stem cells, as in *p63* null epidermis^{4,6,9}. To distinguish between these possibilities, we first examined keratin 10, an early differentiation marker downstream of canonical Notch signalling¹⁶. At E18.5, many keratin-10-positive cells within *K14-miR-203* epidermis were aberrantly juxtaposed to the basement membrane (Fig. 2c). As development progressed, basal progenitor/stem cells

marked by keratin 5 were missing over increasingly larger epidermal stretches that were replaced by flat cells expressing keratin 10 (Fig. 2d). No discernible apoptotic cells were detected as judged by TdT-mediated dUTP nick end labelling (TUNEL) assay and active caspase-3 (data not shown).

The depletion of basal stem cells in *K14-miR-203* epidermis bore a resemblance to *p63* null epidermis^{6,9}. Interestingly, anti-p63 immunolocalization revealed only sporadic, weak expression in E18.5 *K14-miR-203* basal cells (Fig. 2e). Quantification of 5-bromodeoxyuridine (BrdU)-positive cells after a 2-h pulse-labelling revealed a significant reduction in the proliferative pool of E18.5 transgenic progenitors (Fig. 2f).

To address whether miR-203 restricts the proliferative potential of epidermal stem cells, we compared the clonogenic capacity of primary mouse keratinocytes cultured from E18.5 transgenic with wild-type littermates. As expected, wild-type keratinocytes formed typical holoclones composed of small, undifferentiated cells capable of long-term passage^{17,18}. By contrast, transgenic keratinocytes produced mostly paraclones, composed of large, flattened cells^{17,18} (Supplementary Fig. 4b). Moreover, approximately fourfold fewer, and significantly smaller, colonies formed from transgenic compared with wild-type keratinocytes (Fig. 2g and Supplementary Fig. 4c). On passage, only wild-type clones gave rise to colonies. Consistent with a requirement for p63 in maintaining proliferative potential of epidermal stem cells⁹, p63 was markedly diminished in transgenic keratinocytes.

To test whether these dramatic effects of transgene expression are specific to miR-203, we transduced primary mouse keratinocytes with retroviral vectors expressing either miR-203 or miR-203M, a miR-203 mutant harbouring mutations within the 5' seed (Supplementary Fig. 4d), or empty vector alone. Only miR-203 impaired keratinocyte proliferation and colony formation (Fig. 2h and Supplementary Fig. 4e).

To identify early consequences of losing miR-203 expression, we first examined p63 and cell-cycle status in E18.5 *Dicer1* null epidermis. In contrast to the wild type^{2,5,9}, *Dicer1* null epidermis showed frequent p63-positive, BrdU-positive and phospho-histone-H3-positive mitotic suprabasal cells (Supplementary Fig. 5a-c). Fluorescence-activated cell sorting (FACS) quantification revealed an approximately threefold increase in the number of G2/M phase suprabasal cells (Supplementary Fig. 5d).

To test specifically whether miR-203 represses proliferative potential of suprabasal cells *in vivo*, we adapted the antagomir method¹⁹ to repress miR-203 expression in dorsal skin of neonatal mice, in which basal-layer proliferation is still high. We first used a dye-conjugated antagomir to document efficient incorporation in epidermis within 24 h following a single subcutaneous injection (Supplementary Fig. 6a). One day after three dorsal injections, antagomir-203, but not mm-antagomir-203 (a 4-nucleotide mismatched control) or PBS markedly and specifically repressed miR-203 expression in surrounding skin (Fig. 3a and Supplementary Fig. 6b, c).

Similar to E18.5 *Dicer1* null skin, antagomir-treated P4 epidermis showed atypical expansion of p63 expression and BrdU-positive suprabasal cells (Fig. 3b, c). This was further substantiated by the presence of suprabasal cells positive for p63 and the mitotic

marker phospho-histone H3 (Fig. 3d). Finally, quantification of epidermal cells harvested 4 h after BrdU injections revealed that proliferative cells were more numerous in antagomir-203-treated dorsal epidermis than either PBS- or mm-antagomir-203-treated skin (Fig. 3e).

Because these abnormalities occurred early after miR-203 repression, it was unlikely that they arose secondarily from physical perturbations to the skin or mouse. Moreover, the alterations were recapitulated *in vitro* with an antisense oligonucleotide²⁰ that specifically blocked calcium-induced expression of endogenous miR-203 (Supplementary Fig. 6d). BrdU labelling and FACS analysis revealed that anti-miR-203 *in vitro* partially blocked the ability of keratinocytes to exit the cell cycle during calcium-induced differentiation (Fig. 3f).

Having established early consequences of inhibiting miR-203 expression, we examined longer-term effects *in vivo*. Although tail-vein injections of antagomirs afford efficient delivery for some adult tissues¹⁹, it was only partially effective in targeting miR-203 knockdown to skin (Supplementary Fig. 7). Similarly, sustained subcutaneous antagomir injections were not effective once skin matured. To ablate miR-203 efficiently in adult skin, we engineered *K14-CreTm/Dicer1* inducible conditional knockout (iKO) mice and examined long-term consequences of conditionally inducing *Dicer1* ablation. By 30 days, the epidermis had thickened markedly and suprabasal cells were more numerous (Fig. 3g). Atypical p63-positive suprabasal cells were now prevalent in iKO epidermis (Fig. 3h). These results suggested that miR-203 may be required to effectively repress supra-basal p63 and in turn restrict cell proliferation in differentiating epidermis. Importantly, by targeting *Dicer1* after follicles had matured and anchored within the dermis, we could attribute these defects specifically to alterations originating in the epidermis rather than secondarily arising from perturbed hair follicle morphogenesis.

Whereas miR-203 markedly impaired proliferative potential, it had minimal effects on the induction of terminal differentiation markers whether *in vivo* (Fig. 2c, d) or *in vitro* (Supplementary Fig. 8). Thus, of the two key features involved in terminal differentiation, suprabasal miR-203 seemed to act predominantly by restricting proliferative potential of progenitors as they transitioned from basal to suprabasal layers. Although loss of miR-203 resulted in increased suprabasal proliferation, neither *p63* overexpression nor miR-203 reduction are associated with psoriasis^{21,22}, suggesting that neither p63 nor the miRNA are obligatory features of the hyperproliferative state.

Bioinformatics suggest that miRNAs and their predicted targets tend to be mutually exclusive in neighbouring tissues^{23,24}. On the basis of the inverse correlation that we observed between miR-203 and p63 in expression and function, *p63* mRNA surfaced as a possible miR-203 target. Consistent with *in vivo* results (Fig. 2e), p63 was markedly diminished in *K14-miR-203* transgenic primary mouse keratinocytes that were cultured in low calcium (Fig. 4a). Conversely, knocking down endogenous miR-203 in calcium-treated wild-type primary mouse keratinocyte cultures strongly impaired the rapid down-regulation of p63 protein upon terminal differentiation (Fig. 4b), even though *p63* mRNA levels remained appreciable (Supplementary Fig. 9). Conversely, transduction of miR-203 in wild-type primary mouse keratinocytes cultured in low calcium strongly inhibited p63 protein

expression (Fig. 4b), while only slightly decreasing *p63* mRNA (Fig. 4b right-hand graph). Thus, miR-203 seemed to regulate *p63* primarily through translational repression.

To further define the parallels, we examined the status of p21, a well-established *p63* target that functions to restrict proliferative potential of basal epidermal cells²⁵ and is normally repressed in proliferative keratinocytes^{26–28}. Upon miR-203 transduction, however, p21 was upregulated in undifferentiated wild-type primary mouse keratinocytes to an extent that was nearly comparable to its levels in differentiated primary mouse keratinocytes (Fig. 4c). This finding extends the correlation between miR-203 and *p63* to *p63*'s downstream targets, and provides mechanistic insight into how miR-203 may inhibit proliferative potential of epidermal stem cells.

DeltaN*p63α* is the main isoform expressed in epidermis. Within the 3'UTR of *Np63α* mRNA, a hexamer and heptamer match perfectly to miR-203's 5' seed sequence (Fig. 4d). When introduced into the 3'UTR of a luciferase reporter gene, a 599-base-pair mouse *Np63α* 3'UTR fragment encompassing these two putative target sites caused a threefold reduction in activity in *K14-miR-203*-expressing keratinocytes cultured in low calcium (Fig. 4d). Mutations within these two sites together abolished miR-203-mediated repression under the same conditions (Fig. 4e, LoCa). When these assays were performed in high-calcium primary mouse keratinocytes in which endogenous miR-203 was induced (Supplementary Fig. 1b), exogenous miR-203 was not needed to elicit the wild-type *p63* 3'UTR-specific reduction in luciferase activity (Fig. 4e, HiCa). Together, these results indicate that miR-203's effect on *p63* is direct and is mediated through these 3'UTR target sites.

MiR-203 and its suprabasal expression seem to have emerged in vertebrate evolution concomitant with a stratified epidermis. Similarly, the basal expression and function of *p63* in stratified epithelia also seems to be conserved^{4–7,9}. Interestingly, the putative miR-203 recognition sequence exists four times in the *p63* 3'UTR of zebrafish and twice in mouse and human. Moreover, when similarly tested, the *p63* 3'UTR fragments flanking the hexamer motifs from either zebrafish or human reduced luciferase reporter activity comparably to that from mouse (Supplementary Fig. 10). These data suggest that this mechanism for regulating *p63* by miR-203 may be conserved across vertebrates.

Strictly on the basis of bioinformatic studies, the mRNA encoding a zinc-finger protein, *Zfp281*, is the top candidate of miR-203 targets, as it harbours four perfectly conserved miR-203 consensus sites in its 3'UTR²⁹ (Supplementary Fig. 11a). Similarly to *p63*, *Zfp281* is also enriched in basal epidermal stem cells (Supplementary Fig. 11b). In both *K14-miR-203* transgenic mice and miR-203-transduced primary mouse keratinocytes, *Zfp281* mRNA was markedly reduced (Supplementary Fig. 11c). Using a luciferase reporter assay, we showed that the *Zfp281* 3'UTR, but not the mutant form with mutations at four target sites, was specifically regulated by co-transfected and endogenous miR-203 (Supplementary Fig. 11a). Intriguingly, even though *Zfp281* has not been previously studied in skin development, it has been implicated in maintaining the proliferative potential of embryonic stem cells³⁰, and in this regard its function resembles that ascribed to *p63* in keratinocytes.

Taken together, our studies provide evidence that miR-203 acts at least in part by targeting and negatively regulating suprabasal expression of basal genes, thereby acting as a switch between proliferation and differentiation. Although the mechanisms underlying this balance are complex and likely to be dominated by transcriptional controls, the mutually exclusive expression patterns, opposite functions and evolutionarily conserved regulation offer strong support for the intimate relation between miR-203 and its targets in refining the boundary between these two stages in epidermis.

METHODS SUMMARY

Antagomir synthesis and injection

Antagomir-203 and mm-antagomir-203 were designed and synthesized (Dharmacon) as described¹⁹. The antagomir-203 sequence was 5'-UpsCps UAGUGGUCCUAAACAUp sUpsCpsAps-Chol-3' and mm-antagomir-203 was 5'-UpsCpsUcGUGuUCaUAAACAcUpsUpsCpsAps-Chol-3'. Lower-case letters indicate the mismatched nucleotide. Subcutaneous injection was performed on newborn CD-1 mice at a dosage of 80 mg kg⁻¹, according to an established protocol approved by LARC animal facility at the Rockefeller University. Tail-vein injections were performed as described¹⁹. After three initial injections, a booster injection was performed every 4 days. Total treatment was 15 days.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank D. O'Carroll and A. Tarakhovsky for *Dicer1*^{fl/fl} mice; A. Schaefer and P. Greengard for dye-conjugated antagomir-124; Z. Zhang and F. Dietrich for miRNA library sequencing; B. Liu for bioinformatics assistance; A. Giraldez for zebrafish cDNA; N. Stokes and L. Polak for assistance with animals; S. Mazel and X. Fan for assistance in the flow cytometry core facility; J. Racelis for assistance with *in situ* hybridization; and D. Wang and E. Fuchs laboratory members for discussions. R.Y. is supported by the Pathway to Independence Award from the NIH. M.N.P. is supported by the Ruth L. Kirschstein NRSA Fellowship from the NIH. E.F. is an investigator of the Howard Hughes Medical Institute. This work was supported by the HHMI and the NIH.

References

1. Fuchs E. Scratching the surface of skin development. *Nature*. 2007; 445:834–842. [PubMed: 17314969]
2. Lechler T, Fuchs E. Asymmetric cell divisions promote stratification and differentiation of mammalian skin. *Nature*. 2005; 437:275–280. [PubMed: 16094321]
3. Clayton E, et al. A single type of progenitor cell maintains normal epidermis. *Nature*. 2007; 446:185–189. [PubMed: 17330052]
4. Mills AA, et al. p63 is a p53 homologue required for limb and epidermal morphogenesis. *Nature*. 1999; 398:708–713. [PubMed: 10227293]
5. Parsa R, Yang A, McKeon F, Green H. Association of p63 with proliferative potential in normal and neoplastic human keratinocytes. *J Invest Dermatol*. 1999; 113:1099–1105. [PubMed: 10594758]
6. Yang A, et al. p63 is essential for regenerative proliferation in limb, craniofacial and epithelial development. *Nature*. 1999; 398:714–718. [PubMed: 10227294]
7. Lee H, Kimelman D. A dominant-negative form of p63 is required for epidermal proliferation in zebrafish. *Dev Cell*. 2002; 2:607–616. [PubMed: 12015968]

8. Truong AB, Kretz M, Ridky TW, Kimmel R, Khavari P. A p63 regulates proliferation and differentiation of developmentally mature keratinocytes. *Genes Dev.* 2006; 20:3185–3197. [PubMed: 17114587]
9. Senoo M, Pinto F, Crum CP, McKeon F. p63 is essential for the proliferative potential of stem cells in stratified epithelia. *Cell.* 2007; 129:523–536. [PubMed: 17482546]
10. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004; 116:281–297. [PubMed: 14744438]
11. Yi R, et al. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nature Genet.* 2006; 38:356–362. [PubMed: 16462742]
12. Andl T, et al. The miRNA-processing enzyme dicer is essential for the morphogenesis and maintenance of hair follicles. *Curr Biol.* 2006; 16:1041–1049. [PubMed: 16682203]
13. Wienholds E, et al. MicroRNA expression in zebrafish embryonic development. *Science.* 2005; 309:310–311. [PubMed: 15919954]
14. Vasioukhin V, Degenstein L, Wise B, Fuchs E. The magical touch: genome targeting in epidermal stem cells induced by tamoxifen application to mouse skin. *Proc Natl Acad Sci USA.* 1999; 96:8551–8556. [PubMed: 10411913]
15. Grimm D, et al. Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature.* 2006; 441:537–541. [PubMed: 16724069]
16. Blanpain C, Lowry WE, Pasolli HA, Fuchs E. Canonical notch signaling functions as a commitment switch in the epidermal lineage. *Genes Dev.* 2006; 20:3022–3035. [PubMed: 17079689]
17. Barrandon Y, Green H. Three clonal types of keratinocyte with different capacities for multiplication. *Proc Natl Acad Sci USA.* 1987; 84:2302–2306. [PubMed: 2436229]
18. Blanpain C, Lowry WE, Geoghegan A, Polak L, Fuchs E. Self-renewal, multipotency, and the existence of two cell populations within an epithelial stem cell niche. *Cell.* 2004; 118:635–648. [PubMed: 15339667]
19. Krutzfeldt J, et al. Silencing of microRNAs *in vivo* with ‘antagomirs’. *Nature.* 2005; 438:685–689. [PubMed: 16258535]
20. Davis S, Lollo B, Freier S, Esau C. Improved targeting of miRNA with antisense oligonucleotides. *Nucleic Acids Res.* 2006; 34:2294–2304. [PubMed: 16690972]
21. Haider AS, et al. Genomic analysis defines a cancer-specific gene expression signature for human squamous cell carcinoma and distinguishes malignant hyperproliferation from benign hyperplasia. *J Invest Dermatol.* 2006; 126:869–881. [PubMed: 16470182]
22. Sonkoly E, et al. MicroRNAs: novel regulators involved in the pathogenesis of psoriasis? *PLoS ONE.* 2007; 2:e610. [PubMed: 17622355]
23. Farh KK, et al. The widespread impact of mammalian MicroRNAs on mRNA repression and evolution. *Science.* 2005; 310:1817–1821. [PubMed: 16308420]
24. Stark A, Brennecke J, Bushati N, Russell RB, Cohen SM. Animal MicroRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell.* 2005; 123:1133–1146. [PubMed: 16337999]
25. Topley GI, Okuyama R, Gonzales JG, Conti C, Dotto G. P p21(WAF1/Cip1) functions as a suppressor of malignant skin tumor formation and a determinant of keratinocyte stem-cell potential. *Proc Natl Acad Sci USA.* 1999; 96:9089–9094. [PubMed: 10430900]
26. Okuyama R, et al. p53 homologue, p51/p63, maintains the immaturity of keratinocyte stem cells by inhibiting Notch1 activity. *Oncogene.* 2007; 26:4478–4488. [PubMed: 17237812]
27. Nguyen BC, et al. Cross-regulation between Notch and p63 in keratinocyte commitment to differentiation. *Genes Dev.* 2006; 20:1028–1042. [PubMed: 16618808]
28. Westfall MD, Mays DJ, Sniezek JC, Pietenpol JA. The Delta Np63 alpha phosphoprotein binds the p21 and 14–3–3 sigma promoters *in vivo* and has transcriptional repressor activity that is reduced by Hay-Wells syndrome-derived mutations. *Mol Cell Biol.* 2003; 23:2264–2276. [PubMed: 12640112]
29. Grimson A, et al. MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol Cell.* 2007; 27:91–105. [PubMed: 17612493]

30. Wang J, et al. A protein interaction network for pluripotency of embryonic stem cells. *Nature*. 2006; 444:364–368. [PubMed: 17093407]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Spatiotemporal expression of miR-203 during skin development

a, Temporal induction of miR-203 as measured by cloning frequency. **b**, **d**, **e**, *In situ* hybridization reveals restriction of miR-203 to suprabasal, differentiating layers of skin. sb, suprabasal; bl, basal layer; epi, epidermis; der, dermis; sg, sebaceous gland; irs, inner root sheath; hf, hair follicle. Asterisks in **e** represent brown melanin pigment that is not a hybridization signal. In **b**, anti- β 4-integrin co-labelling is in green and miR-203 *in situ* pseudocoloured signal is in red. **c**, qRT-PCR of FACS-purified cells from E15.5 epidermis reveals 25-fold more miR-203 in suprabasal (SB) versus basal layer (BL) cells ($*P < 0.002$). Error bars (s.d.) are derived from three experiments with basal layer level set as 1. Scale bars are 30 μ m.




Figure 2. Premature activation of miR-203 in epidermis restricts its proliferative potential
a, *In situ* hybridization detects precociously expressed miR-203 in basal epidermis and hair germs of *K14-miR-203* Tg skin. WT, wild type. **b**, By P0, Tg epidermis is thinner than the wild type. **c–f**, Signs of basal cell depletion in Tg skin. Arrows denote keratin-10 (K10)-positive, keratin-5 (K5)-negative, β 4-integrin-low cells aberrantly juxtaposed to basement membrane. Note marked reduction of p63 in **e**. In **f**, quantifications reflect $*P < 0.003$, $n = 8$. **g**, Marked reduction in Tg versus wild-type colony-forming efficiency *in vitro* ($**P < 0.0001$, $n = 3$). In **f** and **g**, error bars (s.d.) are derived from eight or three experiments as indicated. **h**, Transduction of primary mouse keratinocytes with miR-203, but not empty vector or mutant miR-203M, results in a marked decline in S-phase cells that incorporate BrdU ($*P < 0.001$, $**P < 0.0001$). Values represent mean \pm s.d. from three experiments. Scale bars are 30 μ m.




Figure 3. Inhibition of miR-203 results in increased epidermal proliferation

a–d, P4 skins analysed after treatment of mice with mm-antagomir-203 (Ctrl) or antagomir-203 (anta). **a**, miR-203 *in situ* hybridizations. **b–d**, Immunofluorescence microscopy. Arrows denote basal-like features in suprabasal cells of antagomir-203-treated epidermis. PH3, phospho-histone H3. **e**, Elevated epidermal proliferation determined by BrdU injections at 4-h intervals and FACS quantification. **f**, The reduction in cycling cells that typically occurs upon calcium-induced differentiation of primary mouse keratinocytes *in vitro* is partially abrogated by anti-miR-203 oligonucleotides. Values represent mean \pm s.d. from three experiments (* $P < 0.005$, ** $P < 0.0001$). **g, h**, Epidermal defects that arise 30 days after tamoxifen-induced *Dicer1* ablation (iKO). Arrows denote suprabasal p63. Scale bars are 30 μ m.




Figure 4. MiR-203 targets p63 mRNA at 3'UTR

a, p63 protein is markedly diminished in Tg versus wild-type primary mouse keratinocytes. **b**, Immunoblot and qRT-PCR quantification of p63 expression. Top panel, calcium-induced (HiCa) p63 downregulation is largely relieved by anti-miR-203 oligonucleotides. Bottom panel, miR-203 transduction is sufficient to repress p63 protein expression in low calcium (LoCa). Right-hand graph, anti-miR-203 treatment slightly recovers *p63* mRNA whereas miR-203 transduction marginally reduces *p63* mRNA ($*P < 0.01$, $n = 3$). **c**, miR-203 transduction in LoCa primary mouse keratinocytes elevates *p21* mRNA level ($**P < 0.005$, $*P < 0.002$, $n = 3$). **d**, Insertion of two miR-203 target sequences within mouse *p63* 3'UTR leads to diminished luciferase (Luc) reporter activity in the presence of miR-203 ($*P < 0.002$, $n = 3$). pGL3 is a luciferase-expressing plasmid from Promega. **e**, Mutation of both target sites (p63M2) abolishes miR-203-mediated repression of luciferase activity; a single-site mutation (p63M) has marginal effects ($**P < 0.0001$, $n = 6$). In HiCa, wild-type but not mutant *p63* 3'UTR represses luciferase activity ($*P < 0.002$, $n = 3$). For all graphs, error bars (s.d.) are derived from the number of experiments as indicated.