



Published in final edited form as:

Nature. 2008 July 03; 454(7200): 104–108. doi:10.1038/nature06969.

A myocardial lineage derives from *Tbx18* epicardial cells

Chen-Leng Cai^{1,4,5,6,*}, Jody C. Martin^{2,*}, Yunfu Sun^{1,*}, Li Cui¹, Lianchun Wang⁷, Kunfu Ouyang³, Lei Yang¹, Lei Bu¹, Xingqun Liang³, Xiaoxue Zhang¹, William B. Stallcup⁸, Christopher P. Denton⁹, Andrew McCulloch², Ju Chen³, and Sylvia M. Evans^{1,3}

¹Skaggs School of Pharmacy, University of California, San Diego, La Jolla, California 92093, USA

²Department of Bioengineering, University of California, San Diego, La Jolla, California 92093, USA

³Department of Medicine, University of California, San Diego, La Jolla, California 92093, USA

⁴Department of Developmental and Regenerative Biology, Mount Sinai School of Medicine, 1 Gustave L. Levy Place, New York, New York 10029, USA

⁵Centre for Molecular Cardiology, Mount Sinai School of Medicine, 1 Gustave L. Levy Place, New York, New York 10029, USA

⁶The Black Family Stem Cell Institute, Mount Sinai School of Medicine, 1 Gustave L. Levy Place, New York, New York 10029, USA

⁷Complex Carbohydrate Research Centre, University of Georgia, Athens, Georgia 30602, USA

⁸Developmental Neurobiology Programme, The Burnham Institute, La Jolla, California 92037, USA

⁹Centre for Rheumatology, Royal Free and University College Medical School, Rowland Hill Street, London NW3 2PF, UK

Abstract

Understanding the origins and roles of cardiac progenitor cells is important for elucidating the pathogenesis of congenital and acquired heart diseases^{1,2}. Moreover, manipulation of cardiac myocyte progenitors has potential for cell-based repair strategies for various myocardial disorders³. Here we report the identification in mouse of a previously unknown cardiac myocyte lineage that derives from the proepicardial organ. These progenitor cells, which express the T-box transcription factor *Tbx18*, migrate onto the outer cardiac surface to form the epicardium, and then make a substantial contribution to myocytes in the ventricular septum and the atrial and ventricular walls. *Tbx18*-expressing cardiac progenitors also give rise to cardiac fibroblasts and coronary smooth muscle cells. The pluripotency of *Tbx18* proepicardial cells provides a theoretical framework for applying these progenitors to effect cardiac repair and regeneration.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to S.M.E. (syevans@ucsd.edu).

*These authors contributed equally to this work.

Full Methods and any associated references are available in the online version of the paper at www.nature.com/nature.

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Emergence of recent data has generated a paradigm shift for our understanding of cardiogenesis, with consequent implications for an understanding of cardiac progenitors and the aetiology of congenital heart disease. It was recognized that cardiac muscle cells derived from precardiac mesoderm subsequently form the primitive heart tube. More recently, the discovery of the secondary or anterior heart field, which contributed cells to the outflow tract and potentially to the right ventricle, suggested the presence of two distinct cardiac lineages⁴. Subsequent lineage studies based on expression of the LIM homeodomain transcription factor *Islet 1* (*Isl1*), and retrospective clonal analysis in the mouse, have confirmed two cardiac lineages, the first and second lineage, based on their timing of entry into the heart and the timing of their differentiation^{4–6}. Here we report a previously unknown myocardial lineage, derived from *Tbx18*-expressing epicardial cells, that makes a substantial contribution to the heart.

Previous studies in avian species have demonstrated that the proepicardium and/or epicardium is a source for coronary vascular progenitors and cardiac fibroblasts^{7–9}. During embryogenesis, cells from the proepicardium emigrate to form the epicardium—the epithelial outer lining of the heart. Epicardial cells undergo an epithelial-to-mesenchymal transition and invade the heart, giving rise to vascular endothelial cells, coronary vascular support cells and adventitial fibroblasts^{7–9}. Recently, regeneration of the zebrafish heart has been shown to be associated with re-activation of an early marker of the proepicardium/epicardium, the T-box transcription factor *Tbx18* (refs 10, 11). *Tbx18*-expressing cells appear to cluster at the wound site of zebrafish heart, with concurrent appearance of neo-vasculature during regeneration¹⁰.

To facilitate visualization of *Tbx18* expression, we generated an *nlacZ* (nuclear *lacZ*) knock-in into the endogenous *Tbx18* locus in mouse (Supplementary Fig. 3), and found that *nlacZ* expression mirrored that of the endogenous *Tbx18* gene (Fig. 1 and Supplementary Fig. 2). Neither *Tbx18* mRNA nor *Tbx18:nlacZ* expression are observed within the heart up to embryonic day (E)11.5 (Fig. 1).

To investigate epicardial lineages in the mouse, we also generated a Cre knock-in into the endogenous *Tbx18* locus (Supplementary Fig. 5), and crossed *Tbx18:Cre* mice with the lineage reporter R26R^{lacZ} (ref. 12) mice. Analysis of *lacZ* expression in embryos from this cross demonstrated early expression consistent with that of endogenous *Tbx18* (ref. 11; Figs 1 and 2 and Supplementary Figs 2 and 7a, b). In contrast to active expression of *Tbx18*, lineage analysis revealed the presence of *Tbx18*-derived lineages within the heart by E10.5 in the region of the forming ventricular septum and in scattered regions within both ventricular walls and atria (Fig. 2d–f and Supplementary Fig. 7b). Co-immunostaining with cardiac troponin T (*cTnT*, also known as *cTnnt*), cardiac troponin I (*cTnI*, also known as *cTnni*), MF20 (a sarcomeric myosin antibody) and the transcription factors *Gata4* and *Nkx2.5* demonstrated that these *Tbx18*-derived cells were cardiomyocytes (Fig. 2g–i and Supplementary Fig. 6). *Tbx18*-lineage-traced, *Nkx2.5*-positive cells could first be observed within the heart at E9.75 (Supplementary Fig. 6a). Complementary lineage studies using an organ explant culture system were consistent with *Tbx18:Cre/R26R^{lacZ}* lineage studies. Outer epicardial cells of embryonic hearts (E11–E13) were selectively labelled with 5-(and 6-)carboxyfluorescein diacetate succinimidyl ester (CFSE)¹³. Hearts harvested at time 0

exhibited specific labelling of the epicardium (Supplementary Fig. 7c). After culture for 18–24 h, fluorescently labelled epicardial cells were observed within the heart, and expressed *cTnT* (Supplementary Fig. 7d–f).

In contrast to earlier stages of heart development, cells actively expressing *Tbx18* are present within the heart at E12.5 (ref. 14). We examined sections from *Tbx18:nlacZ* mice from E10.5 to E17.5 and from neonatal stages using X-gal staining to detect β -galactosidase enzyme, encoded by the *lacZ* gene. (Fig. 1f, g and Supplementary Fig. 4d–f, data not shown for other stages), and by co-immunostaining with β -galactosidase antibody and cell-type-specific markers including *cTnT* and *Nkx2.5* (myocytes), *Pdgfrb* (vascular support cells¹⁵) and *Pecam1* (endothelial cells). Results demonstrated that cells actively expressing *Tbx18* within the heart at or after E12.5 were neither myocytes (*Nkx2.5*-negative, Supplementary Fig. 4a–c, data not shown for other stages) nor endothelial cells (*Pecam1*-negative, data not shown). Using mice containing a fibroblast-specific promoter from the *Col1a2* gene driving expression of a fusion protein in which cre recombinase is fused to a tamoxifen-inducible mutant oestrogen receptor (ER(T))¹⁶, and the *Tbx18*-floxed *nlacZ/nGFP* (nuclear green fluorescent protein) allele (Supplementary Fig. 3), we demonstrated that some *Tbx18*-expressing cells were fibroblasts (Supplementary Fig. 4g). Also, a population of *Tbx18:nlacZ* cells co-expressed *Pdgfrb* (Supplementary Fig. 4h panels 1–6), a marker of vascular support cells (both pericytes and vascular smooth muscle)¹⁵. *Tbx18*-expressing fibroblasts were not stained with the *Pdgfrb* antibody (Supplementary Fig. 4i, panels 1–6), demonstrating that both fibroblasts and vascular support cells actively express *Tbx18* within the heart. Active expression of *Tbx18* was not observed in cardio-myocytes, except within the sinus horns¹⁷.

Adult lineage analysis revealed that *Tbx18* derivatives were observed in the differentiated smooth muscle of the coronary vessels (Fig. 3a, b, c (insets 2 and 3), h, green or white arrows), including the coronary arteries, as evidenced by co-localization of β -galactosidase antibody with neuropilin-1 (*NP-1*, also known as *Nrp1*; Fig. 3i), a specific marker for arteries¹⁸. *Tbx18* expression was maintained in some coronary vascular smooth muscle cells (Fig. 3j, k). At adult stages, a substantial population of ventricular and atrial myocytes were observed to derive from *Tbx18*-expressing progenitors (Fig. 3c–f and Supplementary Fig. 8, black or white notched arrows). Cardiomyocytes isolated from *Tbx18:Cre/R26R^{lacZ}*-lineage-traced hearts exhibited X-gal-positive staining (Fig. 3g). Calcium transient analysis performed on myocytes isolated from *Tbx18:Cre/R26R^{EYFP}*-lineage-traced adult hearts demonstrated no differences between *Tbx18*-labelled myocytes and their non-labelled counterparts (Supplementary Fig. 9). *Tbx18* descendant cells also contributed significantly to the atrioventricular valves (Fig. 3c, inset 4). Co-localization of *Tbx18*-derived cells with endothelial lineages (as marked by *Pecam1*) was not observed in either embryonic or at adult stages (Fig. 3l).

Because previous studies have demonstrated that epicardial cells give rise to coronary endothelial cells^{7–9,19}, we verified immunohisto-chemical results by X-gal staining of affinity-purified endothelial cells isolated from *Tbx18*-lineage-traced hearts (Fig. 3m–o). Purified cardiac endothelial cell fractions did not contain *Tbx18*-lineage-positive cells (Fig. 3o, right panel). We then investigated whether the pro-epicardium contained endothelial

lineages distinct from *Tbx18*-expressing cells. Immunostaining demonstrated a population of *Flk-1* (also known as *Kdr*) cells within the proepicardium that did not express *Tbx18* (Fig. 4a). No *Nkx2.5*-positive myocardial progenitors were observed within the proepicardium (Fig. 4b). We also isolated fibroblasts from *Tbx18*-lineage-traced hearts. Consistent with our *Col1a2*-Cre-ER(T)¹⁶ results (Supplementary Fig. 4g), approximately one-third of cardiac fibroblasts expressed β -galactosidase (Fig. 3p).

Isl1-expressing second heart field progenitors contribute cardio-myocytes that ultimately reside in the outflow tract, the right ventricle, the ventricular septum, the left ventricle, the atria and the atrial septum^{5,20}. *Tbx18*-derived cardiomyocytes contributed significantly to the ventricular septum and atria as well as to small numbers of cells scattered throughout both ventricular walls (Figs 2d–i and 3c–g and Supplementary Figs 7a, b and 8). A comparison of *Tbx18*:Cre, *Isl1*:Cre- and *Isl1*:Cre/*Tbx18*:Cre-lineage-traced hearts indicated complementarity of these two lineages (Fig. 3c and Supplementary Figs 10 and 11a, b), as demonstrated by quantitative analysis of myocytes isolated from lineage-traced hearts (ventricular septum, left ventricle and right ventricle; Supplementary Table 1). Immunostaining and RNA *in situ* hybridization of *Tbx18*, *Isl1* and *MLC2a* (also known as *My17*) revealed that *Tbx18* was not co-expressed with *Isl1* and *MLC2a* during early embryogenesis (Supplementary Figs 2a–e and 11c).

We isolated *Tbx18*:Cre/R26R^{EYFP}-lineage-traced proepicardial cells and cultured them under conditions to favour differentiation into cardiomyocytes or smooth muscle cells²¹. Immunostaining analyses demonstrated efficient conversion of *Tbx18* lineages to myocytes or smooth muscle cells (Supplementary Fig. 12). To evaluate the pluripotency of individual proepicardial cells, single-cell clonal analysis was performed. Out of 336 single proepicardial cells plated on OP9 feeder layers, approximately 37% (124 out of 336) proliferated and formed clones by day seven. Forty clones were picked randomly, each dispersed into two wells, and cultured under conditions to favour either myocyte or smooth muscle cell fates. After five days of culture in differentiation medium, 34% of single-cell clonal derivatives differentiated into cardiomyocytes with an obvious striated cytoarchitecture and expressed *cTnT* (Fig. 4c). Spontaneous contraction was observed in some wells after four days of culture (Supplementary Fig. 13 and Supplementary Video), and myocyte identity was further confirmed by calcium transients (Fig. 4e–g). Each clone (40 out of 40) cultured with smooth muscle culture medium stained with smooth muscle myosin heavy chain (Fig. 4d). No instances were observed where derivatives of a single clone formed only cardiomyocytes and not smooth muscle cells. This demonstrated that a large proportion of proepicardial cells are pluripotent and can adopt either cardiomyocyte or smooth muscle cell fates.

Adult epicardial cells can be activated to migrate *in vitro* and adopt vascular cell fates²². Migratory adult epicardial cells from *Tbx18*:nlacZ mice expressed *Tbx18* (Supplementary Fig. 14). To investigate whether postnatal and adult epicardium retains a similar pluripotency to that of the proepicardium, postnatal and adult epicardial cells isolated from *Tbx18*:Cre/R26R^{EYFP}-lineage-traced mice were cultured on OP9 cells, and then subsequently cultured in myocyte- or smooth-muscle-specific differentiation medium. In contrast to results with proepicardial cells, adult epicardial cells did not convert to *cTnT*-expressing cells under

these culture conditions (data not shown). Understanding the underlying causes of this difference in potential will have significant implications for the possible use of epicardial cells for cardiac repair.

Our data are consistent with a model whereby the first cells to enter the heart from the proepicardium/epicardium give rise to myocyte lineages, first observed at approximately E9.75, whereas subsequent *Tbx18*-positive epicardial lineages that will give rise to vascular support cells and fibroblasts are first observed entering the heart at approximately E12.5. Endothelial lineages within the proepicardium are distinct from those of *Tbx18* cells (Supplementary Fig. 1). *Tbx18* is not actively expressed within cardiac myocyte derivatives, but is observed in a subset of vascular smooth muscle and cardiac fibroblasts, in addition to the epicardium (Fig. 3j, k and Supplementary Fig. 4g–i).

Previous studies in avian embryos have failed to demonstrate that proepicardial lineages give rise to cardiomyocytes within the heart, although myocytic potential has been demonstrated *in vitro*^{7–9,23}. It is unknown whether observed differences reflect distinct experimental approaches or species-specific differences. In mouse, *Tbx18*-expressing cells of the septum transversum that are in continuity with the sinus venosus contribute to the myocardial sleeve of the latter¹⁷. These cells, however, never express *Nkx2.5*, in contrast to the *Tbx18*-derived intracardiac myocytes of the ventricles and atria described here. The distinct embryonic origin of a substantial number of cardiomyocytes from an epicardial lineage provides a new perspective on heart development and congenital or adult heart disease affecting these lineages.

METHODS

Whole-mount RNA *in situ* hybridization and histological analysis

Whole-mount RNA ISH was carried out as described previously²⁴.

Generation of *Tbx18:nlacZ/nGFP*, *Tbx18:Cre* and *Isl1:Cre* knock-in mouse models²⁸

Tbx18 and *Isl1* genomic clones were isolated by screening a mouse genomic lambda library (129/sv, Stratagene). Arms of targeting constructs were amplified on phage DNA templates with high-fidelity DNA polymerase (Pfu, Stratagene, catalogue number 600153). For individual targeting strategy, see Supplementary Figs 3, 5 and 10.

X-gal staining

For the whole-mount staining, mouse embryos and tissues were collected from timed pregnant females. Embryos and tissues were fixed in 4% paraformaldehyde for 15–60 min. After permeabilization (10% Na-deoxycholate, 10% NP40 in PBS), embryos and tissues were stained in X-gal solution (50 mM K-ferricyanide, 50 mM K-ferrocyanide, 200 mM MgCl₂, 100 mg ml⁻¹ X-gal in PBS) for 4–12 h and were then post-fixed with 4% paraformaldehyde. For section staining, embryos and tissues were fixed in 4% paraformaldehyde and were then dehydrated in series sucrose solution. X-gal staining was performed on 6–8- μ m cryosections with previous additional fixation in 4% paraformaldehyde for 6–8 min.

Fluorescent dye lineage tracing of epicardial cells with a heart explant culture system

Embryos were dissected at E11.5 in PBS. About 20 μ l dye solution (30 μ g ml⁻¹) was injected into the pericardial cavity (CFSE, Molecular Probes, C1157). The chest expands under the pressure and it was ensured that it did not shrink back, indicating the dye was not able to leak from the cavity. Embryos were then submerged in media (DMEM) and left at room temperature (23 °C) for 15 min in the dark. Hearts were removed and washed in media for 5 min. Some hearts were selected randomly for fixation in 4% paraformaldehyde. The rest were cultured on a 0.4 μ m polycarbonate membrane insert with 10% embryonic stem cell qualified fetal bovine serum (FBS) (Gibco, 10439024) supplemented DMEM (Gibco, D5796) for 18–24 h. Hearts were fixed and frozen as described previously for immunohistochemistry.

Isolation of cardiomyocytes, endothelial cells, fibroblasts and epicardial cells from adult mouse hearts

For cardiomyocyte isolation, pure adult cardiomyocytes (6–8 weeks) were isolated according to a previously described method²⁹. For Ca²⁺ imaging and *Tbx18/Is11* lineage contribution quantification, different parts of the heart (ventricular septum, left ventricular wall and right ventricular wall) were dissected and separated after retrograde collagenase perfusion via the aorta. After an additional collagenase incubation for 10 min at 37 °C, cardiomyocytes were dispersed mechanically²⁹. Pure populations of myocytes were obtained by centrifugation and cell filtration. Quantification was performed on a number of independent hearts for each sample by cell counts of numerous, random, field-of-view micrographs to determine the percentage of total cardiomyocytes expressing the YFP lineage marker. Statistical analyses were performed on the data sets. For isolation of endothelial cells, adult mouse hearts were minced with a razor blade. Endothelial cells from the hearts were isolated according to a previously described method²⁶. Cardiac fibroblasts were isolated and cultured as described previously²⁵. Pre-plating of cells isolated from the heart digest was shortened to 30 min at 37 °C to increase the purity of fibroblasts. For epicardial cell isolation, adult epicardial cells were isolated from explanted hearts by treatment with collagenase at 37 °C for 10 min (prepared as described for cardiomyocyte isolation). The epicardium was then manually extracted as a monolayer sheet.

Immunostaining

Embryos and tissues were fixed immediately in 4% paraformaldehyde for 10–30 min after dissection. Tissues were embedded and cut by cryosectioning (5–10 μ m). For sections with endogenous YFP, a post-fixation for 5 min on ice was performed before staining. Cells obtained in culture were washed with warm media and then were fixed for 7 min with 10% formalin. Primary antibodies used in this study were: rabbit polyclonal anti- β -galactosidase (Cappel, product number 55978, 1:200), goat polyclonal anti- β -galactosidase (Biogenesis, 4600–1409, 1:200), rabbit polyclonal anti-smooth muscle myosin heavy chain (Biomedical Technologies Inc., BT562, 1:200), rabbit polyclonal anti-NKX2.5 (Santa Cruz Biotechnology, SC14033, 1:50), rat polyclonal anti-Pecam1 (Pharmingen, 550274, 1:100), mouse monoclonal anti-cardiac troponin T (NeoMarkers, MSZ-295-P, 1:200), mouse monoclonal anti- α -smooth muscle actin (Abcam, ab7817, 1:200), mouse monoclonal anti-

α -actinin (sarcomeric) (Sigma-Aldrich, A7811, 1:200), rabbit polyclonal anti-WT1 (Santa Cruz Biotechnology, sc-846, 1:75), goat polyclonal anti-GATA-4 (Santa Cruz Biotechnology, sc-1237, 1:75), mouse polyclonal anti-Isl1 (Developmental Studies Hybridoma Bank, 39.4D5, 1:100). Mouse monoclonal cardiac troponin I was provided by J. Lin (Developmental Studies Hybridoma Bank, 1:100), mouse monoclonal anti-MF20 (1:200) was provided by D. Fischman, and rabbit polyclonal anti-Pdgfrb (1:200) was provided by W. Stallcup. Rabbit polyclonal anti-neuropilin 1 (1:200) was provided by A. L. Kolodkin.

Isolation and differentiation of proepicardial cells

Tbx18-lineage-traced embryos (Tbx18:Cre/R26R^{EYFP}, E9.0–E9.5) were dissected from timed pregnant mice and were quickly washed in cold PBS after dissection. A check was made under the epifluorescent microscope that *Tbx18*-positive cells in the proepicardium have not begun to migrate to the heart. Proepicardial cells were isolated with a pulled-tip glass pipette. The isolated proepicardial cells were cultured on gelatin with a medium designed to induce differentiation into cardiac myocytes or smooth muscle cells²¹.

Single proepicardial cells preparation

Proepicardial cells collected from seven to ten E9.0–E9.5 embryos (Tbx18:Cre/R26R^{EYFP}) were digested with 0.05% trypsin/EDTA (Invitrogen) for 5 min at 37 °C. Single-cell suspension was obtained by pipetting cells in culture medium (85% IMDM (Gibco), 15% FBS (selected batches), 100 μ M non-essential amino acids (Gibco), 2 mM sodium pyruvate (Gibco) and 100 μ M β -mercaptoethanol (Sigma)). Disassociated cells were plated at 0.5 cells per well onto a 96-well plate with a mitomycin C (Sigma)-inactivated OP9 cell feeder layer (YFP signals were examined to assure single-cell colonies derived from Tbx18 lineages). After seven days of culture, clones proliferated from single cells were dispersed to small clumps with 0.05% trypsin/ EDTA for 30 s. These cells were re-plated onto gelatin-coated 16-well chamber slides with one colony split into two wells and then left in stem cell culture medium for two days. Cells were cultured with a medium designed to induce differentiation into cardiac myocytes or smooth muscle cells²¹. Immunostaining was performed after five days of culture.

Ca²⁺ imaging

Cells were loaded with Rhod-2-AM (5 μ M, 15 min; Molecular Probes, R1245MP), and imaged with an Olympus Fluoview 1000 inverted confocal microscope with a $\times 40$ oil immersion lens (numerical aperture 1.3). The line-scan imaging mode was used to measure Ca²⁺ transients of paced cells. The extracellular solution contained 116 mM NaCl, 5.0 mM KCl, 0.8 mM MgSO₄, 1.0 mM NaH₂PO₄, 5.5 mM glucose, 20 mM HEPES and 1.0 mM CaCl₂ (pH 7.4). Image processing and data analysis were performed as previously described²⁷.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank A. Kleckner, J. Lam and M. Zamora for critical technical assistance; D. Bader for discussion; P. Soriano and F. Costantini for providing R26R^{lacZ} and R26R^{EYFP} indicator mice; H. Kubo for providing OP9 cells; and J. Lin, D. Fischman and A. Kolodkin for providing troponin I, Pdgfrb and neuropilin 1 antibodies, respectively. We are also grateful to B. Gelb for his critical comments and revisions of the manuscript. This work was supported by AHA National Scientist Development Grant to C.-L.C. and NIH1RO1 to S.M.E.

References

1. Olson EN. A decade of discoveries in cardiac biology. *Nature Med.* 2004; 10:467–474. [PubMed: 15122248]
2. Srivastava D. Making or breaking the heart: from lineage determination to morphogenesis. *Cell.* 2006; 126:1037–1048. [PubMed: 16990131]
3. Murry CE, Field LJ, Menasche P. Cell-based cardiac repair: reflections at the 10-year point. *Circulation.* 2005; 112:3174–3183. [PubMed: 16286608]
4. Buckingham M, Meilhac S, Zaffran S. Building the mammalian heart from two sources of myocardial cells. *Nature Rev Genet.* 2005; 6:826–835. [PubMed: 16304598]
5. Cai CL, et al. Isl1 identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart. *Dev Cell.* 2003; 5:877–889. [PubMed: 14667410]
6. Kelly, R., Evans, SM. The secondary/anterior heart field. In: Rosenthal, N., Harvey, RP., editors. *Heart Development and Regeneration.* Academic; San Diego: in the press
7. Mikawa T, Fischman DA. Retroviral analysis of cardiac morphogenesis: discontinuous formation of coronary vessels. *Proc Natl Acad Sci USA.* 1992; 89:9504–9508. [PubMed: 1409660]
8. Mikawa T, Gourdie RG. Pericardial mesoderm generates a population of coronary smooth muscle cells migrating into the heart along with in growth of the epicardial organ. *Dev Biol.* 1996; 174:221–232. [PubMed: 8631495]
9. Gittenberger-de Groot AC, Vrancken Peeters MP, Mentink MM, Gourdie RG, Poelmann RE. Epicardium-derived cells contribute a novel population to the myocardial wall and the atrioventricular cushions. *Circ Res.* 1998; 82:1043–1052. [PubMed: 9622157]
10. Lepilina A, et al. A dynamic epicardial injury response supports progenitor cell activity during zebrafish heart regeneration. *Cell.* 2006; 127:607–619. [PubMed: 17081981]
11. Kraus F, Haenig B, Kispert A. Cloning and expression analysis of the mouse T-box gene *Tbx18*. *Mech Dev.* 2001; 100:83–86. [PubMed: 11118889]
12. Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nature Genet.* 1999; 21:70–71. [PubMed: 9916792]
13. Wilm B, Ipenberg A, Hastie ND, Burch JB, Bader DM. The serosal mesothelium is a major source of smooth muscle cells of the gut vasculature. *Development.* 2005; 132:5317–5328. [PubMed: 16284122]
14. Franco D, et al. Left and right ventricular contributions to the formation of the interventricular septum in the mouse heart. *Dev Biol.* 2006; 294:366–375. [PubMed: 16677630]
15. Armulik A, Abramsson A, Betsholtz C. Endothelial/pericyte interactions. *Circ Res.* 2005; 97:512–523. [PubMed: 16166562]
16. Zheng B, Zhang Z, Black CM, de Crombrughe B, Denton CP. Ligand-dependent genetic recombination in fibroblasts: a potentially powerful technique for investigating gene function in fibrosis. *Am J Pathol.* 2002; 160:1609–1617. [PubMed: 12000713]
17. Christoffels VM, et al. Formation of the venous pole of the heart from an Nkx2–5-negative precursor population requires Tbx18. *Circ Res.* 2006; 98:1555–1563. [PubMed: 16709898]
18. You LR, et al. Suppression of Notch signalling by the COUP-TFII transcription factor regulates vein identity. *Nature.* 2005; 435:98–104. [PubMed: 15875024]
19. Perez-Pomares JM, et al. Origin of coronary endothelial cells from epicardial mesothelium in avian embryos. *Int J Dev Biol.* 2002; 46:1005–1013. [PubMed: 12533024]

20. Verzi MP, McCulley DJ, De Val S, Dodou E, Black BL. The right ventricle, outflow tract, and ventricular septum comprise a restricted expression domain within the secondary/anterior heart field. *Dev Biol.* 2005; 287:134–145. [PubMed: 16188249]
21. Moretti A, et al. Multipotent embryonic Isl1⁺ progenitor cells lead to cardiac, smooth muscle, and endothelial cell diversification. *Cell.* 2006; 127:1151–1165. [PubMed: 17123592]
22. Smart N, et al. Thymosin β 4 induces adult epicardial progenitor mobilization and neovascularization. *Nature.* 2007; 445:177–182. [PubMed: 17108969]
23. Kruithof BP, et al. BMP and FGF regulate the differentiation of multipotential pericardial mesoderm into the myocardial or epicardial lineage. *Dev Biol.* 2006; 295:507–522. [PubMed: 16753139]
24. Cai CL, et al. T-box genes coordinate regional rates of proliferation and regional specification during cardiogenesis. *Development.* 2005; 132:2475–2487. [PubMed: 15843407]
25. Subramanian SV, Kelm RJ, Polikandriotis JA, Orosz CG, Strauch AR. Reprogramming of vascular smooth muscle α -actin gene expression as an early indicator of dysfunctional remodeling following heart transplant. *Cardiovasc Res.* 2002; 54:539–548. [PubMed: 12031699]
26. Wang L, Fuster M, Sriramaraio P, Esko JD. Endothelial heparan sulfate deficiency impairs L-selectin- and chemokine-mediated neutrophil trafficking during inflammatory responses. *Nature Immunol.* 2005; 6:902–910. [PubMed: 16056228]
27. Ouyang K, et al. Ca²⁺ sparks and secretion in dorsal root ganglion neurons. *Proc Natl Acad Sci USA.* 2005; 102:12259–12264. [PubMed: 16103366]
28. Hadjantonakis AK, Dickinson ME, Fraser SE, Papaioannou VE. Technicolour transgenics: imaging tools for functional genomics in the mouse. *Nature Rev Genet.* 2003; 4:613–625. [PubMed: 12897773]
29. Zhou YY, et al. Culture and adenoviral infection of adult mouse cardiac myocytes: methods for cellular genetic physiology. *Am J Physiol Heart Circ Physiol.* 2000; 279:H429–H436. [PubMed: 10899083]




Figure 1. LacZ expression in Tbx18:nlacZ knock-in mice recapitulates endogenous *Tbx18* expression

a–d, Section RNA *in situ* hybridization (ISH) of *Tbx18* in mouse embryos (E9.0–E11.5). *Tbx18* is expressed very early in the proepicardium (**a**) and epicardium (**b–d**). For parallel whole-mount RNA ISH, see Supplementary Fig. 2. **e–g**, X-gal staining on cardiac sections from Tbx18:nlacZ embryos at E9.5–E11.5. Tbx18:nlacZ cells are detected in the early epicardium at E9.5 (**e**) and in all epicardial cells covering the heart after E10.5 (**f**, **g**, data not shown for E12.5–E13.5). X-gal staining on Tbx18:nlacZ mouse tissues is consistent with *Tbx18* mRNA ISH. For the Tbx18:nlacZ targeting strategy, see Supplementary Fig. 3. Black arrows indicate *Tbx18* mRNA expression (**a–d**) and Tbx18:nlacZ cells (**e–g**); red arrows in **c** and **d** indicate that *Tbx18* is not expressed in heart between E9.5 and E11.5. For **e–g**, lower panels are high-magnification views of the upper panels in the heart (black arrow regions). LA/RA, left/right atrium; LV/RV, left/right ventricle; OT, outflow tract; and SV, sinus venosus.




Figure 2. Cells derived from *Tbx18*-expressing cells are observed within the heart by E10.5, and exhibit a cardiomyocyte identity

Tbx18:Cre mice were crossed to R26R^{lacZ} indicator mice¹² (*Tbx18*:Cre/R26R^{lacZ}). **a–c**, X-gal staining on whole-mount embryos (E9.5–E11.5) shows *Tbx18*-lineage-traced cells in the proepicardium and epicardium (arrows in **a–c**). **d–f**, X-gal staining on cryosections from E10.5 to E12.5. *Tbx18*-lineage-traced cells are observed in the epicardium (arrows in the upper panels of **d–f**) and within the heart, particularly within the apical region in the developing ventricular septum (arrows in the lower panels of **d–f**). The lower panels of **d–f** show high-magnification views of the ventricles in the upper panel micrographs. For *Tbx18*-lineage-traced cells at E13.5, see Supplementary Fig. 7. **g**, β -galactosidase antibody staining on *Tbx18*-lineage-traced tissue at E11.5. **h**, Cardiac troponin T antibody staining. **i**, Overlay of **g** and **h** revealed that *Tbx18*-lineage-traced cells in heart were cardiac-troponin-T-positive cells. In **g–i**, lower panels are high-magnification views of the upper panels in the ventricular septum region. Arrows in **d–i** indicate *Tbx18*-lineage-traced cells in the ventricular septum region. LA/RA, left/right atrium; LV/ RV, left/right ventricle; and VS, ventricular septum.




Figure 3. *Tbx18* lineage tracing in the adult heart

a, b, Whole-mount X-gal staining of *Tbx18:Cre/R26R^{lacZ}* adult mouse heart (6 weeks). Coronary vasculature (green arrows) is derived from *Tbx18* lineages. Dense staining in the septum is visible (black arrow). **c,** X-gal staining of tissues from *Tbx18:Cre/R26R^{lacZ}* adult mouse heart (6 weeks). *Tbx18* can give rise to: cardiomyocytes within atria (arrows in **c**, inset 1 and Supplementary Fig. 8), the ventricular septum (black arrow in **c**, inset 2) and the ventricular wall (black arrow in **c**, inset 3); coronary vascular support cells (green arrows in **c**, insets 2 and 3); and atrioventricular valves (**c**, inset 4, high magnification for bicuspid valve). Most *Tbx18*-lineage-traced cells within the ventricular septum co-stained with cardiac troponin T (arrows in **d-f**; **f** is a high magnification of the lower panel of **e**), demonstrating they are cardiomyocytes. **g,** A subset of cardiomyocytes isolated from adult *Tbx18*-lineage-traced hearts exhibited X-gal-positive staining. **h,** *Tbx18* lineages give rise to coronary vascular smooth muscle cells (co-localized with smooth muscle myosin heavy chain). **i,** *Tbx18* lineages give rise to coronary artery smooth muscle cells (co-localized with *NP-1*, an artery marker). **j, k,** X-gal staining on *Tbx18:nLacZ* knock-in mice shows that *Tbx18* expression is maintained in some coronary vascular smooth muscle cells from embryonic stages to adulthood. **l-o,** *Tbx18* lineages do not give rise to coronary vascular endothelial cells (not co-localized with *Pecam1*, **l**). **m,** Coronary vascular endothelial cells were isolated from the hearts (green fraction)²⁶ and react with *Pecam1* antibody (**n**, right panel), whereas *Pecam1*⁻ fraction (purple) cells do not (**n**, left panel). **o,** Purified *Pecam1*⁻ cells are Xgal⁻ (**o**, right panel), confirming that *Tbx18*-expressing lineages do not give rise to coronary vascular endothelial cells. A portion of cells in *Pecam1*⁻ fraction are Xgal⁺ (**o**,

left panel). **p**, Cardiac fibroblasts isolated from *Tbx18*-lineage-traced hearts²⁵ demonstrated that approximately 30% of these cells derive from *Tbx18* lineages. Abbreviations are as in Fig. 2.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. *Tbx18*-expressing progenitors within proepicardium are distinct, and retain the capacity to differentiate into cardiomyocytes and smooth muscle cells *in vitro*

a, b, *Tbx18*-expressing progenitors within the proepicardium are not co-stained with antibodies to *Flk-1* (**a**) or *Nkx2.5* (**b**) (E9.0, data not shown for E9.5), suggesting that *Flk-1* endothelial progenitors within the proepicardium are a distinct population from *Tbx18*-expressing cells. *Tbx18* progenitors with myocyte potential do not yet express *Nkx2.5* within the proepicardium, although *Tbx18*-derived myocytes do express *Nkx2.5* within the heart. The second panels in **a** and **b** are high-magnification images of the top panels of **a** and **b**, respectively. **c,** *Tbx18*-lineage-traced proepicardial cells (*Tbx18:Cre/R26R^{EYFP}*) were disassociated and single cells were expanded in proepicardial culture medium. A significant percentage of clones (34%) derived from single proepicardial cells adopt cardiomyocyte fate after incubating in the differentiation medium²¹ (co-stained with cardiac troponin T, bottom panel of **c**). **d,** Every clone (40 out of 40) derived from single *Tbx18* proepicardial cells displayed staining for smooth muscle myosin heavy chain (SM-MHC) antibody after incubating in smooth muscle culture medium. **e–g,** Real-time Ca^{2+} transients of cells derived from single *Tbx18*-lineage-traced proepicardial cells after myocytic differentiation. **e,** Left panel: monitored cell is YFP⁺, suggesting it derived from the *Tbx18:Cre/R26R^{EYFP}*

proepicardium. **e**, Middle panel: differentiated cells were monitored under an inverted confocal microscope after they were preloaded with Ca^{2+} indicator Rhod-2-AM. **e**, Right panel, transmitted light image of the monitored cell. **f**, Line-scanning images and time course (**g**) of spontaneous Ca^{2+} transients obtained from differentiated YFP^{+} cells. PE, proepicardium; SV, sinus venosus; and V, ventricle.