

Title	Genome-scale analyses of health-promoting bacteria: probiogenomics
Authors	Ventura, Marco;O'Flaherty, Sarah;Claesson, Marcus J.;Turroni, Francesca;Klaenhammer, Todd R.;van Sinderen, Douwe;O'Toole, Paul W.
Publication date	2009-01
Original Citation	Ventura M., O'Flaherty S., Claesson M. J., Turroni F., Klaenhammer T.R., van Sinderen D., O'Toole P.W. 2009. Genome-scale analyses of health-promoting bacteria: probiogenomics. Nature Reviews Microbiology 7, pp61-71 doi:10.1038/nrmicro2047
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1038/nrmicro2047
Rights	© 2009 Macmillan Publishers Limited. All rights reserved - http://www.nature.com/authors/editorial_policies/license.html
Download date	2024-05-19 12:28:33
Item downloaded from	https://hdl.handle.net/10468/118



Ventura M., O'Flaherty S., Claesson M. J., Turróni F.,
Klaenhammer T.R., van Sinderen D., O'Toole P. W. 2009.
Genome-scale analyses of health-promoting bacteria:
probiogenomics. *Nature Reviews Microbiology* 7, pp61-71
doi:10.1038/nrmicro2047

Deposited on: 16 Feb 2010

CORA Cork Open Research Archive <http://cora.ucc.ie>

Genome-scale analyses of health-promoting bacteria: probiogenomics

Key words: genomics, functional genomics, probiotic bacteria, intestinal tract, microbiota

Marco Ventura^{*}, Sarah O'Flaherty[‡], Marcus J. Claesson[§], Francesca Turrone^{*}, Todd R. Klaenhammer[‡], Douwe van Sinderen[§], and Paul W. O'Toole[§]

^{*}Department of Genetics, Biology of Microorganisms, Anthropology and Evolution, University of Parma, Italy, [‡]Department of Food, Bioprocessing and Nutrition Sciences, North Carolina State University, Raleigh, North Carolina 27695, [§]Alimentary Pharmabiotic Centre and Department of Microbiology, University College Cork, Western Road, Cork, Ireland

Correspondence to P.W.O
e.mail: pwotoole@ucc.ie

27 **Abstract**

28 The human body is colonized by an enormous population of bacteria (microbiota) that outnumber
29 the human somatic and germ cells and provides the host with additional coding capacity and
30 metabolic activities. Among the human gut microbiota are health-promoting indigenous species,
31 also referred to as probiotic bacteria, which are commonly consumed as live dietary supplements.
32 Although there is a growing list of health benefits provided by the consumption of probiotics, their
33 precise mechanisms of action remain largely unknown. Recent genomics based studies
34 (probiogenomics) are starting to provide insights into the ways probiotic bacteria sense and adapt to
35 the gastrointestinal tract environment. In this review, we will discuss the application of
36 probiogenomics in the elucidation of the molecular basis of probiosis using the well recognized
37 model probiotic bacteria *Bifidobacterium* and *Lactobacillus* as examples.

38

39 The availability of the sequence of the human genome has paved the way for a better understanding
40 of the genetic basis for many aspects of human health and disease. However, fully understanding
41 the human genotype, and its relationship with health and disease susceptibility, requires better
42 information explaining how environmental and developmental factors interact with the genome to
43 influence health status. Human beings are colonized by, or transiently harbour, a wide, complex and
44 dynamic collection of bacteria that outnumber the human somatic and germ cells, and that
45 collectively represent significantly more genetic variety than the genome of their host¹. However, at
46 the present time, the components of the human microbiota remain poorly identified and
47 characterized. Recent culture-independent studies of the microbiota of the human gastrointestinal
48 tract (GIT) have identified more than 1000 phylotypes, representing over 7000 strains and
49 belonging to eight major phyla¹⁻⁴ (see also⁵ for an overview).

50 It has been suggested that the composition of the gut microbiota is the result of selective
51 pressure imposed by the host, and further modulated by competition between constituent bacterial
52 members⁶. The interactions between various bacteria and the human host can be categorized as a
53 continuum ranging from symbiosis to commensalism and through to pathogenesis, where the two
54 former relationships can be grouped as mutualism (Fig. 1). In the human gut environment, the
55 adaptive co-evolution of humans and bacteria may lead to the development of commensal
56 relationships, where neither partner is disadvantaged, or symbiotic relationships where unique
57 metabolic activities or other benefits are provided. The intestinal microbiota contributes to host
58 nutrition^{1, 7, 8} and it impacts on intestinal cell proliferation and differentiation, pH, the development
59 of the immune system and innate and acquired response to pathogens^{1, 9, 10}.

60 Alterations in the composition of the intestinal microbiota have recently been linked to a
61 variety of conditions ranging from Inflammatory Bowel Disease to allergy and obesity^{6, 11-14}.
62 Among the variable constituents of the microbiota are health-promoting indigenous species (or
63 autochthonous microbiota), also known as probiotic bacteria, which are commonly consumed as

live dietary supplements¹⁵. The mechanisms by which probiotic micro-organisms beneficially affect human health (reviewed in^{16, 17}) are typically divided into a number of general categories, including strengthening of the intestinal barrier, modulation of the immune response and antagonism of pathogens either by the production of antimicrobial compounds or through competition for mucosal binding sites^{16, 18}. Although there is suggestive evidence for each of these functional claims, the molecular mechanisms remain largely unknown.

Genomics offers the possibility of accelerating research into probiotic bacteria. In recent years, genome sequencing of gut commensals and symbionts has come to the fore, currently represented by the development of a novel scientific discipline, called probiogenomics¹⁹, which aims to provide insights into the diversity and evolution of commensal/probiotic bacteria and to reveal the molecular basis for their health-promoting activities. The integration of probiogenomics and functional genomic information with data on host gene expression in the human gut will expand our understanding of the roles of (probiotic) microbiota, microbe-microbe and host-microbe interactions. These “omics” approaches allow the simultaneous analysis of very large numbers of genes or proteins²⁰. Probiogenomics is thus one strand of gut systems microbiology. Significantly, when studied in combination with host genome variation, probiogenomics offers a comprehensive systems model, even at individual subject level.

Here we address current developments in analyzing the genome sequences of probiotic bacteria and how these data can be integrated in a global view using omics approaches in order to elucidate genome evolution and genetic adaptation of these bacteria to the human gut ecological niche. We consider the well recognized model probiotic bacteria *Bifidobacterium* spp. and *Lactobacillus* spp. which are phylogenetically distant (although well-characterized; Fig 1), have distinguishing properties, and different depths of biological characterization.

90 **Bifidobacteria genomics**

91 The genus *Bifidobacterium* is relatively small, with 30 species, and a low level of phylogenetic and
92 genomic diversity²¹. Bifidobacteria were originally isolated from a breast-fed infant²³ and since
93 then, 30 species have been isolated from the GIT contents of mammals, birds and insects¹⁹. Those
94 bifidobacteria that may be isolated from the human intestine have attracted the interest of genomic
95 research due to their probiotic properties. However, of the bifidobacterial taxa described to date,
96 genomes of only three strains, which belong to the *B. longum* and *B. adolescentis* groups, have been
97 sequenced to completion (Table 1). The availability of genome sequences provided a genetic basis
98 for the observation that bifidobacteria are extensively prototrophic, indicating that these bacteria are
99 well adapted to grow in an environment such as the human colon, which is poor in certain growth
100 substrates (e.g. vitamins, amino acids and nucleotides)²⁴. In fact, bifidobacterial genome sequences
101 available to date revealed that these organisms harbour genes for the synthesis of at least 19 amino
102 acids and they encode all enzymes needed for the biosynthesis of pyrimidine and purine
103 nucleotides, as well as those required for the synthesis of the B vitamins, folic acid, thiamine and
104 nicotinate (²⁵; Ventura et al., unpublished data; Leahy and D. van Sinderen, unpublished data).
105 Annotation and pathway prediction revealed the presence of all the required genetic information to
106 shunt many monosaccharides or disaccharides into the fructose-6-phosphate pathway²⁴.

107 *Adaptation to the human gut.*

108 The amount and types of “non-digestible” saccharides in the diet (some of which are referred to as
109 prebiotics) has a major influence on the numbers and metabolic activities of different groups of
110 bacteria within the enteric microbiota²⁶. The range of polysaccharide substrates that arrive in the
111 intestine is extremely broad²⁷. This diversity of carbon substrates potentially generates a vast array
112 of ecological roles and niches that may be exploited by gut bacteria. Although some members of the
113 gut microbiota can switch rapidly between different substrates (e.g. derived from diet or of host
114 origin), others (e.g. those associated with insoluble substrates) are much more specialized²⁸. In this
115 context, bifidobacteria have a presumed ecological advantage due to their capacity to metabolize

116 complex sugars derived from the diet as well as from the host²⁹. Genome annotation confirms that
117 genes required for the breakdown of complex sugars are abundant in sequenced bifidobacterial
118 genomes¹⁹. Over 8% of annotated bifidobacterial genes encode enzymes involved in carbohydrate
119 metabolism. These include various glycosyl hydrolases (GH) for utilization of diverse, but in most
120 cases un-identified, plant-derived dietary fibers or complex carbohydrate structures. Most of the
121 bifidobacterial GHs are predicted to be intracellular including those that are thought to hydrolyze
122 arabinogalactans and arabinoxylans, or starch and related polysaccharides^{25, 30, 31}. The genes for
123 these GHs are associated with genetic loci for the uptake of structurally diverse sugar substrates. In
124 fact, about 5% of the total bifidobacterial gene content is dedicated to sugar internalization, through
125 ABC transporters, permeases, and proton symporters rather than phosphoenolpyruvate-
126 phosphotransferase systems (PEP-PTSs)^{25, 32, 33}. Bifidobacteria utilize a kind of docking station to
127 sequester and capture high molecular weight carbohydrates molecules (e.g., xylose- and arabinose-
128 containing polysaccharides; Fig. 2) and bind these to their cell surface^{30, 33}, presumably to avoid
129 losing them to nearby competitors. This is reminiscent of a putative carbohydrate utilization system
130 identified in the genome of *L. plantarum*³⁴, and a system used by *Bacteriodes thetaiotaomicron* for
131 starch utilization³⁵. Enteric bifidobacteria are also able to utilize sialic acid-containing complex
132 carbohydrates in mucin, glycosphingolipids and human milk^{36, 37}. Thus, these bifidobacteria have
133 acquired adaptations to allow them to exploit a rich repertoire of otherwise indigestible components
134 of the human or animal diet.

135 Characterization of the metabolism of prebiotic compounds by bifidobacteria has identified specific
136 transporters and hydrolases for oligosaccharides^{30, 38, 39}. These studies indicated that bifidobacteria
137 ferment different types of fructo-oligosaccharides (FOS); accordingly, the respective FOS
138 metabolism operons possess different genetic architectures⁴⁰, suggesting that these genes were
139 acquired following evolutionary divergence of the species. Prebiotic oligosaccharides are also
140 contained in human milk (e.g., galacto-oligosaccharides), which are hydrolyzed by bifidobacteria
141 through the action of extracellular enzymes encoded by the *galA* gene^{30, 41}. In addition to galacto-

oligosaccharides, human milk consumption provides large amounts of small peptides that are derived from the digestion of milk proteins by the gastric protease pepsin⁴². *Bifidobacterium* genomes encode a rich repertoire of enzymes involved in the breakdown and internalization of peptides such as dipeptidyl aminopeptidases and oligopeptide uptake systems (Ventura et al., unpublished data).

Molecular interaction with the host.

Bacterium-host interactions that result in host benefit can be elucidated by identification and detailed molecular analysis of the bacterial proteins or macromolecules involved. For example a potential probiotic effector molecule, a eukaryotic-type serine protease inhibitor (serpin) was identified in the genome of *B. longum* subsp. *longum*^{25, 43}. Members of the serpin family regulate a wide range of signalling pathways in eukaryotes and some are recognized for their ability to suppress inflammatory responses by inhibiting elastase activity⁴⁴. Recent findings showed that the bifidobacterial serpin-like protein performs an immunomodulatory role in a murine colitis model, by reducing intestinal inflammation⁴³.

Transcriptomic approaches facilitate studies of gene expression profiles and have been successful in studying how individual organisms in bacterial communities affect each other's transcriptome. Recent transcriptomic analyses were performed on bacteria from germ-free mice that had been mono-associated with *B. thetaiotaomicron* —one of the dominant components of the human gut microbiota — and subsequently challenged with *B. longum* subsp. *longum*. The presence of *B. longum* subsp. *longum* provoked an expansion in the diversity of polysaccharides targeted for breakdown by *B. thetaiotaomicron* such as mannose and xylose-containing glycans⁴⁵. The changes in the transcriptional profiles of polysaccharide-utilization related genes by *B. longum* subsp. *longum* and *B. thetaiotaomicron* may imply the existence of symbiosis between these microbial species, where each species possesses a complement of GH activities, which when combined allow both to participate in a synergic harvest of xylose and mannose-containing sugars. This phenomenon has already been described in other microbial communities that degrade

168 cellulose⁴⁶. Alternatively, the shifts in transcription patterns could represent response to competition
169 (see also below for lactobacilli).

170 The elucidation of the molecular impact generated by members of the human microbiota on
171 the human host was also analysed by studying the host epithelium response to co-colonization by *B.*
172 *longum* subsp. *longum* and *B. thetaiotaomicron*⁴⁵. Remarkably, the host response to these two
173 bacterial species was different. In fact, the host response to *B. thetaiotaomicron* was more focused
174 on tumor necrosis factor α and LPS-responsive cytokine produced by natural killer and T
175 macrophages, whereas *B. longum* subsp. *longum* promoted the activation of T-cell-produced
176 cytokine interferon- γ and reduced production by the host of antibacterial proteins such as Reg3 γ
177 (Regenerating islet-derived-3 γ) and Pap (Pancreatitis-associated protein). Thus the host response to
178 enteric bifidobacteria may not only promote their own survival in the human intestine but also affect
179 the composition of the overall human gut microbiota.

180

181 **Comparative genomics of bifidobacteria**

182 Comparisons at the nucleotide level of the fully sequenced bifidobacterial genomes revealed a high
183 degree of conservation and synteny across the entire genomes¹⁹. However, several breakpoint
184 regions were also reported, apparently representing inversions or DNA deletion/insertion points.
185 DNA regions uniquely present in one genome and absent in others were also identified. Most of
186 these correspond to genetic elements presumably acquired by horizontal gene transfer events
187 (HGT), including prophage-like elements, restriction modification systems, integrative plasmids,
188 and genes involved in the biosynthesis of extracellular structures such as exopolysaccharides (EPS)
189 (Fig. 3). Another set of genes disseminated via HGT in bifidobacteria is the CRISPR-related system
190 (CASS) implicated in defence against phages and plasmids⁴⁷, which have been identified in the
191 genome of *B. dentium* Bd1 as well as in the genome of *B. breve* UCC2003 (Ventura et al.,
192 unpublished data; Leahy and D. van Sinderen, unpublished data). Notably these *in silico* analyses
193 were also confirmed by comparative genome hybridization analyses⁴⁸.

194 There is relatively little phylogenetic diversity within the genus *Bifidobacterium* compared
195 to *Lactobacillus* (see below). This is underlined at whole genome level when one compares the oral
196 species (*B. dentium*), which is frequently identified as a component of the microbiota associated
197 with dental caries⁴⁹ with the probiotic species *B. adolescentis* (Fig. 3). Despite the large phenotypic
198 differences, there is a remarkable degree of overall synteny. This reductionist model of genome
199 evolution may be useful for identifying niche-specific genes and genes related to specialized
200 phenotypes.

201

202 **Genomics of *Lactobacillus***

203 The genus *Lactobacillus* has more than 100 species, and is noteworthy for its extreme phylogenetic,
204 phenotypic and ecological diversity²². The microbiological characterization of lactobacilli is
205 historically better developed than that of bifidobacteria, but the genomic analysis is similarly recent.
206 Of the 14 sequenced and published *Lactobacillus* genomes, eight (*L. acidophilus*, *L. casei*, *L.*
207 *fermentum*, *L. gasseri*, *L. johnsonii*, *L. reuteri*, *L. salivarius* and *L. plantarum*) are from
208 cultures/species considered probiotic (Table 1). Interestingly, 11% of the overall coding capacity of
209 the *L. salivarius* genome lies on the first megaplasmid described in lactic acid bacteria; pMP118²².
210 This megaplasmid encodes biologically important features such as a locus for bacteriocin
211 production, a bile salt hydrolase, and two genes that complete the phosphoketolase pathway,
212 officially reclassifying this organism as a facultative heterofermenter²². In fact, plasmids account for
213 15% of the genome of *L. salivarius*, which is not the case with other sequenced probiotic
214 lactobacilli, even though members of this genus are considered relatively replete with plasmids⁹.
215 *Adaptation to the human gut.*

216 The metabolic diversity revealed by the *Lactobacillus* genome sequences available to date is
217 illustrated in Fig 4. Taking the *L. plantarum* WCFS1 genome as reference, it is clear that there is
218 considerable variation in the COG assignments of the gene sets harboured by the respective
219 genomes. Intestinal lactobacilli compensate for their relative degree auxotrophy by being rich in

220 genes for transporters. Their genomes also contain genes that encode acid and bile resistance,
221 capacity for uptake of macromolecules, metabolism of complex carbohydrates, and cell surface
222 proteins that interact with the intestinal mucosa⁶⁰. More strikingly than is evident for bifidobacteria,
223 this adaptation to life in the GIT is further evident when the genome sequences of intestinal isolates
224 are compared with food-adapted lactobacilli such as *L. bulgaricus* and *L. helveticus*. *L. bulgaricus*.
225 The latter, which is widely used as a starter culture in yogurt fermentations, has undergone genome
226 decay to adapt to the milk environment⁵³, and thus harbours numerous degraded or partial
227 carbohydrate pathways and harbours bile salt hydrolase pseudogenes^{53, 60}. In addition, *L. bulgaricus*
228 shows a preference for growth in lactose, further emphasizing its niche adaptation to milk. The
229 genome sequence of *L. helveticus*, a widely used cheese starter culture, has been reported recently⁵².
230 Compared to the closely related *L. acidophilus*, *L. helveticus* has additional genes for fatty acid
231 biosynthesis and specific amino acid metabolism, but notably fewer cell surface proteins and PEP-
232 PTS systems for sugar utilization⁵². Additionally, no functional mucus binding proteins or
233 transporters for complex carbohydrates such as raffinose and fructooligosaccharides are encoded by
234 the *L. helveticus* genome, reflecting the degree of adaptation of *L. helveticus* to a milk environment.
235 In contrast, *L. acidophilus* has adapted to the gut ecological niche by retaining the functional gene
236 sets lacking in *L. helveticus*, emphasizing their importance for probiotic functionality and niche
237 adaptation by autochthonous lactobacilli naturally residing in the GIT.
238 Several studies have examined commensal *Lactobacillus* gene expression in animal model systems.
239 Using a stringent lincomycin-resistance based selection, Walter and colleagues identified
240 surprisingly only three genes that were differentially expressed *in vivo*⁶⁹. Bron *et al.*⁷⁰ used a
241 modified *in vivo* expression technology to identify 72 genes expressed by *L. plantarum* in the
242 mouse GIT, most of which were associated with carbon metabolism, amino acid metabolism, and
243 stress resistance⁷⁰, and many of which were functions previously identified as survival/adaptation
244 factors in pathogens. *L. casei* actively transcribes metabolic genes in the murine intestine, and
245 initiates *de novo* protein synthesis⁷¹. *L. johnsonii* NCC533 expresses different sets of genes

246 depending on its location in the GIT⁷², and surprisingly, 44% of the genome remains untranscribed
247 either *in vitro* or *in vivo*⁷². Interestingly, the prolonged murine gut persistence of NCC533 but not
248 of *L. johnsonii* was recently shown to induce expression of exopolysaccharide synthesis genes,
249 mannose uptake genes and a gene for a putative protease in this strain⁷³. In summary, while there
250 are tantalizing glimpses of commensal *Lactobacillus* gene expression *in vivo*, these are as yet
251 limited to animal models; data from human volunteer studies is keenly awaited.

252 *Molecular basis of the interaction with other commensal bacteria.*

253 Although the biology of commensal bacteria can be investigated in isolation, it must ultimately be
254 understood in the context of the extremely complex intestinal ecosystem⁶¹. *Lactobacillaceae*
255 account for approximately 36 phylotypes among the >1000 phylotypes in the human
256 gastrointestinal microbiota⁵. In the short term, intervention studies in animal models and human
257 subjects provide the key insights into our current understanding of interaction with other
258 commensals.

259 Some lactobacilli may have quite subtle effects on the microbiota. Consumption of *L. rhamnosus*
260 DR20 transiently altered the levels of lactobacilli, bifidobacteria, enterococci, and *Bacteroidetes*,
261 but the variations were generally small⁶² and mechanisms were not investigated. The development
262 of genomic tools facilitated a study⁴⁵, in germ-free mice that were mono-associated with *B.*
263 *thetaiotaomicron*, *B. longum*, *L. casei*, or combinations of these organisms⁴⁵. Presence of *L. casei*
264 resulted in an expanded capacity of *B. thetaiotaomicron* to metabolize polysaccharides, and
265 increased expression of genes for inorganic ion transport and metabolism⁴⁵. The *L. casei*-induced
266 changes in the *Bacteroides* transcriptome were functionally similar to those caused by *B. longum*,
267 but distinct from those induced by administration of *B. animalis* to the mice. Administration of *L.*
268 *paracasei* or *L. rhamnosus* to germ-free mice colonized with human infant microbiota caused
269 modest changes in levels of a limited number of species monitored by culture techniques, but major
270 changes to levels of diverse metabolites including amino acids, methylamines and short-chain fatty

271 acids⁶³. The metabolism of the administered probiotics, coupled with competition for substrates and
272 small molecules, are the likely reasons for the transcriptional and metabolite alterations described in
273 these studies.

274 Numerous studies have reported that administration of probiotics benefits a range of gastrointestinal
275 conditions and infections^{64, 65}, but mechanistic insights are generally lacking. Reduction in vaginal
276 *Lactobacillus* levels that leads to vaginosis has been linked to production of a bacteriocin-like
277 substance by commensal enterococci⁶⁶. From the opposite perspective, the ability of *L. salivarius* to
278 eliminate *Listeria monocytogenes* in a mouse model was dependent on production of the broad
279 spectrum bacteriocin Abp118/salivaricin⁶⁷, and bacteriocin-producing lactobacilli become dominant
280 among strains in a cocktail that reduce *Salmonella* shedding in pigs⁶⁸. Thus bacteriocin production
281 is likely an important general mechanism in the interaction of many lactobacilli and other
282 commensals.

283

284 **Comparative genomics of *Lactobacillus*.**

285 Sequencing of the genomes of twenty lactic acid bacteria (LAB) has demonstrated that loss and
286 decay of ancestral genes has played a key role in the evolution of *Lactobacillales*. *Lactobacillales*
287 diverged from their *Bacillus* ancestor with an estimated loss of 600-1200 genes from a total gene
288 repertoire of 2,100 to 2,200⁵⁰. Many of these genes encoded biosynthetic enzymes or functioned in
289 the sporulation process⁵⁰. However, in addition to major gene losses, gene gains also occurred
290 which appear to reflect the nutrient-rich niches occupied by the LAB, such as milk and the GIT. For
291 example, genes encoding for peptidases, amino acid transport proteins and genes involved in the
292 metabolism and transport of carbohydrates have been duplicated⁵⁰. In addition, comparative
293 analysis between GIT-associated species *L. acidophilus*, *L. gasseri*, and *L. johnsonii* and the dairy
294 species *L. bulgaricus* and *L. helveticus* revealed selective pressure from niche-specific adaptation on
295 the genome evolution of these species⁵¹⁻⁵³.

296 In addition to gene duplication, HGT is also evident in probiotic lactobacilli. For example, the
 297 metabolic diversity of *L. plantarum* is underpinned by the expanded coding capacity afforded by its
 298 larger 3 Mb genome, and a low-GC-content region coding for sugar transport and metabolism genes
 299 which is likely to have been acquired by HGT⁵⁴. Genes encoding cell surface factors in *L. johnsonii*
 300 and the exopolysaccharide cluster in the *L. acidophilus* complex are further examples of HGT in
 301 probiotic lactobacilli^{52, 55}. Moreover, production of reuterin (3-hydroxypropionaldehyde), a potent
 302 broad-spectrum antimicrobial compound⁵⁶, is encoded by a genomic island which is present in some
 303 *L. reuteri* strains⁵⁷⁻⁵⁹, and absent in the sequenced genome of a mouse *L. reuteri* isolate⁵⁸ and the
 304 closely related *L. fermentum*⁵⁹.

305 With genomes of 12 of the 147 recognized species⁷⁴ now fully sequenced, *Lactobacillus* has been
 306 targeted for several comparative whole-genome analyses. Beginning with the report of extreme
 307 diversity between the first two available genomes³⁴, genome sequencing of *L. acidophilus*, *L.*
 308 *gasseri*, *L. delbrueckii* and *L. helveticus* allowed a more focused attention on the ‘acidophilus
 309 complex’^{25, 52, 75}. Large regions of synteny were observed between the species^{25, 52}. Multi-locus
 310 sequence analysis of five housekeeping genes, comparative-genome hybridizations and DNA-
 311 typing showed consistent and stepwise-decreasing levels of similarity within the group, suggesting
 312 a strong role for vertical evolution²⁵. Conversely, differences between trees from 16S rRNA genes
 313 and 401 core genes from *L. acidophilus*, *L. johnsonii* and *L. delbrueckii* indicated a much higher
 314 level (40%) of HGT⁷⁵.

315 In order to infer robust phylogenetic relationships with minimal incongruence, or to elucidate
 316 functional differences between species, a set of carefully selected single-copy ubiquitously-present
 317 genes is necessary. A comparison of 354 core genes from five lactobacilli underscored the
 318 substantial diversification of the genus, and suggested a subgeneric division into three groups²¹.
 319 Furthermore, two overlapping comparative studies, encompassing nine additional *Lactobacillales*
 320 genomes, saw the expansion of the gene core to 567 order-specific genes^{50, 76}. Similarly, the
 321 majority of these encoded information-processing proteins. The finer granularity provided by

LaCOGs (*Lactobacillales*-specific COGs) allowed detection of two genes, whose gene-contexts suggest housekeeping and protein-modification functions. Recently, we extracted 141 core genes from 12 *Lactobacillus* genomes to investigate the case for a single congruent genus phylogeny²². Although this proved impossible at the time, four sub-generic groups were reliably distinguished. These were operationally characterized by absent genes rather than gained/retained genes, consistent with the findings of an earlier study⁷⁶.

Common evolutionary trends in probiotic genomes

Collective analyses of probiotic genome sequences so far available — the probiome — has revealed some generally conserved genetic traits^{22, 25, 52, 54, 55, 59, 76}, which may reflect adaptation to the intestinal niche¹. However, since probiotic bacteria represent diverse and taxonomically heterogeneous groups of microorganisms, the analysis of phyletic (phylogenetic) patterns, i.e. patterns of gene presence/absence in a particular set of genomes, may be overwhelmingly influenced by the evolutionary distance between these two distant phyla. Nevertheless, common trends in the evolution of both *Bifidobacterium* and *Lactobacillus* genomes may be discerned. These include gene loss (e.g. of genes encoding biosynthetic enzymes), gene duplication and HGT. The adaptation of probiotic bacteria to successfully exist and compete in the human gut must have been driven by the occurrence of DNA duplications and genetic acquisitions during their evolution. Many genes involved in sugar metabolism and transport were duplicated or acquired early in the evolution of probiotic bacteria, including those encoding enolase, β -galactosidase, and many other GH⁵⁰. In addition, expansion of peptidases and amino acid transporters has occurred in several lineages of *Lactobacillales* and bifidobacteria. Furthermore, several expanded families include proteins involved in antibiotic resistance in other bacteria, i.e. β -lactamases⁷⁷. Horizontal gene transfer via bacteriophage-mediated or conjugative pathways has been extensively documented in *Lactobacillales* and appears to be important for niche-specific adaptation in probiotic bacteria. In probiotic lactobacilli, HGT played an important role in shaping the common

348 ancestor, in which 84 genes were inferred to be horizontally transferred from different sources⁵⁰. In
349 some cases, the ancestor acquired an additional pseudoparalogous copy of a gene by HGT (e.g.
350 enolase in *Lactobacillales*) while on other occasions, xenologous displacement, i.e., acquisition of
351 genes by HGT followed by the loss of the ancestral orthologous gene⁷⁸ apparently took place.
352 A provocative future challenge will involve the identification of the hypothetical core
353 probiogenome, representing core genome functions of probiotic bacteria. However, only seven
354 genes present in the bifidobacteria but not in the genomes of the other members of the
355 *Actinobacteria* phylum are shared with *Lactobacillales*. Only one of these genes, which encodes a
356 functionally uncharacterized membrane protein, is present in all the *Lactobacillales* genomes so far
357 sequenced⁵⁰.

358

359 **Conclusions and future considerations**

360 Most of the probiotic bacteria marketed today were originally selected on the basis of technological
361 stability or by a variety of easily measurable phenotypes such as ability to tolerate bile salts or
362 survive GIT passage, but not necessarily for their ability to promote health benefits. It is crucial to
363 identify the precise mechanisms by which such probiotic microorganisms influence human health.
364 Such studies should be accelerated by omics approaches involving genomics and functional
365 analyses. Molecular interaction models are being currently developed, although more are required,
366 that monitor the activation of cellular and systemic responses *in vivo* in animal models and in
367 feeding trial participants through the measurement of previously validated biomarkers. The
368 combination of verified molecular models with functional and comparative genomics-based
369 approaches should enable selection of the most appropriate probiotic strain for a particular health
370 benefit or improvement of strain processing and administration regimes that optimize the
371 established health effect. Finally, this might allow the selection of specific probiotics for a particular
372 human genotype, in analogy to personalized genomic medicine efforts.

373 Several issues regarding the sequences of complete probiotic bacterial genomes remain unresolved
374 at present. So far, only a limited number of completed probiotic bacterial genome sequences are
375 available, which only partially represent the total biodiversity of probiotic bacteria residing in the
376 human gut. In this context, understanding of the human gut microbiome will be an important
377 challenge for the future⁷⁹. Furthermore, sequencing the genomes of environmental organisms and
378 carrying out metagenomic surveys of diverse gut environments (human vs. animal GIT) will
379 provide not only an improved understanding of microbial biodiversity but also insights into the
380 evolution of bacterial factors that may be crucial for the commensals (probiotics) establishment in
381 these different gut niches⁸⁰.

382 The first decade of bacterial genomics has afforded unprecedented insights into the evolution of
383 bacterial pathogens (bacterial pathogenomics)⁸¹. The next decade holds the promise of being even
384 more rewarding as the new discoveries about probiotic bacteria provided by probiogenomic efforts
385 are exploited.

386

387 **GLOSSARY**

388 Omics: The integration of genomics methodology and data with functional genomic analyses
389 involving transcriptomics, proteomics, metabolomics and interactomics.

390 Microbiota: The collective microbial community or population resident in a particular locale at a
391 given time-point.

392 Microbiome: The collective genome of the human microbial communities

393 Prebiotics: Growth substrates that are preferentially (or ideally, exclusively) metabolized by a single
394 genus or species, and that may thus be used as dietary supplements to promote growth of a targeted
395 microorganism.

396 Transcriptome: Subsets of genes transcribed in an organism. It represents dynamic links between
397 genomes, proteins and cellular phenotypes.

398 Synteny: Genetic linkage or conservation of gene order.

399 COGs: Clusters of Orthologous Groups are delineated by comparing protein sequences encoded in
400 complete genomes, representing major phylogenetic lineages. Each COG consists of individual
401 proteins or groups of paralogs from at least 3 lineages and thus corresponds to an ancient conserved
402 domain.

403 Neighbour-joining tree: Tree that reconstruct the evolutionary development of organisms based on
404 distances between each pair of taxa.

405 **Acknowledgements**

406 Work in the laboratories of DvS and PWOT is supported by an SFI CSET award to the Alimentary
407 Pharmabiotic Centre, and a DAF/HRB FHRI award to the ELDERMET project. MV was supported
408 by an Italian Award for Outstanding Young Researcher scheme “Incentivazione alla mobilità di
409 studiosi stranieri e italiani residente all’estero” 2005-2009, a Marie Curie Reintegration Grant
410 (MERG-CT-2005-03080) and Parmalat spa, Italy. We thank also S. Leahy, Univ. College Cork and
411 D. John, Trinity College Dublin for the scanning electron microscope images.

412

413 References

- 414 1. Backhed, F., Ley, R. E., Sonnenburg, J. L., Peterson, D. A. & Gordon, J. I. Host-bacterial
415 mutualism in the human intestine. *Science* 307, 1915-20 (2005).
- 416 2. **Eckburg, P. B. et al. Diversity of the human intestinal microbial flora. *Science* 308,**
417 **1635-8 (2005).**
- 418 **This article describes the bacterial diversity occurring in the human gut using 16S rRNA gene**
419 **based libraries.**
- 420 3. Seksik, P. et al. Alterations of the dominant faecal bacterial groups in patients with Crohn's
421 disease of the colon. *Gut* 52, 237-42 (2003).
- 422 4. Turrone, F., Ribbera, A., Foroni, E., van Sinderen, D. & Ventura, M. Human gut microbiota
423 and bifidobacteria: from composition to functionality. *Antonie Van Leeuwenhoek* 94, 35-50
424 (2008).
- 425 5. **Rajilic-Stojanovic, M., Smidt, H. & de Vos, W. M. Diversity of the human**
426 **gastrointestinal tract microbiota revisited. *Environ. Microbiol.* 9, 2125-36 (2007).**
- 427 **This review provides an integrated summary of data from culture independent studies of the**
428 **human gut microbiota.**
- 429 6. Ley, R. E., Peterson, D. A. & Gordon, J. I. Ecological and evolutionary forces shaping
430 microbial diversity in the human intestine. *Cell* 124, 837-48 (2006).
- 431 7. Guarner, F. & Malagelada, J. R. Gut flora in health and disease. *Lancet* 361, 512-9 (2003).
- 432 8. Hooper, L. V. & Gordon, J. I. Commensal host-bacterial relationships in the gut. *Science*
433 292, 1115-8 (2001).
- 434 9. Backhed, F. et al. The gut microbiota as an environmental factor that regulates fat storage.
435 *Proc Natl Acad Sci U S A* 101, 15718-23 (2004).
- 436 10. Samuel, B. S. & Gordon, J. I. A humanized gnotobiotic mouse model of host-archaeal-
437 bacterial mutualism. *Proc Natl Acad Sci U S A* 103, 10011-6 (2006).
- 438 11. Turnbaugh, P. J. et al. An obesity-associated gut microbiome with increased capacity for
439 energy harvest. *Nature* 444, 1027-31 (2006).
- 440 12. Frank, D. N. et al. Molecular-phylogenetic characterization of microbial community
441 imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci U S A* 104, 13780-5
442 (2007).
- 443 13. Kassinen, A. et al. The fecal microbiota of irritable bowel syndrome patients differs
444 significantly from that of healthy subjects. *Gastroenterology* 133, 24-33 (2007).
- 445 14. **Manichanh, C. et al. Reduced diversity of faecal microbiota in Crohn's disease**
446 **revealed by a metagenomic approach. *Gut* 55, 205-11 (2006).**
- 447 **The preceding two references provide evidence for significant microbiota alterations in**
448 **functional bowel disorders.**
- 449 15. Salminen, S., Nurmi, J. & Gueimonde, M. The genomics of probiotic intestinal
450 microorganisms. *Genome Biol* 6, 225 (2005).
- 451 16. Marco, M. L., Pavan, S. & Kleerebezem, M. Towards understanding molecular modes of
452 probiotic action. *Curr Opin Biotechnol* 17, 204-10 (2006).
- 453 17. O'Hara, A. M. & Shanahan, F. Mechanisms of action of probiotics in intestinal diseases.
454 *ScientificWorldJournal* 7, 31-46 (2007).
- 455 18. Saxelin, M., Tynkkynen, S., Mattila-Sandholm, T. & de Vos, W. M. Probiotic and other
456 functional microbes: from markets to mechanisms. *Curr Opin Biotechnol* 16, 204-11 (2005).
- 457 19. Ventura, M. et al. Genomics of Actinobacteria: tracing the evolutionary history of an ancient
458 phylum. *Microbiol Mol Biol Rev* 71, 495-548 (2007).
- 459 20. Joyce, A. R. & Palsson, B. O. The model organism as a system: integrating 'omics' data sets.
460 *Nat Rev Mol Cell Biol* 7, 198-210 (2006).

- 461 21. Ventura, M. et al. Analysis of bifidobacterial evolution using a multilocus approach. *Int J*
462 *Syst Evol Microbiol* 56, 2783-92 (2006).
- 463 22. Claesson, M. J. et al. Multireplicon genome architecture of *Lactobacillus salivarius*. *Proc*
464 *Natl Acad Sci U S A* 103, 6718-23 (2006).
- 465 23. Tissier, H. Traitement des infections intestinales par la methode de la flore bacterienne de
466 l'intestin. *Crit. Rev. Soc. Biol.* 60, 359-361 (1906).
- 467 24. Ventura, M., Canchaya, C., Fitzgerald, G. F., Gupta, R. S. & van Sinderen, D. Genomics as
468 a means to understand bacterial phylogeny and ecological adaptation: the case of
469 bifidobacteria. *Antonie Van Leeuwenhoek* 91, 351-72 (2007).
- 470 25. Schell, M. A. et al. The genome sequence of *Bifidobacterium longum* reflects its adaptation
471 to the human gastrointestinal tract. *Proc Natl Acad Sci U S A* 99, 14422-7 (2002).
- 472 26. Gibson, G. R. & Roberfroid, M. B. Dietary modulation of the human colonic microbiota:
473 introducing the concept of prebiotics. *J Nutr* 125, 1401-12 (1995).
- 474 27. Flint, H. J., Bayer, E. A., Rincon, M. T., Lamed, R. & White, B. A. Polysaccharide
475 utilization by gut bacteria: potential for new insights from genomic analysis. *Nat Rev*
476 *Microbiol* 6, 121-31 (2008).
- 477 28. Sonnenburg, J. L. et al. Glycan foraging in vivo by an intestine-adapted bacterial symbiont.
478 *Science* 307, 1955-9 (2005).
- 479 29. Hooper, L. V., Xu, J., Falk, P. G., Midtvedt, T. & Gordon, J. I. A molecular sensor that
480 allows a gut commensal to control its nutrient foundation in a competitive ecosystem. *Proc*
481 *Natl Acad Sci U S A* 96, 9833-8 (1999).
- 482 30. Hinz, S. W., Verhoef, R., Schols, H. A., Vincken, J. P. & Voragen, A. G. Type I
483 arabinogalactan contains beta-D-Galp-(1-->3)-beta-D-Galp structural elements. *Carbohydr*
484 *Res* 340, 2135-43 (2005).
- 485 31. Ryan, S. M., Fitzgerald, G. F. & van Sinderen, D. Screening for and identification of starch-,
486 amylopectin-, and pullulan-degrading activities in bifidobacterial strains. *Appl Environ*
487 *Microbiol* 72, 5289-96 (2006).
- 488 32. Maze, A., O'Connell-Motherway, M., Fitzgerald, G. F., Deutscher, J. & van Sinderen, D.
489 Identification and characterization of a fructose phosphotransferase system in
490 *Bifidobacterium breve* UCC2003. *Appl Environ Microbiol* 73, 545-53 (2007).
- 491 **33. van den Broek, L. A., Hinz, S. W., Beldman, G., Vincken, J. P. & Voragen, A. G.**
492 ***Bifidobacterium* carbohydrases-their role in breakdown and synthesis of (potential)**
493 **prebiotics. *Mol Nutr Food Res* 52, 146-63 (2008).**
- 494 **This paper is providing the state of the art in the area of enzymes encoded by bifidobacteria**
495 **involved in the hydrolysis of carbohydrates**
- 496 34. Siezen, R. et al. *Lactobacillus plantarum* gene clusters encoding putative cell-surface protein
497 complexes for carbohydrate utilization are conserved in specific gram-positive bacteria.
498 *BMC Genomics* 7, 126 (2006).
- 499 35. Hooper, L. V., Midtvedt, T. & Gordon, J. I. How host-microbial interactions shape the
500 nutrient environment of the mammalian intestine. *Annu Rev Nutr* 22, 283-307 (2002).
- 501 36. Hoskins, L. C. et al. Mucin degradation in human colon ecosystems. Isolation and properties
502 of fecal strains that degrade ABH blood group antigens and oligosaccharides from mucin
503 glycoproteins. *J Clin Invest* 75, 944-53 (1985).
- 504 37. Ruas-Madiedo, P., Gueimonde, M., Fernandez-Garcia, M., de los Reyes-Gavilan, C. G. &
505 Margolles, A. Mucin degradation by *Bifidobacterium* strains isolated from the human
506 intestinal microbiota. *Appl Environ Microbiol* 74, 1936-40 (2008).
- 507 38. Ehrmann, M. A., Korakli, M. & Vogel, R. F. Identification of the gene for beta-
508 fructofuranosidase of *Bifidobacterium lactis* DSM10140(T) and characterization of the
509 enzyme expressed in *Escherichia coli*. *Curr Microbiol* 46, 391-7 (2003).

39. Katayama, T. et al. Molecular cloning and characterization of *Bifidobacterium bifidum* 1,2-alpha-L-fucosidase (AfcA), a novel inverting glycosidase (glycoside hydrolase family 95). *J Bacteriol* 186, 4885-93 (2004).
40. Ryan, S. M., Fitzgerald, G. F. & van Sinderen, D. Transcriptional regulation and characterization of a novel beta-fructofuranosidase-encoding gene from *Bifidobacterium breve* UCC2003. *Appl Environ Microbiol* 71, 3475-82 (2005).
41. Gonzalez, R., Klaassens, E. S., Malinen, E., de Vos, W. M. & Vaughan, E. E. Differential transcriptional response of *Bifidobacterium longum* to human milk, formula milk and galactooligosaccharide. *Appl Environ Microbiol* (2008).
42. Liepke, C. et al. Human milk provides peptides highly stimulating the growth of bifidobacteria. *Eur J Biochem* 269, 712-8 (2002).
43. Ivanov, D. et al. A serpin from the gut bacterium *Bifidobacterium longum* inhibits eukaryotic elastase-like serine proteases. *J Biol Chem* 281, 17246-52 (2006).
44. Potempa, J., Korzus, E. & Travis, J. The serpin superfamily of proteinase inhibitors: structure, function, and regulation. *J Biol Chem* 269, 15957-60 (1994).
45. **Sonnenburg, J. L., Chen, C. T. & Gordon, J. I. Genomic and metabolic studies of the impact of probiotics on a model gut symbiont and host. *PLoS Biol* 4, e413 (2006).**
This paper describes the crosstalk existing between bifidobacteria and *Bacteroides* in the murine intestine as well as between these bacteria and their host.
46. Kato, S., Haruta, S., Cui, Z. J., Ishii, M. & Igarashi, Y. Stable coexistence of five bacterial strains as a cellulose-degrading community. *Appl Environ Microbiol* 71, 7099-106 (2005).
47. Barrangou, R. et al. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* 315, 1709-12 (2007).
48. Klijn, A., Mercenier, A. & Arigoni, F. Lessons from the genomes of bifidobacteria. *FEMS Microbiol Rev* 29, 491-509 (2005).
49. Aas, J. A. et al. Bacteria of dental caries in primary and permanent teeth in children and young adults. *J Clin Microbiol* 46, 1407-17 (2008).
50. Makarova, K. S. & Koonin, E. V. Evolutionary genomics of lactic acid bacteria. *J Bacteriol* 189, 1199-208 (2007).
51. Makarova, K. et al. Comparative genomics of the lactic acid bacteria. *Proc. Natl. Acad. Sci. U S A* 103, 15611-6 (2006).
- This landmark study provided a large tranche of genomic data to allow studies of genome evolution in lactic acid bacteria.**
52. Altermann, E. et al. Complete genome sequence of the probiotic lactic acid bacterium *Lactobacillus acidophilus* NCFM. *Proc Natl Acad Sci U S A* 102, 3906-12 (2005).
53. van de Guchte, M. et al. The complete genome sequence of *Lactobacillus bulgaricus* reveals extensive and ongoing reductive evolution. *Proc. Natl. Acad. Sci. U S A* 103, 9274-9 (2006).
54. **Kleerebezem, M. et al. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc Natl Acad Sci U S A* 100, 1990-5 (2003).**
This is the first article describing the genome sequence of a member of the genus *Lactobacillus*.
55. **Pridmore, R. D. et al. The genome sequence of the probiotic intestinal bacterium *Lactobacillus johnsonii* NCC 533. *Proc Natl Acad Sci U S A* 101, 2512-7 (2004).**
This paper describes the genome contents of a common used probiotic bacterium belonging to the genus *Lactobacillus*.
56. Talarico, T. L., Casas, I. A., Chung, T. C. & Dobrogosz, W. J. Production and isolation of reuterin, a growth inhibitor produced by *Lactobacillus reuteri*. *Antimicrob. Agents Chemother.* 32, 1854-8 (1988).
57. Santos, F. et al. The complete coenzyme B12 biosynthesis gene cluster of *Lactobacillus reuteri* CRL1098. *Microbiology* 154, 81-93 (2008).

- 561 58. Sriramulu, D. D. et al. *Lactobacillus reuteri* DSM 20016 produces cobalamin-dependent diol
562 dehydratase in metabolosomes and metabolizes 1,2-propanediol by disproportionation. *J.*
563 *Bacteriol.* 190, 4559-67 (2008).
- 564 59. Morita, H. et al. Comparative Genome Analysis of *Lactobacillus reuteri* and *Lactobacillus*
565 *fermentum* Reveal a Genomic Island for Reuterin and Cobalamin Production. *DNA Res* 15,
566 151-61 (2008).
- 567 60. Pfeiler, E. A. & Klaenhammer, T. R. The genomics of lactic acid bacteria. *Trends Microbiol*
568 15, 546-53 (2007).
- 569 61. Whitman, W. B., Coleman, D. C. & Wiebe, W. J. Prokaryotes: the unseen majority. *Proc.*
570 *Natl. Acad. Sci. U S A* 95, 6578-83 (1998).
- 571 62. Tannock, G. W. et al. Analysis of the fecal microflora of human subjects consuming a
572 probiotic product containing *Lactobacillus rhamnosus* DR20. *Appl. Environ. Microbiol.* 66,
573 2578-88 (2000).
- 574 63. Martin, F. P. et al. Probiotic modulation of symbiotic gut microbial-host metabolic
575 interactions in a humanized microbiome mouse model. *Mol. Syst. Biol.* 4, 157 (2008).
- 576 64. Hickson, M. et al. Use of probiotic *Lactobacillus* preparation to prevent diarrhoea associated
577 with antibiotics: randomised double blind placebo controlled trial. *Brit. Med. J.* 335, 80
578 (2007).
- 579 65. Sullivan, A. & Nord, C. E. Probiotics and gastrointestinal diseases. *J. Intern. Med.* 257, 78-
580 92 (2005).
- 581 66. Kelly, M. C., Mequio, M. J. & Pybus, V. Inhibition of vaginal lactobacilli by a bacteriocin-
582 like inhibitor produced by *Enterococcus faecium* 62-6: potential significance for bacterial
583 vaginosis. *Infect. Dis. Obstet. Gynecol.* 11, 147-56 (2003).
- 584 **67. Corr, S. C. et al. Bacteriocin production as a mechanism for the antiinfective activity**
585 **of *Lactobacillus salivarius* UCC118. *Proc. Natl. Acad. Sci. U S A* 104, 7617-7621 (2007).**
586 **This study identified the first molecular mechanism whereby probiotic bacteria modulate the**
587 **microbiota *in vivo*.**
- 588 68. Casey, P. G. et al. A five-strain probiotic combination reduces pathogen shedding and
589 alleviates disease signs in pigs challenged with *Salmonella enterica* Serovar Typhimurium.
590 *Appl. Environ. Microbiol.* 73, 1858-63 (2007).
- 591 **69. Walter, J. et al. Identification of *Lactobacillus reuteri* genes specifically induced in the**
592 **mouse gastrointestinal tract. *Appl. Environ. Microbiol.* 69, 2044-51 (2003).**
- 593 **70. Bron, P. A., Grangette, C., Mercenier, A., de Vos, W. M. & Kleerebezem, M.**
594 **Identification of *Lactobacillus plantarum* genes that are induced in the gastrointestinal**
595 **tract of mice. *J Bacteriol* 186, 5721-9 (2004).**
596 **This manuscript provides an insight into the molecular interactions between a commensal**
597 **microorganism and a murine-model host.**
- 598 71. Oozeer, R. et al. Differential activities of four *Lactobacillus casei* promoters during bacterial
599 transit through the gastrointestinal tracts of human-microbiota-associated mice. *Appl.*
600 *Environ. Microbiol.* 71, 1356-63 (2005).
- 601 72. Denou, E. et al. Gene expression of commensal *Lactobacillus johnsonii* strain NCC533
602 during in vitro growth and in the murine gut. *J. Bacteriol.* 189, 8109-19 (2007).
- 603 73. Denou, E. et al. Identification of genes associated with the long-gut-persistence phenotype
604 of the probiotic *Lactobacillus johnsonii* strain NCC533 using a combination of genomics
605 and transcriptome analysis. *J. Bacteriol.* 190, 3161-8 (2008).
- 606 74. Euzèby, J. P. (2007). List of Bacterial names with Standing in Nomenclature. *Int J Syst*
607 *Bacteriol.* 47, 590-592 (1997).
- 608 75. Nicolas, P., Bessieres, P., Ehrlich, S. D., Maguin, E. & van de Guchte, M. Extensive
609 horizontal transfer of core genome genes between two *Lactobacillus* species found in the
610 gastrointestinal tract. *BMC Evol. Biol.* 7, 141 (2007).

611 76. Makarova, K. et al. Comparative genomics of the lactic acid bacteria. Proc Natl Acad Sci U
612 S A 103, 15611-6 (2006).

613 77. Teuber, M., Meile, L. & Schwarz, F. Acquired antibiotic resistance in lactic acid bacteria
614 from food. *Antonie Van Leeuwenhoek* 76, 115-37 (1999).

615 78. Koonin, E. V., Makarova, K. S. & Aravind, L. Horizontal gene transfer in prokaryotes:
616 quantification and classification. *Annu Rev Microbiol* 55, 709-42 (2001).

617 79. Turnbaugh, P. J. et al. The human microbiome project. *Nature* 449, 804-10 (2007).

618 80. **Ley, R. E. et al. Evolution of mammals and their gut microbes. *Science* 320, 1647-51**
619 **(2008).**

620 **This paper describes the bacterial diversity present in the gut of numerous mammals.**

621 81. Pallen M.J., & Wren, B.M. Bacterial pathogenomics. *Nature* 449, 835-842 (2007).

622

623 82. Lee, J. H. et al. Comparative genomic analysis of the gut bacterium *Bifidobacterium longum*
624 reveals loci susceptible to deletion during pure culture growth. *BMC Genomics* 9, 247
625 (2008).

626 83. Leahy, S. C., Higgins, D. G., Fitzgerald, G. F. & van Sinderen, D. Getting better with
627 bifidobacteria. *J Appl Microbiol* 98, 1303-15 (2005).

628

629

630

631

632

633

634

635

636

637

Table 1: General features of sequenced *Bifidobacterium* and *Lactobacillus* genomes.

Species	Genome size (bp)	%GC	Gene numbers	Proteins	Source	Accession number	Reference
<i>B. longum</i> subsp. <i>longum</i> NCC2705	2,256,640	60%	1798	1727	Human GIT	NC_004307	²⁵
<i>B. longum</i> subsp. <i>longum</i> DJ010A	2,375,286	59%	1908	1908	Human GIT	NC_010816	⁸²
<i>B. breve</i> UCC2003	2422668	59%	1868		Infant feces	Project ID: 13487	⁸³
<i>B. adolescentis</i> ATCC15703	2,089,645	59%	1701	1631	Human GIT	NC_008618	-
<i>B. adolescentis</i> L2-32	2,385,710	59%	2499	2428	Infant feces	NZ_AAXD000000000	-
<i>B. animalis</i> subsp. <i>lactis</i> HN019	1,915,892	60%	1632	1578	-	NZ_ABOT000000000	-
<i>Lactobacillus acidophilus</i> NCFM	1,993,560	34%	1936	1862	Human GIT	NC_006814	⁵²
<i>Lactobacillus casei</i> ATCC334	2,895,264	46%	2909	2751	Emmental cheese	NC_008526	⁷⁶
<i>Lactobacillus gasseri</i> ATCC33323	1,894,360	35%	1898	1755	Human GIT	NC_008530	⁵⁰
<i>Lactobacillus jonsonii</i> NCC533	1,992,676	34%	1918	1821	Human GIT	NC_005362	⁵⁵
<i>Lactobacillus plantarum</i> WCFS1	3,308,274	44%	3135	3007	Human saliva	NC_004567	⁵⁴
<i>Lactobacillus reuteri</i> F275	1,999,618	38%	2027	1900	Human GIT	NC_009513	⁶⁰
<i>Lactobacillus fermentum</i> IFO 3956	2,098,685	51%	1912	1843	-	NC_010610	⁶⁰
<i>Lactobacillus salivarius</i> susp. <i>salivarius</i> UCC118	1,827,111	32%	1864	1717	Human GIT	NC_007929	²²

LEGENDS

Figure 1: Ecological, evolutionary and morphological overview of bifidobacteria and lactobacillae.

[A] Schematic representation of the biological relationships between bacteria and the human body.

Commensalisms or symbiosis is a consequence of the co-evolution of host-bacterial relationships.

B| Evolutionary relationships between the main GIT commensal bacterial groups (bifidobacteria on the left and lactobacillae on the right) based on neighbour-joining tree of 16S rRNA genes

sequences. Bar indicates scale for computed distances. Bacterial taxa for which the whole genome sequences is available are shaded in blue, whereas for those that is still on progress are shaded in grey.

C| electron micrographs illustrating the cell morphology of bifidobacteria (e.g., *B. breve* UCC2003) (right panel) and lactobacillae (e.g., *L. salivarius* UCC118) (Left panel). Both scanning electron microscope images were prepared by. S. Leahy, Univ. College Cork and D. John, Trinity College Dublin. Magnification ca. 20,000 fold; scale bar is 2 micrometres.

Figure 2: Putative strategy adopted by bifidobacteria to secure sugar nutrients for their own benefit.

Bifidobacteria use a kind of docking station to capture complex sugars (e.g., xylan and arabino based molecules) and bind these to the bacterial cell surface, without loosing them to nearby

competitors. In the latter case the docking station is a complex of modular glycanases, which are anchored at the cell surface by a transmembrane domain. The enzymatic activities degrade the

arabinoxylan molecules to oligosaccharides that are subsequently transported across the bacterial membrane by a transporter protein; the presence of the bacterial cell wall may prohibit diffusion of these nutrients away from the transporter.

Figure 3. Comparative analysis of *Bifidobacterium* genomes. A| A comparison of the *B. dentium* Bd1 and *B. adolescentis* ATCC15703 genomes. B| Comparison of gene order conservation between two genome pairs, illustrating different forms of bifidobacterial genome evolution. X and Y axes represent the linearised chromosomes of *B. dentium* Bd1 and *B. adolescentis* ATCC15703, respectively.

Figure 4. Comparative analysis of *Lactobacillus* genomes. Circular genome atlas of *L. plantarum* WCFS1 with mapped orthologs (defined as reciprocal best FastA hits with more than 30% identity over at least 80% of both protein lengths) in 13 publicly available *Lactobacillus* genomes. The outer circle shows *L. plantarum* followed, inwards, by *L. salivarius*, *L. brevis*, *L. reuteri* F275, *L. reuteri* F275 (Japanese), *L. fermentum*, *L. acidophilus*, *L. helveticus*, *L. johnsonii*, *L. gasseri*, *L. bulgaricus* ATCC 11842, *L. bulgaricus* ATCC BAA-365, *L. casei*, *L. sakei*, G+C percentage, and GC skew (window-sizes 10,000 bp). Red colour represents COG categories in Metabolism, green - Information Storage and Processing, blue - Cellular Processes and Signalling, and grey - poorly or not categorised.





Figure 2

a)



b)




Figure 3




Figure 4

Table 1: General features of sequenced *Bifidobacterium* and *Lactobacillus* genomes.

Species	Genome size (bp)	%GC	Gene numbers	Proteins	Source	Accession number	Reference
<i>B. longum</i> subsp. <i>longum</i> NCC2705	2,256,640	60%	1798	1727	Human GIT	NC_004307	25
<i>B. longum</i> subsp. <i>longum</i> DJ010A	2,375,286	59%	1908	1908	Human GIT	NC_010816	82
<i>B. breve</i> UCC2003	2422668	59%	1868		Infant feces	Project ID: 13487	83
<i>B. adolescentis</i> ATCC15703	2,089,645	59%	1701	1631	Human GIT	NC_008618	-
<i>B. adolescentis</i> L2-32	2,385,710	59%	2499	2428	Infant feces	NZ_AAXD00000000	-
<i>B. animalis</i> subsp. <i>lactis</i> HN019	1,915,892	60%	1632	1578	-	NZ_ABOT00000000	-
<i>Lactobacillus acidophilus</i> NCFM	1,993,560	34%	1936	1862	Human GIT	NC_006814	52
<i>Lactobacillus casei</i> ATCC334	2,895,264	46%	2909	2751	Emmental cheese	NC_008526	76
<i>Lactobacillus gasseri</i> ATCC33323	1,894,360	35%	1898	1755	Human GIT	NC_008530	50
<i>Lactobacillus jonsonii</i> NCC533	1,992,676	34%	1918	1821	Human GIT	NC_005362	55
<i>Lactobacillus plantarum</i> WCFS1	3,308,274	44%	3135	3007	Human saliva	NC_004567	54
<i>Lactobacillus reuteri</i> F275	1,999,618	38%	2027	1900	Human GIT	NC_009513	60
<i>Lactobacillus fermentum</i> IFO 3956	2,098,685	51%	1912	1843	-	NC_010610	60
<i>Lactobacillus salivarius</i> susp. <i>salivarius</i> UCC118	1,827,111	32%	1864	1717	Human GIT	NC_007929	22

