

The versatility and adaptation of bacteria from the genus *Stenotrophomonas*

Robert P. Ryan^{*}, Sebastien Monchy[†], Massimiliano Cardinale[§], Safiyh Taghavi[‡], Lisa Crossman^{||}, Matthew B. Avison[¶], Gabriele Berg[§], Daniel van der Lelie[‡] and J. Maxwell Dow^{*}

Abstract | The genus *Stenotrophomonas* comprises at least eight species. These bacteria are found throughout the environment, particularly in close association with plants. Strains of the most predominant species, *Stenotrophomonas maltophilia*, have an extraordinary range of activities that include beneficial effects for plant growth and health, the breakdown of natural and man-made pollutants that are central to bioremediation and phytoremediation strategies and the production of biomolecules of economic value, as well as detrimental effects, such as multidrug resistance, in human pathogenic strains. Here, we discuss the versatility of the bacteria in the genus *Stenotrophomonas* and the insight that comparative genomic analysis of clinical and endophytic isolates of *S. maltophilia* has brought to our understanding of the adaptation of this genus to various niches.

The genus *Stenotrophomonas*, which is phylogenetically placed in the Gammaproteobacteria, was first described with the type species *Stenotrophomonas maltophilia*¹. This species was originally named *Pseudomonas maltophilia* by Hugh and Ryschenko in 1961, but was later transferred to the genus *Xanthomonas*² before it was given its own genus¹. The genus name (from the Greek 'stenos', meaning narrow, 'trophus', meaning one who feeds and 'monas', meaning unit) was intended to highlight the limited nutritional range of the bacterium¹. However, several studies subsequently demonstrated that the genus is capable of great metabolic versatility and intraspecific heterogeneity^{3–8} (FIG. 1). The genus currently comprises eight species, *S. maltophilia*, *Stenotrophomonas nitritireducens*⁹, *Stenotrophomonas rhizophila*¹⁰, *Stenotrophomonas acidaminiphila*¹¹, *Stenotrophomonas koreensis*¹², *Stenotrophomonas chelatiphaga*¹³, *Stenotrophomonas terrae*¹⁴ and *Stenotrophomonas humi*¹⁴. *Stenotrophomonas dokdonensis* was described in 2006 (REF. 15) but was transferred in 2008 to the genus *Pseudoxanthomonas*¹⁶. Phenotypic and genotypic studies as well as analysis of the ecological and metabolic diversity of these bacteria have revealed further differentiation at the species level.

Although *Stenotrophomonas* spp. occur ubiquitously in the environment, soil and plants are their main environmental reservoirs. *S. maltophilia* is a typical, often dominant member of the microbial communities that are found on or in plants and has a worldwide distribution (reviewed in REFS 17, 18). Members of the genus *Stenotrophomonas* have an important ecological role in the nitrogen and sulphur cycles^{19–21} and several *Stenotrophomonas* species, especially *S. maltophilia* and *S. rhizophila*, can engage in beneficial interactions with plants (FIG. 1). *S. maltophilia* is also an emerging human pathogen that is responsible for fatal infections in humans (reviewed in REFS 18, 22). In contrast to the phylogenetically closely related genera *Xanthomonas* and *Xylella*, no *Stenotrophomonas* species are known to be phytopathogenic (FIG. 2).

Various molecular tools, including transposon mutagenesis, allelic exchange and reporter fusions, have been developed to facilitate the genetic analysis of the diverse activities of *Stenotrophomonas* spp. Furthermore, the full genome sequence of an environmental isolate, *S. maltophilia* R551-3, and a clinical isolate, *S. maltophilia* K279a, are now available. In this Review, we describe the versatility of bacteria from the genus *Stenotrophomonas*, using a comparison of these

^{*}BIOMERIT Research Centre, Department of Microbiology, BioSciences Institute, University College Cork, Cork, Ireland.

[†]Brookhaven National Laboratory, Biology Department, Upton, New York 11973-5000, USA.

[§]Institute for Environmental Biotechnology, Graz University of Technology, Graz, Austria.

[‡]Pathogen Sequencing Unit, The Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK.

[¶]Department of Cellular and Molecular Medicine, University of Bristol, School of Medical Sciences, University Walk, Bristol, BS8 1TD, UK. Correspondence to R.P.R. e-mail: r.ryan@ucc.ie doi:10.1038/nrmicro2163




Figure 1 | Biotechnological uses of *Stenotrophomonas* spp. *Stenotrophomonas* spp. have many traits that could be used in different biotechnological processes. Some *Stenotrophomonas* spp. can produce antimicrobial compounds that protect plants, as well as generate factors that can promote plant growth. Also, many *Stenotrophomonas* spp. have a high level of intrinsic resistance to heavy metals and antibiotics and have been shown to degrade a wide range of compounds, including pollutants, and could potentially be used in bioremediation and phytoremediation. *Stenotrophomonas maltophilia* is also known to cause human disease as a result of its ability to colonize immunocompromised patients, and has been shown to be virulent in a nematode model.

recently sequenced genomes to highlight the possible genetic basis of adaptation to different niches.

Associations of *Stenotrophomonas* with plants

Stenotrophomonas species, especially *S. maltophilia* and *S. rhizophila*, are often found in association with plants. These bacteria can be isolated from the rhizosphere^{23,24} or from internal plant tissues, particularly from the vascular tissues of the root and stem (FIG. 3; [Supplementary information S1](#) (movie)). Endophytic strains of *S. maltophilia* have been isolated from the roots of many plant species, including cucumber (*Cucumis sativus*)²⁵, oilseed rape (*Brassica napus*)²⁶, potato (*Solanum tuberosum*)²⁶, strawberry (*Fragaria x ananassa*)²⁶, alfalfa (*Medicago sativa*)²⁷, sunflower (*Helianthus annuus*)²⁷, maize (*Zea mays*)²⁸, rice (*Oryza sativa*)²⁹, wheat (*Triticum aestivum*)³⁰, various weeds, willow (*Salix herbacea*)³¹ and poplar (*Populus*)³².

Factors involved in plant colonization. Most *Stenotrophomonas* spp. are highly adaptable to hostile and nutrient-limited environments. Several factors have a known or suggested influence on the ability of *Stenotrophomonas* spp. to colonize and survive on plant surfaces. The establishment of interactions between plants and microorganisms in the rhizosphere is preceded by the movement of free-living microorganisms towards the plant roots and can involve chemotaxis towards attractants that are present in plant root exudates. In addition to flagellar motility, other contributing factors to the colonization of plant tissues include a high bacterial growth rate, vitamin B1 synthesis, the exudation of NADH dehydrogenase and bacterial lipopolysaccharides (LPSs), particularly the O antigen (reviewed in REFS 17, 18). Furthermore, many *Stenotrophomonas* spp.

can produce extracellular enzymes (proteases, lipases, nucleases, chitinases and elastases) that have been shown to be important in plant colonization by other rhizosphere microorganisms.

S. maltophilia produces various pili or fimbriae that have been implicated both in adhesion to surfaces³³ and in the formation of complex biofilms. Both adhesion and biofilm formation might contribute to the ability of *S. maltophilia* to compete with other microorganisms on the surface of plant roots, and are certainly important for the colonization of medical devices that leads to infection in humans³⁴. de Oliveira-Garcia and colleagues³³ characterized SMF1 fimbriae from *S. maltophilia* strains SMDP92 and ATCC 1363. SMF1 fimbriae are composed of a 17 kDa fimbrin subunit that shares significant amino-terminal amino acid sequence similarity to the CupA fimbriae of *Pseudomonas aeruginosa* and to several fimbriae from pathogenic *Escherichia coli*. All of the clinical *S. maltophilia* isolates that were tested produced the 17 kDa fimbrin. The genomes of *S. maltophilia* strains K279a and R551-3 contain genes that encode type I pili (Sterm_0582-85, Sterm_1304-09 and Sterm_2358-66; based on R551-3 genome annotation), which have been implicated in adhesion and the early stages of biofilm formation, and type IV pili (Sterm_1417-22 and Sterm_3223-26), which have been implicated in adherence, auto-aggregation, twitching motility and biofilm formation. These gene clusters are distributed throughout each genome of the sequenced strains in a similar manner, which may indicate that there are some similarities in the plant and animal colonization strategies. Both *S. maltophilia* SMDP92 and ATCC 1363 carry *manA*, *manB*, *rmlB*, *rmlA*, *rmlC* and *rmlD* (also known as Stemr_0515-30), which encode enzymes involved in the biosynthesis of LPS and exopolysaccharides. Mutations in *manA*, *rmlA* and *rmlC* affect biofilm formation and twitching motility in *S. maltophilia* WR-C³⁵.

Epiphytic *Stenotrophomonas* bacteria can also alter the properties of the leaf surface to which they attach. Schreiber *et al.*³⁶ have demonstrated that *S. maltophilia* SaO5sm can increase the water permeability of *Hedera* and *Prunus* cuticles, which in turn should increase the availability of water and dissolved compounds in the phyllosphere, thereby enhancing the environmental growth conditions for the bacteria. The molecular mechanism (or mechanisms) responsible for the observed effects is as yet unknown, although it has been proposed that extracellular enzymes that degrade the cutin polymer and/or plasticizers, such as biosurfactants, could be responsible³⁷.

The synthesis of compatible solutes by bacteria contributes to their survival under the changing osmolarities that occur in the rhizosphere³⁸. *S. maltophilia* accumulates trehalose as the only compatible solute, whereas *S. rhizophila* produces glucosylglycerol in addition to trehalose^{39,40}. These sugars often accumulate intracellularly and protect against various stresses⁴¹. *S. maltophilia* K279a encodes the Smlt2757 and Smlt2759 proteins, which are involved in the biosynthesis of trehalose through the degradation of

Intraspecific heterogeneity

The quality of being diverse within a single species.

Rhizosphere

The zone around roots that is influenced by the plant and is a region of high microbial activity.

Endophytic

A microorganism that lives within a plant for at least part of its life cycle without causing apparent disease.

Epiphytic

Describes a bacterium that grows on or attaches to the surface of a living plant.

Phyllosphere

The micro-environment on the leaf surface of a plant.

Compatible solute

An organic compound that acts as a cytoplasmic solute to regulate water content for bacterial cells growing in environments of high osmolarity.




Figure 2 | Phylogenetic analysis of the eight validly described *Stenotrophomonas* species and related taxa. This neighbour-joining tree illustrates the maximum likelihood based on 16S rDNA sequences of *Stenotrophomonas* species and related taxa. Bootstrap values for >50% of 100 replicates (maximum likelihood/neighbour joining) are shown at the corresponding nodes. The scale bar represents 0.01 changes per nucleotide position.

glycogen by the TreY–TreZ pathway⁴². Genes encoding the enzymes in a second pathway for trehalose biosynthesis that involves the conversion of maltose (the trehalose synthase pathway), are carried on another region of the *S. maltophilia* K279a genome, which is absent from *S. maltophilia* R551-3 (Supplementary information S2 (table)). Both *S. maltophilia* K279a and *S. maltophilia* R551-3 can also produce trehalose from glucose through the trehalose-6-phosphate synthase–trehalose-6-phosphate phosphatase pathway (encoded by *otsA* and *otsB*). However, unlike the clinical isolate *S. maltophilia* K279a, the environmental isolate *S. maltophilia* R551-3 does not encode the pathway to use trehalose and therefore cannot use this sugar as a sole carbon source³².

S. maltophilia might also have the capacity to protect itself from predation by soil protozoa, which could confer a selective advantage over other bacteria. The genome of the environmental isolate *S. maltophilia* R551-3 contains a cluster of six genes (Stemr_2139-44), which is absent from the *S. maltophilia* K279a isolate (Supplementary information S3 (table)). The *rebA–C* genes encode refractile inclusion bodies, known as R bodies, which

are toxic to sensitive species of *Paramecium*⁴³, a genus of unicellular ciliate protozoa that live in freshwater environments. As similar *rebA* and *rebB* gene clusters are found in *Xanthomonas axonopodis* pv. *citri* str. 306 and in *Shewanella denitrificans* OS217, these proteins could conceivably have a role in the defence of bacteria against predation by protozoa in the rhizosphere or bulk soil.

Biotechnological uses of *Stenotrophomonas* spp.

Stenotrophomonas spp. are promising candidates for biotechnological applications in agriculture; treatment with *Stenotrophomonas* spp. can result in enhanced plant growth and can influence plant development on marginal soil (FIG. 1). For example, plant growth promotion of up to 180% was observed for wheat, tomato, lettuce, sweet pepper, melon, celery and carrot in the highly salinated soils of Uzbekistan¹⁷. *Stenotrophomonas* spp. also have promising applications in bioremediation and phytoremediation⁴⁴, as these bacteria can metabolize a large range of organic compounds that are present in the rhizosphere, including phenolic compounds found in plant root exudates. Accordingly, *S. maltophilia* can degrade *p*-nitrophenol and 4-chlorophenol⁴⁵, polycyclic aromatic hydrocarbons⁴⁶,

R body

A bacterial inclusion consisting of long proteinaceous ribbons rolled up inside the bacterial cell.

Phenolic compound

A chemical compound that is characterized by the presence of a hydroxyl group attached to a six-membered aromatic ring; many plants produce phenolic compounds.




Figure 3 | Root and endosphere colonization by *Stenotrophomonas rhizophila*. Volume renderings of confocal laser scanning micrographs of DsRed-labelled *Stenotrophomonas rhizophila* DSM 14405^T cells in the rhizosphere of tomato plants. **a** | Colonization of 1-week-old tomato roots. The colonization pattern is characterized by small colonies that mainly grow inside the epidermal cells and the root hairs. **b, c** | Magnifications of the regions selected by the continuous line and dotted line, respectively, in part **a**, observed from a different angle. **d** | Seven *S. rhizophila* DSM 14405^T cells forming a cluster inside a new tomato root hair. **e–g** | Cutting planes of the image in panel **d** on the x, y and z axes, respectively. The scale bars represent 40 μm (**a**), 3 μm (**b**), 20 μm (**c**) and 5 μm (**d–g**). Images courtesy of C. Zachow, Graz University of Technology, Austria.

selenium compounds⁴⁷, benzene, toluene, ethylbenzene⁴⁸ and xenobiotics^{44,48}. These broad metabolic properties could provide plants with protection against the phytotoxic effects of these compounds. *Stenotrophomonas* spp. can enhance plant productivity by several mechanisms, including the production of the plant growth hormone indole-3-acetic acid (IAA)⁴⁹, nitrogen fixation^{21,50} and the oxidation of elemental sulphur, which in turn provides sulphate for the plants¹⁹. Examples of how *Stenotrophomonas* spp. have been used in such applications are provided in TABLE 1. In the following sections, we detail some of the traits of *Stenotrophomonas* spp. that make them amenable to such applications.

Metal tolerance. Many *S. maltophilia* strains have intrinsic resistance to various heavy metals^{51,52}. For example, the *S. maltophilia* strains Sm777 and D457R have been shown to tolerate various toxic metals, such as cadmium, lead, cobalt, zinc, mercury and silver. When tested in tenfold diluted tryptic soy broth, strain Sm777 is additionally tolerant to 50 mM selenite, 25 mM tellurite and 50 mM uranyl salts. These properties of *S. maltophilia* have the potential to be exploited for bioremediation purposes or to aid phytoremediation. Furthermore, the tolerance of *S. maltophilia* strains to heavy metals could be useful in the bioremediation of soils that are polluted with heavy metals and xenobiotics.

A different complement of genes that specify metal tolerance has been identified in the *S. maltophilia* R551-3 and K279a genomes (Supplementary information S4 (table)). Genes coding for copper and mercury resistance, located on a genomic island in *S. maltophilia* K279a, were absent from the *S. maltophilia* R551-3 genome. Conversely, loci coding for arsenic resistance (*ars*, Stemr_2020–2024) and two tellurium resistance proteins (Stemr_2893–94) were identified in the *S. maltophilia* R551-3 genome but were absent from the *S. maltophilia* K279a genome.

Biocontrol properties. The potential of *Stenotrophomonas* spp. for the biocontrol of oomycete, fungal and bacterial pathogens has been demonstrated in several systems that include both monocotyledonous and dicotyledonous crops as hosts^{53–61} (examples of these are provided in TABLE 1). *Stenotrophomonas* spp. can prevent the growth or activity of plant pathogens by several different mechanisms. *In vitro*, most *S. maltophilia* isolates have antifungal activity⁷, and the new antifungal compounds maltophilin⁶² and xanthobaccin⁶⁰ have been described. *Stenotrophomonas* spp. also produce volatile organic compounds (VOCs) with antifungal activity^{63–65} (see below). *S. maltophilia* strains have an extraordinarily high hydrolytic potential; they produce diverse proteases, chitinases, glucanases, DNases, RNases, lipases and laccases^{23,66,67}. Both chitinolytic and proteolytic activities contribute to the biocontrol activity of *S. maltophilia*^{61,68–70}. Chitinases might protect plants against fungal pathogens through fungal cell wall lysis but might also have a role in triggering plant defence mechanisms⁷¹. A chitinase from *S. maltophilia* strain C5 was shown to suppress summer patch disease (caused by *Magnaporthe poae* Lanschoot and Jackson) in Kentucky bluegrass by the activation of disease resistance genes⁷². The precise roles of the other exoenzymes in the antagonistic activity of *S. maltophilia* remain to be elucidated and it is therefore likely that further bacterial products also control plant disease by inducing plant defences.

Another factor that is important for the control of fungal infection is competition for iron. *Stenotrophomonas* cells can efficiently capture siderophores that are produced by other microorganisms⁷³, such as ferrichrome, which is produced by fungi of several genera, including the phytopathogenic genus *Ustilago*⁷⁴. Both of the sequenced *S. maltophilia* genomes encode many TonB-dependent receptors (TBDRs), outer membrane proteins that

Xenobiotic

A chemical that is only man-made, and otherwise is not found in the environment.

Biocontrol

The control of harmful pests and pathogens through the use of microorganisms.

Siderophore

A small organic molecule that is produced by bacteria to sequester iron.

Table 1 | Applications of *Stenotrophomonas* spp. in biocontrol

Strain	Pathogen and/or plant species	Refs
<i>Stenotrophomonas maltophilia</i>	Soil-borne pathogen	55
<i>S. maltophilia</i>	<i>Rhizoctonia solani</i> damping-off in bark compost media	58
<i>S. maltophilia</i> 3089	<i>Verticillium dahliae</i> ; oilseed rape	54
<i>S. maltophilia</i> 34S1	Summer patch disease of turf grass	57,72
<i>S. maltophilia</i> C3*	Biocontrol of brown patch disease	56
<i>Stenotrophomonas</i> sp. strain SB-K88	Sugar beet damping-off disease	60
<i>S. maltophilia</i> C3*	<i>Bipolaris sorokiniana</i> ; tall fescue	61
<i>S. maltophilia</i> W81 mutant with overproduction of an extracellular serine protease	Biological control of <i>Pythium ultimum</i>	54
<i>S. maltophilia</i> 18	<i>Fusarium graminearum</i> ; wheat	155
<i>S. maltophilia</i> RAY132	Sulphur-oxidizing PGPR; canola	19
<i>S. maltophilia</i> PD3533	<i>Ralstonia solanacearum</i> race 3 biovar 2; potato	59
<i>S. maltophilia</i> R551-3	Endophytic strain isolated from poplar	32

*In a series of papers, the potential of *S. maltophilia* to control plant pathogens was described^{56,61}. Later, a taxonomic study published by Sullivan *et al.* showed that this strain belongs to *Lysbacter enzymogenes*¹⁵⁶. PGPR, plant growth promoting rhizobacteria.

are primarily known for the active transport of iron–siderophore complexes in Gram-negative bacteria. In *Xanthomonas campestris*, different subsets of TBDRs mediate iron uptake and the use of plant carbohydrates⁷⁵. There are major differences, however, in the complement of genes encoding TBDRs; the *S. maltophilia* R551-3 genome carries 82 such genes compared with 65 in the *S. maltophilia* K279a genome. Many of the genes that are present in *S. maltophilia* R551-3 but absent from *S. maltophilia* K279a are linked to genes encoding proteins that are involved in iron transport and siderophore uptake. Such an overrepresentation of TBDRs is found in only a limited number of organisms, but is common in *Xanthomonas* spp. and in aquatic bacteria that scavenge complex carbohydrates⁷⁵. Although both *S. maltophilia* R551-3 and K279a produce the siderophore enterobactin, this additional capacity for iron uptake suggests that iron competition with other organisms for endophytic (or rhizospheric) growth is important.

Bioactive natural products. Many plant-associated bacteria are well known for their diverse range of secondary metabolic products, including antibiotics, anticancer compounds, VOCs and antifungal, antiviral, insecticidal and immunosuppressant agents. *S. maltophilia* strains also produce bioactive compounds, including antibiotics and enzymes^{76,77}. Several proteases produced by *Stenotrophomonas* spp. are so much more effective than those currently in use in industry that it is thought these proteases could ‘revolutionize’ washing agents⁷⁸. A selection of these compounds has been highlighted in BOX 1. Although a wide range of biologically active compounds has been isolated from *Stenotrophomonas* spp., these organisms still remain an untapped source of novel natural products.

VOCs. The VOCs that are produced by *Stenotrophomonas* spp. can also negatively influence fungal growth and serve as inter- and intra-organismic communication signals^{64,65}. Many different VOCs that are produced by *S. maltophilia* and *S. rhizophila* inhibit mycelial growth of the soil-borne pathogen *Rhizoctonia solani* by more than 90%⁶³. Two of these VOCs have been characterized as β -phenylethanol and dodecanal. The precise mode of action of these secondary metabolites on the target organism is not known. By contrast, VOCs such as acetone, 2-methyl-1-butanol, heptanal and octanal, which are produced by *Trichoderma* spp., are known to reduce fungal growth by inhibiting protein synthesis⁷⁸.

Antimicrobial resistance

Many strains of *S. maltophilia* are also well known for their multiple antibiotic resistance phenotypes⁷, which is consistent with the elevated antibiotic and bactericidal selection pressure that is found in their biotopes⁷⁹. Multiple antibiotic resistance could help *S. maltophilia* to compete in the rhizosphere^{3,79,80}, which supports intense microbiological activity and competition in comparison to the nutrient-limited bulk soil. Comparative genomic analysis and experimental testing of the clinical and endophytic *S. maltophilia* isolates K279a and R551-3 showed that many of the genes that encode antimicrobial drug resistance and resistance-nodulation-division (RND) family transporters (tripartite efflux pumps that are involved in antibiotic resistance) are conserved (Supplementary information S4 (table)). In addition, the *S. maltophilia* R551-3 genome carries two additional RND operons (Stemr_2065-67 and Stemr_951-52) that are putatively involved in antibiotic efflux, a macrolide ATP-binding cassette-type transporter (Stemr_2509-10) and three putative major facilitator superfamily antibiotic transporters (Stemr_3598, Stemr_3605 and Stemr_3603, the last of which is putatively involved in chloramphenicol transport). The presence of these genes indicates that the endophytic and clinical strains have a similar level of antibiotic resistance, with possibly an even broader resistance spectrum for the endophytic strain *S. maltophilia* R551-3. In both strains, most antibiotic resistance genes are not associated with mobile genetic elements, such as phages or transposons, which makes it unlikely that *S. maltophilia* K279a acquired its antibiotic resistance genes in the clinical environment.

Adaptation and metabolic versatility

Stenotrophomonas spp. can efficiently colonize such different biotopes as plants, humans and marine environments. Comparative studies of the recently determined genome sequence of the endophytic *S. maltophilia* strain R551-3 with that of the clinical isolate *S. maltophilia* K279a have provided insight into functions that could be associated with adaptation to these different niches. Approximately 85% of the 4,175 *S. maltophilia* R551-3 genes are homologous to genes from *S. maltophilia* K279a, and have the same organization in both strains (FIG. 4). This indicates that the ancestral *S. maltophilia* core genome had to be well equipped to allow survival, colonization and competition for resources in such a

Biotope

The natural environment of a microorganism.

Box 1 | *Stenotrophomonas* species as a source of useful compounds and activities

Stenotrophomonas spp. have promising applications in bioremediation and phytoremediation⁴⁴ as a result of their ability to metabolize many of the compounds that are present in the rhizosphere, including xenobiotics. A selection of the xenobiotic compounds that *Stenotrophomonas* spp. can degrade is provided in the table. Strains of *Stenotrophomonas maltophilia* could be exploited in the bioremediation of polluted soils owing to their intrinsic resistance to heavy metals. *Stenotrophomonas* spp. are also capable of biocatalysis, including the conversion of linoleic acid to 10-hydroxy-12(Z)-octadecenoic acid.

Useful compound or activity	Refs
Degradation of xenobiotic compounds	
Biodegradation of <i>p</i> -nitrophenol, 4-chlorophenol and 4-nitroaniline	139
<i>N</i> -demethylation of the insecticide acetamiprid	140
Biodegradation of nonylphenol	141
Degradation of polypropylene glycols	142
Decolourization of synthetic dyes	67
Degradation of keratin	143
Degradation of herbicides 4-(2,4-dichlorophenoxy) butyric acid and 4-(4-chloro-2-methylphenoxy) butyric acid	144
Heavy metal tolerance and remediation	
Accumulation of intracellular uranium	145
Tolerance of heavy metals	52
Removal of Au(III) from contaminated waste water	146
Cr(VI) resistance	147
Biocatalysis and synthesis of novel compounds	
Conversion of linoleic acid to 10-hydroxy-12(Z)-octadecenoic acid	148
7 α -OH epimerization of bile acids	149
Hydroxysteroid dehydrogenase	150
Cholylglycine hydrolase	150
Aminopeptidase activity against 4-hydroxyproline-containing peptides	151
Synthesis of glucosylglycerol	39
Saccharification of alginate	152
Synthesis of terephthalic acid	153
Production of the glycosidase inhibitor valienamine from validomycin A	154
Biocontrol and antimicrobial activity	
Production of antifungal xanthobaccins and the macrocyclic lactam antibiotic maltophilin	23,61

wide range of biotopes. Divergence in the gene complement of the two organisms might reflect adaptation to specific niches, and in the following sections we highlight some of these differences.

The *S. maltophilia* R551-3 genome contains 27 regions that are absent from *S. maltophilia* K279a; the size of these regions ranges from 1.7 kb to 61.5 kb (Supplementary information S2,S3 (tables)). Six of these regions represent putative genomic islands, as characterized by different codon usage from the rest of the genome, location next to a tRNA gene and/or the presence of a flanking integrase gene. By contrast, the genome of *S. maltophilia* K279a contains 19 regions that are absent from *S. maltophilia* R551-3 (Supplementary information S2,S3 (tables)), five of which are putative genomic islands⁸⁰. These islands are often inserted at the same positions in both the *S. maltophilia* R551-3 and K279a genomes, which points to the existence of

insertion hotspots in the core genome of this species (FIG. 5). In addition to genomic islands, the *S. maltophilia* R551-3 genome carries 14 complete insertion sequence (IS) elements: five intact copies of the IS481 family, three intact copies of the IS3 (also known as IS150) family (identical to ISPsy9 from K279a) and six intact copies of the IS110 family. Two additional truncated transposases of the IS110 family were also identified. In addition, two phages were found, one of which (Stemr_910-36) also occurs in *S. maltophilia* K279a.

Filamentous haemagglutinin proteins are adhesins that are secreted by the type V protein secretion pathway and are involved in cell–cell aggregation. These filamentous haemagglutinins have also been shown to mediate contact between the phytopathogen *Xylella fastidiosa* and plant cells⁸¹ and are important virulence factors that are involved in the adhesion of *Bordetella pertussis* to mammalian host cells⁸². *S. maltophilia* R551-3




Figure 4 | Genome maps of *Stenotrophomonas maltophilia* R551-3 and K279a. Genome maps of the poplar endophyte *S. maltophilia* R551-3 (a) and of the opportunistic pathogen *S. maltophilia* K279a (b) are shown. From the outside in, the circles represent coordinates in kilobase pairs (kbp), %GC content, predicted open reading frames (ORFs) in the clockwise and anticlockwise orientations, GC skew ((G-C and G+C) in a 1,000-bp window), transposable elements (pink) and pseudogenes (grey), and the putative *S. maltophilia* K279a genomic islands (red).

encodes three filamentous haemagglutinins (Stemr_0113, Stemr_2248 and Stemr_2356). One of these (Stemr_0113) has a low level of amino acid sequence similarity (only 53% identity) with its *S. maltophilia* K279a homologue (*shlA*, which encodes *Smlt1390*) compared with the average similarity of proteins that are shared by the two strains. The *S. maltophilia* K279a haemagglutinin is located with five haemagglutinin open reading frame fragments. Another region that is specific to *S. maltophilia* K279a and is located on a putative complex transposon carries genes that encode an adhesin and a type IV pilus as well as a peptidase (Supplementary information S2 (table)). All these differences between *S. maltophilia* K279a and *S. maltophilia* R551-3 could be related to niche adaptation or host preference.

A key component of the bacterial outer membrane is LPS, and changes in LPS structure have been correlated with changes in resistance to various antimicrobial agents. Many studies have examined the LPS in *S. maltophilia* in an effort to assess its contribution to antimicrobial resistance in this organism^{83,84} and as a basis for serotyping. As outlined above, mutations in *manA*, *rmlA* and *rmlC* affect LPS structure in *S. maltophilia* WR-C³⁵. Adjacent to these genes in the sequenced genomes is a locus that is also probably involved in LPS biosynthesis and that shows considerable variability between the two *S. maltophilia* strains (Supplementary information S5 (figure)). This locus is flanked by the genes encoding cystathionine gamma lyase (*metB*) at one end and an electron transport flavoprotein (*etfA*) at the other. An equivalently located, highly variable LPS locus has been described in a range of xanthomonads that infect rice, citrus and crucifers⁸⁵ and was probably acquired by horizontal gene transfer⁸⁵. Differences in the complement and nature of the genes in this locus are indicative of alterations in the structure of LPS, particularly the O antigen moiety. Serotyping of heat-stable O antigens from *S. maltophilia* has revealed a considerable level of variation between isolates, with 31 defined serotypes. The structure of the O antigen polysaccharides has been described for a number of these serotypes. Most of the polymers have branched repeating units, often with lateral pentosyl substitutions⁸⁶.

Variation in LPS biosynthetic gene clusters between strains is common in bacterial pathogens of animals, for which it might have a role in evading the host immune system. The role of differences in O antigen or LPS structure between related plant pathogenic or plant-associated bacteria is less certain, although involvement in host-range selection and specificity has been proposed⁸⁷. Serotype analysis of a range of *S. maltophilia* strains, however, shows no clear delineation between clinical and environmental isolates. Furthermore, other roles for variations in O antigen structure, such as promoting insensitivity to phage infection, should not be overlooked.

The endophytic strain *S. maltophilia* R551-3, which has no direct plant growth promoting effects on its poplar host, does not carry genes for the plant growth promotion mechanisms that have been described for other endophytes, such as metabolism of the plant signal molecules γ -amino butyric acid and phenyl acetic acid, degradation of the ethylene precursor 1-aminocyclopropane-1-carboxylic acid and acetoin production. In addition, only low levels of IAA are produced by *S. maltophilia* R551-3 (REF. 32).

Although *S. maltophilia* is related to plant pathogenic *Xanthomonas* spp., it is not a phytopathogen and accordingly neither the *S. maltophilia* K279a nor R551-3 genomes encode genes that are known to contribute to *Xanthomonas* virulence, including those encoding the type III secretion system and genes encoding certain plant cell wall degrading enzymes, such as pectinases. These sequences also do not seem to encode the virulence factor type VI secretion system. In addition, neither genome carries the *zonula occludens* toxin (*zot*) gene that was identified in several clinical isolates of *S. maltophilia*⁸⁸. This gene is similar to the *zot* gene in *Vibrio cholerae*, which encodes the major virulence factor enterotoxin⁸⁸. However, genes that are involved in phytopathogenesis, including those encoding enzymes such as cellulase (glycosyl hydrolase family 5) and those encoding type I, type II (*Sec*), type IV, type V and the twin arginine transporter (TAT) secretion systems, are present. Functional genomic analysis will allow an




Figure 5 | **Comparison of gene content and organization in the genomes of *Stenotrophomonas maltophilia* R551-3 and K279a.** The genomes were compared using LinePlot from [MaGe](#) (see Further information). Syntenic regions between *S. maltophilia* R551-3 and K279a are displayed in purple, and the location and putative functions of selected genes within regions that are specific to each strain are indicated. The green boxes represent putative genomic islands whereas the blue boxes represent virulence genes. LPS, lipopolysaccharide.

investigation of the factors that are important for the association of *Stenotrophomonas* spp. with plants.

***S. maltophilia*: an opportunistic pathogen**

S. maltophilia is the only species of *Stenotrophomonas* that is known to cause human disease⁵, but there is considerable phylogenetic and phenotypic variability among *S. maltophilia* isolates, including those from patients in a single hospital^{89–91}. This is probably the result of the many environmental niches of this bacterium; most infections are likely to reflect contact with separate environmental sources. Indeed, there are few instances of outbreaks of *S. maltophilia*, and those that occur are caused by a single contaminated source, such as a water source^{92–95}. Despite this, there is evidence that certain phylogenetic groups are better able to cause infection than others⁹⁶.

S. maltophilia is almost exclusively a hospital-acquired pathogen and has been associated with bacteraemic infections and pneumonia, both with a high rate of mortality⁹⁷, in immunocompromised patients. This reflects a requirement for three major risk factors for infection: severe debilitation and/or neutropenia; the presence of indwelling devices such as ventilator tubes and/or intravenous catheters for prolonged periods; and multiple and/or prolonged courses of broad-spectrum antimicrobial drugs^{98–100}. There are also examples of soft tissue, ocular, wound and burn infections, and endocarditis¹⁸.

The bacteraemia isolate *S. maltophilia* K279a carries several genes that encode factors that could allow this strain to adhere to surfaces and form biofilms, which are both key factors in the colonization of indwelling devices. Studies have shown that *S. maltophilia* isolates have cytotoxic effects *in vitro* against HEp-2, Vero and

HeLa cell lines after 24 hours¹⁰¹, and *S. maltophilia* K279a killed almost all of the N2 *Caenorhabditis elegans* in the assay within 24 hours¹⁰². However, as one might expect of a true opportunist, *S. maltophilia* has no type III secretion system⁸⁰.

A recent study by Waters and colleagues¹⁰³ tried to address the lack of documentation of the potential of *S. maltophilia* for virulence by investigating the immunostimulatory properties of 24 *S. maltophilia* clinical respiratory and non-respiratory isolates (from blood, skin and soft tissue). In this study, which involved a neonatal mouse model of pneumonia and macrophage cell lines, they determined the rates of pneumonia, bacteraemia and mortality, as well as the inflammatory response that is elicited by *S. maltophilia* infection. They demonstrated that the respiratory and non-respiratory *S. maltophilia* isolates were highly immunostimulatory but weakly invasive, which indicates that these isolates could contribute to airway inflammation.

Whether *S. maltophilia* clinical isolates are colonizers or true pathogens is still controversial in some cases. This is particularly the case for pneumonia, because it is rare to culture pure *S. maltophilia* from the lungs, and severely debilitated patients are often colonized asymptotically^{104,105}. Nonetheless, a recent study showed that 4.5% of nosocomial pneumonia in patients in intensive care units and 6% of ventilator-associated pneumonias are caused by *S. maltophilia*¹⁰⁶. The attributable mortality for *S. maltophilia* pneumonia (as far as this can be accurately quantified in patients with multiple and complex pathologies) is estimated at 20–30%¹⁰⁷. As many as 25% of adult patients with cystic fibrosis carry *S. maltophilia* in their lungs at any one time, although these numbers are highly variable¹⁰⁸. Also, there is no evidence of any reduction in lung function associated

with *S. maltophilia* colonization¹⁰⁹. Although the rates are variable, approximately 1% of all nosocomial bacteraemias are caused by *S. maltophilia* and the attributed mortality has been estimated at approximately 25%¹¹⁰. These bacteraemias are usually caused by an indwelling device, and the prognosis is usually good following the removal of the device^{111–113}.

The primary reason for the increase in *S. maltophilia* infections is the intrinsic resistance of this species to many front-line antimicrobials, such as β -lactams, including carbapenems¹¹⁴, aminoglycosides (except gentamicin)^{48,80,115,116}, macrolides, tetracycline, chloramphenicol and older quinolones^{80,117–119}. Furthermore, *S. maltophilia* isolates can rapidly develop resistance to newer fluoroquinolones, gentamicin and minocycline through mutation; the underlying mechanisms are not certain, but are likely to be the result of the overproduction of intrinsic efflux pumps⁸⁹. Typically, empiric therapy for *S. maltophilia* is trimethoprim–sulphamethoxazole (TMP–SMX), to which >95% of isolates are sensitive^{120,121}. However, resistance is increasing as a result of the spread of acquired mobile resistance determinants^{122,123} and, in many patients, TMP–SMX therapy is contra-indicated¹²⁴. Alternatives include ticarcillin–clavulanate, respiratory quinolones such as gatifloxacin and moxifloxacin, minocycline and possibly tigecycline^{120,125}. For how long these drugs would remain clinically efficacious if they were used as front-line therapy is uncertain, however, as resistant mutants arise spontaneously at high rates *in vitro*⁹⁶ (V. C. Gould and M.B.A., unpublished observations).

The *de novo* mobilization of antibiotic resistance genes from environmental bacteria or opportunistic pathogens has contributed greatly to the increase in antibiotic resistance in more common and important pathogens. Examples include the mobilization of the *SHV* and *CTX-M* β -lactamase genes from *Klebsiella pneumoniae* and *Kluyvera* spp., respectively, on composite transposons that are likely to have formed since the clinical use of antibiotics began^{126,127}. One potential consequence of the increased colonization of patients by *S. maltophilia* is the mobilization of its L1 metallo- β -lactamase gene onto a plasmid that can then confer broad-spectrum β -lactam resistance, including to the last-line carbapenems, in *P. aeruginosa* and even the Enterobacteriaceae. Another possibility is the mobilization of a potential quinolone resistance determinant, *qnr*, that was identified in the *S. maltophilia* K279a¹²⁸ and R551-3 genome sequences (Supplementary information S4 (table)). Resistance plasmid and transposon carriage has been demonstrated in *S. maltophilia*, as has sharing of these elements with *E. coli in vitro*^{123,129–131}, suggesting that it might be only a matter of time before L1 mobilization occurs, which would be a hugely significant event.

Polymicrobial infections and interspecies signalling.

There is an increasing appreciation of the polymicrobial nature of human infections and of the potentially important role for interspecies interactions in bacterial virulence and the response to therapy. *S. maltophilia* can be found together with the opportunistic pathogen *P. aeruginosa* in diverse niches, including the lungs of patients

with cystic fibrosis, where *S. maltophilia* can protect antibiotic-sensitive strains of *P. aeruginosa* by degrading antibiotics¹³². Interspecies signalling between these organisms can also influence *P. aeruginosa*. *S. maltophilia* possesses a cell–cell signalling system that is mediated by a diffusible signal factor (DSF) that was first identified in the related plant pathogen *X. campestris*^{102,133}. In *S. maltophilia* strain WR-C the DSF can be one of eight structurally related fatty acids that include *cis*-11-methyl-2-dodecenoic acid, the DSF signal from *X. campestris*^{133,134}. Genome analyses revealed that *S. maltophilia* does not synthesize *N*-acyl homoserine lactones (*N*-AHLs) or autoinducer 2 (AI2), which are signal molecules that are commonly found in other Gram-negative organisms, and *N*-AHLs have not been detected in *S. maltophilia* cultures^{135,136}. In the clinical isolate *S. maltophilia* K279a, DSF signalling controls several functions, including the production of an extracellular protease, aggregative behaviour and virulence in a nematode model¹⁰². Although *P. aeruginosa* does not synthesize DSF, it can respond to the signal that is produced by *S. maltophilia* by altering biofilm architecture and increasing tolerance to the cationic antimicrobial peptides polymyxin B and colistin^{137,138}. This response of *P. aeruginosa* to DSF requires PA1396, a sensor kinase with an input domain that is related to the sensory input domain of RpfC, which is responsible for DSF perception in xanthomonads. Homologues of PA1396 occur in other pseudomonads, some of which are plant pathogenic or plant associated, as well as in unrelated bacteria, including the poplar endophyte *Enterobacter* sp. 638. These observations indicate that modulation of bacterial behaviour through DSF-mediated interspecies signalling with *S. maltophilia* is a phenomenon that could also occur in a non-pathogenic context in rhizospheric or endophytic communities.

Implications for biotechnological uses. The application of rhizospheric and endophytic *S. maltophilia* strains to control plant pathogens⁵⁹ or promote plant health should be carefully considered. Not only do these strains present a risk as reservoirs for antibiotic resistance genes, but they also have potential as opportunistic pathogens. It is known that clinical isolates of *S. maltophilia* and other opportunistic pathogens can actively multiply in the rhizosphere⁴⁹. This has led to suggestions that the rhizosphere is an environmental reservoir for such opportunists. Furthermore, the mechanisms responsible for the colonization of plants and for the antagonistic activity of *S. maltophilia* strains against plant pathogens are similar to those that are responsible for the colonization of human tissues and for pathogenicity⁷⁹. Such considerations should perhaps be translated into appropriate measures, such as banning plants from hospitals.

Concluding remarks

A fuller understanding of the versatility, adaptation and potential uses of this fascinating group of organisms presents both considerable challenges and opportunities for the future. Determination of the genome sequences of the clinical and endophytic *S. maltophilia* strains forms the basis for functional genomic analyses

Riboswitch

A conformational switch in RNA molecules that is induced by small metabolites and leads to a switch in gene regulatory function.

to test the contribution of specific functions to the tenacity of these bacteria in colonization, their broad resistance to antibiotics and their ability to enter into close endophytic associations with plants. These studies might also be expected to reveal alternative targets for antimicrobial action that could include interference with quorum sensing or cell–cell signalling, interference with the action of riboswitches and inhibition of adherence to surfaces and biofilm formation. Given the ubiquitous nature of *Stenotrophomonas* spp., the impact of interactions with other organisms in polymicrobial communities, which may be associated with

infections or endophytic colonization, merits attention. Further insights into the role of *Stenotrophomonas* spp. in the nitrogen cycle should emerge from analysis of the genome of *Stenotrophomonas* sp. SKA14, a nitrogen-fixing bacterium that was isolated from the Baltic Sea. The sequencing of other *Stenotrophomonas* spp., such as *S. rhizophila*, that have potential uses in promoting plant growth, biocontrol or bioremediation but are not pathogenic are also warranted. Finally, *Stenotrophomonas* spp. should continue to be a source of useful or novel enzymatic capabilities, reflecting their metabolic versatility.

- Palleroni, N. J. & Bradbury, J. F. *Stenotrophomonas*, a new bacterial genus for *Xanthomonas maltophilia* (Hugh 1980) Swings. *et al.* 1983. *Int. J. Syst. Bacteriol.* **43**, 606–609 (1993).
- Swings, J., Devos, P., Vandenmooter, M. & Deley, J. Transfer of *Pseudomonas maltophilia* Hugh 1981 to the genus *Xanthomonas maltophilia* (Hugh 1981) comb. nov. *Int. J. Syst. Bacteriol.* **33**, 409–413 (1983).
- Berg, G., Roskot, N. & Smalla, K. Genotypic and phenotypic relationships between clinical and environmental isolates of *Stenotrophomonas maltophilia*. *J. Clin. Microbiol.* **37**, 3594–3600 (1999).
- Chatelut, M., Dournes, J. L., Chabanon, G. & Marty, N. Epidemiologic typing of *Stenotrophomonas (Xanthomonas) maltophilia* by PCR. *J. Clin. Microbiol.* **33**, 912–914 (1995).
- Coenye, T., Vanlaere, E., Falsen, E. & Vandamme, P. *Stenotrophomonas africana* Drancourt. *et al.* 1997 is a later synonym of *Stenotrophomonas maltophilia* (Hugh 1981) Palleroni and Bradbury 1993. *Int. J. Syst. Evol. Microbiol.* **54**, 1235–1237 (2004).
- Hauben, L., Vauterin, L., Moore, E. R. B., Hoste, B. & Swings, J. Genomic diversity of the genus *Stenotrophomonas*. *Int. J. Syst. Bacteriol.* **49**, 1749–1760 (1999).
- Minkwitz, A. & Berg, G. Comparison of antifungal activities and 16S ribosomal DNA sequences of clinical and environmental isolates of *Stenotrophomonas maltophilia*. *J. Clin. Microbiol.* **39**, 139–145 (2001).
- Nesme, X., Vanechoutte, M., Orso, S., Hoste, B. & Swings, J. Diversity and genetic relatedness within genera *Xanthomonas* and *Stenotrophomonas* using restriction-endonuclease site differences of PCR-amplified 16S ribosomal-RNA gene. *Syst. Appl. Microbiol.* **18**, 127–135 (1995).
Highlights the complexity of the *Stenotrophomonas* species and details useful methods to discern them from other xanthomonads.
- Finkmann, W., Altendorf, K., Stackebrandt, E. & Lipski, A. Characterization of N₂O-producing *Xanthomonas*-like isolates from biofilters as *Stenotrophomonas nitritireducens* sp. nov., *Luteimonas mephitis* gen. nov., sp. nov. and *Pseudoxanthomonas broegbemensis* gen. nov., sp. nov. *Int. J. Syst. Evol. Microbiol.* **50**, 273–282 (2000).
- Wolf, A., Fritze, A., Hagemann, M. & Berg, G. *Stenotrophomonas rhizophila* sp. nov., a novel plant-associated bacterium with antifungal properties. *Int. J. Syst. Evol. Microbiol.* **52**, 1937–1944 (2002).
Defines the plant-associated species *S. rhizophila*.
- Assih, E. A. *et al.* *Stenotrophomonas acidaminiphila* sp. nov., a strictly aerobic bacterium isolated from an upflow anaerobic sludge blanket (UASB) reactor. *Int. J. Syst. Evol. Microbiol.* **52**, 559–568 (2002).
- Yang, H. C., Im, W. T., Kang, M. S., Shin, D. Y. & Lee, S. T. *Stenotrophomonas koreensis* sp. nov., isolated from compost in South Korea. *Int. J. Syst. Evol. Microbiol.* **56**, 81–84 (2006).
- Kaparullina, E., Doronina, N., Chistyakova, T. & Trotsenko, Y. *Stenotrophomonas chelatiphaga* sp. nov., a new aerobic EDTA-degrading bacterium. *Syst. Appl. Microbiol.* **32**, 157–162 (2009).
- Heylen, K., Vanparrys, B., Peirsegaale, F., Lebbe, L. & De Vos, P. *Stenotrophomonas terrae* sp. nov. and *Stenotrophomonas humii* sp. nov., two nitrate-reducing bacteria isolated from soil. *Int. J. Syst. Evol. Microbiol.* **57**, 2056–2061 (2007).
- Yoon, J. H., Kang, S. J., Oh, H. W. & Oh, T. K. *Stenotrophomonas dokdonensis* sp. nov., isolated from soil. *Int. J. Syst. Evol. Microbiol.* **56**, 1363–1367 (2006).
- Lee, D. S. *et al.* *Pseudoxanthomonas sacceonensis* sp. nov., isolated from BTEX-contaminated soil in Korea, transfer of *Stenotrophomonas dokdonensis* Yoon *et al.* 2006 to the genus *Pseudoxanthomonas* as *Pseudoxanthomonas dokdonensis* comb. nov. and emended description of the genus *Pseudoxanthomonas*. *Int. J. Syst. Evol. Microbiol.* **58**, 2235–2240 (2008).
- Berg, G., Egamberdieva, D., Lugtenberg, B. & Hagemann, M. in *Symbiosis and Stress* (eds Seckbach, J. & Grube, M.) (in the press).
- Denton, M. & Kerr, K. G. Microbiological and clinical aspects of infection associated with *Stenotrophomonas maltophilia*. *Clin. Microbiol. Rev.* **11**, 57–80 (1998).
- Banerjee, M. & Yesmin, L. Sulfur-oxidizing plant growth promoting rhizobacteria for enhanced canola performance. US Patent 07491535 (2002).
- Ikemoto, S., Suzuki, K., Kaneko, T. & Komagata, K. Characterization of strains of *Pseudomonas maltophilia* which do not require methionine. *Int. J. Syst. Bacteriol.* **30**, 437–447 (1980).
- Park, M. *et al.* Isolation and characterization of diazotrophic growth promoting bacteria from rhizosphere of agricultural crops of Korea. *Microbiol. Res.* **160**, 127–133 (2005).
- Lockhart, S. R. *et al.* Antimicrobial resistance among Gram-negative bacilli causing infections in intensive care unit patients in the United States between 1993 and 2004. *J. Clin. Microbiol.* **45**, 3352–3359 (2007).
- Berg, G., Marten, P. & Ballin, G. *Stenotrophomonas maltophilia* in the rhizosphere of oilseed rape — occurrence, characterization and interaction with phytopathogenic fungi. *Microbiol. Res.* **151**, 19–27 (1996).
- Juhnke, M. E. & Desjardins, E. Selective medium for isolation of *Xanthomonas maltophilia* from soil and rhizosphere environments. *Appl. Environ. Microbiol.* **55**, 747–750 (1989).
- Mahaffee, W. F. & Klopper, J. W. Temporal changes in the bacterial communities of soil, rhizosphere, and endorhiza associated with field-grown cucumber (*Cucumis sativus* L.). *Microb. Ecol.* **34**, 210–223 (1997).
- Berg, G. *et al.* Plant-dependent genotypic and phenotypic diversity of antagonistic rhizobacteria isolated from different *Verticillium* host plants. *Appl. Environ. Microbiol.* **68**, 3328–3338 (2002).
- Schwieger, F. & Tebbe, C. C. Effect of field inoculation with *Sinorhizobium meliloti* L33 on the composition of bacterial communities in rhizospheres of a target plant (*Medicago sativa*) and a non-target plant (*Chenopodium album*) — linking of 16S rRNA gene-based single-strand conformation polymorphism community profiles to the diversity of cultivated bacteria. *Appl. Environ. Microbiol.* **66**, 3556–3565 (2000).
- Chelius, M. K. & Triplett, E. W. Immunolocalization of dinitrogenase reductase produced by *Klebsiella pneumoniae* in association with *Zea mays* L. *Appl. Environ. Microbiol.* **66**, 783–787 (2000).
- Mehnaz, S. *et al.* Isolation and 16S rRNA sequence analysis of the beneficial bacteria from the rhizosphere of rice. *Can. J. Microbiol.* **47**, 110–117 (2001).
- Germida, J. J. & Siciliano, S. D. Taxonomic diversity of bacteria associated with the roots of modern, recent and ancient wheat cultivars. *Biol. Fertil. Soils* **33**, 410–415 (2001).
- Sturz, A. V., Matheson, B. G., Arsenault, W., Kimpinski, J. & Christie, B. R. Weeds as a source of plant growth promoting rhizobacteria in agricultural soils. *Can. J. Microbiol.* **47**, 1013–1024 (2001).
- Taghavi, S. *et al.* Mechanisms underlying the beneficial effects of endophytic bacteria on growth and development of poplar. *Appl. Environ. Microbiol.* **75**, 748–757 (2009).
- de Oliveira-Garcia, D. *et al.* Fimbriae and adherence of *Stenotrophomonas maltophilia* to epithelial cells and to abiotic surfaces. *Cell. Microbiol.* **5**, 625–636 (2003).
- Elvers, K. T., Leeming, K. & Lappin-Scott, H. M. Binary culture biofilm formation by *Stenotrophomonas maltophilia* and *Fusarium oxysporum*. *J. Ind. Microbiol. Biotechnol.* **26**, 178–183 (2001).
- Huang, T. P., Somers, E. B. & Wong, A. C. L. Differential biofilm formation and motility associated with lipopolysaccharide/exopolysaccharide-coupled biosynthetic genes in *Stenotrophomonas maltophilia*. *J. Bacteriol.* **188**, 3116–3120 (2006).
- Schreiber, L. *et al.* Plant–microbe interactions: identification of epiphytic bacteria and their ability to alter leaf surface permeability. *New Phytol.* **166**, 589–594 (2005).
- Riederer, M. & Schonherr, J. Effects of surfactants on water permeability of isolated plant cuticles and on the composition of their cuticular waxes. *Pesticide Sci.* **29**, 85–94 (1990).
- Miller, K. J. & Wood, J. M. Osmoadaptation by rhizosphere bacteria. *Annu. Rev. Microbiol.* **50**, 101–136 (1996).
- Hagemann, M. *et al.* The plant-associated bacterium *Stenotrophomonas rhizophila* expresses a new enzyme for the synthesis of the compatible solute glucosylglycerol. *J. Bacteriol.* **190**, 5898–5906 (2008).
- Roder, A., Hoffmann, E., Hagemann, M. & Berg, G. Synthesis of the compatible solutes glucosylglycerol and trehalose by salt-stressed cells of *Stenotrophomonas* strains. *FEMS Microbiol. Lett.* **243**, 219–226 (2005).
- Elbein, A. D., Pan, Y. T., Pastuszak, I. & Carroll, D. New insights on trehalose: a multifunctional molecule. *Glycobiology* **13**, 17R–27R (2003).
- Avonce, N., Mendoza-Vargas, A., Morett, E. & Iturriaga, G. Insights on the evolution of trehalose biosynthesis. *BMC Evol. Biol.* **6**, 109 (2006).
- Heruth, D. P., Pond, F. R., Dilts, J. A. & Quackenbush, R. L. Characterization of genetic determinants for R-body synthesis and assembly in *Caedibacter taeniospiralis* 47 and 116. *J. Bacteriol.* **176**, 3559–3567 (1994).
- Binks, P. R., Nicklin, S. & Bruce, N. C. Degradation of hexahydro-1, 3, 5-triazine-1, 3, 5-triazine (RDx) by *Stenotrophomonas maltophilia* PB1. *Appl. Environ. Microbiol.* **61**, 1318–1322 (1995).
- Zhang, J. F., Zheng, Y. G., Liu, Z. Q. & Shen, Y. C. Preparation of 3-ketovalidoxylamine A C-N lyase substrate: *N*-p-nitrophenyl-3-ketovalidamine by *Stenotrophomonas maltophilia* CCTCC M 204024. *Appl. Microbiol. Biotechnol.* **73**, 1275–1281 (2007).
- Juhasz, A. L., Stanley, G. A. & Britz, M. L. Microbial degradation and detoxification of high molecular weight polycyclic aromatic hydrocarbons by *Stenotrophomonas maltophilia* strain VUN 10,003. *Lett. Appl. Microbiol.* **30**, 396–401 (2000).

47. Dungan, R. S., Yates, S. R. & Frankenberger, W. T. Transformations of selenate and selenite by *Stenotrophomonas maltophilia* isolated from a seleniferous agricultural drainage pond sediment. *Environ. Microbiol.* **5**, 287–295 (2003).
48. Lee, E. Y., Jun, Y. S., Cho, K. S. & Ryu, H. W. Degradation characteristics of toluene, benzene, ethylbenzene, and xylene by *Stenotrophomonas maltophilia* T3-c. *J. Air Waste Manage. Assoc.* **52**, 400–406 (2002).
49. Suckstorff, I. & Berg, G. Evidence for dose-dependent effects on plant growth by *Stenotrophomonas* strains from different origins. *J. Appl. Microbiol.* **95**, 656–663 (2003).
50. Liba, C. M. *et al.* Nitrogen-fixing chemo-organotrophic bacteria isolated from cyanobacteria-deprived lichens and their ability to solubilize phosphate and to release amino acids and phytohormones. *J. Appl. Microbiol.* **101**, 1076–1086 (2006).
51. Alonso, A., Sanchez, P. & Martinez, J. L. *Stenotrophomonas maltophilia* D457R contains a cluster of genes from Gram-positive bacteria involved in antibiotic and heavy metal resistance. *Antimicrob. Agents Chemother.* **44**, 1778–1782 (2000).
52. Pages, D. *et al.* Heavy metal tolerance in *Stenotrophomonas maltophilia*. *PLoS One* **3**, e1539 (2008).
53. Berg, G., Knaape, C., Ballin, G. & Seidel, D. Biological control of *Verticillium dahliae* Kleb. by natural occurring rhizosphere bacteria. *Arch. Phytopathol. Plant Protection* **29**, 249–262 (1994).
54. Dunne, C., Moenne-Loccoz, Y., de Bruijn, F. J. & O'Garra, F. Overproduction of an inducible extracellular serine protease improves biological control of *Pythium ultimum* by *Stenotrophomonas maltophilia* strain W81. *Microbiology* **146**, 2069–2078 (2000).
55. Elad, Y., Chet, I. & Baker, R. Increased growth-response of plants induced by rhizobacteria antagonistic to soilborne pathogenic fungi. *Plant Soil* **98**, 325–330 (1987).
56. Giesler, L. J. & Yuen, G. Y. Evaluation of *Stenotrophomonas maltophilia* strain C3 for biocontrol of brown patch disease. *Crop Prot.* **17**, 509–513 (1998).
57. Kobayashi, D. Y., Guglielmoni, M. & Clarke, B. B. Isolation of the chitinolytic bacteria *Xanthomonas maltophilia* and *Serratia marcescens* as biological control agents for summer patch disease of turfgrass. *Soil Biol. Biochem.* **27**, 1479–1487 (1995).
58. Kwok, O. C. H., Fahy, P. C., Hoihtink, H. A. J. & Kuter, G. A. Interactions between bacteria and *Trichoderma hamatum* in suppression of rhizoctonia damping-off in bark compost media. *Phytopathology* **77**, 1206–1212 (1987).
59. Messiha, N. A. S. *et al.* *Stenotrophomonas maltophilia*: a new potential biocontrol agent of *Ralstonia solanacearum*, causal agent of potato brown rot. *Eur. J. Plant Pathol.* **118**, 211–225 (2007).
Describes the first example of the release of *S. maltophilia* for disease control.
60. Nakayama, T., Homma, Y., Hashidoko, Y., Mizutani, J. & Tahara, S. Possible role of xanthobaccins produced by *Stenotrophomonas* sp strain SB-K88 in suppression of sugar beet damping-off disease. *Appl. Environ. Microbiol.* **65**, 4334–4339 (1999).
61. Zhang, Z. & Yuen, G. Y. Biological control of *Bipolaris sorokiniana* on tall fescue by *Stenotrophomonas maltophilia* strain C3. *Phytopathology* **89**, 817–822 (1999).
62. Jakobi, M. *et al.* Maltophilin: a new antifungal compound produced by *Stenotrophomonas maltophilia* R3089. *J. Antibiotics* **49**, 1101–1104 (1996).
63. Kai, M., Effmert, U., Berg, G. & Piechulla, B. Volatiles of bacterial antagonists inhibit mycelial growth of the plant pathogen *Rhizoctonia solani*. *Arch. Microbiol.* **187**, 351–360 (2007).
64. Stotzyk, G. & Schenck, S. Volatile organic compounds and microorganisms. *Crit. Rev. Microbiol.* **4**, 333–382 (1976).
65. Wheatley, R. E. The consequences of volatile organic compound mediated bacterial and fungal interactions. *Antonie Van Leeuwenhoek* **81**, 357–364 (2002).
66. Debette, J. Isolation and characterization of an extracellular proteinase produced by a soil strain of *Xanthomonas maltophilia*. *Curr. Microbiol.* **22**, 85–90 (1991).
67. Galai, S., Limam, F. & Marzouki, M. N. A new *Stenotrophomonas maltophilia* strain producing laccase. Use in decolorization of synthetic dyes. *Appl. Biochem. Biotechnol.* 17 Oct 2008 (doi: 10.1007/s12010-008-8369-y).
68. Zhang, Z. & Yuen, G. Y. Effects of culture fluids and preinduction of chitinase production on biocontrol of *Bipolaris leaf spot* by *Stenotrophomonas maltophilia* C3. *Biol. Control* **18**, 277–286 (2000).
69. Zhang, Z., Yuen, G. Y., Sarath, G. & Penheiter, A. R. Chitinases from the plant disease biocontrol agent, *Stenotrophomonas maltophilia* C3. *Phytopathology* **91**, 204–211 (2001).
70. Zhang, Z. G. & Yuen, G. Y. The role of chitinase production by *Stenotrophomonas maltophilia* strain C3 in biological control of *Bipolaris sorokiniana*. *Phytopathology* **90**, 384–389 (2000).
71. Mastretta, C. *et al.* Endophytic bacteria and their potential application to improve the phytoremediation of contaminated environments. *Biotechnol. Genetic Eng. Rev.* **23**, 175–207 (2006).
72. Kobayashi, D. Y., Reedy, R. M., Bick, J. & Oudemans, P. V. Characterization of a chitinase gene from *Stenotrophomonas maltophilia* strain 3451 and its involvement in biological control. *Appl. Environ. Microbiol.* **68**, 1047–1054 (2002).
73. Jurkevitch, E., Hadar, Y. & Chen, Y. Differential siderophore utilization and iron uptake by soil and rhizosphere bacteria. *Appl. Environ. Microbiol.* **58**, 119–124 (1992).
74. Ardon, O. *et al.* Iron uptake in *Ustilago maydis*: studies with fluorescent ferrichrome analogues. *Microbiology* **143**, 3625–3631 (1997).
75. Blanvillain, S. *et al.* Plant carbohydrate scavenging through TonB-dependent receptors: a feature shared by phytopathogenic and aquatic bacteria. *PLoS ONE* **2**, e224 (2007).
76. Cao, Z. J. *et al.* Characterization of a novel *Stenotrophomonas* isolate with high keratinase activity and purification of the enzyme. *J. Ind. Microbiol. Biotechnol.* **36**, 181–188 (2009).
77. Siegert, P. *et al.* Medium/means containing proteases from *Stenotrophomonas maltophilia*. Patent DE 102007033104 20070713 (2007).
78. Humphris, S. N., Wheatley, R. E. & Bruce, A. The effects of specific volatile organic compounds produced by *Trichoderma* spp. on the growth of wood decay basidiomycetes. *Holzforschung* **55**, 233–237 (2001).
79. Berg, G., Eberl, L. & Hartmann, A. The rhizosphere as a reservoir for opportunistic human pathogenic bacteria. *Environ. Microbiol.* **7**, 1673–1685 (2005).
An insightful review that describes the rhizosphere as a reservoir for opportunistic pathogenic bacteria.
80. Crossman, L. C. *et al.* The complete genome, comparative and functional analysis of *Stenotrophomonas maltophilia* reveals an organism heavily shielded by drug resistance determinants. *Genome Biol.* **9**, R74 (2008).
The first published genome sequence of a *Stenotrophomonas* species, the clinical isolate *S. maltophilia* K279a.
81. Guilhaert, M. R. & Kirkpatrick, B. C. Identification of *Xylella fastidiosa* antiviralure genes: hemagglutinin adhesins contribute to *X. fastidiosa* biofilm maturation and colonization and attenuate virulence. *Mol. Plant Microbe Interact.* **18**, 856–868 (2005).
82. Colombi, D. *et al.* Haemagglutination induced by *Bordetella pertussis* filamentous haemagglutinin adhesin (FHA) is inhibited by antibodies produced against FHA(430–873) fragment expressed in *Lactobacillus casei*. *Curr. Microbiol.* **53**, 462–466 (2006).
83. McKay, G. A., Woods, D. E., MacDonald, K. L. & Poole, K. Role of phosphoglucosyltransferase of *Stenotrophomonas maltophilia* in lipopolysaccharide biosynthesis, virulence, and antibiotic resistance. *Infect. Immun.* **71**, 3068–3075 (2003).
84. Rahmati-Bahram, A., Magee, J. T. & Jackson, S. K. Temperature-dependent aminoglycoside resistance in *Stenotrophomonas (Xanthomonas) maltophilia*: alterations in protein and lipopolysaccharide with growth temperature. *J. Antimicrob. Chemother.* **37**, 665–676 (1996).
85. Patil, P. B., Bogdanove, A. J. & Sonti, R. V. The role of horizontal transfer in the evolution of a highly variable lipopolysaccharide biosynthesis locus in xanthomonads that infect rice, citrus and crucifers. *BMC Evol. Biol.* **7**, 243 (2007).
86. Winn, A. M. & Wilkinson, S. G. Structure of the O16 antigen of *Stenotrophomonas maltophilia*. *Carbohydrate Res.* **330**, 279–283 (2001).
87. da Silva, A. C. R. *et al.* Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* **417**, 459–463 (2002).
88. Hagemann, M., Hasse, D. & Berg, G. Detection of a phage genome carrying a zonula occludens like toxin gene (*zot*) in clinical isolates of *Stenotrophomonas maltophilia*. *Arch. Microbiol.* **185**, 449–458 (2006).
89. Gould, V. C. & Avison, M. B. SmeDEF-mediated antimicrobial drug resistance in *Stenotrophomonas maltophilia* clinical isolates having defined phylogenetic relationships. *J. Antimicrob. Chemother.* **57**, 1070–1076 (2006).
90. Roscetto, E. *et al.* PCR-based rapid genotyping of *Stenotrophomonas maltophilia* isolates. *BMC Microbiol.* **8**, 202 (2008).
91. Valdezate, S. *et al.* High genetic diversity among *Stenotrophomonas maltophilia* strains despite their originating at a single hospital. *J. Clin. Microbiol.* **42**, 693–699 (2004).
92. Gulcan, H., Kuzucu, C. & Durmaz, R. Nosocomial *Stenotrophomonas maltophilia* cross-infection: three cases in newborns. *Am. J. Infect. Control* **32**, 365–368 (2004).
93. Park, Y. S. *et al.* Pseudooutbreak of *Stenotrophomonas maltophilia* bacteremia in a general ward. *Am. J. Infect. Control* **36**, 29–32 (2008).
94. Sakhni, E., Weissmann, A. & Oren, I. Fulminant *Stenotrophomonas maltophilia* soft tissue infection in immunocompromised patients: an outbreak transmitted via tap water. *Am. J. Med. Sci.* **323**, 269–272 (2002).
95. Squier, C., Yu, V. L. & Stout, J. E. Waterborne nosocomial infections. *Curr. Infect. Dis. Rep.* **2**, 490–496 (2000).
96. Gould, V. C., Okazaki, A. & Avison, M. B. β -Lactam resistance and β -lactamase expression in clinical *Stenotrophomonas maltophilia* isolates having defined phylogenetic relationships. *J. Antimicrob. Chemother.* **57**, 199–203 (2006).
97. Paez, J. I. G. & Costa, S. F. Risk factors associated with mortality of infections caused by *Stenotrophomonas maltophilia*: a systematic review. *J. Hosp. Infect.* **70**, 101–108 (2008).
98. Ansari, S. R. *et al.* Risk factors for infections with multidrug-resistant *Stenotrophomonas maltophilia* in patients with cancer. *Cancer* **109**, 2615–2622 (2007).
99. Cheong, H. S. *et al.* Risk factors for mortality and clinical implications of catheter-related infections in patients with bacteraemia caused by *Stenotrophomonas maltophilia*. *Int. J. Antimicrob. Agents* **32**, 538–540 (2008).
100. Hanes, S. D. *et al.* Risk factors for late-onset nosocomial pneumonia caused by *Stenotrophomonas maltophilia* in critically ill trauma patients. *Clin. Infect. Dis.* **35**, 228–235 (2002).
101. Figueiredo, P. M. S. *et al.* Cytotoxic activity of clinical *Stenotrophomonas maltophilia*. *Lett. Appl. Microbiol.* **43**, 443–449 (2006).
102. Fouhy, Y. *et al.* Diffusible signal factor-dependent cell–cell signaling and virulence in the nosocomial pathogen *Stenotrophomonas maltophilia*. *J. Bacteriol.* **189**, 4964–4968 (2007).
103. Waters, V. J. *et al.* Immunostimulatory properties of the emerging pathogen *Stenotrophomonas maltophilia*. *Infect. Immun.* **75**, 1698–1703 (2007).
104. Kwa, A. L. H. *et al.* The impact of multidrug resistance on the outcomes of critically ill patients with Gram-negative bacterial pneumonia. *Diagnostic Microbiol. Infect. Dis.* **58**, 99–104 (2007).
105. Pathmanathan, A. & Waterer, G. W. Significance of positive *Stenotrophomonas maltophilia* culture in acute respiratory tract infection. *Eur. Resp. J.* **25**, 911–914 (2005).
106. Weber, D. J. *et al.* Microbiology of ventilator-associated pneumonia compared with that of hospital-acquired pneumonia. *Infect. Control Hosp. Epidemiol.* **28**, 825–831 (2007).
107. Aisenberg, G. *et al.* *Stenotrophomonas maltophilia* pneumonia in cancer patients without traditional risk factors for infection, 1997–2004. *Eur. J. Clin. Microbiol. Infect. Dis.* **26**, 13–20 (2007).
108. Steinkamp, G. *et al.* Prospective evaluation of emerging bacteria in cystic fibrosis. *J. Cystic Fibrosis* **4**, 41–48 (2005).
109. Goss, C. H., Mayer-Hamblett, N., Aitken, M. L., Rubenfeld, G. D. & Ramsey, B. W. Association between *Stenotrophomonas maltophilia* and lung function in cystic fibrosis. *Thorax* **59**, 955–959 (2004).
110. Senol, E., DesJardin, J., Stark, P. C., Barefoot, L. & Snyderman, D. R. Attributable mortality of *Stenotrophomonas maltophilia* bacteremia. *Clin. Infect. Dis.* **34**, 1653–1656 (2002).
111. Boktour, M. *et al.* Central venous catheter and *Stenotrophomonas maltophilia* bacteremia in cancer patients. *Cancer* **106**, 1967–1973 (2006).

112. Friedman, N. D., Korman, T. M., Fairley, C. K., Franklin, J. C. & Spelman, D. W. Bacteremia due to *Stenotrophomonas maltophilia*: an analysis of 45 episodes. *J. Infect.* **45**, 47–53 (2002).
113. Lai, C. H. et al. Central venous catheter-related *Stenotrophomonas maltophilia* bacteraemia and associated relapsing bacteraemia in haematology and oncology patients. *Clin. Microbiol. Infect.* **12**, 986–991 (2006).
114. Avison, M. B. et al. Differential regulation of L1 and L2 β -lactamase expression in *Stenotrophomonas maltophilia*. *J. Antimicrob. Chemother.* **49**, 387–389 (2002).
115. Lambert, T., Ploy, M. C., Denis, F. & Courvalin, P. Characterization of the chromosomal aac(6)-Iz gene of *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* **43**, 2366–2371 (1999).
116. Okazaki, A. & Avison, M. B. Aph(3)-IIC, an aminoglycoside resistance determinant from *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* **51**, 359–360 (2007).
117. Alonso, A. & Martinez, J. L. Cloning and characterization of SmeDEF, a novel multidrug efflux pump from *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* **44**, 3079–3086 (2000).
118. Alonso, A. & Martinez, J. L. Expression of multidrug efflux pump SmeDEF by clinical isolates of *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* **45**, 1879–1881 (2001).
119. Zhang, L., Li, X. Z. & Poole, K. SmeDEF multidrug efflux pump contributes to intrinsic multidrug resistance in *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* **45**, 3497–3503 (2001).
120. Falagas, M. E., Valkimadi, P. E., Huang, Y. T., Matthaiou, D. K. & Hsueh, P. R. Therapeutic options for *Stenotrophomonas maltophilia* infections beyond co-trimoxazole: a systematic review. *J. Antimicrob. Chemother.* **62**, 889–894 (2008).
121. Livermore, D. M. et al. Non-susceptibility trends among *Pseudomonas aeruginosa* and other non-fermentative Gram-negative bacteria from bacteraemias in the UK and Ireland, 2001–06. *J. Antimicrob. Chemother.* **62**, 1155–1163 (2008).
122. Barbolla, R. et al. Class 1 integrons increase trimethoprim-sulfamethoxazole MICs against epidemiologically unrelated *Stenotrophomonas maltophilia* isolates. *Antimicrob. Agents Chemother.* **48**, 666–669 (2004).
123. Toleman, M. A., Bennett, P. M., Bennett, D. M. C., Jones, R. N. & Walsh, T. R. Global emergence of trimethoprim/sulfamethoxazole resistance in *Stenotrophomonas maltophilia* mediated by acquisition of *sul* genes. *Emerg. Infect. Dis.* **13**, 559–565 (2007).
124. Lawson, D. H. & Paice, B. J. Adverse reactions to trimethoprim-sulfamethoxazole. *Rev. Infect. Dis.* **4**, 429–433 (1982).
125. Nicodemo, A. C. & Paez, J. I. G. Antimicrobial therapy for *Stenotrophomonas maltophilia* infections. *Eur. J. Clin. Microbiol. Infect. Dis.* **26**, 229–237 (2007).
126. Ford, P. J. & Avison, M. B. Evolutionary mapping of the SHV β -lactamase and evidence for two separate IS26-dependent bla(SHV) mobilization events from the *Klebsiella pneumoniae* chromosome. *J. Antimicrob. Chemother.* **54**, 69–75 (2004).
127. Poirer, L., Lartigue, M. F., Decusser, J. W. & Nordmann, P. ISEcp1B-mediated transposition of bla(CTX-M) in *Escherichia coli*. *Antimicrob. Agents Chemother.* **49**, 447–450 (2005).
128. Sanchez, M. B., Hernandez, A., Rodriguez-Martinez, J. M., Martinez-Martinez, L. & Martinez, J. L. Predictive analysis of transmissible quinolone resistance indicates *Stenotrophomonas maltophilia* as a potential source of a novel family of Qnr determinants. *BMC Microbiol.* **8**, 148 (2008).
129. Al Naiemi, N., Duim, B. & Bart, A. A CTX-M extended-spectrum β -lactamase in *Pseudomonas aeruginosa* and *Stenotrophomonas maltophilia*. *J. Med. Microbiol.* **55**, 1607–1608 (2006).
130. Avison, M. B., von Heldreich, C. J., Higgins, C. S., Bennett, P. M. & Walsh, T. R. A TEM-2 β -lactamase encoded on an active Tn1-like transposon in the genome of a clinical isolate of *Stenotrophomonas maltophilia*. *J. Antimicrob. Chemother.* **46**, 879–884 (2000).
131. De Gelder, L., Williams, J. J., Ponciano, J. M., Sota, M. & Top, E. M. Adaptive plasmid evolution results in host-range expansion of a broad-host-range plasmid. *Genetics* **178**, 2179–2190 (2008).
132. Kataoka, D. et al. The indirect pathogenicity of *Stenotrophomonas maltophilia*. *Int. J. Antimicrob. Agents* **22**, 601–606 (2003).
133. Huang, T. P. & Wong, A. C. L. A cyclic AMP receptor protein-regulated cell-communication system mediates expression of a FecA homologue in *Stenotrophomonas maltophilia*. *Appl. Environ. Microbiol.* **73**, 5034–5040 (2007).
134. Wang, L. H. et al. A bacterial cell-cell communication signal with cross-kingdom structural analogues. *Mol. Microbiol.* **51**, 903–912 (2004).
135. Veselova, M. et al. Production of *N*-acylhomoserine lactone signal molecules by Gram-negative soil-borne and plant-associated bacteria. *Folia Microbiologica* **48**, 794–798 (2003).
136. Zhu, H., Thuruthyl, S. J. & Willcox, M. D. P. Production of *N*-acyl homoserine lactones by Gram-negative bacteria isolated from contact lens wearers. *Clin. Exp. Ophthalmol.* **29**, 150–152 (2001).
137. Ryan, R. P. & Dow, J. M. Diffusible signals and interspecies communication in bacteria. *Microbiology* **154**, 1845–1858 (2008).
138. Ryan, R. P. et al. Interspecies signalling via the *Stenotrophomonas maltophilia* diffusible signal factor influences biofilm formation and polymyxin tolerance in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **68**, 75–86 (2008).
- Reports signalling between *S. maltophilia* and *P. aeruginosa*, another bacterial species occupying the same niches, and discusses the consequences for bacterial behaviour.**
139. Liu, Z., Yang, C. & Qiao, C. L. Biodegradation of *p*-nitrophenol and 4-chlorophenol by *Stenotrophomonas* sp. *FEMS Microbiol. Lett.* **277**, 150–156 (2007).
140. Chen, T., Dai, Y. J., Ding, J. F., Yuan, S. & Ni, J. P. N-demethylation of neonicotinoid insecticide acetamiprid by bacterium *Stenotrophomonas maltophilia* CGMCC 1.1788. *Biodegradation* **19**, 651–658 (2008).
141. Soares, A., Guieysse, B., Delgado, O. & Mattiasson, B. Aerobic biodegradation of nonylphenol by cold-adapted bacteria. *Biotechnol. Lett.* **25**, 731–738 (2003).
142. Tachibana, S., Kuba, N., Kawai, F., Duine, J. A. & Yasuda, M. Involvement of a quinoprotein (PQQ-containing) alcohol dehydrogenase in the degradation of polypropylene glycols by the bacterium *Stenotrophomonas maltophilia*. *FEMS Microbiol. Lett.* **218**, 345–349 (2003).
143. Cao, F., Ren, Y. & Hua, W. Cyclomaltotriose mixed esters of anti-inflammatory drugs and short-chain fatty acids and study of their enzymatic hydrolysis *in vitro*. *Carbohydrate Res.* **344**, 526–530 (2009).
144. Smejkal, C. W., Seymour, F. A., Burton, S. K. & Lappin-Scott, H. M. Characterisation of bacterial cultures enriched on the chlorophenoxyalkanoic acid herbicides 4-(2,4-dichlorophenoxy) butyric acid and 4-(4-chloro-2-methylphenoxy) butyric acid. *J. Ind. Microbiol. Biotechnol.* **30**, 561–567 (2003).
145. Merroun, M. L. & Selenska-Pobell, S. Bacterial interactions with uranium: an environmental perspective. *J. Contaminant Hydrology* **102**, 285–295 (2008).
146. Song, H. P., Li, X. G., Sun, J. S., Xu, S. M. & Han, X. Application of a magnetotactic bacterium, *Stenotrophomonas* sp. to the removal of Au(III) from contaminated wastewater with a magnetic separator. *Chemosphere* **72**, 616–621 (2008).
147. Morel, M. A. et al. Cellular and biochemical response to Cr(VI) in *Stenotrophomonas* sp. *FEMS Microbiol. Lett.* **291**, 162–168 (2009).
148. Yu, L., Liu, Y. & Wang, J. Identification of novel denitrifying bacteria *Stenotrophomonas* sp. ZZ15 and *Oceanimonas* sp. YC13 and application for removal of nitrate from industrial wastewater. *Biodegradation* **20**, 391–400 (2009).
149. Dean, M. et al. Characterization of cholyglycine hydrolase from a bile-adapted strain of *Xanthomonas maltophilia* and its application for quantitative hydrolysis of conjugated bile salts. *Appl. Environ. Microbiol.* **68**, 3126–3128 (2002).
150. Pedrini, P. et al. *Xanthomonas maltophilia* CBS 897.97 as a source of new 7 β - and 7 α -hydroxysteroid dehydrogenases and cholyglycine hydrolase: improved biotransformations of bile acids. *Steroids* **71**, 189–198 (2006).
151. Nakajima, Y. et al. Dipeptidyl aminopeptidase IV from *Stenotrophomonas maltophilia* exhibits activity against a substrate containing a 4-hydroxyproline residue. *J. Bacteriol.* **190**, 7819–7829 (2008).
152. Choi, S.-K. et al. Designing selective, high affinity ligands of 5-HT_{1D} receptor by covalent dimerization of 5-HT_{1D} ligands derived from 4-fluoro-N-[3-(1-methyl-4-piperidinyl)-1H-indol-5-yl]benzamide. *J. Medicinal Chem.* **51**, 3609–3616 (2008).
153. Ping, Z. et al. Enhancement of leaching copper by electro-oxidation from metal powders of waste printed circuit board. *J. Hazard. Mater.* **166**, 746–750 (2009).
154. Xue, Y.-P. & Zheng, Y.-G. Production of valienamine using a two-step process with *Stenotrophomonas maltophilia*. *J. Biotechnol.* **136**, S367–S368 (2008).
155. Dal Bello, G. M., Monaco, C. I. & Simon, M. R. Biological control of seedling blight of wheat caused by *Fusarium graminearum* with beneficial rhizosphere microorganisms. *World J. Microbiol. Biotechnol.* **18**, 627–636 (2002).
156. Sullivan, R. F., Holtman, M. A., Zylstra, G. J., White, J. F. & Kobayashi, D. Y. Taxonomic positioning of two biological control agents for plant diseases as *Lysobacter enzymogenes* based on phylogenetic analysis of 16S rDNA, fatty acid composition and phenotypic characteristics. *J. Appl. Microbiol.* **94**, 1079–1086 (2003).

Acknowledgements

G.B. thanks C. Zachow, M. Hagemann, B. Lugtenberg and D. Egamberdieyeva; her research was supported by the Deutsche Forschungsgemeinschaft, the Austrian Science Foundation FWF and by the INTAS project 04-82-6969. Research by S.M., S.T. and D.v.d.L. was supported by the US Department of Energy, Office of Science, BER, project number KP1102010 under contract DE-AC02-98CH10886, by Laboratory Directed Research and Development funds (LDRD05-063 and LDRD09-005) and by Royalty Funds at the Brookhaven National Laboratory under contract with the US Department of Energy. M.B.A.'s work on *S. maltophilia* has been funded by the Wellcome Trust and the British Society for Antimicrobial Chemotherapy. R.P.R. and J.M.D. are indebted to Y. McCarthy for helpful discussions. R.P.R. and J.M.D. were supported in part by grants awarded by the Science Foundation of Ireland (SFI 03/IN3/B373 and 07/IN.1/B955 to J.M.D.). Sequencing of *S. maltophilia* R551-3 was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract number DE-AC02-05CH11231, by Lawrence Livermore National Laboratory under contract number DE-AC52-07NA27344 and by Los Alamos National Laboratory under contract number DE-AC02-06NA25396.

DATABASES

Entrez Gene:

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene&term=1234>

Entrez Genome Project:

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=genomeprj>
[Bordetella pertussis](#) | [Caenorhabditis elegans](#) | [Escherichia coli](#) | [Klebsiella pneumoniae](#) | [Medicago sativa](#) | [Pseudomonas aeruginosa](#) | [Rhizoctonia solani](#) | [Shewanella denitrificans](#) | [Solanum tuberosum](#) | [Stenotrophomonas maltophilia K279a](#) | [Stenotrophomonas maltophilia R551-3](#) | [Triticum aestivum](#) | [Vibrio cholerae](#) | [Xanthomonas axonopodis](#) | [Xanthomonas campestris](#) | [Xylella fastidiosa](#) | [Zea mays](#)

UniProtKB:

<http://www.uniprot.org>
 CupA | PA1396 | SMe1 | Smlt1390 | Smlt2757 | Smlt2759 | Zot

FURTHER INFORMATION

MaGe: <http://www.genoscope.cns.fr/agg/mage>
Stenotrophomonas maltophilia K279a genome: http://www.sanger.ac.uk/Projects/S_maltophilia
Stenotrophomonas maltophilia 5513 genome: http://genome.jgi-psf.org/finished_microbes/stema/stema_home.html

SUPPLEMENTARY INFORMATION

See online article: S1 (movie) | S2 (table) | S3 (table) | S4 (table) | S5 (figure)

ALL LINKS ARE ACTIVE IN THE ONLINE PDF