

Historic Perspectives on Annonaceous Acetogenins from the Chemical Bench to Preclinical Trials

Authors

Chih-Chuang Liaw^{1,2}, Tung-Ying Wu³, Fang-Rong Chang³, Yang-Chang Wu^{3,4}

Affiliations

¹ Graduate Institute of Pharmaceutical Chemistry, College of Pharmacy, China Medical University, Taichung, Taiwan

² Department of Life Sciences, National Chung-Hsing University, Taichung, Taiwan

³ Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung, Taiwan

⁴ Cancer Center, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

Key words

- Annonaceae
- acetogenins
- mitochondria complex I inhibitor
- pesticidal activity
- anticancer activity
- apoptosis
- acetogenin mimics

Abstract

Studies on the Annonaceous acetogenins began after the first cytotoxic acetogenin, uvaricin, was isolated in 1982. This attractive finding made many medicinal and natural product chemists direct their efforts on the isolation and identification of these classes of compounds. As more Annonaceous acetogenins were isolated, more information about them was uncovered. From their structural identification to the total synthesis of natural product analogues and from cell-based screening and molecular-based targeting to animal testing, the mechanisms of action of the Annonaceous acetogenins became clearer. The purpose of this review is to give an account of recent studies on this class of compounds and their analogues, which will aid us not only in clarifying how the Annonaceous acetogenins act but also in establishing principles for the further development of this class of compounds.

Abbreviations

AGE:	Annonaceous acetogenin	H23:	non-small cell human lung cancer cells
APICID:	atmospheric pressure in-source collision-induced dissociation	H8:	HPV16 subgenes-immortalised human endocervical cells
CC:	column chromatography	HCT-8:	intestinal adenocarcinoma cells
CCC:	countercurrent chromatography	HSCC:	high-speed countercurrent chromatography
CD:	circular dichroism	HT-29:	human colon cancer cells
Complex I:	NADH:ubiquinone oxidoreductase	IC ₅₀ :	the half maximal inhibitory concentration
Dansyl-NH:	5-dimethylaminonaphthalen-1-yl-sulfonamide	ITC:	isothermal titration calorimetry
DIP:	direct-inlet probe	K562:	human immortalised myelogenous leukaemia line
EI-MS:	electron-impact mass spectrometry	KB 3-1:	human epidermoid carcinoma cells
ESI-MS:	electron-spray ionisation mass spectrometry	LC/MS:	liquid chromatography/mass spectrometry
FITC:	fluorescein isothiocyanate	LT ₅₀ :	median lethal time
H125:	lung adenocarcinoma cells	M17/Adr:	adriamycin-resistant murine mammary cells
		MCF-7/Adr:	adriamycin-resistant human mammary adenocarcinoma cells
		MCF-7/wt:	nonresistant human mammary adenocarcinoma cell wild type
		MDA-MB-468:	human mammary adenocarcinoma cells
		MDR:	multiple-drug resistant
		NADH:	nicotinamide adenine dinucleotide, reduced form
		NBD-NH:	7-nitrobenzo[c][1,2,5]oxadiazol-4-yl-amino
		NCI:	National Cancer Institute
		NMA:	naphthylmethoxyacetic acid
		P388:	murine lymphoblastoid cells
		P-gp:	P-glycoprotein mediated pumps
		PI:	propidium iodide
		PO:	oral administration
		PO3:	pancreatic adenocarcinoma
		PS:	PS system (P-388 lymphocytic leukaemia in mice)
		SAR:	structure-activity relationship

received Nov. 30, 2009

revised April 26, 2010

accepted April 28, 2010

Bibliography

DOI <http://dx.doi.org/10.1055/s-0030-1250006>
Published online June 24, 2010
Planta Med 2010; 76:
1390–1404 © Georg Thieme
Verlag KG Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Dr. Yang-Chang Wu
Graduate Institute of Natural
Products
Kaohsiung Medical University
100 Shih-Chuan 1st Road
703 Kaohsiung
Taiwan
Republic of China
Phone: + 886 73 12 11 01
ext. 2197
Fax: + 886 73 11 47 73
yachwu@kmu.edu.tw

Sf9 cell:	pupal ovarian tissue of the Fall armyworm <i>Spodoptera frugiperda</i>	THP:	tetrahydropyran
T/C:	test/control	TLC:	thin-layer chromatography
T24:	bladder cancer cells	TMS:	trimethylsilyl
THF:	tetrahydrofuran	YEM:	yellow fever mosquito larvae assay

Introduction

Plants have a long history of use in the treatment of human diseases. Botanical extracts have long been regarded as a source of new and useful pharmaceuticals. According to Cragg's investigation, approximately 62% of commercially available drugs have natural product origins [1]. These natural products also play important roles as direct treatments or as templates (lead compounds) that are modified for the treatment of human diseases. The commonly known example from folkloric medicinal plants is the anticancer agent paclitaxol, a diterpene from *Taxus brevifolia* (Taxaceae) discovered by Wall and Wani in 1971 [2] that is now used for the treatment of advanced ovarian cancer. Other examples of anticancer drugs derived from natural products, such as camptothecin from *Camptotheca acuminata* (Nyssaceae) [3], podophyllotoxin from *Podophyllum peltatum* [4], vincristine and vinblastine from *Vinca rosea* [5], and adriamycin from *Streptomyces peucetius* [6], have encouraged pharmaceutical chemists to search for new drugs from medicinal plant sources.

Annonaceous acetogenins (AGEs) are a unique class of C₃₅ or C₃₇ secondary metabolites of Annonaceous plants derived from the polyketide pathway. Extensive studies on AGEs have indicated that these naturally occurring compounds possess a broad spectrum of bioactivity, including anticancer, antiparasitic, insecticidal and immunosuppressive effects. In 27 years, more than 500 AGEs were isolated from various parts of the plants of this family [7–10]. These bioactive AGEs became more important, particularly in pharmaceutical research [11, 12]. A great interest in investigating the mechanisms of action of a series of AGEs emerged from the leaps in knowledge about the processes involved in tumour cell death. Members of this class of natural compounds are regarded as "potential" candidates for future generations of anticancer drugs. AGEs have become one of the most interesting classes of natural products at present.

Annonaceous plants are important economic crops in Asia. There is an especially abundant biomass of Annonaceous plants in Taiwan. In our studies, AGEs from Formosan Annonaceous plants showed significant cytotoxicity against ovarian cancer cells, 1A9, with more potential than paclitaxol [13]. The purpose of this review is to give a short historical introduction as well as an account of the recent studies on AGEs and their analogues. This brief outline begins with a description of the sources, isolation, chemistry and biological activities of this class of natural compounds. Achievements in the studies of AGEs are noted by their significant cytotoxicity. Recent studies on the mechanisms of action of the pesticidal and antitumour AGEs are reviewed within the individual sections. In addition, modified analogues of AGEs and AGE mimics were made to verify hypotheses regarding the modes of action of the AGEs. Historically following the studies of the AGEs from the abundant natural biomass, it will be helpful for us to not only clarify how AGEs act but also establish the principles that will guide the further development of these natural products.

Previous Studies on the AGEs before the 2000s

The isolation of uvaricin (1) from the roots of *Uvaria accuminata* Oliv. by Jolad et al. in 1982 and its excellent bioactivity in the PS test system initiated the studies on AGEs as a hot topic in drug discovery [14]. Following the experimental and major biochemical conceptual advances, the studies on AGEs during the early 1980s to about 2000 could be summarised in three stages, including the initial stage (before 1990), the middle stage (from 1991–1995), and modern stage (1996–2000). In the initial stage (before 1990), several groups, including McLaughlin's (Purdue, USA), Cave's (CNRS, France), Fujimoto's (Tokyo, Japan), Pettit's (Arizona, USA), and Sneden's (Virginia, USA), began efforts toward the isolation and structural identification of these types of bioactive compounds.

In the isolation procedure, AGEs were first concentrated via solvent partitioning driven by their amphiphathic properties. Methanol is usually the solvent of choice for the extraction of these AGEs from plant materials. By partitioning the crude extract with chloroform and water, the chloroform layer will become enriched with the AGEs. Chromatographic techniques, such as gradient elution column chromatography (CC), flash chromatography and preparative TLC, were generally utilised for the isolation and purification of these compounds. During this period of time the major population of AGEs was discovered. For example, the first mono-tetrahydrofuran (THF) AGE, annonacin (2), was isolated from *Annona densicoma* Mart [15]; the adjacent bis-THF AGE, squamocin (3), was isolated from *A. squamosa* L. [16]. The first nonadjacent bis-THF AGE, bullatalicin (4) [17], was isolated from *A. bullata* A. Rich. and was also isolated from *A. cherimolia* Mill as cherimoline (5) [18] (renamed by Cortes et al. as cherimolins-1 and 2 [19]). The first AGE with the saturated γ -lactone moiety, laherradurine (6), was isolated from *A. cherimolia* seeds, of which the structure was revised by the same group later [19, 20]. The primary structures of the AGEs were determined by ¹H- and ¹³C-NMR and mass spectroscopy, in which the former method (NMR) could confirm the presence of functional group substituents (the γ -lactone ring moiety and the oxygen-bearing moieties) and the latter (MS) could determine the placement of substituents along the carbon skeleton. Rupprecht et al. published the first review on AGEs in 1990 [7], reporting the strategies for their structural elucidation and revising some structures of the published compounds. At that time, three main types of AGEs, mono-THF, adjacent bis-THF and nonadjacent bis-THF, were classified based on the presence of the THF ring moieties.

The stereochemistry of the substituents of AGEs was mainly determined by organic syntheses of the partial structures and X-ray crystallography of the AGEs. Hoyer et al. (Minnesota, USA) first proposed a valuable method for the quantitative correlation of the ¹H-NMR chemical shift data for a series of model diastereomeric bis-tetrahydrofurans [21, 22]. Pettit et al. first elucidated the relative stereochemistry of the eight chiral centres of rolliniastatin 1 (7), by X-ray structure determination of its 15-O-*p*-bromophenylurethane derivative, as 4S*,15S*,16S*,19R*,20R*,23S*,24R*,36R* [23]. However, the absolute configuration of rolliniastatin

1 (7) should be the mirror image of the proposed structure because of its CD data [$n-\pi^*$ $\Delta\epsilon$ (235) < 0 and $\pi-\pi^*$ $\Delta\epsilon$ (220 nm) > 0] [24, 25], which indicates a 36S configuration. In 1990, Born et al. proposed a systematic comparison of the ^1H - and ^{13}C -NMR chemical shifts of the diagnostic protons for two diastereomers, which became a popular way to confirm the *erythro* and *threo* effects on the protons of the tetrahydrofuran rings [26]. Meanwhile, Hoyer et al. proposed a symmetry-assisted synthesis of the bis-THF moiety [27]. Biosynthetically, the AGEs are regarded to originate from polyhydroxy C-32 or C-34 fatty acids, to which a 2-propanol unit is added to form the methylated α,β -unsaturated γ -lactone. Hoyer's proposed idea really matched the biosynthetic hypothesis for the construction of these compounds (● Fig. 1).

Following the rapid development of chromatographic techniques in the early 1990s, studies on the AGEs in the middle stage (from 1991 to 1995) focused on the isolation and purification of various types of AGEs and the identification of the stereochemistry of AGEs by chemical methods. More laboratories worldwide became active in research related to the isolation, structural elucidation, and even total synthesis and mechanisms of anticancer action of these compounds. Our group also began studying the Formosan AGEs at this time.

Since the 1970s, our group has mainly studied the secondary metabolites from plants of the Lauraceae. Because of the close phylogenic relationships of both the Lauraceae and Annonaceae families, we started to study the secondary metabolites of the Formosan Annonaceous plants (● Table 1). In Taiwan, there are 21 species (8 genera) of Annonaceous plants, of which three, *Fissistigma glaucescens* (Hance) Merr., *F. oldhami* (Hemsl.) Merr. and *Goniothalamus amuyon* (Blanco) Merr., are native to the Taiwan island [28, 29]. Our early studies on the Formosan Annonaceous plants focused on alkaloids, which showed diverse biological functions, including cytotoxicity [30], antiplatelet aggregation activity [31], cardiovascular activity [32, 33] and antimicrobial activity [34]. Afterwards, diterpenoids, styrylpyrones [12, 35, 36] and some linear AGEs [37], mono-THF AGEs [38, 39], and adjacent bis-THF AGEs [39] were isolated from various parts of the *Annona* plants in our laboratory. Two groups in China also started related research on AGEs in China during that period [40, 41], one of which – Dr. Yang's group in Yun-Nan – cooperated with us to elucidate the structures of the AGEs from *A. muricata* [42].

To improve the efficiency of the chromatography, repeated open column chromatography and high-performance liquid chromatography (HPLC) were introduced for the isolation of AGEs [43–46] such that AGEs with minor structural differences could be isolated more easily and quickly. Based on the isolations of different AGEs, their general structural features can be divided into two classes: 1) the moiety of the γ -lactone rings: the α,β -unsaturated γ -lactone ring (normal form) or the ketolactone (isoform) (see ● Fig. 2); and 2) the oxygen-bearing moieties (see ● Fig. 3) [9, 47]. In addition, the structure of (–)-muricatacin (8), the first shortened AGE with only a terminal γ -lactone moiety and an aliphatic chain, was reported in 1991 [48].

Due to the difficulty in crystallising aliphatic AGEs, the determination of the absolute stereochemistry of AGEs became the main challenge of the research work. Hoyer et al. first used spectroscopic methodologies to determine the absolute configurations of stereogenic carbinol centres in the five adjacent bis-THF AGEs and four mono-THF AGEs with the refined Mosher method in 1992 [43, 44]. They also validated the configuration at the C-4 carbinol centre in these bioactive AGEs as the *R* configuration [49].




Fig. 1 Structures 1–7.

Fujimoto et al. designed a series of mono-THF compounds with various conformations and reported their ^{13}C -NMR resonances, which are useful in determining the conformations of the THF ring moieties [50]. Yu et al. made a single crystal of (+)-gigantecin (9) from *A. coriacea* Mart for an X-ray study to determine its absolute conformation [51]. These research findings accelerated the process for the structural identification of the AGEs.

On the other hand, Duret et al. suspected that AGEs with terminal ketolactones (isoforms) were artefacts of the translactonisation of 4-hydroxy-AGEs. They further confirmed this hypothesis by performing the extraction and characterisation of the initial AGEs from fresh crude materials under the effects of alkaloids, basic media, and alcohols. These reagents affected the kinetics of the translactonisation [52], a result that was later supported by Duret et al. work showing that 4-hydroxylated AGEs led to iso-AGEs under basic conditions [53].

The studies on AGEs in the modern stage (from 1996 to 2000) were concerned with the efficient identification of AGEs by hyphenated techniques and other spectroscopic methods. Although normal and reverse-phase HPLC are powerful tools for the isolation of natural products, they still have some limitations, such as the amount of sample that can be purified per unit time, the cost of the solvent, the size of the columns. Searching for new technologies to facilitate the chromatographic work is very important. Hopp et al. used countercurrent chromatography (CCC) to isolate

	Seed or unripe fruit	Fruit	Leaves	Stem	Bark	Root
Annona						
<i>A. cherimola</i> Mill.		*	*	*		
<i>A. glabra</i> L.		*		*		
<i>A. artemoya</i> (<i>A. cherimola</i> X <i>A. squamosa</i>)	*					
<i>A. montana</i> Macf.	*		*	*	*	
<i>A. muricata</i> L.	*		*			
<i>A. reticulata</i> L.	*		*			
<i>A. squamosa</i> L.	*		*	*		
<i>A. purpurea</i> L.			*	*		
Artabotrys						
<i>A. hexaptalus</i> (L. f.) Bhandqri						
<i>A. uncinatus</i> (Lam.) Merr.		*	*	*		*
Cananga						
<i>Cananga odorata</i> (Lam.) Hook. F. & Thomas	*		*			
Fissistigma						
<i>F. glaucescens</i> (Hance) Merr. #			*	*		
<i>F. oldhami</i> (Hemsl.) Merr. #	*		*	*		
Goniiothalamus						
<i>G. amuyon</i> (Blanco) Merr. #			*	*		
Polyalthia						
<i>P. longifolia</i> Benth. et Hook. F.			*			
<i>P. longifolia</i> Benth. et Hook. F. 'pendula'			*			
<i>P. suberosa</i> Hook. f.						
<i>P. liukuensis</i> Hatusima #						
Rollinia						
<i>R. mucosa</i> Baill	*		*	*		
Uvaria						
<i>U. rufa</i> Bl.				*		

Table 1 Formosan Annonaceous plants collected for investigation of their secondary metabolites

* Chemical components of the material have been studied in our laboratory. # Annonaceous plants that are native in Taiwan island




Fig. 2 γ -Lactone subunits in AGEs.

four AGEs, (2,4-*cis*- and *trans*)-9-hydroxyasimicinone (**10**), (2,4-*cis*- and *trans*)-squamoxinone B (**11**), (2,4-*cis*- and *trans*)-squamoxinone C (**12**), and isoannoreticuin (**13**), from the bark of *A. squamosa* [54]. Duret et al. also applied high-speed countercurrent chromatography (HSCCC) to the separation of AGEs from *A. atemoya* to give two major AGEs, squamocin (**3**) and bullatacin (rolliniastatin-2, **14**), and six other known AGEs [55]. Moreover, to separate mixtures of AGEs that cannot be easily purified by regular HPLC methods, some Japanese scholars and our laboratory introduced the recycle-HPLC system for isolating AGEs.

To develop a convenient spectral methodology to determine the stereochemistry of AGEs, Gawronski and Wu provided a far more reliable way of determining the absolute configuration of the γ -lactone ring moiety through the analysis of the CD spectra of butenolides [25]. Duret et al. modified the Mosher method and de-

termined the stereochemistry of asimicin (**15**) by the long-range anisotropic effect of 2-NMA (naphthylmethoxyacetic acid), [56] (● Fig. 4).

The other key tool for determining the structures of AGEs is mass spectrometry (MS). Generally, electron-impact mass spectrometry (EI-MS) is the preferred technique for determining the placement of the tetrahydrofuran rings and functional groups (hydroxy, ketone, acetoxy and double bond) along the hydrocarbon chain. The derivatised AGEs, such as TMS and acetyl derivatives, are helpful in the elucidation of these structures. In addition, the direct-inlet probe technique (DIP) and lower evaporator energy (e.g., 30 eV) have been suggested for use with EI-MS scanning because AGEs easily decompose thermally. The structure of squamocin (**3**) from *A. squamosa* was characterised by a combination of chemical derivatisation and precursor-ion scanning mass spectrometry. The lactone portion of squamocin (**3**) was modified with *N,N*-dimethylethylenediamine in the vapour phase to afford a strong positive charge at one end of the skeleton [57]. In 1997, Gu et al. (XenoBiotic Laboratories), cooperating with the McLaughlin group, analysed AGEs from *R. mucosa* that were amenable to liquid chromatography/mass spectrometry (LC/MS) with ionisation source-atmospheric pressure in-source collision-induced dissociation (APICID) to detect the presence of 40 known AGEs, in addition to four new AGEs of diverse structures, in a bioactive crude methanol-soluble fraction from this plant extract [58]. They also observed a unique fragmentation rule for AGEs with a hydroxy group at C-4, which had a characteristic loss of a terminal γ -lactone (112 amu) during ESI-MS scanning [58]. This




Fig. 3 Tetrahydrofuran (THF), tetrahydropyran (THP), and other oxygen-bearing subunits in Annonaceous AGEs.

rapid and relatively uncomplicated selective ionisation procedure also provided a convenient and useful method for identifying AGEs with or without a hydroxy group at C-4.

The Extensive Studies on AGEs after 2000

After 2000, we continued investigating the AGEs from Formosan plants of *Annona* species [59–62]. Among them, two epimeric AGEs, muricins A (**16**) and B (**17**) [59], were isolated and their absolute configurations were determined by the modified Mosher method. Muricin B (**17**) is the first Annonaceous acetogenin to possess a hydroxy group with the *S*-configuration at C-4, where the typical configuration of the hydroxy group was *R*. In 2003, we reported a novel skeleton of abridged AGE, rollicosin (**18**), from the unripe fruits of *Rollinia mucosa* Baill, which was the first identified compound that contained lactone moieties on both sides of an aliphatic chain [62]. Soon after this isolation, Chinese scholars reported the second abridged AGE, squamostolide (**19**), from *A. squamosa* [63] (● Fig. 5).

During this time, the function and mechanism of action of AGEs were investigated. The link between the mitochondrial respiratory chain and cell apoptosis was clarified [64]. The latter, namely programmed cell death, is a normal physiological process that selectively and desirably destroys cells and tissues without an inflammatory response, as opposed to a necrotic cell death. Instead of focusing on the inhibition of mitochondria complex I, we found




Fig. 4 Structures 8–15.

that bullatacin (**14**) could induce cell death via apoptosis based on an analysis of the morphological changes of bullatacin (**14**)-treated Hep 2.2.15, as determined by double staining with fluorescein isothiocyanate (FITC)-labelled annexin V and propidium iodide (PI) [65]. This finding also opened a new window for exploring the mechanism of action of AGEs. Herein, we summarise the studies on the cytotoxicity and pesticidal activity of AGEs in the last decade.

Biological activities and the mechanisms of action of the AGEs

Although AGEs were reported with high potential and diverse biological activities, including antibacterial [66], insecticidal, cytotoxic and immunosuppressive effects, pharmaceutical scientists were interested in how AGEs worked in cells (what the mechanism of the anticancer action of AGEs is) and whether the compounds could work *in vivo*. Combined with the traditional uses of the Annonaceous plants in North America and South-East Asia, scientists noticed the pesticidal activities of AGEs and proposed a possible mechanism of pesticidal action involved to ATP levels in pests. Thereafter, they found that both the cytotoxic and pesticidal activities should be related to ATP generation and NADH oxi-




Fig. 5 Structures 16–19.

dation in mitochondria, which directed many ongoing studies about the interaction of AGEs with the mitochondrial complex I.

Anticancer activity: Jolad et al. first reported the significant *in vivo* cytotoxic activity of uvaricin (**1**) by the PS screening system [14]. Ahammadsahib et al. used normal mice bearing L1210 murine leukaemia and athymic mice bearing A2780 conventional ovarian cancer xenografts to study the cytotoxic action of bullatacin (**14**) and analogues in 1993; meanwhile these compounds also have potential as insecticides in insect-derived Sf9 cells. The toxicity of AGEs in both cases probably arises from the strong inhibitory ability of mitochondrial electron transport with specific action at complex I [67]. Degli Esposti et al. (France) first used mammalian mitochondria to study the action of AGEs toward the NADH-ubiquinone reductase (complex I) and reported that bullatacin (**14**) inhibited the proton pumping function of complex I with similar efficiency under steady-state and non-steady-state conditions, as compared to the action of rotenone and piericidin [68]. Because of the ability to inhibit the mitochondrial complex I, the main gate of the energy production in the cells, AGEs have been regarded as candidates for future generations of antitumour drugs with different mechanisms.

Besides blocking the NADH:ubiquinone oxidoreductase (complex I) in the electron transport system, AGEs are also powerful inhibitors of the NADH oxidases peculiar to the plasma membranes of cancer cells. Both mechanisms of action result in the inhibition of ATP production and may account for the observation that AGEs are more effective at killing multiple-drug resistant (MDR) tumours than their nonresistant counterparts since the MDR pumps on the cell membranes require ATP to function. In addition, Oberlies et al. observed that AGEs could selectively inhibit the cell growth of cancerous cells by *in vitro* cell inhibition assays against three murine (P388, PO3, and M17/Adr) and two human (H8 and H125) cancer cell lines [69]. Interestingly, the work of Oberlies et al. proposed that this class of natural products showed a certain biological activity against some drug-resistant cancers. Currently, multidrug-resistant cancers are hard to cure because the cancer cells have developed a mechanism to overcome the anticancer agents. Based on the biochemical differences between MDR and parental cancer cells, such as the ATP-depen-


dent P-glycoprotein-mediated pumps (P-gp) and the higher demand for ATP in the MDR cancer cells, Oberlies et al. used bullatacin (**14**) to test two cell lines, MDR human mammary adenocarcinoma (MCF-7/Adr) cells and the parental, nonresistant wild-type (MCF-7/wt) cells [70]. Therefore, ATP depletion could be another mode of action of AGEs that offers a special advantage in the chemotherapeutic treatment of MDR tumours.

Shimada et al. also proposed a model for explaining the action of AGEs [71]. They suspected that the lactone ring alone could directly interact with the binding to complex I, and the THF rings with flanking OH groups function just as hydrophilic anchors at the membrane surface that allow lateral diffusion (or random distribution) of the lactone ring in the membrane interior. To verify the model, Kuwabara et al. synthesised a series of analogues with two terminal γ -lactone rings [72]. However, the bioassay results did not show that these analogues worked twice as well as AGEs did.

To clarify the mechanism of action of AGEs, we cooperated with biochemists and pharmacologists in Taiwan due to its abundant amount of naturally occurring Formosan AGEs. Yuan et al. found that annonacin (**2**) could arrest T24 bladder cancer cells at the G1 phase and cause cytotoxicity in a Bax- and caspase-3-related pathway [73]. In addition, squamocin (**3**) was also observed to arrest the same cancer cells at the G1 phase and cause a selective cytotoxicity in S-phase-enriched T24 cells via the same pathway of cleaving the functional protein of PARP and inducing cell apoptosis [74]. Squamocin (**3**) was also found to inhibit the proliferation of K562 cells via G2/M arrest in association with the induction of p21, p27 and the reduction of Cdk1 and Cdc25C kinase activities [75]. These works connected the AGEs with cell apoptosis, which exploited the multiple functions of AGEs as new anticancer candidates.

Pesticidal activity: Following the North American folk use of Annonaceous plants as pesticides, Rupprecht et al. first noted this application. They determined the pesticidal potencies of the extracts from the paw paw tree (*Asimina triloba*) by the brine shrimp test (*Artemia salina* larvae), which paralleled the significant activities seen against the striped cucumber beetle (*Acalymma vittatum* F.), Mexican bean beetle (*Epilachna varivestis* Mulsant), mosquito larvae (*Aedes aegypti* L.), blowfly larvae (*Calliphora vicina* Meigen), melon aphid (*Aphis gossypii* Glover), two-spotted-spider mite (*Tetranychus urticae* Koch) and free-living nematode (*Caenorhabditis elegans*). During the preliminary screening, asimicin (**15**) was isolated and its pesticidal action evaluated [76]. Moreover, by the same bioactivity-guided isolation/fractionation method, bullatacin (**14**) was isolated and its pesticidal effects observed at concentrations as low as 1 ppm, whereas bullatacinone (**20**) lacked pesticidal activities [24]. Meanwhile, Ratnayake et al. conducted a controlled study on the pesticidal potencies of extracts from various plant parts of the paw paw tree (*Asimina triloba*) using the brine shrimp test [77] (● Fig. 6).

He et al. further evaluated the pesticidal properties of 44 AGEs using the yellow fever mosquito larvae (YFM) assay [78]. The results clearly demonstrated that most AGEs had pesticidal properties. In addition, they indicated that the adjacent bis-THF AGEs with three hydroxy groups, for example, bullatacin (**14**) and trilobin (**21**), were the most potent. They further made AGEs, mono-THF, adjacent bis-THF, and nonadjacent bis-THF types as insecticidal baits to test the potent toxicity of these compounds against insecticide-susceptible and -resistant German cockroaches, compared to the activities of some conventional synthetic insecti-



cides [79]. Ohsawa et al. evaluated the insecticidal activities of AGEs from the seeds of the pond apple, *A. glabra* L. with a micro-sprayer on the cabbage leaf or the filter paper [80]. Meanwhile, Guadano et al. also found that annonacin (2) showed anti-feedant effects on *L. decemlineata* and squamocin (3) was toxic to *L. decemlineata* and *M. persicae*. They also proved that both AGEs were not mutagenic but were toxic in the absence of a metabolic activation system [81].

Londershausen et al. also noticed that extracts of ground seeds from *A. squamosa* revealed interesting insecticidal properties. AGEs were determined to be the active components through an activity-monitored fractionation. The investigation of ATP-levels (at the LT_{50} value) in *Plutella xylostella* under treatment with squamocin (3) and antimycin A revealed values of 1.45 and 1.35 $\mu\text{mol/g}$ fresh weight, respectively. Further studies revealed that squamocin (3) showed an inhibitory effect on NADH-cytochrome *c*-reductase and complex I of insect mitochondria with IC_{50} values of 4–8 $\mu\text{mol/g}$ protein and 0.8 μM , respectively. Similar results for squamocin (3) were observed for the inhibition of complex I from bovine heart muscle (IC_{50} : <0.1 μM) or *Neurospora crassa* cells (IC_{50} : 0.3 μM), but no effects on other coupling sites of mitochondrial complexes were observed [43]. These assembled experimental results were the work of Lewis et al. in 1993 [82]. Friedrich et al. and Hollingworth et al. simultaneously reported the insecticidal action of AGEs to be a result of the inhibition of mitochondria complex I [83,84]. Friedrich et al. found that the inhibition of mitochondrial and bacterial NADH:ubiquinone oxidoreductase (complex I) by AGEs was not purely competitive [83]. They demonstrated that AGEs should affect the electron-transfer step from the high-potential iron-sulfur cluster to ubiquinone by directly acting at the ubiquinone-catalytic site of complex I.

The Studies on Modifications and Analogues of the AGEs

The structure-activity relationship (SAR) studies of AGEs are always interesting for medicinal and natural product chemists. Miyoshi et al. noticed that the alkyl spacer between the γ -lactone and hydroxylated THF ring moieties elicited potent inhibitory activities on the NADH oxidase [85]. They summarised the SAR rules of AGEs as follows: 1) the adjacent bis-THF ring moiety is not an essential structural factor for inhibition, and the mono-

THF ring compounds can maintain potent activities; 2) this stereochemical factor was also not essential for potent activity irrespective of the number (one or two) of THF rings; 3) the THF rings of the AGEs had strong interactions with the interface of lipid bilayers irrespective of the stereochemistry in the THF region; and 4) the spacer moiety is very important for potent activity [86]. Takada et al. also tested the NADH oxidase activity of two naturally occurring AGEs, bullatacin (14) and diepomuricanin (22), and several synthesised analogues in a comparison with that of piericidin A [87]. They concluded that both ring moieties, the γ -lactone ring and the tetrahydrofuran ring, acted in a cooperative manner on the enzyme and that the optimal length of the alkyl spacer was 13 carbon atoms. These results supported the above hypothesis that Miyoshi et al. offered.

To consider solely the role of the THF ring moieties, Murai et al. synthesised Δlac -AGE (23) (AGE without the α,β -unsaturated γ -lactone ring), which was also shown to be a novel type of inhibitor that acts at the terminal electron transfer step of mitochondrial NADH-ubiquinone oxidoreductase (complex I). They also synthesised a photolabile Δlac -AGE (24) connected to a biotin probe to trace the labelled peptide without the use of a radioisotope. This photolabile Δlac -AGE (24) elicited potent inhibition of bovine heart mitochondrial complex I at nanomolar levels [88]. Ichimaru et al. further synthesised a series of Δlac -AGEs, in which the stereochemistry around the hydroxylated tetrahydrofuran (THF) ring moiety was systematically modified, and examined their inhibitory effects on complex I. The results revealed that the bis-THF ring analogues are much more potent than the mono-THF ring analogues and that the stereochemistry around the bis-THF ring moiety played a significant role in the inhibitory effects on complex I [89]. Compound 25 showed a similar IC_{50} value as was observed for bullatacin (14) in the reduction of NADH oxidase activity (0.60–0.65 mmol NADH/min/mg of protein) in submitochondrial particles. Intriguingly, Ichimaru et al. demonstrated that the inhibitory site of complex I on which Δlac -AGEs acted might be different from that at which natural AGEs did.

On the other hand, these featured structures, such as the γ -lactone ring moiety, one to three THF/THP rings with multiple chiral centres, and an alkyl side chain make AGEs difficult and challenging synthetic targets. Because substantial amounts of pure samples are required for further biological and clinical studies, a number of total syntheses of AGEs have been reported in the literature since the 1990s. Recent advances in the total syntheses of AGEs include mono-THF AGEs: murisolin (26) [90,91], longicin (27) [92,93], and *cis*-solamin (28) [94,95], adjacent bis-THF AGEs: bullatacin (14) [96], rolliniastatin 1 (7) [23,97], rollimembrin (29) [97,98], 10-hydroxyasimicin (30) [99,100], membranacin (31) [97,101], asimicin (15) [76,102], longimicin D (32) [103–104], and mucoxin (33) [105,106], non-adjacent bis-THF AGEs: *cis*-sylvaticin (34) [107,108] and gigantecin (9) [109,110], and others, jimenezin (35) [111,112], mucocin (36) [113,114], pyranicin (37) [115,116], pyragoncin (38) [115,117,118], rollicosin (18) [62,119] and squamostolide (19) [63,120] (● Fig. 7).

In addition to the total syntheses of various AGEs, some special analogues were designed to improve the bioactivities through, for example, modifications of the γ -lactone ring, the THF ring, and hydroxy moieties on the aliphatic chain.




Fig. 7 Structures 22–38.

Modifications of the γ -lactone ring moiety

Hoppen et al. designed and prepared quinone-mucocin (**39**) and quinone-squamocin D (**40**) to elucidate the mechanisms of action of the AGEs. The IC_{50} values of **39** and **40** in the inhibition of the mitochondrial NADH-ubiquinone oxidoreductase complex were 3.6 and 1.7 nM, respectively [121]. These results supported their hypothesis that AGEs are competitive inhibitors at the ubiquinone binding site of complex I based on the structural similarity between the butenolide and the quinone. Arndt et al. synthesised a systematic variation of featured structures, the butenolide and the ether components, to evaluate the critical factors for the interaction of the AGEs with complex I. Their results and data from the smaller substructures indicated that the substructures of the AGEs, the polyether component and the lipophilic side chain, would be necessary for strong binding of the AGEs to complex I [122].

In addition, aromatic heterocycles are commonly found as base structures of potent complex I inhibitors. Duval et al. tried to replace the α,β unsaturated γ -lactone moiety of squamocin (**3**) with benzimidazole via an unusual condensation-oxidative decarboxylation reaction with 1,2-diamines in the presence of acetic acid and oxygen. Although they did not clarify the inhibitory ability of the modified squamocin toward complex I, one of the benzimidazole analogues (**41**) showed cytotoxicity (KB 3–1) with an IC_{50} value of 2.2×10^{-3} μ M and induced a 61% accumulation of the G1 phase of the KB 3–1 cell cycle at concentrations of 1–5 nM, with apoptosis above 10 nM [123]. In 2006, Duval et al. semisynthesised a series of heterocyclic analogues of squamocin (**3**). Their results suggested that the binding of this hybrid inhibitor (**41**) was responsible for a negative allosteric effect at the level of the first ubiquinone-binding site of mitochondrial complex I [124].

Duval et al. also prepared a small library of the γ -keto ester derivatives of squamocin (**3**) and screened their biological activities,

including their cytotoxicity against KB 3–1 cells, inhibition of mitochondrial complex I and of complex III. However, these modified analogues with an open γ -lactone ring did not show better activity than that of the parent compound, squamocin (**3**) [125]. Except for the adjacent bis-THF and nonadjacent bis-THF AGEs, Kojima et al. made a series of α,β -unsaturated- γ -lactone-free, nitrogen-containing heterocyclic analogues of solamin (**42**), a mono-THF acetogenin. The cytotoxicities of the compounds were investigated against 39 tumour cell lines. One of them, a 1-methylpyrazol-5-yl derivative (**43**), showed a selective increase in cytotoxicity against NCI-H23 with a potency 80 times higher than that of solamin [126] (● Fig. 8).

Modification of the THF ring moiety

The AGEs are a large class of naturally occurring polyketides that exhibit potent anticancer activities. In 2000, based on both the difficulty associated with total syntheses of AGEs and the straightforward means by which their structures can be simplified, Chinese and French scholars both proposed to replace the ethylene bridge in the THF rings with normal and iso-terminal lactone moieties, respectively [127, 128].

Yao and coworkers further studied simplified AA005 (**44**) and its analogues, which showed potent antitumour activities and significant selectivity between normal cells and cancer cells (≥ 7 , [129]). Zeng et al. designed and synthesised (4*R*)-hydroxylated analogue **45** based on the structure of bullatacin (**14**). The preliminary screenings showed that the IC_{50} values of **45** were 1.6×10^{-3} and 8×10^{-2} μ g/mL against HT-29 and HCT-8 cells, respectively. A remarkable enhancement of cytotoxic effect for the 4(*R*)-hydroxy analogue (**45**) was observed [130]. The results support a very interesting piece of information, namely that both the butenolide and ethylene glycol subunits play essential roles in the cytotoxicities of the compounds against tumour cell lines. Their precise




Fig. 8 Structures 39–43.

role is not yet clear. Additionally, the presence of the hydroxy group at C-10 and the absolute configuration of the methyl group on the butenolide moiety (**46**) are less important for their activity [131].

Rodier et al. tried to introduce a benzoyl group to fix the moiety between the ether linkage; these analogues display interesting cell cycle effects but are less potent than bullatacinone (**20**), a compound with the same terminal lactone [132]. Fujita et al. also tried to replace the bis-adjacent THF ring by a 1,2-cyclopentenediol bis-ether skeleton to obtain simplified mimics **47–50**. Based on the evaluation of the inhibitory effects on mitochondrial NADH:ubiquinone oxidoreductase (complex I), the 1,2-cyclopentenediol bis-ether motif also showed very potent inhibitory activity at the nanomolar level [133].

For the study of AGE mimics, Liu et al. further designed, synthesised and evaluated a new series of mimics containing a terminal lactam [134]. They found that the *N*-methylated lactam-containing compounds **51**, **52**, and **53** exhibited comparable potencies to that of AA005 (**44**) and similar selectivity among cancer cells. *N*-Methyl compound **51** shows comparable activities to that of AA005 (**44**) and retains similar cell selectivity. It was also revealed that the stereogenic centre on the lactam is not essential for antitumour activity. Recently, Liu et al. synthesised a series of analogues by replacing the acyclic bis-ether functionality of AA005 (**44**) with certain conformationally constrained fragments. Interestingly, most newly synthesised mimetics were found to exhibit potent activities against breast cancer cells and showed satisfactory selectivities between cancerous and non-cancerous cells. Among them, an *N,N'*-dimethyl bis-amide compound (**54**) exhibits more potency against MDA-MB-468 cells than does its parent molecule AA005 (**44**). Studies by Lin et al. indicate that bisamide analogues of AA005 make this unique class of anticancer agents much simpler and allow more flexibility for their future development [135] (● Fig. 9).




Fig. 9 Structures 44–54.

Replacement of the hydroxy moiety on the aliphatic chain

Some scholars have studied the replacement of the hydroxy moieties on the aliphatic chain of AGEs. Ye et al. obtained halogen-substituted AGEs, 4(*S*)-chloro-4-deoxygigantetrocin A and 4(*S*)-18-dichloro-4,18-dideoxy-asimilobin, by treating gigantetrocin A with triphenylphosphine and CCl_4 . The chlorinated compounds showed decreased bioactivities in the brine shrimp lethality test and against human tumour cell lines [136]. Kojima et al. made C4-fluorinated analogue (**55**) of solamin and evaluated its anti-tumour activities against 39 tumour cell lines. They found that C4-fluorinated solamin (**55**) showed more potent growth inhibitory activity against cancer cell lines than did solamin [137]. On the other hand, Gallardo et al. made 10-oximeganacone (**56**), the first bioactive nitrogenated acetogenin, which showed potent inhibition towards complex I by the titration of the NADH oxidase and NADH:ubiquinone oxidoreductase activities [138]. Duret et al. semisynthesised amino derivatives from two natural AGEs, rolliniastatin-1 and squamocin. Although it is noteworthy that these amino-AGEs still retain some activity, more studies are required to confirm the potencies of these derivatives as new specific and efficient anticancer agents [139].

A variety of chemical strategies has been applied to investigating biological processes. Recently, fluorescent modifications became powerful tools for visualising the distribution of bioactive natural

products in cells and investigating their targeting. In 2005, Derbre et al. synthesised hybrids consisting of an AGE tail connected to a fluorescent tag. Using fluorescent microscopy, both **57** and **58** were initially observed in Jurkat cell mitochondria, but they diffused into the cytosol of apoptotic cells, supporting the conclusion that squamocin (**3**) passes through the plasma membrane and targets the mitochondria. Indeed, both semisynthesised fluorescent derivatives were shown to be potent apoptosis inducers that were directed to this organelle. Besides, they proposed that the lactone moiety seems not to interfere with the mitochondrial targeting but apparently influenced the bioactivity of AGEs [140]. Alexander et al. attached ethyl 7-dimethylaminocoumarin 4-acetate to the diols of (–)-mucocin (**59**) through amide coupling chemistry. Although coumarin-labelled mucocin can also induce fluorescently coded morphogenic responses, no expected response was found [141]. This result might be due to the occupation of the mitochondrial recognition site by the fluorescent coumarin group. To overcome the above disadvantage, Maezaki et al. and Kojima et al. labelled the terminal aliphatic chain of solamin (**60**, **61**) with fluorescent groups, 7-nitrobenzo [c][1,2,5]oxadiazol-4-yl-amino (NBD-NH-) and 5-dimethylaminonaphthalen-1-yl-sulfonamide (dansyl-NH-), in 2007 and 2009, respectively [142,143]. It was anticipated that these compounds would be used to explore the targeting of AGEs. Among AGE mimics, Liu et al. tried to modify the C-10 hydroxy group of the AGE mimic to introduce a label based on the results of the anticancer-activity screenings of parallel synthetic analogues. Fluorescent-imaging studies revealed that AA005-flu's (**62** and **63**) distribution in normal human cells was significantly different from that in cancer cells. AA005-flu accumulated in the mitochondria of the cancer cells. This direct and visible evidence suggests that membrane recognition of AA005 (**44**) is involved in its selective bioactivity [144] (► Fig. 10).

AGEs as cation ionophores

Although many research results mentioned the mechanisms of AGEs, for instance, the inhibitory action of mitochondria complex I (NADH:ubiquinone oxidoreductase) [145], induction of programmed cell death by the expression of the pro-apoptotic proteins Bax, Bad, caspase-3 [74] and the structure-activity relationships of either natural, semisynthesised or synthesised compounds, the diverse bioactivities of the various types of AGEs still seem difficult to explain. Some researchers notice the chemophysical features of various AGEs and more direct evidence was provided as a new structure-activity relationship for AGEs. Sasaki et al. first reported the ionophore activity of the AGEs. It was revealed by NMR studies that the structurally-related analogues of AGEs form supramolecular complexes with metal cations [146]. These studies indicated that hydroxylated bis-THF derivatives, structural components of the potent antitumour AGEs, formed supramolecular complexes with metal cations. In particular, some formed 2 : 1 ligand : metal complexes with calcium cations with high selectivity [146]. Although Araya et al. evaluated the ion-transport and ion-binding activities of AGEs using apparatus W-08 in 1995 and did not find any special activity [147], in 1998 Sasaki et al. indicated that two AGEs, bullatacin (**14**) and asimicin (**15**), and their structurally related analogues binding bivalent cations such as Ca^{2+} and Mg^{2+} [148,149]. Peyrat et al. evaluated the ^{13}C -NMR longitudinal relaxation times (T_1) of both annonacin (**2**) and squamocin (**3**) in the absence and presence of Ca^{2+} ions to assess the structural changes that accompany complexations. They thought that the interesting cytotoxic activities




Fig. 10 Structures **55**–**63**.

of the THF- γ -lactone derivatives could be explained by their ionophoric ability. Their results also show differences in the stoichiometry of the complexes for mono-THF AGE and bis-THF AGE with Ca^{2+} ions [150].

In biological studies with living cells, we assumed that AGEs play a role in the bioavailability of the cations in the cell membranes due to their amphiphilic nature. While culturing smooth muscle cells of the human coronary artery with squamocin (**3**) in our study, we observed that squamocin (**3**) (an adjacent bis-THF acetogenin) could induce a transient but strong increase in the large-conductance Ca^{2+} -activated K^{+} channels [151]. In a whole-cell configuration, squamocin (**3**, 0.3–100 μM) induced a Ca^{2+} -activated K^{+} current [$I_{\text{K}(\text{Ca})}$] in a concentration-dependent manner with an EC_{50} value of 4 μM . When cells were exposed to a Ca^{2+} -free solution, squamocin (**3**, 3 μM) induced a transient increase in $I_{\text{K}(\text{Ca})}$. In the continued presence of squamocin (**3**), an additional increase in ex-




Fig. 11 Structures **64** and **65**.

tracellular Ca^{2+} (1 mM) caused a significant increase in $I_{K(\text{Ca})}$. In the cell-attached configuration of the single-channel recordings, squamocin (**3**) applied to the bath increased the activity of large-conductance Ca^{2+} -activated K^+ (BK_{Ca}) channels without altering the single-channel conductance. These findings provide evidence that squamocin (**3**) can activate $I_{K(\text{Ca})}$ in coronary arterial smooth muscle cells. The initial transient activation of $I_{K(\text{Ca})}$ may reflect squamocin-induced Ca^{2+} release from intracellular Ca^{2+} stores, whereas the sustained activation of $I_{K(\text{Ca})}$ may arise from the squamocin-induced Ca^{2+} influx across the cell membrane. The stimulatory effects of squamocin (**3**) on these channels should affect the functional activity of vascular smooth muscle cells [151]. We speculate that AGEs could use their hydrophilic centres (THF rings with flanking hydroxy groups) to bind cations like Ca^{2+} and surround the ion core by its peripheral hydrophobic regions (long chains). This arrangement allows the molecules to dissolve effectively in the membrane and diffuse transversely into cells as ionophores. We clarified the interaction between mono-THF AGEs and Ca^{2+} by isothermal titration calorimetry (ITC), which is an extremely powerful and highly sensitive technique for measuring the heats of interaction of reacting species in dilute solution. Interestingly, we found that the mono-THF AGEs annonacin (**2**) and uvariamicin-I (**64**) interacted with Ca^{2+} by an exothermic process, indicating the formation of AGE-calcium complexes [152] (● Fig. 11).

From the Chemical Bench to Preclinical Trials

In 1976, Ratnayake et al. found that extracts of the leaves and twigs of the native paw paw tree, *Asimina triloba*, were bioactive in the antitumour screens of the U.S. National Cancer Institute (NCI). Following the sound phytochemical studies on AGEs [77], Gu et al. used the three most active AGEs, bullatacin (**14**), asimicidin (**15**) and trilobacin (**65**), to establish quality control over AGE extracts of the paw paw tree by LC/MS/MS [153]. In this study they identified that small twigs from the months of May and June were the optimum plant sources for the commercial harvest of biomass for extraction. They further tried to develop some useful commercial products containing the AGEs, including head lice shampoo (in 2001), ointment, lotion, spray for plant pests, and paw paw capsules (in 2003) for human administration. The entire process from the safety and toxicology of the AGEs to the success of commercial products was described in McLaughlin's 2008 review [154]. More recently, Cuendet et al. reported the potential of the standardised extract from the twigs of *A. triloba* to mediate a cancer chemopreventative effect in the *N*-methyl-*N*-nitrosourea-induced mammary carcinogenesis model. As McLaughlin et al. did, they used three potent bioactive AGEs, bullatacin (**14**), asimicidin

(**15**), and trilobacin (**65**), in their standardised extract. Mammary tumour latency was increased from 55 to 66 days in Sprague-Dawley rats given a diet containing paw paw extract (1250 and 2500 mg/kg diet; based on maximum tolerated dose studies) [155].

In Taiwan, plants of the genus *Annona* are important economic crops for their edible fruits. The abundant material obtained from the seeds and the excellent cytotoxicities of the AGEs from this material attracted us to further develop pharmaceutical products. In addition to the aforementioned achievements, oral gavage (PO) animal studies have been performed by MDS Pharma Services and us. The extract from *A. muricata*, WYC-AA07, was applied to the xenograft tumour model of human MCF-7 breast tumour cells on SCID mice (assay 580000). WYC-AA07 at 10 mg/kg was administered daily by PO for a total of 10 doses. The tumour size, body weight and signs of overt animal toxicities after dosing were monitored and recorded for 25 days. WYC-AA07 at 10 mg/kg PO caused a significant decrease in the tumour weights from day 13 to day 25. However, it also caused a significant decrease in body weight on days 9, 13, 17, and 21. The other experiment was done on the xenograft tumour model of human HT-29 colon tumours on SCID mice (assay 580100). WYC-AA07 at 20 mg/kg PO caused death in one half of the animals and a significant decrease in body weight on day 8. Recently, we tested the toxicity of squamocin (**3**) on nude mice at 20 mg/kg PO. Although most of the animals showed an abnormal neuron function that caused the mice to move uncontrollably in a lateral direction, the mice recovered after we stopped the administration of squamocin (**3**). On the other hand, an ethnopharmacological investigation reported the similar side effect of AGEs that a neurodegenerative tauopathy endemic to the Caribbean island of Guadeloupe was suspected to be linked to the consumption of Annonaceous plants. Escobar-Khondiker et al. further found that annonacin (**2**) induced the retrograde transport of mitochondria to decrease ATP levels, which induces changes in the intracellular distribution of tau in a way that shares characteristics with some neurodegenerative diseases [156]. The possible adverse effect of AGEs on neuron cells should be a concern in advanced pharmaceutical applications for alternative therapy.

Perspectives

The AGEs are one of the most interesting classes of natural products appearing in the past two decades. They exhibit a wide variety of biological activities, and, impressively, some of them have comparable cytotoxic to taxol against various cancer cells. Both featured structures, the hydroxylated THF and γ -lactone ring moieties, are thought to be the pharmacophores that block the electron transport system of mitochondrial complex I. Much effort has been dedicated to elucidating the underlying cytotoxic mechanisms of the AGEs and to synthesising AGE analogues by altering the spacing between two moieties, removing either one of two critical featured structures (Δ Lac AGEs or muricatacin), or mimicking the THF rings by ether linkages. Although none of the modified AGEs obtained thus far have demonstrated activities comparable to those of the naturally occurring AGEs, studies on the synthesis and mechanisms of action of these compounds established solid fundamental knowledge and drug discovery experience. For example, the studies on analogues of AGEs created a series of compounds that contain completely different skeletons, of which some also show excellent bioactivities against various can-

cer cells. In addition, *in vivo* tests in mice showed the antitumor effects and even some possible adverse effects of AGEs. However, there was no released data about the *in vivo* pharmacokinetic study of AGEs. Thus, more *in vivo* studies are necessary. Combined with the above information, it will be substantially helpful to further develop these types of compounds as new drugs.

References

- Cragg GM, Newman DJ, Snader KM. Natural products in drug discovery and development. *J Nat Prod* 1997; 60: 52–60
- Wani MC, Taylor HL, Wall ME, Coggon P, McPhail AT. Plant antitumor agents. VI. Isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J Am Chem Soc* 1971; 93: 2325–2327
- Wall ME, Wani MC, Cook CE, Palmer KH, McPhail AT, Sim GA. Plant antitumor agents. I. The isolation and structure of camptothecin, a novel alkaloidal leukemia and tumor inhibitor from *Camptotheca acuminata*. *J Am Chem Soc* 1966; 88: 3888–3890
- Hartwell JL, Schrecker AW. Components of podophyllin. V. The constitution of podophyllotoxin. *J Am Chem Soc* 1951; 73: 2909–2916
- Neuss N, Gorman M, Hargrove W, Cone NJ, Biemann K, Buechi G, Manning RE. Vinca alkaloids. XXI. The structures of the oncolytic alkaloids vinblastine (VLB) and vincristine (VCR). *J Am Chem Soc* 1964; 86: 1440–1442
- Arcamone F, Franceschi G, Penco S, Selva A. Adriamycin (14-hydroxy-daunomycin), a novel antitumor antibiotic. *Tetrahedron Lett* 1969; 13: 1007–1010
- Rupprecht JK, Hui YH, McLaughlin JL. Annonaceous acetogenins: a review. *J Nat Prod* 1990; 53: 237–278
- Zeng L, Ye Q, Oberlies H, Shi G, Gu ZM, He K, McLaughlin JL. Recent advances in annonaceous acetogenins. *Nat Prod Rep* 1996; 13: 275–306
- Cave A, Figadere B, Laurens A, Cortes D. Acetogenins from Annonaceae. Progress in the chemistry of organic natural products, vol. 70. Berlin, Heidelberg: Springer Publisher; 1997: 81–288
- Bermejo A, Figadere B, Zafra-Polo MC, Barrachina I, Estornell E, Cortes D. Acetogenins from Annonaceae. Recent progress in isolation, synthesis, and mechanisms of action. *Nat Prod Rep* 2005; 22: 263–303
- Leboeuf M, Cave A, Bhaumik PK, Mukherjee B, Mukherjee R. The phytochemistry of the Annonaceae. *Phytochemistry* 1982; 21: 2783–2813
- Wu YC. New research and development on the Formosan annonaceous plants. In: Atta-ur-Rahman, editor. *Studies in natural products chemistry*, vol. 33. Amsterdam: Elsevier B.V.; 2006: 957–1024
- Nakanishi Y, Chang FR, Liaw CC, Wu YC, Bastow KF, Lee KH. Acetogenins as selective inhibitors of the human ovarian 1A9 tumor cell line. *J Med Chem* 2003; 46: 3185–3188
- Jolad SD, Hoffmann JJ, Schram KH, Cole JR. Uvaricin, a new antitumor agent from *Uvaria accuminata* (Annonaceae). *J Org Chem* 1982; 47: 3151–3153
- McCloud TG, Smith DL, Chang C-J, Cassidy JM. Annonacin, a novel, biologically active polyketide from *Annona densicoma*. *Experientia* 1987; 43: 947–949
- Fujimoto Y, Eguchi T, Kakinuma K, Ikekawa N, Sahai M, Gupta YK. Squamocin, a new cytotoxic bis-tetrahydrofuran containing acetogenin from *Annona squamosa*. *Chem Pharm Bull* 1988; 36: 4802–4806
- Hui YH, Rupprecht JK, Anderson JE, Liu YM, Smith DL, Chang CJ, McLaughlin JL. Bullatalicin, a novel bioactive acetogenin from *Annona bullata* (Annonaceae). *Tetrahedron* 1989; 45: 6941–6948
- Cortes D, Rios JL, Villar A, Valverde S. Cherimoline et dihydrocherimoline: deux nouvelles γ -lactones bis-tetrahydrofuranniques possédant une activité antimicrobienne. *Tetrahedron Lett* 1984; 25: 3199–3202
- Cortes D, Myint SH, Dupont B, Davoust D. Acetogenins of the Annonaceae. Part 20. Bioactive acetogenins from seeds of *Annona cherimolia*. *Phytochemistry* 1993; 32: 1475–1482
- Rios JL, Cortes D, Valverde S. Acetogenins, aporphinoids, and azaanthraquinone from *Annona cherimolia* seeds. *Planta Med* 1989; 55: 321–323
- Hoye TR, Suhadolnik JC. On the stereochemistry of the bistetrahydrofuran moiety of uvaricin. Proton chemical shifts can play a crucial role in complex structure determination. *J Am Chem Soc* 1987; 109: 4402–4403
- Hoye TR, Zhuang ZP. Validation of the proton NMR chemical shift method for determination of stereochemistry in the bistetrahydrofuran moiety of uvaricin-related acetogenins from Annonaceae: rolliniastatin 1 (and asimicin). *J Org Chem* 1988; 53: 5578–5580
- Pettit GR, Cragg GM, Polonsky J, Herald DL, Goswami A, Smith CR, Moretti C, Schmidt JM, Weisleder D. Isolation and structure of rolliniastatin 1 from the South American tree *Rollinia mucosa*. *Can J Chem* 1987; 65: 1433–1435
- Hui YH, Rupprecht JK, Liu YM, Anderson JE, Smith DL, Chang CJ, McLaughlin JL. Bullatacin and bullatacinone: two highly potent bioactive acetogenins from *Annona bullata*. *J Nat Prod* 1989; 52: 463–477
- Gawronski J, Wu YC. A note on the determination of absolute configuration of acetogenins by circular dichroism. *Polish J Chem* 1999; 73: 241–243
- Born L, Lieb F, Lorentzen JP, Moeschler H, Nonfon M, Soellner R, Wendisch D. The relative configuration of acetogenins isolated from *Annona squamosa*: annonin I (squamocin) and annonin VI. *Planta Med* 1990; 56: 312–316
- Hoye TR, Suhadolnik JC. Symmetry-assisted synthesis of triepoxide stereoisomers of *E,Z,E*-dodeca-2,6,10-trien-1,12-diol and their cascade reactions to 2, 5-linked bistetrahydrofurans. *J Am Chem Soc* 1985; 107: 5312–5313
- Huang TS. Flora of Taiwan, vol. 2. Taipei: Lungwei Printing Co. Ltd.; 1996: 415–419
- Liao JC. A list of scientific names of woody plants in Taiwan. Taipei: College of Agriculture, National Taiwan University; 1993
- Wu YC, Chang GY, Ko FN, Teng CM. Bioactive constituents from the stems of *Annona montana*. *Planta Med* 1995; 61: 146–149
- Chen KS, Ko FN, Teng CM, Wu YC. Antiplatelet and vasorelaxing actions of some aporphinoids. *Planta Med* 1996; 62: 133–136
- Lin CH, Yang CM, Ko FN, Wu YC, Teng CM. Antimuscarinic action of lirioidenine, isolated from *Fissistigma glaucescens*, in canine tracheal smooth muscle. *Br J Pharmacol* 1994; 113: 1464–1470
- Chang GJ, Wu MH, Wu YC, Su MJ. Electrophysiological mechanisms for antiarrhythmic efficacy and positive inotropy of lirioidenine, a natural aporphine alkaloid from *Fissistigma glaucescens*. *Br J Pharmacol* 1996; 118: 1571–1583
- Ko FN, Yu SM, Su MJ, Wu YC, Teng CM. Pharmacological activity of (–)-discretamine, a novel vascular α -adrenoceptor and 5-hydroxytryptamine receptor antagonist, isolated from *Fissistigma glaucescens*. *Br J Pharmacol* 1993; 110: 882–888
- Wu YC, Chang FR, Duh CY, Wang SK, Wu TS. Cytotoxic styrylpyrrolones of *Goniothalamus amuyon*. *Phytochemistry* 1992; 31: 2851–2853
- Wu YC, Duh CY, Chang FR, Chang GY, Wang SK, Chang JJ, McPhail DR, McPhail AT, Lee KH. The crystal structure and cytotoxicity of goniodiol-7-monoacetate from *Goniothalamus amuyon*. *J Nat Prod* 1991; 54: 1077–1081
- Wu YC, Chang FR, Chen KS, Liang SC, Lee MR. Studies on acetogenins from Formosan Annonaceous plants. 3. Diepoxymontin, a novel acetogenin from *Annona montana*. *Heterocycles* 1994; 38: 1475–1478
- Wu YC, Chang FR, Duh CY, Wang SK. Anoreticuin and isoanoreticuin: two new cytotoxic acetogenins from *Annona reticulata*. *Heterocycles* 1992; 34: 667–674
- Chang FR, Wu YC, Duh CY, Wang SK. Studies on the acetogenins of Formosan Annonaceous plants. II. Cytotoxic acetogenins from *Annona reticulata*. *J Nat Prod* 1993; 56: 1688–1694
- Zhang L, Yang R, Wu X. Chemical compositions of *Goniothalamus howii* (I). *Zhiwu Xuebao* 1993; 35: 390–396
- Yu J, Luo X, Sun L, Liu C, Hong S, Ma L. Squamostatin-B, a new polyketide from *Annona squamosa* (Annonaceae). *Chin Chem Lett* 1993; 4: 423–426
- Li CM, Mu Q, Hao XJ, Sun HD, Zheng HL, Wu YC. Three new bioactive Annonaceous acetogenins from *Annona muricata*. *Chin Chem Lett* 1994; 5: 747–750
- Londershausen M, Leicht W, Lieb F, Moeschler H, Weiss H. Molecular mode of action of annonins. *Pestic Sci* 1991; 33: 427–438
- Gu ZM, Fang XP, Miesbauer LR, Smith DL, McLaughlin JL. 30-, 31-, and 32-hydroxybullatacinones: bioactive terminally hydroxylated Annonaceous acetogenins from *Annona bullata*. *J Nat Prod* 1993; 56: 870–876
- Gu ZM, Fang XP, Rieser MJ, Hui YH, Miesbauer LR, Smith DL, Wood KV, McLaughlin JL. New cytotoxic Annonaceous acetogenins: bullatanocin and cis- and trans-bullatanocinone, from *Annona bullata* (Annonaceae). *Tetrahedron* 1993; 49: 747–754
- Gu ZM, Fang XP, Zeng L, Wood KV, McLaughlin JL. Bullacin: a new cytotoxic Annonaceous acetogenin from *Annona bullata*. *Heterocycles* 1993; 36: 2221–2228
- Zafra-Polo MC, Figadere B, Gallardo T, Tormo JR, Cortes D. Natural acetogenins from Annonaceae, synthesis and mechanisms of action. *Phytochemistry* 1998; 48: 1087–1117

- 48 Rieser MJ, Kozlowski JF, Wood KV, McLaughlin JL. Muricatacin: a simple biologically active acetogenin derivative from the seeds of *Annona muricata* (Annonaceae). *Tetrahedron Lett* 1991; 32: 1137–1140
- 49 Rieser MJ, Hui YH, Rupprecht JK, Kozlowski JF, Wood KV, McLaughlin JL, Hanson PR, Zhuang Z, Hoye TR. Determination of absolute configuration of stereogenic carbinol centers in Annonaceous acetogenins by proton and fluorine-19 NMR analysis of Mosher ester derivatives. *J Am Chem Soc* 1992; 114: 10203–10213
- 50 Fujimoto Y, Murasaki C, Shimada H, Nishioka S, Kakinuma K, Singh S, Singh M, Gupta YK, Sahai M. Annonaceous acetogenins from the seeds of *Annona squamosa*. Non-adjacent bis-tetrahydrofuranic acetogenins. *Chem Pharm Bull* 1994; 42: 1175–1184
- 51 Yu JG, Hu XE, Ho DK, Bean MF, Stephens RE, Cassady JM, Brinen LS, Clardy J. Absolute stereochemistry of (+)-gigantecin from *Annona coriacea* (Annonaceae). *J Org Chem* 2002; 59: 1598–1599
- 52 Duret P, Laurens A, Hocquemiller R, Cortes D, Cave A. Acetogenins of Annonaceae. 34. Isoacetogenins, artifacts issued from translocationization from annonaceous acetogenins. *Heterocycles* 1994; 39: 741–749
- 53 Duret P, Figadere B, Hocquemiller R, Cave A. Epimerization of Annonaceous acetogenins under basic conditions. *Tetrahedron Lett* 1997; 38: 8849–8852
- 54 Hopp DC, Conway WD, McLaughlin JL. Using countercurrent chromatography to assist in the purification of new Annonaceous acetogenins from *Annona squamosa*. *Phytochem Anal* 1999; 10: 339–347
- 55 Duret P, Waechter AI, Margraff R, Foucault A, Hocquemiller R, Cave A. High-speed countercurrent chromatography: a promising method for the separation of the Annonaceous acetogenins. *J Liq Chromatogr Relat Technol* 1997; 20: 627–635
- 56 Duret P, Waechter A-I, Figadere B, Hocquemiller R, Cave A. Determination of absolute configurations of carbinols of Annonaceous acetogenins with 2-naphthylmethoxyacetic acid esters. *J Org Chem* 1998; 63: 4717–4720
- 57 Hirayama K, Akashi S, Yuji R, Niitsu U, Fujimoto Y. Structural studies of polyhydroxybis(tetrahydrofuran) acetogenins from *Annona squamosa* using the combination of chemical derivatization and precursor-ion scanning mass spectrometry. *Org Mass Spectrom* 1993; 28: 1516–1524
- 58 Gu ZM, Zhou D, Wu J, Shi G, Zeng L, McLaughlin JL. Screening for Annonaceous acetogenins in bioactive plant extracts by liquid chromatography/mass spectrometry. *J Nat Prod* 1997; 60: 242–248
- 59 Chang FR, Wu YC. Novel cytotoxic Annonaceous acetogenins from *Annona muricata*. *J Nat Prod* 2001; 64: 925–931
- 60 Chang FR, Liaw CC, Lin CY, Chou CJ, Chiu HF, Wu YC. New adjacent bis-tetrahydrofuran Annonaceous acetogenins from *Annona muricata*. *Planta Med* 2003; 69: 241–246
- 61 Liaw CC, Chang FR, Wu CC, Chen SL, Bastow Kenneth F, Hayashi KI, Nozaki H, Lee KH, Wu YC. Nine new cytotoxic monotetrahydrofuranic Annonaceous acetogenins from *Annona montana*. *Planta Med* 2004; 70: 948–959
- 62 Liaw CC, Chang FR, Wu MJ, Wu YC. A novel constituent from *Rollinia mucosa*, rollicosin, and a new approach to develop Annonaceous acetogenins as potential antitumor agents. *J Nat Prod* 2003; 66: 279–281
- 63 Xie HH, Wei XY, Wang JD, Liu MF, Yang RZ. A new cytotoxic acetogenin from the seeds of *Annona squamosa*. *Chin Chem Lett* 2003; 14: 588–590
- 64 Wolvetang EJ, Johnson KL, Krauer K, Ralph SJ, Linnane AW. Mitochondrial respiratory chain inhibitors induce apoptosis. *FEBS Lett* 1994; 339: 40–44
- 65 Chih HW, Chiu HF, Tang KS, Chang FR, Wu YC. Bullatacin, a potent anti-tumor Annonaceous acetogenin, inhibits proliferation of human hepatocarcinoma cell line 2.2.15 by apoptosis induction. *Life Sci* 2001; 69: 1321–1331
- 66 Cartagena E, Colom OA, Neske A, Valdez JC, Bardon A. Effects of plant lactones on the production of biofilm of *Pseudomonas aeruginosa*. *Chem Pharm Bull* 2007; 55: 22–25
- 67 Ahammadsahib KI, Hollingworth RM, McGovern JP, Hui YH, McLaughlin JL. Mode of action of bullatacin: a potent antitumor and pesticidal Annonaceous acetogenin. *Life Sci* 1993; 53: 1113–1120
- 68 Degli Esposti M, Ghelli A, Ratta M, Cortes D, Estornell E. Natural substances (acetogenins) from the family Annonaceae are powerful inhibitors of mitochondrial NADH dehydrogenase (complex I). *Biochem J* 1994; 301: 161–167
- 69 Oberlies NH, Jones JL, Corbett TH, Fotopoulos SS, McLaughlin JL. Tumor cell growth inhibition by several Annonaceous acetogenins in an *in vitro* disk diffusion assay. *Cancer Lett* 1995; 96: 55–62
- 70 Oberlies NH, Croy VL, Harrison ML, McLaughlin JL. The Annonaceous acetogenin bullatacin is cytotoxic against multidrug-resistant human mammary adenocarcinoma cells. *Cancer Lett* 1997; 115: 73–79
- 71 Shimada H, Grutzner JB, Kozlowski JF, McLaughlin JL. Membrane conformations and their relation to cytotoxicity of asimicin and its analogs. *Biochemistry* 1998; 37: 854–866
- 72 Kuwabara K, Takada M, Iwata J, Tatsumoto K, Sakamoto K, Iwamura H, Miyoshi H. Design syntheses and mitochondrial complex I inhibitory activity of novel acetogenin mimics. *Eur J Biochem* 2000; 267: 2538–2546
- 73 Yuan SSF, Chang HL, Chen HW, Yeh YT, Kao YH, Lin KH, Wu YC, Su JH. Annonacin, a mono-tetrahydrofuran acetogenin, arrests cancer cells at the G1 phase and causes cytotoxicity in a Bax- and caspase-3-related pathway. *Life Sci* 2003; 72: 2853–2861
- 74 Yuan SSF, Chang HL, Chen HW, Kuo FC, Liaw CC, Su JH, Wu YC. Selective cytotoxicity of squamocin on T24 bladder cancer cells at the S-phase via a Bax-, Bad-, and caspase-3-related pathways. *Life Sci* 2006; 78: 869–874
- 75 Lu MC, Yang SH, Hwang SL, Lu YJ, Lin YH, Wang SR, Wu YC, Lin SR. Induction of G2/M phase arrest by squamocin in chronic myeloid leukemia (K562) cells. *Life Sci* 2006; 78: 2378–2383
- 76 Rupprecht JK, Chang CJ, Cassady JM, McLaughlin JL, Mikolajczak KL, Weisleder D. Asimicin, a new cytotoxic and pesticidal acetogenin from the pawpaw, *Asimina triloba* (Annonaceae). *Heterocycles* 1986; 24: 1197–1201
- 77 Ratnayake S, Rupprecht JK, Potter WM, McLaughlin JL. Evaluation of various parts of the paw paw tree, *Asimina triloba* (Annonaceae), as commercial sources of the pesticidal Annonaceous acetogenins. *J Econ Entomol* 1992; 85: 2353–2356
- 78 He K, Zeng L, Ye Q, Shi G, Oberlies NH, Zhao GX, Njoku CJ, McLaughlin JL. Comparative structure-activity relationship evaluations of Annonaceous acetogenins for pesticidal activity. *Pestic Sci* 1997; 49: 372–378
- 79 Alali FQ, Kaakeh W, Bennett GW, McLaughlin JL. Annonaceous acetogenins as natural pesticides: potent toxicity against insecticide-susceptible and -resistant German cockroaches (Dictyoptera: Blattellidae). *J Econ Entomol* 1998; 91: 641–649
- 80 Ohsawa K, Atsuzawa S, Mitsui T, Yamamoto I. Isolation and insecticidal activity of three acetogenins from seeds of pond apple, *Annona glabra* L. *Nippon Noyaku Gakkaishi* 1991; 16: 93–96
- 81 Guadano A, Gutierrez C, De la Pena E, Cortes D, Gonzalez-Coloma A. Insecticidal and mutagenic evaluation of two Annonaceous acetogenins. *J Nat Prod* 2000; 63: 773–776
- 82 Lewis MA, Arnason JT, Philogene BJR, Rupprecht JK, McLaughlin JL. Inhibition of respiration at site I by asimicin, an insecticidal acetogenin of the pawpaw, *Asimina triloba* (Annonaceae). *Pestic Biochem Physiol* 1993; 45: 15–23
- 83 Friedrich T, Ohnishi T, Forche E, Kunze B, Jansen R, Trowitzsch W, Hoefle G, Reichenbach H, Weiss H. Two binding sites for naturally occurring inhibitors in mitochondrial and bacterial NADH: ubiquinone oxidoreductase (complex I). *Biochem Soc Trans* 1994; 22: 226–230
- 84 Hollingworth RM, Ahammadsahib KI, Gadelhak G, McLaughlin JL. New inhibitors of complex I of the mitochondrial electron transport chain with activity as pesticides. *Biochem Soc Trans* 1994; 22: 230–233
- 85 Miyoshi H, Ohshima M, Shimada H, Akagi T, Iwamura H, McLaughlin JL. Essential structural factors of Annonaceous acetogenins as potent inhibitors of mitochondrial complex I. *Biochim Biophys Acta* 1998; 1365: 443–452
- 86 Miyoshi H. Inhibitors of mitochondrial respiratory enzymes. *J Pestic Sci* 2005; 30: 120–121
- 87 Takada M, Kuwabara K, Nakato H, Tanaka A, Iwamura H, Miyoshi H. Definition of crucial structural factors of acetogenins, potent inhibitors of mitochondrial complex I. *Biochim Biophys Acta* 2000; 1460: 302–310
- 88 Murai M, Ichimaru N, Abe M, Nishioka T, Miyoshi H. Synthesis of photo-labile Δ lac-acetogenin for photoaffinity labeling of mitochondrial complex I. *J Pestic Sci* 2006; 31: 156–158
- 89 Ichimaru N, Yoshinaga N, Nishioka T, Miyoshi H. Effect of stereochemistry of Δ lac-acetogenins on the inhibition of mitochondrial complex I (NADH-ubiquinone oxidoreductase). *Tetrahedron* 2007; 63: 1127–1139
- 90 Miynt SH, Laurens A, Hocquemiller R, Cave A, Davoust D, Cortes D. Murisolin: a new cytotoxic mono-tetrahydrofuran- γ -lactone from *Annona muricata*. *Heterocycles* 1990; 31: 861–867

- 91 Hattori Y, Kimura Y, Moroda A, Konno H, Abe M, Miyoshi H, Goto T, Makabe H. Synthesis of murisolin, (15R,16R,19R,20S)-murisolin A, and (15R,16R,19S,20S)-16,19-cis-murisolin and their inhibitory action with bovine heart mitochondrial complex I. *Chem Asian J* 2006; 1: 894–904
- 92 Ye Q, Zeng L, Zhang Y, Zhao GX, McLaughlin JL, Woo MH, Evert DR. Longicin and goniothalamycinone: novel bioactive monotetrahydrofuran acetogenins from *Asimina longifolia*. *J Nat Prod* 1995; 58: 1398–1406
- 93 Hanessian S, Giroux S, Buffat M. Total synthesis and structural confirmation of (+)-longicin. *Org Lett* 2005; 7: 3989–3992
- 94 Gleye C, Duret P, Laurens A, Hocquemiller R, Cave A. cis-Monotetrahydrofuran acetogenins from the roots of *Annona muricata*. *J Nat Prod* 1998; 61: 576–579
- 95 Goksel H, Stark CBW. Total synthesis of cis-solamin: exploiting the RuO₄-catalyzed oxidative cyclization of dienes. *Org Lett* 2006; 8: 3433–3436
- 96 Zhao H, Gorman JST, Pagenkopf BL. Advances in Lewis acid controlled carbon-carbon bond-forming reactions enable a concise and convergent total synthesis of bullatacin. *Org Lett* 2006; 8: 4379–4382
- 97 Keum G, Hwang CH, Kang SB, Kim Y, Lee E. Stereoselective syntheses of rolliniastatin 1, rollimembrin, and membranacin. *J Am Chem Soc* 2005; 127: 10396–10399
- 98 Gonzalez MC, Tormo JR, Bermejo A, Zafra-Polo MC, Estornell E, Cortes D. Rollimembrin, a novel acetogenin inhibitor of mammalian mitochondrial complex I. *Bioorg Med Chem Lett* 1997; 7: 1113–1118
- 99 He K, Shi G, Zhao GX, Ye Q, Schwedler JT, Wood KV, McLaughlin JL. Three new adjacent bis-tetrahydrofuran acetogenins with four hydroxyl groups from *Asimina triloba*. *J Nat Prod* 1996; 59: 1029–1034
- 100 Natrass GL, Diez E, McLachlan MM, Dixon DJ, Ley SV. The total synthesis of the Annonaceous acetogenin 10-hydroxyasimicin. *Angew Chem Int Ed Engl* 2005; 44: 580–584
- 101 Saez J, Sahpaz S, Villaescusa L, Hocquemiller R, Cave A, Cortes D. Acetogenins of the Annonaceae. 18. Rioclarin and membranacin, two new bis-tetrahydrofuran acetogenins of the seeds of *Rollinia membranacea*. *J Nat Prod* 1993; 56: 351–356
- 102 Marshall JA, Sabatini JJ. An outside-in approach to adjacent bistetrahydrofuran Annonaceous acetogenins with C₂ core symmetry. Total synthesis of asimicin and a C32 analogue. *Org Lett* 2006; 8: 3557–3560
- 103 Ye Q, He K, Oberlies NH, Zeng L, Shi G, Evert D, McLaughlin JL. Longimicins A–D: novel bioactive acetogenins from *Asimina longifolia* (Annonaceae) and structure-activity relationships of asimicin type of annonaceous acetogenins. *J Med Chem* 1996; 39: 1790–1796
- 104 Tominaga H, Maezaki N, Yanai M, Kojima N, Urabe D, Ueki R, Tanaka T. First total synthesis of longimicin D. *Eur J Org Chem* 2006; 2006: 1422–1429
- 105 Shi G, Kozlowski JF, Schwedler JT, Wood KV, MacDougall JM, McLaughlin JL. Muconin and mucocin: additional nonclassical bioactive acetogenins from *Rollinia mucosa*. *J Org Chem* 1996; 61: 7988–7989
- 106 Narayan RS, Borhan B. Synthesis of the proposed structure of mucocin via regio- and stereoselective tetrahydrofuran ring-forming strategies. *J Org Chem* 2006; 71: 1416–1429
- 107 Shi G, Zheng L, Gu Z-M, MacDougall JM, McLaughlin JL. Absolute stereochemistries of sylvaticin and 12,15-cis-sylvaticin, bioactive C-20,23-cis nonadjacent bistetrahydrofuran Annonaceous acetogenins, from *Rollinia mucosa*. *Heterocycles* 1995; 41: 1785–1796
- 108 Donohoe TJ, Harris RM, Burrows J, Parker J. Total synthesis of (+)-cis-sylvaticin: double oxidative cyclization reactions catalyzed by osmium. *J Am Chem Soc* 2006; 128: 13704–13705
- 109 Alkofahi A, Rupprecht JK, Liu YM, Chang CJ, Smith DL, McLaughlin JL. Gigantecin: a novel antimitotic and cytotoxic acetogenin, with nonadjacent tetrahydrofuran rings, from *Goniothalamus giganteus* (Annonaceae). *Experientia* 1990; 46: 539–541
- 110 Hoye TR, Eklov BM, Jeon J, Khoroosi M. Sequencing of three-component olefin metatheses: total synthesis of either (+)-gigantecin or (+)-14-deoxy-9-oxygigantecin. *Org Lett* 2006; 8: 3383–3386
- 111 Chavez D, Acevedo LA, Mata R, Jimenezin, a novel Annonaceous acetogenin from the seeds of *Rollinia mucosa* containing adjacent tetrahydrofuran-tetrahydropyran ring systems. *J Nat Prod* 1998; 61: 419–421
- 112 Bandur NG, Bruckner D, Hoffmann RW, Koert U. Total synthesis of jimenezin via an intramolecular allylboration. *Org Lett* 2006; 8: 3829–3831
- 113 Shi G, Alfonso D, Fatope MO, Zeng L, Gu ZM, Zhao GX, He K, MacDougall JM, McLaughlin JL. Mucocin: a new Annonaceous acetogenin bearing a tetrahydropyran ring. *J Am Chem Soc* 1995; 117: 10409–10410
- 114 Crimmins MT, Zhang Y, Diaz FA. Total synthesis of (–)-mucocin. *Org Lett* 2006; 8: 2369–2372
- 115 Alali FQ, Rogers L, Zhang Y, McLaughlin JL. Unusual bioactive Annonaceous acetogenins from *Goniothalamus giganteus*. *Tetrahedron* 1998; 54: 5833–5844
- 116 Strand D, Rein T. Total synthesis of pyranicin. *Org Lett* 2004; 7: 199–202
- 117 Strand D, Rein T. Synthesis of pyragonicin. *Org Lett* 2005; 7: 2779–2781
- 118 Takahashi S, Ogawa N, Koshino H, Nakata T. Total synthesis of the proposed structure for pyragonicin. *Org Lett* 2005; 7: 2783–2786
- 119 Quinn KJ, Isaacs AK, DeChristopher BA, Szklarz SC, Arvary RA. Asymmetric total synthesis of rollicocin. *Org Lett* 2005; 7: 1243–1245
- 120 Makabe H, Kimura Y, Higuchi M, Konno H, Murai M, Miyoshi H. Synthesis of (4R,15R,16R,21S)- and (4R,15S,16S,21S)-rollicocin, squamostolide, and their inhibitory action with bovine heart mitochondrial complex I. *Bioorg Med Chem* 2006; 14: 3119–3130
- 121 Hoppen S, Emde U, Friedrich T, Grubert L, Koert U. Natural-product hybrids: design, synthesis, and biological evaluation of quinone-Annonaceous acetogenins. *Angew Chem Int Ed Engl* 2000; 39: 2099–2102
- 122 Arndt S, Emde U, Baurle S, Friedrich T, Grubert L, Koert U. Quinone-Annonaceous acetogenins: synthesis and complex I inhibition studies of a new class of natural product hybrids. *Chem Eur J* 2001; 7: 993–1005
- 123 Duval R, Lewin G, Hocquemiller R. Semisynthesis of heterocyclic analogues of squamocin, a cytotoxic annonaceous acetogenin, by an unusual oxidative decarboxylation reaction. *Bioorg Med Chem* 2003; 11: 3439–3446
- 124 Duval RA, Lewin G, Peris E, Chahboune N, Garofano A, Droese S, Cortes D, Brandt U, Hocquemiller R. Heterocyclic analogues of squamocin as inhibitors of mitochondrial complex I. On the role of the terminal lactone of Annonaceous acetogenins. *Biochemistry* 2006; 45: 2721–2728
- 125 Duval RA, Poupon E, Romero V, Peris E, Lewin G, Cortes D, Brandt U, Hocquemiller R. Analogues of cytotoxic squamocin using reliable reactions: new insights into the reactivity and role of the α,β -unsaturated lactone of the Annonaceous acetogenins. *Tetrahedron* 2006; 62: 6248–6257
- 126 Kojima N, Fushimi T, Maezaki N, Tanaka T, Yamori T. Synthesis of hybrid acetogenins, α,β -unsaturated- γ -lactone-free nitrogen-containing heterocyclic analogues, and their cytotoxicity against human cancer cell lines. *Bioorg Med Chem Lett* 2008; 18: 1637–1641
- 127 Zeng BB, Wu Y, Yu Q, Wu YL, Li Y, Chen XG. Enantiopure simple analogues of Annonaceous acetogenins with remarkable selective cytotoxicity towards tumor cell lines. *Angew Chem Int Ed* 2000; 39: 1934–1937
- 128 Rodier S, Le Huerou Y, Renoux B, Doyon J, Renard P, Pierre A, Gesson JP, Gree R. Synthesis and cytotoxic activity of acetogenin analogues. *Bioorg Med Chem Lett* 2000; 10: 1373–1375
- 129 Yao ZJ, Wu HP, Wu YL. Polyether mimics of naturally occurring cytotoxic Annonaceous acetogenins. *J Med Chem* 2000; 43: 2484–2487
- 130 Jiang S, Liu ZH, Sheng G, Zeng BB, Cheng XG, Wu YL, Yao ZJ. Mimicry of Annonaceous acetogenins: enantioselective synthesis of a (4R)-hydroxy analogue having potent antitumor activity. *J Org Chem* 2002; 67: 3404–3408
- 131 Zeng BB, Wu Y, Jiang S, Yu Q, Yao ZJ, Liu ZH, Li HY, Li Y, Chen XG, Wu YL. Studies on mimicry of naturally occurring Annonaceous acetogenins: non-THF analogues leading to remarkable selective cytotoxicity against human tumor cells. *Chem Eur J* 2003; 9: 282–290
- 132 Rodier S, Le Huerou Y, Renoux B, Doyon J, Renard P, Pierre A, Gesson JP, Gree R. New cytotoxic analogues of Annonaceous acetogenins. *Anticancer Drug Des* 2001; 16: 109–117
- 133 Fujita D, Ichimaru N, Abe M, Murai M, Hamada T, Nishioka T, Miyoshi H. Synthesis of non-THF analogs of acetogenin toward simplified mimics. *Tetrahedron Lett* 2005; 46: 5775–5779
- 134 Liu HX, Huang GR, Zhang HM, Wu JR, Yao ZJ. Annonaceous acetogenin mimics bearing a terminal lactam and their cytotoxicity against cancer cells. *Bioorg Med Chem Lett* 2007; 17: 3426–3430
- 135 Liu HX, Shao F, Li GQ, Xun GL, Yao ZJ. Tuning the acyclic ether moiety of anticancer agent AA005 with conformationally constrained fragments. *Chem Eur J* 2008; 14: 8632–8639
- 136 Ye Q, Shi G, He K, McLaughlin JL. Chlorinated Annonaceous acetogenins and their bioactivities. *J Nat Prod* 1996; 59: 994–996

- 137 Kojima N, Hayashi H, Suzuki S, Tominaga H, Maezaki N, Tanaka T, Yamori T. Synthesis of C4-fluorinated solamins and their growth inhibitory activity against human cancer cell lines. *Bioorg Med Chem Lett* 2008; 18: 6451–6453
- 138 Gallardo T, Saez J, Granados H, Tormo JR, Velez ID, Brun N, Torres B, Cortes D. 10-Oximeguanacone, the first nitrogenated acetogenin derivative found to be a potent inhibitor of mitochondrial complex I. *J Nat Prod* 1998; 61: 1001–1005
- 139 Duret P, Hocquemiller R, Gantier JC, Figadere B. Semisynthesis and cytotoxicity of amino acetogenins and derivatives. *Bioorg Med Chem* 1999; 7: 1821–1826
- 140 Derbre S, Roue G, Poupon E, Susin Santos A, Hocquemiller R. Annonaceous acetogenins: the hydroxyl groups and THF rings are crucial structural elements for targeting the mitochondria, demonstration with the synthesis of fluorescent squamocin analogues. *Chembiochem* 2005; 6: 979–982
- 141 Alexander MD, Burkart MD, Leonard MS, Portonovo P, Liang B, Ding X, Joullie MM, Gullledge BM, Aggen JB, Chamberlin AR, Sandler J, Fenical W, Cui J, Charpure SJ, Polosukhin A, Zhang HR, Evans PA, Richardson AD, Harper MK, Ireland CM, Vong BG, Brady TP, Theodorakis EA, La Clair JJ. A central strategy for converting natural products into fluorescent probes. *Chembiochem* 2006; 7: 409–416
- 142 Maezaki N, Urabe D, Yano M, Tominaga H, Morioka T, Kojima N, Tanaka T. Synthesis of fluorescent solamin for visualization of cell distribution. *Heterocycles* 2007; 73: 159–164
- 143 Kojima N, Morioka T, Yano M, Suga Y, Maezaki N, Tanaka T. Convergent synthesis of fluorescence labeled solamin. *Heterocycles* 2009; 79: 387–393
- 144 Liu HX, Huang GR, Zhang HM, Jiang S, Wu JR, Yao ZJ. A structure-activity guided strategy for fluorescent labeling of Annonaceous acetogenin mimetics and their application in cell biology. *Chembiochem* 2007; 8: 172–177
- 145 Zafra-Polo MC, Gonzalez MC, Estornell E, Sahpaz S, Cortes D. Acetogenins from Annonaceae, inhibitors of mitochondrial complex I. *Phytochemistry* 1996; 42: 253–271
- 146 Sasaki S, Naito H, Maruta K, Kawahara E, Maeda M. Novel calcium ionophores: supramolecular complexation by the hydroxylated-bistetrahydrofuran skeleton of potent antitumor annonaceous acetogenins. *Tetrahedron Lett* 1994; 35: 3337–3340
- 147 Araya H, Fujimoto Y, Hirayama K. Structural elucidation of tetrahydrofuranic acetogenins by means of precursor-ion scanning method. *Yuki Gosei Kagaku Kyokaishi* 1994; 52: 765–777
- 148 Sasaki S, Maruta K, Naito H, Sugihara H, Hiratani K, Maeda M. New calcium-selective electrodes based on Annonaceous acetogenins and their analogs with neighboring bistetrahydrofuran. *Tetrahedron Lett* 1995; 36: 5571–5574
- 149 Sasaki S, Maruta K, Maemura R, Kawahara E, Maeda M. Novel acyclic ligands for metal cations based on the adjacent bistetrahydrofuran as analogs of natural Annonaceous acetogenins. *Tetrahedron* 1998; 54: 2401–2410
- 150 Peyrat JF, Mahuteau J, Figadere B, Cave A. NMR studies of Ca^{2+} complexes of Annonaceous acetogenins. *J Org Chem* 1997; 62: 4811–4815
- 151 Wu SN, Chiang HT, Chang FR, Liaw CC, Wu YC. Stimulatory effects of squamocin, an Annonaceous acetogenin, on Ca^{2+} -activated K^+ current in cultured smooth muscle cells of human coronary artery. *Chem Res Toxicol* 2003; 16: 15–22
- 152 Liaw CC, Yang YL, Chen M, Chang FR, Chen SL, Wu SH, Wu YC. Mono-tetrahydrofuran Annonaceous acetogenins from *Annona squamosa* as cytotoxic agents and calcium ion chelators. *J Nat Prod* 2008; 71: 764–771
- 153 Gu ZM, Zhou D, Lewis NJ, Wu J, Johnson HA, McLaughlin JL, Gordon J. Quantitative evaluation of Annonaceous acetogenins in monthly samples of paw paw (*Asimina triloba*) twigs by liquid chromatography/electrospray ionization/tandem mass spectrometry. *Phytochem Anal* 1999; 10: 32–38
- 154 McLaughlin JL. Paw paw and cancer: Annonaceous acetogenins from discovery to commercial products. *J Nat Prod* 2008; 71: 1311–1321
- 155 Cuendet M, Oteham CP, Moon RC, Keller WJ, Peaden PA, Pezzuto JM. Dietary administration of *Asimina triloba* (Paw Paw) extract increases tumor latency in N-methyl-N-nitrosourea-treated rats. *Pharm Biol* 2008; 46: 3–7
- 156 Escobar-Khondiker M, Hoellerhage M, Muriel MP, Champy P, Bach A, Depienne C, Respondek G, Yamada ES, Lannuzel A, Yagi T, Hirsch Etienne C, Oertel WH, Jacob R, Michel PP, Ruberg M, Hoeglinger GU. Annonacin, a natural mitochondrial complex I inhibitor, causes Tau pathology in cultured neurons. *J Neurosci* 2007; 27: 7827–7837