

# Dual function of an *Arabidopsis* transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression

Yoh Sakuma<sup>\*†</sup>, Kyonoshin Maruyama<sup>†</sup>, Feng Qin<sup>†</sup>, Yuriko Osakabe<sup>\*†</sup>, Kazuo Shinozaki<sup>‡§</sup>, and Kazuko Yamaguchi-Shinozaki<sup>\*†§¶</sup>

<sup>\*</sup>Graduate School of Agricultural and Life Sciences, University of Tokyo, Tokyo 113-8657, Japan; <sup>†</sup>Biological Resources Division, Japan International Research Center for Agricultural Sciences, Tsukuba, Ibaraki 305-8686, Japan; <sup>‡</sup>RIKEN Plant Science Center, Yokohama, Kanagawa 230-0045, Japan; and <sup>§</sup>Core Research for Evolutional Science and Technology, Japan Science and Technology Agency, Kawaguchi, Saitama 332-0012, Japan

Edited by Diter von Wettstein, Washington State University, Pullman, WA, and accepted August 16, 2006 (received for review July 5, 2006)

**Transcription factor DREB2A interacts with a cis-acting dehydration-responsive element (DRE) sequence and activates expression of downstream genes involved in drought- and salt-stress response in *Arabidopsis thaliana*. Intact DREB2A expression does not activate downstream genes under normal growth conditions. A negative regulatory domain exists in the central region of DREB2A, and deletion of this region transforms DREB2A to a constitutive active form (*DREB2A CA*). We carried out microarray analysis of transgenic *Arabidopsis*-overexpressing *DREB2A CA* and found that the overexpression of *DREB2A CA* induces not only drought- and salt-responsive genes but also heat-shock (HS)-related genes. Moreover, we found that transient induction of the *DREB2A* occurs rapidly by HS stress, and that the sGFP-DREB2A protein accumulates in nuclei of HS-stressed cells. DREB2A up-regulated genes were classified into three groups based on their expression patterns: genes induced by HS, genes induced by drought stress, and genes induced by both HS and drought stress. DREB2A up-regulated genes were down-regulated in DREB2A knockout mutants under stress conditions. Thermotolerance was significantly increased in plants overexpressing *DREB2A CA* and decreased in *DREB2A* knockout plants. Collectively, these results indicate that DREB2A functions in both water and HS-stress responses.**

AP2-type transcription factor | drought-stress tolerance | heat-stress tolerance | knockout mutants | microarray analysis

Environmental stresses such as heat, cold, drought, and high salinity influence plant growth and productivity. Plants respond and adapt to these stresses to survive under stress conditions at physiological and biochemical levels. These stresses have been shown to induce the expression of genes with various functions in a variety of plants (1–4). The cis- and trans-acting elements that function in stress-responsive gene expression have been precisely analyzed to elucidate the molecular mechanisms of gene expression in response to drought stress (5). The dehydration-responsive element (DRE) with the core sequence A/GCCGAC was identified as a cis-acting promoter element that regulates gene expression in response to drought, salt, and cold stresses in *Arabidopsis* (6). A similar motif was identified as the C-repeat and low-temperature-responsive element in cold-inducible genes (7, 8).

*Arabidopsis* cDNAs encoding the ethylene-responsive element-binding factor/APETALA2 (ERF/AP2)-type DRE-binding proteins (DREB) *CBF1*, *DREB1A*, and *DREB2A* have been isolated with a yeast one-hybrid screening (9, 10). They specifically bind to the DRE/C-repeat sequence and activate the transcription of genes driven by the DRE sequence in *Arabidopsis*. Three DREB1/C-repeat-binding factor (CBF) proteins are encoded by genes that lie in tandem on chromosome 4 in the order of *DREB1B/CBF1*, *DREB1A/CBF3*, and *DREB1C/CBF2* (10, 11). *Arabidopsis* also contains two DREB2 proteins, DREB2A and DREB2B (10). Although the expression of all three *DREB1/CBF* genes is induced

by cold stress but not by drought and salt stress, both *DREB2* genes are induced by drought and salt stress (10). Both DREB1/CBF and DREB2 proteins bind to DRE, but DREB1/CBFs are thought to function in cold-responsive gene expression, whereas DREB2s are involved in drought-responsive gene expression.

Overexpression of DREB1/CBF driven by the 35S *CaMV* promoter causes growth retardation under normal growth conditions and increases stress tolerance of drought, high-salinity, and freezing in transgenic *Arabidopsis* (10, 12, 13). More than 40 downstream targets of DREB1/CBF have been identified by using microarray analysis (14–17). Many of their protein products, such as RNA-binding proteins, sugar transport proteins, desaturases, late embryogenesis-abundant (LEA) proteins, and osmoprotectant biosynthesis-proteins, are known to function against stresses (4) and are probably responsible for the stress tolerance of the transgenic plants. Transcription factors were also downstream targets, which suggests the existence of further regulation of gene expression downstream of the DRE/DREB regulon (16–18). Conserved sequences in the promoter regions of the genes directly downstream of *DREB1A* were analyzed, and A/GCCGACNT was found in their promoter regions between –51 and –450 as a consensus DRE (16).

DREB2 also has a conserved ERF/AP2 DNA-binding domain and recognizes the DRE sequence. A genome search of *Arabidopsis* revealed at least six *DREB2* homologues other than *DREB2A* and *DREB2B*. Because *DREB2A* and *DREB2B* are the only genes induced strongly by drought and high salinity among the eight genes of DREB2-type proteins, they are thought to be major transcription factors functioning under drought- and salt-stress conditions (19, 20). Domain analysis of DREB2A by using *Arabidopsis* protoplasts revealed that a negative regulatory domain exists in the central region of DREB2A, and deletion of this region transforms DREB2A to a constitutive active form (21). Overexpression of the constitutive active form of DREB2A (*35S:DREB2A CA*) significantly increased tolerance to drought

This paper results from the Arthur M. Sackler Colloquium of the National Academy of Sciences, "From Functional Genomics of Model Organisms to Crop Plants for Global Health," held April 3–5, 2006, at The National Academy of Sciences in Washington, DC. The complete program is available on the NAS web site at [www.nasonline.org/functional.genomics](http://www.nasonline.org/functional.genomics).

Author contributions: Y.S., K.S., and K.Y.-S. designed research; Y.S., K.M., F.Q., and Y.O. performed research; K.M. analyzed data; and Y.S. and K.Y.-S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS direct submission.

Abbreviations: DRE, dehydration-responsive element; HS, heat shock; HSP, HS protein; HSF, HS transcription factor; CBF, C-repeat-binding factor.

Data deposition: The complete set of microarray data has been deposited in the European Bioinformatics Institute ArrayExpress database, [www.ebi.ac.uk/arrayexpress](http://www.ebi.ac.uk/arrayexpress) (accession nos. E-MEXP-601, MEXP-602, and MEXP-603).

<sup>†</sup>To whom correspondence should be addressed. E-mail: [akys@mail.ecc.u-tokyo.ac.jp](mailto:akys@mail.ecc.u-tokyo.ac.jp).

© 2006 by The National Academy of Sciences of the USA

stress but only slightly increased tolerance to freezing. Overexpression of this gene also resulted in growth retardation in transgenic *Arabidopsis* plants. Microarray analyses of the 35S:*DREB2A CA* plants revealed that *DREB2A* regulates the expression of many drought-inducible genes. However, some genes regulated by *DREB2A* are not regulated by *DREB1A*, which also recognizes DRE/C-repeat but functions in cold-stress-responsive gene expression (21). Promoter analysis of the *DREB1A*- and *DREB2A*-regulated genes and gel mobility-shift assay by using both recombinant proteins indicated that *DREB1A* and *DREB2A* exhibit different DNA-binding specificities. *DREB1A* has the highest affinity to A/GCCGACNT, whereas *DREB2A* preferentially binds ACCGAC (16, 21). This difference controls the induction of different downstream genes between *DREB1A* and *DREB2A*. Although the *DREB2A*-regulated genes play important roles in drought-stress tolerance, they are not sufficient to withstand freezing stress.

In contrast to drought- and cold-stress responses, the acquisition of tolerance to heat-shock (HS) stress is correlated to the induction of HS proteins (HSPs). The expression of some *HSP* genes is reported to be induced not only by HS but also by drought stress (22). HSPs act as molecular chaperones by maintaining homeostasis of protein folding and thus help to maintain the metabolic and structural integrity of cells (22–24). HSP expression is regulated at the transcriptional level by HS transcription factors (HSFs), which recognize the conserved HS element in the promoter regions of *HSP* genes (24–26). *Arabidopsis* contains 21 different *HSF* genes (27), and this high multiplicity suggests the importance of backup and diversification of HSFs. However, microarray and genetic data indicate that different regulatory systems contribute to heat tolerance at different stages of the plant life cycle, and that different gene sets may control the complex multigenic process for acquiring thermotolerance (28, 29).

In this study, we carried out a microarray analysis by using 35S:*DREB2A CA* plants and found that many HS-inducible genes were up-regulated. We showed that the expression of *DREB2A* was rapidly and transiently induced by HS and dehydration treatments, and that the *DREB2A* protein is accumulated by HS stress and localized in nuclei. Two T-DNA-tagged *dreb2a* mutants were used for further functional analysis. Many up-regulated genes in the 35S:*DREB2A CA* plants were down-regulated in the *dreb2a* mutants. Here we report the HS stress tolerance of the 35S:*DREB2A CA* transgenic plants and HS sensitivity of the *dreb2a* mutants and discuss the role of *DREB2A* in both HS and drought stress responses.

## Results

**Microarray Analysis of *DREB2A* Up-Regulated Genes in Transgenic *Arabidopsis* Plants Overexpressing the Constitutive Active Form of *DREB2A*.** We have shown that the *DREB2A* protein has a negative regulatory domain in the region between amino acid residues 136 and 165. Consequently, deletion of this region transforms *DREB2A* to a constitutive active form (*DREB2A CA*) (21). Furthermore, overexpression of *DREB2A CA* with the *CaMV 35S* promoter induces expression of many stress-responsive genes (21). To understand the entire transcriptional network of *DREB2A*, we analyzed up-regulated genes in the transgenic plants overexpressing *DREB2A CA* (35S:*DREB2A CA*) by using an Agilent *Arabidopsis* 2 Oligo Microarray (Agilent Technologies, Palo Alto, CA), covering >21,000 genes. We used 3-wk-old *Arabidopsis* plants of two independent lines overexpressing *DREB2A CA*, 35 S:*DREB2A CA-a*, and 35S:*DREB2A CA-b* (21). Two experiments were performed for each RNA sample by using different labels, cy3 or cy5. In comparison to vector control plants, 483 genes with the average of expression ratios greater than two times in the 35S:*DREB2A CA* plants are listed in Table 1, which is published as supporting information on the PNAS

web site. Among the 483 up-regulated genes, 68 and 107 genes were induced by drought and salt stress, respectively, and 120 genes were induced by drought and/or salt more than two times (K.M. and K.Y.-S., unpublished data; Fig. 5, which is published as supporting information on the PNAS web site).


We subsequently focused on 36 *DREB2A* up-regulated genes that exhibited remarkable expression ratios greater than eight times, as summarized in Table 2, which is published as supporting information on the PNAS web site. Among the 36 genes, 14 and 19 are induced by drought and high salinity, respectively. The DRE sequence (A/GCCGAC), which is a recognition sequence of *DREB2A*, is found in the 1,000-bp promoter region of 29 genes among these 36 up-regulated genes. Expression for several of the identified up-regulated genes was confirmed by using quantitative RT-PCR and shown to increase in the 35S:*DREB2A CA* plants. With the exception of *At1g52690* (LEA protein), their expression levels directly to *DREB2A* expression (see Fig. 6, which is published as supporting information on the PNAS web site). Among the 36 up-regulated genes in 35S:*DREB2A CA*, genes for LEA proteins, dehydrins, and *COR15A*, which function in acquisition of stress tolerance to drought and high salinity, were identified (5). Interestingly, a gene for a HSF, *AtHsfA3*, was included in the up-regulated genes, and its expression ratio was highest in the 35S:*DREB2A CA* plants (except *DREB2A*). The *HSF* gene family consists of genes that encode transcription factors involved in HS-inducible gene expression (27). In addition, genes for chloroplast-localized small HSP (CPsHSP, *At1g52560*), DnaJ-like protein (*At5g37440*), HSP70 (*At3g12580*), and HSP18.2 (*At5g59720*), which likely function in HS stress response, were also overexpressed in the transgenic plants. These results suggest that *DREB2A* functions not only in drought- and salt-stress responsive gene expression but also in HS stress-responsive gene expression.

To analyze the relationship between HS stress-inducible genes and the up-regulated genes in the 35S:*DREB2A CA* plants, a similar microarray analysis was performed. Total RNA samples were prepared from 3-wk-old plants treated at 37°C for 0.5 or 5 h and nontreated control plants (Tables 3 and 4, which are published as supporting information on the PNAS web site). Heat-inducible genes (268 and 778) with expression ratios greater than two times were identified at 0.5 and 5 h, respectively. Among the 36 up-regulated genes in the 35S:*DREB2A CA* plants listed in Table 2, nine genes were shown to be induced by heat. Expression of *DREB2A* is also induced by HS stress. Interestingly, induction for four of these nine genes, such as *At1g52560* (CPsHSP), *HSP18.2*, *At4G36010* (thaumatin-like protein), and *AtCYP18-1*, was heat-specific, whereas the other five genes and *DREB2A* were induced by drought, salt, and heat stress (Table 2).

**Expression Analysis of *DREB2A* and Its Up-Regulated Genes Under High-Temperature Stress Condition.** We performed quantitative RT-PCR to confirm the expression of *DREB2A* during heat stress (Fig. 1A). The expression of *DREB2A* was rapidly and transiently induced by HS treatment (37°C) and peaked ≈30 min. Then, the expression of *DREB2A* drastically decreased to a relatively low level that was maintained until 10 h after the treatment. In contrast, *DREB2A* was gradually induced by drought and salt stress during 10 h. We also analyzed the effect of H<sub>2</sub>O<sub>2</sub>-induced oxidative stress and found that, although *DREB2A* was also gradually induced by H<sub>2</sub>O<sub>2</sub>, the level of its induction was lower than those by drought and salt stress. We then analyzed the effects of various plant hormone treatments. *DREB2A* was not induced by any plant hormones such as abscisic acid, cytokinin, ethephon, auxin, methyl jasmonate, and salicylic acid (data not shown). We analyzed expression of a *DREB2A* homolog and found a similar rapid induction of *DREB2B* in response to HS stress (Fig. 1B). Similarly, the *DREB2B* gene was also gradually induced by drought, high salinity, and H<sub>2</sub>O<sub>2</sub> (Fig. 1B).

To ascertain that HS stress induces accumulation of the






**Fig. 3.** Analysis of *DREB2A* knockout mutants. (A) A scheme of the *Arabidopsis DREB2A* gene. Exons (open boxes) and an intron (line) are indicated. The positions of the T-DNA insertions are shown (not to scale). (B–E) Expression of *DREB2A* (B), *DREB2A* downstream genes induced by both HS and drought (C), *DREB2A* downstream genes induced by HS (D), and *DREB2A* downstream genes induced by drought (E) was analyzed in wild-type and *dreb2a* plants by quantitative RT-PCR (the highest expression level was set to 100). Data represent means and standard errors of three replications.

3C shows the expression of *RD17*, *At4g36010* (thaumatin-like protein), and *AtHsfA3* that are induced by both HS and drought stress. In contrast to wild-type plants, HS-inducible expression of these genes was decreased remarkably in both *dreb2a-1* and *dreb2a-2*. Drought-inducible expression of the genes was also decreased, but their low-level expression was still detected even under stress conditions. Expression of *HSP70*, *HSP18.2*, and *AtCYP18-1* that are induced by HS stress but not drought stress was also decreased in *dreb2a-1* and *dreb2a-2* (Fig. 3D). Although a significant decrease of drought-stress-inducible expression of *At1g52690* (LEA protein) was observed in knockout mutants, the response of *RD29A* was unclear (Fig. 3E).

Then, we compared the expression profiles in the two *DREB2A* mutants of 3-wk-old plants under drought- or HS-stress conditions with that in the wild-type plants by using the Agilent *Arabidopsis* 2 Oligo Microarray. Genes with expression ratios  $<0.5$  on the average of two mutant lines are listed in Tables 5 and 6, which are published as supporting information on the PNAS web site. Eleven and 50 genes showed decreased expression in the *DREB2A* mutants under drought- and HS-stress conditions, respectively. Among these down-regulated genes, 31 were over-expressed more than two times in the *35S:DREB2A CA* plants (Table 7, which is published as supporting information on the PNAS web site). We searched for conserved sequences in the 31 promoter regions of the 36 *DREB2A* up-regulated genes by using the MEME program (30). The 1,000-bp promoter regions upstream of putative transcriptional initiation sites that are the 5' end of the full-length cDNA clones were used for this search (31), and we found a highly conserved sequence (A/GCCGAC) that included the DRE-core motif. We previously reported that *DREB2A* has a higher affinity to ACCGAC than GCCGAC (21). Therefore, the frequency of ACCGAC and GCCGAC in the promoter region of the 35 genes was verified. Twenty-eight genes had DRE, A/GCCGAC, sequence(s) within 1,000-bp promoter regions. Among the 28 genes, 24 (86%) have at least one ACCGAC sequence in their promoter regions.

**HS-Stress Tolerance of the *35S:DREB2A* Plants and the *DREB2A* Knockout Mutants.** Expression of *DREB2A* was induced by HS transiently and significantly. Moreover, microarray analysis of the

*35S:DREB2A CA* plants showed that *DREB2A* up-regulates the expression of many HS-inducible genes in the plants. We were interested to determine whether the *35S:DREB2A* plants had an improved tolerance to heat stress. Seven-day-old vector control and transgenic plants, germinating on a filter paper premoistened by liquid germination medium, were grown at 22°C and then subjected to HS treatment at 45°C. As shown in Fig. 4A, two



**Fig. 4.** HS-stress tolerance of the *35S:DREB2A CA* and *dreb2a* plants. (A) One-week-old seedlings of the vector control or the *35S:DREB2A CA* plants were treated at 22°C or 45°C for 1 h. (B) One-week-old seedlings of wild-type or *dreb2a* plants were preincubated at 37°C for 1 h and then treated at 22°C or 49°C for 1 h. After treatment, plants were grown under normal conditions for 1 wk. Percentages of surviving plants and numbers of surviving plants per total numbers of tested plants are indicated around the photographs. More than 15 plants were used per test, and each test was repeated three times. \*, These plants had significantly lower survival rates than the wild-type plants ( $\chi^2$  test,  $P < 0.05$ ). \*\*, These plants had significantly higher or lower survival rates than the vector control or the wild-type plants ( $\chi^2$  test,  $P < 0.01$ ).

transgenic lines of the *35S:DREB2A CA-a*, *35S:DREB2A CA-b*, and vector control plants grew well under 22°C. However, when the plants were exposed at 45°C for 1 h, only 13.4% of the vector control plants survived, whereas 88.0% of *35S:DREB2A CA-a* and 85.4% of *35S:DREB2A CA-b* survived during the subsequent 7-day recovery at 22°C. These results clearly indicate augmented thermotolerance of the *35S:DREB2A CA* plants.

We then compared the thermotolerance of the *DREB2A* knockout mutants to that of wild-type plants. Seven-day-old plants grown on a filter paper were subjected to 37°C for 1 h, then followed by 49°C for 1 h. Wild-type plants (78.4%) survived during the subsequent 7-day recovery at 22°C, whereas 62.2% and 51.4% of *dreb2a-1* and *dreb2a-2* survived, respectively (Fig. 4B). Thus, the phenotype of these mutants in survival under HS stress contrasted with those of the transgenic plants overexpressing *DREB2A CA*, indicating that *DREB2A* functions in improving HS-stress tolerance.

## Discussion

The DRE contains the core sequence A/GCCGAC and has been identified as a cis-acting promoter element in regulating gene expression in response to drought, salt, and cold stresses in *Arabidopsis* (6). *DREB2A* was subsequently isolated as a cDNA encoding a DREB protein by using a yeast one-hybrid screening (10). Therefore, the function of *DREB2A* under these stresses has been well studied (10, 21), but the effect of HS stress has not been previously studied. However, in our current study, we clearly identified the up-regulation of many HS-related genes such as *AtHsfA3* and *HSPs*, as a result of overexpressing *DREB2A CA* (Table 2 and Fig. 6). We subsequently confirmed that these genes were induced by HS stress (Table 2 and Fig. 2). Moreover, we found that the *DREB2A* gene is also strongly induced by HS stress (Fig. 1A), and that the sGFP-*DREB2A* protein is stabilized in the nuclei in plant cells by HS stress (Fig. 1C). We hypothesized that *DREB2A* is important for the regulation of gene expression in response to heat stress as well as drought and salt stress. We later confirmed this hypothesis by using the *dreb2a* mutants and the *35S:DREB2A CA* plants. Elimination of the *DREB2A* gene decreased HS- and drought-stress-inducible expression of *DREB2A* up-regulated genes (Fig. 3). Finally, we found that the *35S:DREB2A CA* plants showed enhanced thermotolerance, and that the *dreb2a* mutants displayed reduced survival under HS condition (Fig. 4).

Four of five HS-related genes, *AtHsfA3* and three *HSPs*, that are up-regulated in the *35S:DREB2A CA* plants with remarkable expression ratios greater than eight times, showed strong inducible expression by HS stress (Table 2). *AtHsfA3* is a member of the *Hsf* gene family, which consists of 21 genes belonging to three major classes that are based on structural differences, HsfA, HsfB, and HsfC. HsfAs, have been shown to be the major transcription factors that function in heat-induced expression of *HSP* genes (24). *AtHsfA3* might act as a transcription factor in HS-stress-responsive gene expression, but detailed function of *AtHsfA3* has not yet been identified. *HSP70* (*At3g12580*) encodes one of the *HSP70* chaperones that have an essential function in preventing aggregation and in assisting refolding of nonnative proteins under stress conditions. They are also involved in the degradation of unstable proteins by targeting the proteins to lysosomes or proteasomes (22). Overexpression of *HSP70* genes in transgenic plants has been shown to correlate positively with the acquisition of thermotolerance and results in enhanced tolerance to drought, salt, and heat stress (22). Two genes, *At1g52560* (CPsHSP) and *HSP18.2*, encode small HS proteins (sHSPs). Although the sHSPs alone are not able to refold nonnative proteins, they probably bind to unfolded proteins and facilitate further refolding by HSP complexes (22). These HSPs probably contribute to the acquisition of stress tolerance not only to heat stress but also to drought and salt stress

in the *35S:DREB2A CA* plants. Five more genes such as *RD17* encoding a group two LEA protein (dehydrin), *AtGolS1* encoding a galactinol synthase (32), *MT2A* encoding a putative detoxification-related gene (33), *AT4g36010* (thaumatin-like protein) encoding a PR-5 like protein related to biotic stress, and *AtCYP18-1* encoding an unknown protein, were shown to be HS-inducible. These genes may have important roles to increase HS stress tolerance in the transgenic plants. On the other hand, thermotolerance of the *dreb2a* mutants was not dramatically reduced (Fig. 4). Our microarray analysis revealed that 956 genes showed HS-responsive expression, and 110 of them are up-regulated by *DREB2A* (Tables 3 and 4). Functionally redundant proteins are encoded by *DREB2A*-regulated and -unregulated HS-responsive genes. It appears that the latter genes are capable of complementing the *DREB2A* mutation.

Our microarray analysis of *35S:DREB2A CA* plants confirmed that many drought-stress- or HS-inducible genes are potentially downstream of *DREB2A* (Table 2 and Fig. 2). However, an analysis with a gain-of-function method with constitutive and ectopic gene expression sometimes creates false responses. Therefore, we confirmed the expression of these *DREB2A* up-regulated genes in the *dreb2a* mutants by using quantitative RT-PCR and microarray analyses. Six *DREB2A* up-regulated genes showed clear down-regulation of stress-inducible expression in both *dreb2a-1* and -2 plants (Fig. 3 C and D). Similar to *DREB2A*, most of the genes such as *RD17*, *HSP70*, *AtCYP18-1*, and *HSP18.2* showed rapid and transient expression in response to HS stress, but their expression was slightly delayed. Although their expression peaked 1 h after stress treatment (Fig. 2 A and B), HS-inducible *DREB2A* expression peaked between 30 min and 1 h (Fig. 1A). This time lag supports the hypothesis that these genes are downstream targets of *DREB2A*. Drought-stress-inducible expression of *At1g52690* (LEA protein) also decreased in the *dreb2a* mutants (Fig. 3E). In addition, stress-inducible expression of *At1g01470* (LEA14), *COR15A*, *At3g17520*, *XERO2*, and *RD29A* was reduced in the microarray analysis of the *dreb2a* mutants (Table 7). All of these genes have DRE in their promoter regions, suggesting they are direct downstream genes of *DREB2A*.

Among 483 *DREB2A* up-regulated genes with the expression ratios greater than two times, 68 and 107 genes showed drought- and salt-responsive gene expression, respectively (Fig. 5). Fifty-five (80%) of the 68 drought-stress-inducible genes are also induced by salt stress. This extensive overlap suggests that a common regulatory mechanism exists between drought- and salt-induced gene expression of the *DREB2A* up-regulated genes. One hundred ten genes showed HS-responsive expression, and only 38 of them showed drought- and/or salt stress-responsive expression (Fig. 5). HS- and drought-specific expression of the *DREB2A* up-regulated genes was also detected by quantitative RT-PCR. Genes shown in Fig. 2B responded to HS but not to drought stress. In contrast, genes shown in Fig. 2C responded to drought but not to HS stress. These results suggest that distinct regulatory mechanisms exist between HS- and drought-/salt-induced gene expression of the *DREB2A* up-regulated genes. Recently, Diaz-Martin *et al.* (34) reported that a sunflower *DREB2*-type protein, HaDREB2, interacts with a heat-stress factor A9 (HaHSFA9), and these proteins synergistically induce expression of *HaHSP17.6*. Interaction of HS or drought-/salt-stress specific cofactors such as HaAHSFA9 may be necessary to activate *DREB2A*. Otherwise, HS- or drought-/salt stress-specific repressors may inhibit the gene expression activated by *DREB2A*.

When sGFP was fused to full-length (FL) *DREB2A* (sGFP-*DREB2A*) and expressed in *Arabidopsis*, only weak green fluorescence was observed in the nucleus under unstressed control conditions. In contrast, strong sGFP signals were observed in the nucleus of the transgenic plants expressing chimeric genes with

sGFP fused to DREB2A CA. DREB2A CA lacks the negative regulatory domain that contains a PEST sequence (21). The FL DREB2A protein containing the PEST sequence may be degraded rapidly by the ubiquitin–proteasome system (21). However, strong GFP fluorescence was detected in the nuclei of transgenic plants expressing *sGFP-DREB2A* under HS-stress conditions (Fig. 1C). These data indicate that the DREB2A protein is stabilized by HS stress and nuclear-localized during stress conditions. The PEST sequence of DREB2A (RSDASEVTSTSSQSEVCTVETPGCV) contains many phosphorylation target sites of protein kinases such as PKC and CKII. Because the phosphorylation of the PEST sequence has been reported to be important for protein degradation (35), it is plausible that the modification of the DREB2A protein by these kinases might be important for its stabilization under stress conditions.

In summary, DREB2A is shown to play an important role in crosstalk between drought- and heat-stress responses. *DREB2A* is HS-stress-inducible and contains an HS element in its promoter. Moreover, DREB2A controls HSF as one of the downstream genes. This complex cascade of transcription may enable plants to properly respond to both heat and drought stress.

### Materials and Methods

**Plant Materials.** *Arabidopsis* plants were grown, transformed, and treated as described (21). The *dre2a-1* (line 379F02) and *dre2a-2* (line 179C04) mutants were obtained from the GABI-Kat (36). For HS-stress treatment, *Arabidopsis* seedlings were grown on agar plates at 22°C for 3 wk and then transferred to 37°C.

**Quantitative RT-PCR and Microarray Analysis.** Total RNA was extracted with TriZol Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions, and quantitative RT-PCR was performed as described (21). The primer pairs used in the real-time PCR are shown in Table 8, which is published as supporting information on the PNAS web site. Microarray analysis and data mining were carried out as described (37).

**Observation of Subcellular Localization of Green Fluorescent Signals in Transgenic Plants.** The DNA sequence of sGFP fused with the DREB2A coding sequence and Nos terminator (NosT) of 35S:*GFP-DREB2A* plasmid (21) was amplified by PCR and cloned into the pGreenII0029 vector (38) by PstI and SalI sites. The 1.8 kbp of the DREB2A promoter sequence (19) was subsequently inserted in front of *sGFP-DREB2A-NosT* by the PstI site. Plant transformation was carried out as described (10). GFP fluorescence was analyzed by a confocal laser-scanning microscope LSM5 PASCAL (Zeiss, Oberkochen, Germany).

**HS Tolerance Test.** *Arabidopsis* seeds were germinated and grown on two layers of filter paper containing 4 ml of germination medium (6) in 90-mm plastic Petri dishes. Petri dishes were sealed with Micropore surgical tape (3M Health Care, St. Paul, MN) and incubated at 22°C. Twenty-five to 30 seeds were sown for each experiment, and seedlings that germinated within 7 days after sowing were subjected to the HS-tolerance test. Throughout the incubation period, 2 ml of sterilized water was added to Petri dishes every 3 days. Seven-day-old seedlings were subjected to HS treatment on a paper box (width 13 mm × depth 130 mm × height 50 mm) that was placed in a hybridization oven (HB-80, TAITEC, Saitama, Japan). After HS treatment, plants were grown under normal conditions for 1 wk, and the number of surviving plants was counted.

We thank Dr. Y. Niwa (Shizuoka Prefectural University, Shizuoka, Japan) for providing the *sGFP* gene. We are grateful for the excellent technical support provided by Kyoko Yoshiwara, Hiroko Sado, Ekuko Ohgawara, and Kaori Amano, and for skillful editorial assistance by Masami Toyoshima of the Japan International Research Center for Agricultural Sciences. We also thank the National Institute of Agrobiological Sciences for supporting the 22k microarray analysis. This work was supported in part by the Program for Promotion of Basic Research Activities for Innovative Biosciences, Core Research for Evolutional Science and Technology, and by a project grant from the Ministry of Agriculture, Forestry, and Fisheries, Japan.

- Thomashow MF (1999) *Annu Rev Plant Physiol Plant Mol Biol* 50:571–599.
- Zhu JK (2002) *Annu Rev Plant Biol* 53:247–273.
- Shinozaki K, Yamaguchi-Shinozaki K (2003) *Curr Opin Plant Biol* 6:410–417.
- Bartels D, Sunkar R (2005) *Crit Rev Plant Sci* 24:23–58.
- Yamaguchi-Shinozaki K, Shinozaki K (2006) *Annu Rev Plant Biol* 57:781–803.
- Yamaguchi-Shinozaki K, Shinozaki K (1994) *Plant Cell* 6:251–264.
- Baker SS, Wilhelm KS, Thomashow MF (1994) *Plant Mol Biol* 24:701–713.
- Jiang C, Lu B, Singh J (1996) *Plant Mol Biol* 30:679–684.
- Stockinger EJ, Gilmour SJ, Thomashow MF (1997) *Proc Natl Acad Sci USA* 94:1035–1040.
- Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1998) *Plant Cell* 10:1391–1406.
- Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM, Thomashow MF (1998) *Plant J* 16:433–442.
- Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF (1998) *Science* 280:104–106.
- Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1999) *Nat Biotechnol* 17:287–291.
- Seki M, Narusaka M, Abe H, Kasuga M, Yamaguchi-Shinozaki K, Carninci P, Hayashizaki Y, Shinozaki K (2001) *Plant Cell* 13:61–72.
- Fowler S, Thomashow MF (2002) *Plant Cell* 14:1675–1690.
- Maruyama K, Sakuma Y, Kasuga M, Ito Y, Seki M, Goda H, Shimada Y, Yoshida S, Shinozaki K, Yamaguchi-Shinozaki K (2004) *Plant J* 38:982–993.
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF (2005) *Plant J* 41:195–211.
- Sakamoto H, Maruyama K, Sakuma Y, Meshi T, Iwabuchi M, Shinozaki K, Yamaguchi-Shinozaki K (2004) *Plant Physiol* 136:2734–2746.
- Nakashima K, Shinwari ZK, Sakuma Y, Seki M, Miura S, Shinozaki K, Yamaguchi-Shinozaki K (2000) *Plant Mol Biol* 42:657–665.
- Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K, Yamaguchi-Shinozaki K (2002) *Biochem Biophys Res Commun* 290:998–1009.
- Sakuma Y, Maruyama K, Osakabe K, Qin F, Seki M, Shinozaki K, Yamaguchi-Shinozaki K (2006) *Plant Cell* 18:1292–1309.
- Wang W, Vinocur B, Shoseyov O, Altman A (2004) *Trends Plant Sci* 9:244–252.
- Vierling E (1991) *Annu Rev Plant Physiol Plant Mol Biol* 42:579–620.
- Sung DY, Guy CL (2003) *Plant Physiol* 132:979–987.
- Wu C (1995) *Annu Rev Cell Dev Biol* 11:441–469.
- Miller G, Mittler R (2006) *Ann Bot* 98:279–288.
- Nover N, Bharti N, Doring P, Mishra SK, Ganguli A, Scharf KD (2001) *Cell Stress Chaperones* 6:177–189.
- Larkindale J, Hall JD, Knight MR, Vierling E (2005) *Plant Physiol* 138:882–897.
- Busch W, Wunderlich M, Schoffl F (2006) *Plant J* 41:1–14.
- Bailey TL, Elkan C (1995) *Proc Int Conf Intell Syst Mol Biol* 3:21–29.
- Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, et al. (2002) *Plant J* 31:279–292.
- Taji T, Ohsumi C, Iuchi S, Seki M, Kasuga M, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K (2002) *Plant J* 29:417–426.
- Zhou J, Goldsbrough P (1994) *Plant Cell* 6:875–884.
- Diaz-Martin J, Alguera C, Prieto-Dapena P, Espinosa JM, Jordano J (2005) *Plant Physiol* 139:1483–1494.
- Salmeron J, Janzen J, Soneji Y, Bump N, Kamens J, Allen H, Ley SC (2001) *J Biol Chem* 276:22215–22222.
- Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B (2003) *Plant Mol Biol* 53:247–259.
- Furihata T, Maruyama K, Fujita Y, Umezawa T, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K (2006) *Proc Natl Acad Sci USA* 103:1988–1993.
- Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM (2000) *Plant Mol Biol* 42:819–832.