

A membrane-tethered transcription factor defines a branch of the heat stress response in *Arabidopsis thaliana*

Hongbo Gao*, Federica Brandizzi*[†], Christoph Benning[‡], and Robert M. Larkin*^{‡§}

*Michigan State University–Department of Energy Plant Research Laboratory, [†]Department of Biochemistry and Molecular Biology, and [‡]Department of Plant Biology, Michigan State University, East Lansing, MI 48824

Communicated by Michael F. Thomashow, Michigan State University, East Lansing, MI, August 28, 2008 (received for review December 14, 2007)

In plants, heat stress responses are controlled by heat stress transcription factors that are conserved among all eukaryotes and can be constitutively expressed or induced by heat. Heat-inducible transcription factors that are distinct from the “classical” heat stress transcription factors have also been reported to contribute to heat tolerance. Here, we show that *bZIP28*, a gene encoding a putative membrane-tethered transcription factor, is up-regulated in response to heat and that a *bZIP28* null mutant has a striking heat-sensitive phenotype. The heat-inducible expression of genes that encode BiP2, an endoplasmic reticulum (ER) chaperone, and HSP26.5-P, a small heat shock protein, is attenuated in the *bZIP28* null mutant. An estradiol-inducible *bZIP28* transgene induces a variety of heat and ER stress-inducible genes. Moreover, heat stress appears to induce the proteolytic release of the predicted transcription factor domain of *bZIP28* from the ER membrane, thereby causing its redistribution to the nucleus. These findings indicate that *bZIP28* is an essential component of a membrane-tethered transcription factor–based signaling pathway that contributes to heat tolerance.

BiP2 | *bZIP28* | endoplasmic reticulum | signal transduction | thermotolerance

As sessile organisms, plants must complete their life cycle in an ever-changing environment. To help cope with these fluctuating environmental conditions, plants have evolved particular responses for individual stresses and various combinations of stresses (1). The heat shock response involves the accumulation of molecular chaperones or heat shock proteins (HSPs) that stabilize partially unfolded proteins (2, 3). The induction of HSPs by heat stress is controlled by heat stress transcription factors (HSFs) that bind heat stress elements (5'-AGAAAnnTTCT-3') and are conserved among eukaryotes (2, 4). Plants are unusual in having 20 or more genes that encode HSFs, whereas other eukaryotes contain 1 to 3 of these genes. The expansion of HSF genes in plants is thought to result from a combination of diversity in activity and expression patterns of individual family members (2, 4). When temperatures are optimal, constitutively expressed HSFs are down-regulated by HSPs that bind and retain HSFs in the cytosol. During heat stress, these HSP-HSF complexes dissociate, allowing HSFs to redistribute to the nucleus and regulate gene expression (2, 5–7). The transcriptional activity of HSFs is also dependent on their phosphorylation state (7, 8). The heat stress response is attenuated by HSPs and the heat shock factor binding protein 1 (HSBP1), which binds to and inhibits HSFs (7).

Transcription factors that are induced by heat and are not related to these “classical” HSFs have recently been reported to contribute to heat tolerance in plants. DREB2A, an ethylene-responsive element-binding factor/Apetala 2-type transcription factor that contributes to drought and heat tolerance (9, 10), contributes to heat tolerance by inducing *HsfA3* expression (11, 12). Also, an *Arabidopsis* relative of the human nuclear transcription factor X-box binding 1 that contributes to salt and

defense responses (13, 14) was recently shown to be heat inducible and to contribute to heat tolerance (15). These findings give evidence of cross-talk between heat stress and other stress signaling pathways.

Membrane-tethered transcription factors (MTTFs) are maintained in an inactive state by associating with membranes through one or more transmembrane domains (TMDs). In response to specific signals, an MTTF fragment that contains the transcription factor domain but lacks a TMD, is released from membranes by regulated intramembrane proteolysis (RIP), is redistributed to the nucleus, and regulates the expression of particular nuclear genes. MTTFs regulate diverse processes in prokaryotes and eukaryotes (16–20). Here, we show that *bZIP28*, which encodes a putative basic leucine zipper (bZIP) type MTTF, contributes to the up-regulation of heat-responsive genes and heat tolerance.

Results

The *bZIP28* gene is predicted to encode an MTTF with one (bZIP) domain, one TMD, and a site 1 protease (S1P) cleavage site and site 2 protease (S2P) cleavage site that contribute to RIP (refs. 16, 21, 22; [supporting information \(SI\) Fig. S1](#)). Genome expression data publicly available from AtGenExpress indicated that *bZIP28* is up-regulated in response to heat, a result that we confirmed ([Fig. S2](#)). These data lead us to postulate that *bZIP28* might be an MTTF with a role in the heat shock response.

To test the membrane association and topology of AtbZIP28, we carried out cell fractionation experiments using *Arabidopsis* protoplasts that transiently expressed a fusion protein in which YFP was fused to the amino terminus of AtbZIP28 (YFP-bZIP28). YFP-bZIP28 accumulated in the pellet fraction, which contains cellular membranes as did the endoplasmic reticulum (ER) membrane marker GFP-calnexin (ref. 23; [Fig. 1A](#)). We did not observe an AtbZIP28 signal in either of the supernatant fractions, which contain soluble cytosolic and vacuolar proteins (site 1) and soluble microsomal proteins (site 2). An ER-targeted and -retained form of GFP (GFP-HDEL; ref. 24) was extracted in site 2 and pellet fractions, as expected for a soluble protein that localizes to the lumen of the ER (25). To determine the membrane topology of *bZIP28*, we digested fraction P with proteinase K. We observed that *bZIP28*, but not GFP-HDEL, was susceptible to proteinase K digestion in the absence of detergent ([Fig. 1B](#)). These data suggest that *bZIP28* is oriented


Author contributions: H.G., F.B., C.B., and R.M.L. designed research; H.G. and F.B. performed research; H.G., F.B., C.B., and R.M.L. analyzed data; and H.G. and R.M.L. wrote the paper.

The authors declare no conflict of interest.

[§]To whom correspondence should be addressed at: Michigan State University–Department of Energy Plant Research Laboratory, Room S206 Plant Biology Laboratories, East Lansing, MI 48824. E-mail: larkinr@msu.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0808463105/DCSupplemental.


© 2008 by The National Academy of Sciences of the USA



the mutant and was similar to the *bZIP28* mutant that was stably transformed with a transgene expressing a fusion protein in which YFP was fused to the amino terminus of bZIP28 (YFP-bZIP28) (Fig. 4). A comprehensive RT-PCR analysis of the *bZIP28* T-DNA insertion allele indicates that bZIP28 mRNA does not accumulate in the bZIP28 mutant (Fig. S4C) and that partial RNAs can be transcribed from this allele upstream and downstream of the T-DNA insertion site (Fig. S4D). The observation that the heat-sensitive phenotype can be rescued by a transgene expressing the full-length bZIP28 fused to YFP indicates that the heat-sensitive phenotype results from a loss of bZIP28 function rather than from a gain of function like the transcription of aberrant mRNAs from the T-DNA insertion allele.

From these data, we hypothesized that bZIP28 is likely an MTF that regulates the expression of heat-inducible genes. To identify candidate genes regulated by bZIP28, we searched publicly available data sets for genes that are up-regulated by heat and are coexpressed with *bZIP28*. Two genes identified during this search, At1g52560 and At5g42020, which encode the HSP26.5-P and the ER-localized chaperone BiP2, respectively, were up-regulated by heat less in the *bZIP28* mutant than in WT (Fig. 5A). The heat induction of *HSP17.4-CIII* (At1g54050) was indistinguishable in the *bZIP28* mutant compared with WT, as are 13 other genes that are heat induced and coexpressed with *bZIP28* (Fig. 5A and Table S1).

Our model predicts that a heat-induced signal releases bZIP28 from its membrane tether by triggering RIP and that the free bZIP domain contributes to the heat induction of genes such as *BiP2*. To test whether the bZIP domain of bZIP28 is sufficient to induce the *BiP2* gene, we used an estradiol-inducible promoter (28) to drive expression of a truncated version of bZIP28 that contained the bZIP domain but lacked the predicted TMD and the entire carboxy-terminus (i.e., bZIP28 Δ 301–675). Estradiol treatments specifically induced expression of the bZIP domain and caused *BiP2* mRNA to accumulate above the levels observed in the *bZIP28* mutant; accumulation of *BiP2* mRNA levels correlated with the accumulation of bZIP28 mRNA levels (Fig. 5B). Elevated levels of *bZIP28* Δ 301–675 expression were sufficient to induce nine other heat-inducible genes that are also induced by ER stress or encode ER-localized proteins with diverse functions (ref. 29; Fig. S5 and Tables S2 and S3). *HSP26.5-P* was not similarly induced by estradiol-inducible deletion construct (H.G., unpublished data), which indicates that although bZIP28 is necessary for optimal expression of *HSP26.5-P* (Fig. 5A), bZIP28 Δ 301–675 alone is not sufficient for heat induction of *HSP26.5-P*. Although low levels of *BiP2* expression appeared to be induced by estradiol in the mutant, low levels of *BiP2* expression are induced by a variety of



manipulations similar to those used in this experiment (30). Low levels of induced expression were observed for *BiP2* regardless of whether seedlings were treated with estradiol (Fig. S6). These data indicate that low levels of *BiP2* expression are induced by experimental manipulations, as has been reported previously (30); that bZIP28 $\Delta 301$ –675 is responsible for the remainder of this induction; and that estradiol does not affect *BiP2* expression independent of bZIP28 $\Delta 301$ –675.

Discussion

Here, we show that bZIP28 is an ER-localized MTF that contributes to heat tolerance. Heat appears to induce a signal that triggers the cleavage of bZIP28, followed by the redistribution of bZIP28 from the ER to the nucleoplasm. Once in the nucleoplasm, bZIP28 up-regulates heat-regulated genes. To our knowledge, such an MTF-based signaling system has not been previously shown to contribute to heat tolerance in any system. Such a membrane-tethering mechanism suggests that bZIP28 may be tethered to the ER to respond to heat-induced ER stress. Consistent with this idea, unfolded protein stress within the ER was reported to trigger the redistribution of bZIP28 from the ER to the nucleus at optimal growth temperatures (22).

The redistribution of soluble transcription factors from the cytosol to the nucleus during heat stress is well established. HSFs are maintained in a dormant state at optimal temperatures by

forming a complex with HSPs in the cytosol. During heat stress, HSPs bind and stabilize misfolded proteins, thereby liberating HSFs, which redistribute to the nucleus and regulate gene expression (2, 3). Subsequent to the heat stress response, the activity of HSFs is attenuated and HSFs redistribute to the cytosol. In plants, HSBP1, HSP70, and HSP17.4-CII contribute to this down-regulation of HSF activity (31–34). Membrane tethering of bZIP28 and the formation of HSF-HSP complexes would appear to be distinct mechanisms that allow plants to respond rapidly to heat stress that is localized in the ER and cytosol, respectively. Proper attenuation of HSF activity appears essential in plants. For instance, null alleles of a gene that encodes HSBP1 in maize are embryo lethal, apparently attributable to an unattenuated heat stress response during embryogenesis (32). The tethering of bZIP28 to the ER suggests that bZIP28 is similar to HSFs in that the activity of bZIP28 is detrimental when temperatures are optimal. Indeed, driving bZIP28 expression by a strong and constitutive promoter severely inhibits plant growth at optimal temperatures (22).

In animal cells, MTFs such as ATF6 and SREBP are retained in the ER by protein–protein interactions that can be disrupted by particular signals. Once these interactions are disrupted, the MTF can be transported into the Golgi, where it is cleaved by RIP. This mechanism involves two distinct proteases known as S1P and S2P that cleave the MTF between the transcription factor domain and the TMD (16, 19). In plants, bZIP17 and a plant protease that is related to S1P contribute to salt tolerance (21). Therefore, a similar signaling mechanism appears to be at least partially conserved between plants and animals. Such a signaling mechanism also seems likely for bZIP28, because bZIP28 appears to contain a canonical S1P cleavage site (21).

bZIP28 is related to three other MTFs in *Arabidopsis*: bZIP17, bZIP49, and bZIP60. bZIP28 and bZIP60 regulate genes that are responsive to unfolded protein stress (22, 35), which is consistent with some redundancy among bZIP28 and its relatives. However, our analysis of a bZIP28 mutant during a heat stress treatment not only indicates that bZIP28 is required for heat tolerance but that if other MTFs contribute to this heat stress response, they are at most only partially redundant with bZIP28. Because bZIP17, but not bZIP28, was reported to promote salt tolerance (21, 22), at least this one bZIP28 relative is regulated by specific stresses and might affect distinct regulons, as has been suggested for MTFs in animal cells (36). A system of ER-localized MTFs that is regulated by overlapping and specific ER stresses may be beneficial for optimal regulation of ER-related processes in diverse environmental conditions.

Our findings not only suggest that bZIP28 is activated by a mechanism such as RIP during heat stress but indicate that bZIP28 expression is induced during heat stress. These findings suggest that the bZIP28 activity may be required for an immediate heat stress response and that sustained bZIP28 activity may also contribute to heat tolerance. Our results also show that bZIP28 is required for the optimal expression of two heat-inducible genes that contribute to proper protein folding and thermotolerance in plants (2, 3). Additionally, high levels of the ER-localized BiP2 promote protein secretion in *Arabidopsis* (37). Therefore, a role for bZIP28 in the regulation of these genes is consistent with a role in protecting the ER from heat stress and promoting a proper unfolded protein response. Our initial gene expression analysis indicates that bZIP28 can induce several heat and ER stress-responsive genes that encode proteins of diverse functions when this factor is liberated from its ER tether. These initial results are consistent with a broad influence of the ER and bZIP28 on heat-responsive gene expression. Because the severe chlorotic phenotype of the bZIP28 mutant developed over a period of 5 days after a brief heat stress treatment, the bZIP28-mediated transcriptome responses that

are most important for heat tolerance may occur at any point during the heat stress treatment or during the subsequent recovery period at an optimal temperature.

Materials and Methods

Plant Materials and Growth Conditions. Plants were grown at 22°C under 12-h light and 12-h dark cycles. Light was provided by broad-spectrum fluorescent tube lamps at 120 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$. For heat stress experiments, 2–4 h after dawn, plants were transferred to a growth chamber with these same light conditions but with the temperature set at 42°C. The bZIP28 T-DNA insertion mutant Salk_132285 (27) was obtained from the *Arabidopsis* Biological Resource Center (Ohio State University, Columbus, OH). The T-DNA insertion was verified by amplifying the insertion site using the following oligonucleotides: LbB1, 5'-GCGTGGACCGCTTGCTGCAACT-3'; 132285-LP, 5'-CACCATTAAATTC-TAACCAAGC-3'; and 132285-RP, 5'-GTTGCCTTAAAGCGACATTCTC-3', in addition to Phusion DNA polymerase (New England BioLabs). The PCR product that contained the insertion site could be amplified using primers LbB1 and 132285-RP. Sequence data generated at the Research Technology Support Facility (Michigan State University, East Lansing, MI) with LbB1 confirmed that a T-DNA disrupted the first exon in Salk_132285.

Phenotypic analyses were conducted on 0.5 X Murashige-Skoog (MS) media (Sigma) containing 0.6% phytagar (Caisson Laboratories). The leaf tissue for immunoblotting was obtained from 4-week-old plants grown on soil.

Transient expression in *Arabidopsis* protoplasts was performed as previously described (38) and analyzed 12 h after transformation. Transient expression in tobacco was performed as previously described (39) and analyzed 2 days after infiltration.

Protoplast Fractionation. *Arabidopsis* protoplasts expressing YFP-bZIP28 were fractionated as previously described (40). Protoplasts were pelleted in 250 mM NaCl, gently resuspended in ice-cold extraction buffer (0.1 M Tris-HCl, (pH 7.8), 0.2 M NaCl, 1 mM EDTA), and incubated on ice for 10 min. The resuspended protoplasts were centrifuged for 15 min at 16,000 $\times g$ at 4°C in a microcentrifuge. The first supernatant, which contains cytosolic and vacuolar proteins, was removed and the pellet was resuspended by gentle sonication. The resuspended pellet was centrifuged for 15 min at 16,000 $\times g$ at 4°C in a microcentrifuge. The second supernatant, which contains soluble microsomes, was removed, and the pellet fraction, containing larger cellular membrane fragments and organelles, was resuspended in extraction buffer. Equal volumes of each fraction were analyzed by means of SDS/PAGE and immunoblotting using anti-GFP antibodies, which recognize GFP and YFP (Abcam).

Topology Analysis. Topology analysis was performed with a proteinase K protection assay, as previously described (40). Briefly, microsomes of *Arabidopsis* protoplasts expressing YFP-bZIP28 were prepared by osmotic shock in 100 mM Tris-HCl (pH 7.5), 12% sucrose, and 2 mM CaCl_2 . The extracts were centrifuged at 1000 $\times g$ for 5 min, and the middle phase containing microsomes was removed. Microsomes were treated with 0.2 mg/ml proteinase K with or without 1% (vol/vol) Triton X-100 for 20 min on ice. Inactivation of the proteinase K was achieved by boiling for 10 min. One percent (vol/vol) Triton X-100 was added to samples that had not previously been treated with the detergent to release proteins from the membranes. Equal volumes of microsomes were analyzed by immunoblotting, as described in *Cell Fractionation*.

Bioinformatic Analysis. BLAST searches of the protein database were done at The *Arabidopsis* Information Resource (TAIR; <http://arabidopsis.org/Blast/>) and the National Center for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov/BLAST/>). The prediction of the TMD was obtained from the ARAMEMNON database (41). N-glycosylation sites were predicted using the NetNGlyc server (<http://www.cbs.dtu.dk/services/NetNGlyc/>). To identify genes that are coexpressed with bZIP28, we downloaded microarray data from AtGenExpress through TAIR (www.arabidopsis.org) and analyzed these data using Microsoft Excel. Genes that were expressed at least twofold more in heat stress conditions with a maximum value larger than 30 were chosen for further analysis. The list of coexpressed genes was narrowed further by comparing the response of these genes to bZIP28 during other abiotic stress treatments, namely, cold, salt, osmotic, genotoxic, drought, oxidative, wounding, and UV stress. We chose genes that, like bZIP28, are induced more by heat than these other treatments. ER-localized genes were identified using the SUBA database (42).

Construction of YFP Fusion Genes. For transient expression experiments, a YFP-bZIP28 fusion gene was constructed that was driven by the cauliflower

mosaic virus 35S promoter. For this gene fusion, the coding region of the *bZIP28* gene, including the intron, was amplified using 5'-ATCGACTAGTGAATCAACATCCGTGG-3', 5'-TCAGGTGGCTACGAGATGGAGACC-3', and Phusion DNA polymerase, digested with *SpeI* and inserted between the *SpeI* and *StuI* sites in a modified pCambia vector with an *EYFP* gene and a *BAR* gene (Cambia). The resulting plasmid was sequenced with gene-specific oligonucleotides.

For expression in stable transgenic plants, a *YFP-bZIP28* fusion gene was constructed that was driven by the *bZIP28* promoter. For this gene fusion, all DNA fragments were amplified as described above, except that the 1 kb of the promoter region was amplified using 5'-GGATTCTAGAAGATGCCAGCAAACA-3' and 5'-CATGCCATGGTCATCGTCGGAGGATTTCG-3' and digested with *XbaI* and *NcoI*; the *YFP* gene was amplified using 5'-GATGTCATGATGACAAGGGCGAGGAGC-3' and 5'-CGGTTCTAGACAAGACCGGCAACAGGATTTC-3' and digested with *BspHI* and *MluI*; the coding region and a 454-bp downstream region of *bZIP28* were amplified using 5'-GACGACGCGTGAATCAACATCCGTGGTTGCTCC-3' and 5'-GCTCGGTACACCTGATAGTGGCCTGCTTC-3' and digested with *MluI* and *BstEII*. These three fragments were ligated into a derivative of the transformation vector pCambia3302 (Cambia) cut by *XbaI* and *BstEII*.

To express an estradiol-inducible truncated *bZIP28* protein lacking the TMD and residues carboxy-terminal to the TMD, the *bZIP28* coding sequence was amplified as described above using 5'-ATTCCTCGAGATGGAATCAACATCCGTGGTTGC-3' and 5'-ATCCACTAGTTCAGACAGGCTTAGGATTAACTTAGG-3', digested with *XhoI* and *SpeI*, and inserted into pER8 (28).

Microscopy. Fluorescence imaging was performed using a Zeiss Laser Scanning Confocal Microscope 510 META. A 488-nm line of an argon laser was used to excite GFP, and a 514-nm line was used to excite YFP. Emission filters of 505–550 nm and 530–600 nm were used for GFP and YFP, respectively, with appropriate main dichroics. Samples were viewed with an 63 $\times$ oil-immersion objective. Postacquisition image handling was done with Zeiss AIM software and Paint Shop Pro (Corel, Ottawa, Canada).

RT-PCR Analysis. For the heat induction experiment, plants were grown for 2 weeks on 0.5 $\times$ MS media, as described above. Mutant and WT plants were placed

in a 42°C growth chamber for 0, 30, 70, and 150 min. For estradiol-mediated induction of the *bZIP* domain, plants were grown for 1 week on MS media, as described above, and subsequently placed in a tube containing 0.5 $\times$ MS liquid medium and 2 μ M estradiol (Sigma Chemical Co.) for 1, 2, and 4 h. The plants were then harvested and frozen in liquid nitrogen. Total RNA was isolated by using an RNeasy kit (Qiagen). To compare gene expression levels among the samples, equal amounts of total RNA were used for semiquantitative RT-PCR analysis with a SuperScript One-step RT-PCR kit (Invitrogen). The primers used for each gene were as follows: At3g10800, 5'-CAACATCCGTGGTGTCTCTC-3' and 5'-ACTAGCAACCTTCTTGAGCTTAC-3'; At1g54050, 5'-TCAGATATCCAGGTTACAGTGAGG-3' and 5'-GTCTTAGGTTTCGGCGGCTG-3'; At1g52560, 5'-CTCCACCTAATGAGTCTTCC-3' and 5'-CTCCGCTTAATGTCCTCAACC-3'; At5g42020, 5'-AGGCTACGAAGTTAGGATCAGTTATTG-3' and 5'-CTCTCAGGATTAAACAGCGCC-3'; and At4g05320, 5'-TCAATTCTCTACCGTGATCAAGATGCA-3' and 5'-GGTGTCAAGTCTCCACCTCAAGAGTA-3'. mRNA levels were quantitated using the Quantity One 1-D Analysis Software (Bio-Rad) and normalized to UBQ10, which had been quantitated using the same software.

For the comprehensive RT-PCR expression analysis of the *bZIP28* T-DNA insertion allele (Fig. S4), RT-PCR was as described above, except that ACGAGTACCGATGGAATCAACATCCGTGGTTGCTC and GAAATCTCCGTCTCATCGTCGAG were used to monitor expression of this allele upstream of the insertion site and CAACGAAGGTGATGATGACGACG and AGACAAACCGCAAAGCTTCC were used to monitor the expression of this allele downstream of the T-DNA insertion site.

ACKNOWLEDGMENTS. This work was supported by a Michigan State University-Department of Energy Plant Research Laboratory group project (Grant DE-FG02-91ER20021). We thank Min Zhang (from Capital Normal University, Beijing, China), Hong Cai, and Yaojian Liu for technical assistance. We thank Neil Adhikari from the Michigan State University-Department of Energy Plant Research Laboratory and the Michigan State University Genetics Program for helping prepare figures. We thank the following colleagues for helpful advice: ShengYang He, Jianping Hu, and Beronda Montgomery-Kaguri from the Michigan State University-Department of Energy Plant Research Laboratory and Lee Kroos from the Michigan State University Department of Biochemistry and Molecular Biology.

- Mittler R (2006) Abiotic stress, the field environment and stress combination. *Trends Plants Sci* 11:15–19.
- Baniwal SK, et al. (2004) Heat stress response in plants: A complex game with chaperones and more than twenty heat stress transcription factors. *J Biosci* 29:471–487.
- Kotak S, et al. (2007) Complexity of the heat stress response in plants. *Curr Opin Plant Biol* 10:310–316.
- von Koskull-Döring P, Scharf KD, Nover L (2007) The diversity of plant heat stress transcription factors. *Trends Plants Sci* 12:452–457.
- Duina AA, Kalton HM, Gaber RF (1998) Requirement for Hsp90 and a CyP-40-type cyclophilin in negative regulation of the heat shock response. *J Biol Chem* 273:18974–18978.
- Yamada K, et al. (2007) Cytosolic HSP90 regulates the heat shock response that is responsible for heat acclimation in *Arabidopsis thaliana*. *J Biol Chem* 282:37794–37804.
- Morimoto RI (1998) Regulation of the heat shock transcriptional response: Cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes Dev* 12:3788–3796.
- Reindl A, Schöffl F, Schell J, Koncz C, Bakó L (1997) Phosphorylation by a cyclin-dependent kinase modulates DNA binding of the *Arabidopsis* heat-shock transcription factor HSF1 in vitro. *Plant Physiol* 115:93–100.
- Sakuma Y, et al. (2006) Functional analysis of an *Arabidopsis* transcription factor, DREB2A, involved in drought-responsive gene expression. *Plant Cell* 18:1292–1309.
- Sakuma Y, et al. (2006) Dual function of an *Arabidopsis* transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression. *Proc Natl Acad Sci USA* 103:18822–18827.
- Schramm F, et al. (2008) A cascade of transcription factor DREB2A and heat stress transcription factor HsfA3 regulates the heat stress response of *Arabidopsis*. *Plant J* 53:264–274.
- Yoshida T, et al. (2008) Functional analysis of an *Arabidopsis* heat-shock transcription factor HsfA3 in the transcriptional cascade downstream of the DREB2A stress-regulatory system. *Biochem Biophys Res Commun* 368:515–521.
- Asano T, et al. (2008) AtNF1L1, an *Arabidopsis* homologue of the human transcription factor NF-X1, functions as a negative regulator of the trichothecene phytoxin-induced defense response. *Plant J* 53:450–464.
- Lisso J, Altmann T, Müssig C (2006) The AtNF1L1 gene encodes a NF-X1 type zinc finger protein required for growth under salt stress. *FEBS Lett* 580:4851–4856.
- Larkindale J, Vierling E (2008) Core genome responses involved in acclimation to high temperature. *Plant Physiol* 146:748–761.
- Ron D, Walter P (2007) Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 8:519–529.
- Hoppe T, Rape M, Jentsch S (2001) Membrane-bound transcription factors: Regulated release by RIP or RUP. *Curr Opin Cell Biol* 13:344–348.
- Selkoe D, Kopan R (2003) Notch and presenilin: Regulated intramembrane proteolysis links development and degeneration. *Annu Rev Neurosci* 26:565–597.
- Goldstein JL, DeBose-Boyd RA, Brown MS (2006) Protein sensors for membrane sterols. *Cell* 124:35–46.
- Makinoshima H, Glickman MS (2006) Site-2 proteases in prokaryotes: Regulated intramembrane proteolysis expands to microbial pathogenesis. *Microbes Infect* 8:1882–1888.
- Liu JX, Srivastava R, Che P, Howell SH (2007) Salt stress responses in *Arabidopsis* utilize a signal transduction pathway related to endoplasmic reticulum stress signaling. *Plant J* 51:897–909.
- Liu JX, Srivastava R, Che P, Howell SH (2007) An endoplasmic reticulum stress response in *Arabidopsis* is mediated by proteolytic processing and nuclear relocation of a membrane-associated transcription factor, bZIP28. *Plant Cell* 19:4111–4119.
- Irons SL, Evans DE, Brandizzi F (2003) The first 238 amino acids of the human lamin B receptor are targeted to the nuclear envelope in plants. *J Exp Bot* 54:943–950.
- Batoko H, Zheng HQ, Hawes C, Moore I (2000) A rab1 GTPase is required for transport between the endoplasmic reticulum and Golgi apparatus and for normal Golgi movement in plants. *Plant Cell* 12:2201–2218.
- Brandizzi F, et al. (2003) ER quality control can lead to retrograde transport from the ER lumen to the cytosol and the nucleoplasm in plants. *Plant J* 34:269–281.
- Staehelin LA (1997) The plant ER: A dynamic organelle composed of a large number of discrete functional domains. *Plant J* 11:1151–1165.
- Alonso JM, et al. (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301:653–657.
- Zuo J, Niu QW, Chua N-H (2000) Technical advance: An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. *Plant J* 24:265–273.
- Urade R (2007) Cellular response to unfolded proteins in the endoplasmic reticulum of plants. *FEBS J* 274:1152–1171.
- Martinez IM, Chrispeels MJ (2003) Genomic analysis of the unfolded protein response in *Arabidopsis* shows its connection to important cellular processes. *Plant Cell* 15:561–576.
- Lee J, Schöffl F (1996) An Hsp70 antisense gene affects the expression of HSP70/HSC70, the regulation of HSF, and the acquisition of thermotolerance in transgenic *Arabidopsis thaliana*. *Mol Gen Genet* 252:11–19.
- Fu S, Meeley R, Scanlon MJ (2002) Empty pericarp2 encodes a negative regulator of the heat shock response and is required for maize embryogenesis. *Plant Cell* 14:3119–3132.
- Port M, et al. (2004) Role of Hsp17.4-CII as coregulator and cytoplasmic retention factor of tomato heat stress transcription factor HsfA2. *Plant Physiol* 135:1457–14570.
- Kim B, Schöffl F (2002) Interaction between *Arabidopsis* heat shock transcription factor 1 and 70 kDa heat shock proteins. *J Exp Bot* 53:371–375.

35. Iwata Y, Koizumi N (2005) An *Arabidopsis* transcription factor, AtbZIP60, regulates the endoplasmic reticulum stress response in a manner unique to plants. *Proc Natl Acad Sci USA* 102:5280–5285.
36. Bailey D, O'Hare P (2007) Transmembrane bZIP transcription factors in ER stress signaling and the unfolded protein response. *Antioxid Redox Signal* 9:2305–2321.
37. Wang D, Weaver ND, Kesarwani M, Dong X (2005) Induction of protein secretory pathway is required for systemic acquired resistance. *Science* 308:1036–1040.
38. He P, Shan L, Sheen J (2007) The use of protoplasts to study innate immune responses. *Methods Mol Biol* 354:1–9.
39. Voinnet O, Rivas S, Mestre P, Baulcombe D (2003) An enhanced transient expression system in plants based on suppression of gene silencing by the p19 protein of tomato bushy stunt virus. *Plant J* 33:949–956.
40. Hanton SL, et al. (2005) Diacidic motifs influence the export of transmembrane proteins from the endoplasmic reticulum in plant cells. *Plant Cell* 17:3081–3093.
41. Schwacke R, et al. (2003) ARAMEMNON, a novel database for *Arabidopsis* integral membrane proteins. *Plant Physiol* 131:16–26.
42. Heazlewood JL, Verboom RE, Tonti-Filippini J, Small I, Millar AH (2007) SUBA: The *Arabidopsis* subcellular database. *Nucleic Acids Res* 35:D213–D218.