

## Soil fertility in flooded and non-flooded irrigated rice systems

Kanwar Lal Sahrawat<sup>a\*</sup>

Volume 58, Issue 4, Pages 423-436, 2012

DOI: <http://dx.doi.org/10.1080/03650340.2010.522993>

This is author version post print archived in the official Institutional Repository of  
ICRISAT [www.icrisat.org](http://www.icrisat.org)

---

### Soil fertility in flooded and non-flooded irrigated rice systems

**Kanwar Lal Sahrawat**

**ICRISAT, Patancheru 502 324, India**

#### Abstract

Lowland rice system in Asia makes a major contribution to the global rice supply and is often cited as an example of a sustainable system in which two or three crops of rice are grown in sequence under submerged conditions. However, reports indicate that water shortage may become critical in some regions for lowland rice cultivation; and there is high potential in exploring rice cultivation under moisture regimes that save water and also increase productivity. The objective of this paper is to analyze the consequences of switching the growing of rice from flooded to non-flooded conditions on soil fertility management. Fertility advantages of submerged

27 rice include amelioration of chemical fertility, preferential accumulation of organic  
28 matter and improved availability of major, secondary and selected micronutrients,  
29 which contribute to the long-term maintenance of soil fertility and sustainability of  
30 lowland rice system. On the other hand, the fertility problems in aerobic rice have to  
31 be dealt through diversification of the production systems. Continuous growing of  
32 aerobic rice on a plot might not be sustainable due to chemical and biological  
33 constraints.

---

34

35 **Keywords:** aerobic rice; chemical and biological fertility; nutrient availability; paddy  
36 rice; soil health; sustainability

37

38

### 39 **Introduction**

40 Lowland rice or paddy systems in Asia make a major contribution to the global rice  
41 supply (Cassman and Pingali 1995). The lowland rice system is often cited as an  
42 example of a sustainable system (De Datta 1981; Kyuma 2004). Growing of rice in  
43 submerged soils is an integral component of the traditional, age-old technology in  
44 monsoon Asia (Kyuma 2004). This method of rice cultivation involves land  
45 preparation by cultivating the land in the flooded or saturated state (termed  
46 puddling), followed by transplanting rice seedlings into the puddled paddies, and  
47 growing of rice in submerged soils until two to three weeks prior to the harvest of  
48 the crop.

49 Over 70% of Asia's rice is produced in irrigated lowland fields with high  
50 irrigation requirements to maintain a layer of standing water on the soil surface

51 during most of the growing season (Bouman and Tuong 2001).**The water**  
52 **requirement of flooded rice in the Philippines determined during six seasons**  
53 **including both rainy and post-rainy seasons, varied from 1240 to 1889 mm in the**  
54 **flooded fields and 790-1430 mm in aerobic fields (Bouman et al. 2005). Actually,**  
55 irrigation water shortage being experienced in some regions of Asia, is threatening  
56 the traditional system of lowland rice cultivation (Bouman et al. 2007). **Hence, there**  
57 **is a need for exploring alternate water management practices that save water and**  
58 **at the same time enhance water productivity. Water productivity here is defined as**  
59 **the weight of economic yield or grains produced per unit of water input including**  
60 **rainfall plus irrigation (Bouman et al. 2005).**

61 **For growing rice under non-flooded conditions, there is a need to develop a**  
62 **strategy that includes the development of rice cultivars that are adapted to aerobic**  
63 **conditions. Also, management practices need to be developed in an integrated**  
64 **nutrient and pest management approach so that adapted rice cultivars sustain**  
65 **high yields. It is however, realized that rice yields under aerobic conditions are**  
66 **influenced by complex site-specific and ecological constraints and they need to be**  
67 **considered while developing management practices to go with adopted rice**  
68 **cultivars.**

69 Indeed, aerobic rice is seen as an emerging option to produce rice with less  
70 water than the submerged rice system (Tuong et al. 2005; Xue et al. 2008; Kato et al.  
71 2009). The aerobic rice uses cultivars that maintain high productivity under aerobic  
72 non-submerged or non-saturated soils (Atlin et al. 2006; Peng et al. 2006; Kato et al.  
73 2009). There is an overlap in the definitions of traditional upland rice and aerobic  
74 rice (Bouman et al. 2007). **Generally, rice grown on fertile uplands using high-**  
75 **yielding cultivars with adequate water supply can be regarded as aerobic rice and**

76 **non-irrigated or rain-fed rice with lower productivity expectations is regarded as**  
77 **upland rice (Kato et al. 2009).**

78 The present water shortage situation for growing rice (Bouman et al. 2007)  
79 indeed presents an opportunity to review and critically analyze the effects on soil  
80 fertility of non-flooded rice systems relative to those that accrue in flooded or  
81 submerged rice systems. Therefore, the main objective of this paper is to provide a  
82 setting for nutrient disorder-related problems that constrain aerobic rice as  
83 compared to flooded rice system with emphasis on soil fertility maintenance. Future  
84 research in this important area should focus in diagnosing and managing nutrient  
85 disorders and pest and diseases for the adoption and adaptation of aerobic or non-  
86 flooded rice systems by rice growers especially in the regions facing critical  
87 irrigation water shortages.

88

### 89 **Basic principles of fertility in rice soils**

90 Rice is a sub-aquatic plant, well adapted to flooded soils, and thus is able to derive  
91 the benefits following flooding of the soil. However, upland rice is also grown in  
92 well-drained soils. **Lowland rice is perhaps** the only food crop, which thrives in  
93 submerged soils in monsoon Asia and other regions prone to seasonal or prolonged  
94 flooding (Kamoshita 2007). The adaptation of the lowland rice to flooded conditions  
95 is due to the presence of aerenchyma or pore space in the rice plant that conduit air  
96 from leaves to roots (Reddy and De Laune 2008).

97 **Submerged soils benefit the rice crop by providing a more conducive**  
98 **environment for nutrient availability and uptake as a result of the adjustment of**  
99 **soil pH in the neutral range (Ponnamperuma 1972, 1984). The presence of free**  
100 **water on the soil eliminates water stress and minimizes weed competition to the**

101 **rice crop growth (Ponnamperuma 1975; DeDatta 1981; Rao et al. 2007). Also, the**  
102 **forms and availability of nutrients are related to moisture supply and flooded**  
103 **condition improves both availability (favorable soil pH) and the accessibility**  
104 **(nutrients delivery to rice plant roots improved both by mass flow and diffusion**  
105 **mechanisms) (Ponnamperuma 1975, 1984). Moreover, soil physical properties**  
106 **related to structure, which are important under arable or drained conditions, are**  
107 **not as important, as long as the soil is submerged under water (Ponnaperuma**  
108 **1984). In general, soil chemical properties are improved following submergence of**  
109 **the soil (Ponnamperuma 1984; Narteh and Sahrawat 1999). Paddy soils also provide**  
110 a congenial environment for biological nitrogen fixation by a range of aerobic,  
111 facultative anaerobes and anaerobic bacteria (Ponnamperuma 1972; Sahrawat 1998;  
112 Kyuma 2004; Sahrawat 2004 b).

113         The most important effect of submerging a soil under water is to cut the  
114 supply of oxygen. As a result, the entrapped oxygen is quickly exhausted and the  
115 soil becomes devoid of free oxygen. The lack of free oxygen or anaerobiosis causes  
116 soil reduction and sets in motion a series of physical, chemical, and biological  
117 processes that profoundly influence the quality of a soil as medium for growing rice  
118 or any other wetland crop (e.g., see Ponnamperuma 1972, 1984; Kyuma 2004).  
119 Submerging an aerobic soil in the water decreases its redox potential, which drops  
120 and stabilizes at a fairly stable range of + 200 mV to - 300 mV depending on the soil,  
121 especially on the content of organic matter and reducible nutrient elements – nitrate  
122 N ( $\text{NO}_3^-$ ), manganic manganese ( $\text{Mn}^{4+}$ ), sulfate ( $\text{SO}_4^{2-}$ ) and ferric iron ( $\text{Fe}^{3+}$ ),  
123 especially iron (Table 2). However, the redox potential of the surface water and first  
124 few mm of top soil in contact with the surface water remains relatively oxidized in

125 the redox potential range of + 300 to + 500 mV (Gambrell and Patrick 1978; Patrick  
126 and Reddy 1978; Fiedler et al. 2007).


127 The Eh controls the stability of various oxidized components [oxygen, nitrate,  
128 manganese (Mn IV), ferric (Fe III) iron, sulfate (SO<sub>4</sub><sup>2-</sup>), and carbon dioxide (CO<sub>2</sub>)] in  
129 submerged soils and sediments (Patrick and Reddy 1978; Fiedler et al. 2007, Table 3).  
130 Soil reduction is influenced by the quality of the decomposable organic matter (OM)  
131 and the capacity of reduction is controlled by the quantity of easily reducible iron or  
132 active iron (Sahrawat 2004 a).

133

#### 134 **Fertility advantages of flooded rice system**

135 *Convergence of pH in the neutral range and the implications for nutrient availability*

136 Under submerged condition, the pH of soils is generally stabilized in the neutral  
137 range (6.5-7.5). Following submergence, the pH of alkaline soils decrease because  
138 under prevailing anaerobic condition ferric iron is used as an electron acceptor for  
139 oxidizing organic matter and during this process acidity is consumed  
140 (Ponnamperuma 1972; Narteh and Sahrawat 1999):



142 In these redox reactions, ferric iron (from amorphous ferric hydroxides)  
143 serves as an electron acceptor and OM (CH<sub>2</sub>O) as the electron donor. This reaction  
144 results in the consumption of acidity and raising the soil pH.

145 A decrease in pH of alkali or calcareous soils is the result of accumulation of  
146 carbon dioxide in flooded soil, which neutralizes alkalinity. Moreover, carbon  
147 dioxide produced is retained in the flooded soil due to restricted diffusion through  
148 standing water on the soil surface. This allows large quantities of carbon dioxide to  
149 accumulate and form mild acid, which helps in neutralizing alkalinity in the soil-

150 floodwater system (see equations 2 and 3). Moreover, the submerged soil system  
151 provides an ideal environment for reaction between carbon dioxide (carbonic acid)  
152 and alkalinity.



155 Thus, iron reduction and carbon dioxide concentration in the submerged reduced  
156 soil system, play key role in controlling the pH of submerged soils. The above  
157 reactions require an optimum temperature (between 25 and 35°C) and the  
158 availability of easily decomposable organic matter, reducible iron and other electron  
159 acceptors such as sulfate and carbon dioxide (Ponnamperuma 1972; Sahrawat  
160 2004a).

161 The convergence of soil pH in the neutral range following submerging of soils  
162 benefits the lowland rice crop through better availability of nutrients such as  
163 ammonium, phosphorus (P), potassium (K) and exchangeable cations, which are  
164 mobilized in soil solution. Aluminum (Al) toxicity and other acid soil-related  
165 nutrient problems prevalent in the upland soils (for review see Sahrawat 2009) are  
166 reduced or alleviated by soil submergence of flooding (Ponnamperuma 1972; Narteh  
167 and Sahrawat 1999). A summary of results, gleaned from various sources in the  
168 literature, on the influence of flooding on nutrient availability is provided in Table 1.

169 Equally importantly, the growing of wetland rice in submerged soil is  
170 recognized as a component of technology for the reclamation of salt-affected (saline,  
171 saline-alkali and alkali soils) soils (Gupta and Abrol 1990; van Asten 2003); because  
172 the growing of a lowland rice crop keeps the salt-affected soils productive even  
173 during the reclamation phase. During the reclamation of salt-affected soils, growing

174 of a lowland rice crop allows ponding of water to facilitate the leaching of salts after  
175 the application of amendments such as gypsum and organic matter. The application  
176 of carbonaceous materials (e.g., rice straw from previous harvest and or compost) to  
177 salt-affected soils prior to submergence and growing of the lowland rice, can further  
178 catalyze the amelioration of these soils. Production of extra carbon dioxide helps to  
179 neutralize the alkalinity of alkali soils. In the case of saline soils, ponding of water on  
180 the soil surface facilitates the leaching of salts (Gupta and Abrol 1990; Rao and  
181 Pathak 1996; Qadir et al. 2007).

182

### 183 *Organic matter accumulation*

184 In addition to the favorable effects of soil submergence on fertility in general and N fertility  
185 in particular, lowland rice cultivation maintains or in some cases improves the OM status of  
186 paddy soils. A review of recent global literature showed that organic matter status of soils  
187 under continuous rice (two or three crops per year) is either maintained or even increased  
188 compared with soils under upland rice or in wetland rice-upland crop sequence where a  
189 general decline in soil organic matter has been reported (Witt et al. 2000; Sahrawat 2004 b;  
190 Pampolino et al. 2008; Cheng et al. 2009).

191 Witt et al. (2000) showed that the sequestration of organic C and total N in  
192 wetland soils was significant during two years of cropping under flooded condition.  
193 An experiment was conducted on a clay soil at the International Rice Research  
194 Institute in Los Baños, Laguna, Philippines where five successive croppings (1993-  
195 1995) involving rice-rice or maize-rice were grown. Surface (0-15 cm) soil samples  
196 were taken at the start of the experiment in 1993 (wet season) and again in 1995 after  
197 harvest of the fifth crop in the wet season. There was a net gain in soil organic C and  
198 total N under the rice-rice system and a net decline under the maize-rice system.

199 Replacement of dry season flooded rice crop by maize caused a reduction in C and N  
200 sequestration in the soil. The results demonstrated the capacity of continuous  
201 irrigated lowland rice system to sequester C and N during relatively short time  
202 periods (Table 4) and were in accord with those reported by other researchers on the  
203 long-term benefits of flooding on soil OM accumulation (Zhang and He 2004;  
204 Shrestha et al. 2006; Pampolino et al. 2008; Cheng et al. 2009 ; Nayak et al. 2009).

205 Results reported from long-term experiments suggest that soil organic matter  
206 (SOM) levels under rice-wheat system in the Indo-Gangetic Plains have declined  
207 (Bhandari et al., 2002; Regmi, et al. 2002). On the other hand, prolonged submerged  
208 soil conditions stimulate SOM accumulation and C sequestration in wetland soils  
209 and sediments (Sahrawat 2004 b; Sahrawat et al. 2005; Pampolino et al. 2008; Nayak  
210 et al. 2009). In a long-term (32 years) study of SOM sequestration in the rice-wheat  
211 and maize-wheat systems in Punjab (India) showed that in both rice-wheat and  
212 maize-wheat cropping systems the application of farm yard manure or balanced  
213 fertilization resulted in higher C sequestration. However, the rice-wheat system  
214 (mean value of 260 kg C ha<sup>-1</sup> year<sup>-1</sup> in 0-60 cm depth) had a greater capacity to  
215 sequester C as compared to the maize-wheat (mean value of 70 kg C ha<sup>-1</sup> year<sup>-1</sup> in  
216 0-60 cm depth) system because of greater C input through enhanced productivity  
217 (Kukul et al. 2009).

218 Nishimura et al. (2008) studied the effects of land use change from paddy  
219 cultivation to upland crop cultivation on soil C budget and found that the drainage  
220 of paddy fields for upland cultivation caused significant C loss from crop land soil.  
221 These results are in accord with those of earlier studies which showed that the  
222 drainage of paddy fields for upland crop cultivation causes loss of SOM due to

223 enhanced decomposition of OM under aerobic condition (Mitsuchi 1974; Koizumi et  
224 al. 1993; Hu et al. 2004) The benefits of OM accumulation under long-term paddy  
225 rice cultivation were reversed by bringing the land under upland crop culture  
226 (Sahrawat 2004 b). Prolonged cultivation of lowland rice permits the accumulation of  
227 SOM. For a detailed discussion of OM accumulation in submerged soils and  
228 sediments, readers are referred to a review by Sahrawat (2004 b).

229         The decomposition of OM in aerobic soils is rapid in the presence of oxygen,  
230 which is the most efficient electron acceptor. On the other hand, in the absence of  
231 oxygen in flooded soils and sediments, decomposition of OM depends on the  
232 availability of alternate electron acceptors such as  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$  or  $\text{Fe}^{2+}$ . Since iron is  
233 present in high amounts in rice soils, the ferric-ferrous iron redox reaction plays a  
234 dominant role in the oxidation of OM and its mineralization in submerged soils and  
235 sediments (Sahrawat 2004a, 2010). Compared with arable soils, the decomposition of  
236 organic materials in submerged soils is slower, incomplete and inefficient, leading to  
237 net accumulation of OM (Sahrawat 2004 b; Reddy and De Laune 2008).

238         The deficiencies of nutrients such as N, phosphorus (P), and sulfur (S) affect  
239 the growth of bacteria, which in turn influence C fixation, storage, and release in  
240 wetland ecosystems. The formation of recalcitrant complexes stabilizes OM, making  
241 it less accessible for decomposition by microbial activity and hence its accumulation.  
242 In addition, the productions of compounds in submerged soils and sediments, which  
243 are toxic to microbial population, also retard soil OM decomposition (Sahrawat  
244 2004b).

245         The most important factor responsible for net accumulation of OM in wetland  
246 soils and sediments is the high net primary productivity of these systems (Neue et al.

247 1997; Sahrawat 2004 b). In essence, slow decomposition of OM and higher net  
248 primary productivity of submerged rice soils lead to net accumulation of organic  
249 matter and N in submerged soils and sediments.

250

### 251 *Plant nutrient availability*

252 Pre-flooding of the soil for about four weeks prior to transplanting of the rice  
253 seedlings leads to the release of ammonium, P, K and other exchangeable ions in soil  
254 solution, which is good for growth of the rice plant (Ponnamperuma 1984; Sahrawat  
255 and Narteh 2002). This may allow the farmer to skip the basal application of N and  
256 reduce the application rates of P and K in some cases. The extent and release of  
257 ammonium and other cations and anions depends on soil chemical characteristics  
258 including pH, OM and texture (Ponnamperuma 1972; Narteh and Sahrawat 1999;  
259 Sahrawat 2010).

260 Flooding soil is a great pH neutralizer in problem soils. This is brought about  
261 by the neutralization of acidity in acid soils and alkalinity in alkaline soils following  
262 flooding, thereby generally influencing favorably, the release and the availability of  
263 plant nutrients. Soils with moderate to high content of inherent or added organic  
264 matter can help bring soil pH to a neutral range, favoring nutrient uptake by  
265 wetland rice. Submergence of soil improves the availability of ammonium-N, P, K,  
266 calcium (Ca), magnesium (Mg), iron (Fe), manganese (Mn), and silicon (Si). Toxic  
267 concentrations of Al and Mn in soil solution are minimized with reduced solubility  
268 of these metals as a result of increased pH. On the other hand, the availability of S  
269 may be reduced due to sulfate reduction to sulfide in flooded soils. The supply of  
270 micronutrients such as copper (Cu) and molybdenum (Mo) is generally adequate.  
271 The availability of zinc (Zn) is reduced in submerged soils (Ponnamperuma 1972;

272 Sahrawat 1998; Narteh and Sahrawat, 1999). Also see Table 1 for the summary of  
273 results on nutrient availability under flooded vs. aerobic conditions.

274         There are other disadvantages associated with certain tropical soils in the  
275 humid regions that adversely influence the growth and production of lowland rice  
276 crop. For example, reducing conditions following flooding of iron-rich, acid soils  
277 lead to accumulation of excessive concentrations of ferrous iron in soil solution. The  
278 accumulation of excessive amount of ferrous iron in soil solution could cause iron  
279 toxicity to lowland rice. For detailed discussion of the various aspects of the topic,  
280 the readers are referred to reviews by Becker and Asch (2005) and Sahrawat (2004c).  
281 Also, submerged soils with high amount of OM or with added as fresh crop and  
282 organic residues may lead to the production of organic acids and sulfide, which can  
283 be toxic to the rice plant (Kyuma 2004).

284         A summary of results on the comparative status of soil organic matter,  
285 availability of plant nutrients and other factors that influence rice plant growth and  
286 development under flooded and non-flooded conditions (Table 1) show that the  
287 fertility and other associated advantage under submerged conditions outweigh those  
288 in the aerobic rice conditions. In the longer these fertility advantages or  
289 disadvantages have a cumulative effect on soil quality for growing rice.

290

291

292

### 293 **Aerobic rice and soil fertility**

294 The soil reduction-driven moderation or amelioration in soil pH observed in  
295 submerged rice soils is not observed in aerobic soils. This is the major difference in  
296 the submerged and aerobic soils, which has an overwhelming influence on the

297 general soil health and the availability of major and micro-nutrient elements in rice  
298 soils (Sahrawat 1998). The extreme soil pH acidic or alkaline in aerobic soils indeed  
299 causes complex nutrient imbalances in the root zone of plants because of high  
300 concentrations of nutrients such as Al, sodium (Na) and calcium (Foy 2002; Hiradate  
301 et al. 2007; Fageria and Baligar 2008; Sahrawat 2009). For example, Ponnampereuma  
302 (1975) reported that Fe and P deficiencies were prevalent in mineral soils with pH in  
303 the neutral range, while on strongly acid soils, P deficiency and Mn toxicity were the  
304 likely toxicities to aerobic rice. The deficiencies of Fe and Zn to aerobic rice on the  
305 calcareous soils is well established (Ponnampereuma 1975; Yoshida 1981). The absence  
306 of free water on soil surface under aerobic rice soil conditions greatly influences  
307 weed control, and the availability and accessibility of nutrients by plant roots  
308 (Ponnampereuma 1975; Rao et al. 2007; Mahajan et al. 2009; Sahrawat 2009).

309 Soil water regime especially in high pH calcareous soils, greatly influences the  
310 availability of micronutrients such as Fe and Mn; and the rice crop grown under  
311 upland aerobic conditions frequently suffers from Fe and at times Mn deficiency  
312 (Ponnampereuma 1975; Yoshida 1981; Takkar et al. 1989; Takkar 1996; Fageria et al.  
313 2002; Maruyama et al. 2005; Gao et al. 2006; Tao et al. 2007). On the other hand, in  
314 acid upland soils, the acid-soil-related infertility is a major constraint in the humid  
315 tropical regions. These soil infertility problems result from low pH, Al toxicity, P  
316 deficiency and low base saturation, and the interactions between various deficiencies  
317 and toxicities (Fageria and Baligar 2008; Sahrawat 2009).

318

319 Yield decline resulting from continuous cropping of non-flooded or aerobic  
320 rice is a constraint to the widespread adoption of the aerobic rice technology in Asia.

321 Shift in water regime from flooded to non-flooded conditions influence the  
322 availability N and requirement of the crop. For example, Belder et al. (2005) reported  
323 from a field study that the amount of N accounted for was higher under aerobic  
324 irrigated conditions than under flooded conditions. It was concluded that there is a  
325 need for optimizing the rate and timing of N application efficient N nutrition of the  
326 rice crop. Similar results were reported by Nie et al. (2009) from a pot culture study.

327 It was suggested that the form, rate and time of N application need to be  
328 optimized for satisfactory growth the rice crop. Also, fertilizer N source that  
329 acidified the soil was found to be more efficient in N nutrition of the rice plants (Nie  
330 et al. 2009). Further studies showed that soil acidification (by application of dilute  
331 sulfuric acid) followed by application of N improved both plant growth of rice  
332 plants grown in aerobic conditions in pots (Xiang et al. 2009). Nitrogen application  
333 was more effective than in increasing plant growth and N uptake than soil  
334 acidification. It was indicated that a reduction in soil N availability and plant N  
335 uptake following an increase in the soil pH probably contributed to the declines  
336 observed in the growth and yield of monocropped aerobic rice (Xiang et al. 2009).

337 Thus, the issues relating to the sustainability of the aerobic rice, the crop need  
338 to be a part of the diversified system to maintain soil quality and health and control  
339 pests and diseases, e.g., through the use of legumes in the cropping systems.  
340 Without diversification of the system, there is a potential threat to the sustainability  
341 of the system due to biological and soil chemical fertility constraints. For example,  
342 research showed that although aerobic rice can yield 3-6 t ha<sup>-1</sup> under tropical climatic  
343 conditions, the crop can suffer from immediate yield failure or a drastic yield  
344 decline. Several factors have been suggested as the causal agents, which vary from

345 the involvement of chemicals, nutrient imbalance and biological agents especially  
346 nematodes and root pathogens (Ventura and Watanabe 1978; George et al. 2002;  
347 Fageria and Baligar 2003; Coyne et al. 2004; Atlin et al. 2006; Kreye et al. 2009;  
348 Sahrawat 2009).

349 A critical analysis of the results reported from diverse sites would suggest  
350 that under aerobic rice on soils with pH in the alkaline range, micronutrient  
351 imbalances especially caused by the deficiency of Fe and Mn seem the potential  
352 causal factors and need further research (Fageria et al. 2002). Under acidic conditions  
353 in aerobic soils, low base saturation leads to nutrient imbalance (Fageria and Baligar  
354 2008; Sahrawat 2009). Another major constraint that influences the growth and yield  
355 of aerobic rice is the establishment of root knot nematode that invade the rice plant  
356 roots (Coyne et al. 2004; Kreye et al. 2009). On the other hand, soil submergence  
357 precludes the invasion of new roots of the rice plant by root knot nematode (e.g., see  
358 Coyne et al. 2004; Bridge et al. 2005). Moreover, under extended aerobic condition  
359 during the season, these problems related to micronutrient imbalance and root knot  
360 nematode invasion may occur and need to be addressed to help the rice growers  
361 cope with the problems associated with water-shortage. There is an obvious need to  
362 diagnose the causal factors that lead to yield decline in aerobic rice and to develop  
363 suitable management options to alleviate or avoid such constraints (Fageria and  
364 Baligar 2003; Kreye et al. 2009; Sahrawat 2009).

### 365 **Perspectives**

366 The benefits of growing rice in submerged soils are well documented and it is  
367 known that growing rice in submerged state not only imparts stability to rice  
368 production by alleviating water shortage, effective weed control (Ponnaperuma  
369 1972, 1984; DeDatta 1981), but also form the basis of soil fertility and organic matter

370 conservation and maintenance in the longer-term (Sahrawat 2004b; Pampolino et al.  
371 2008). Soil erosion is not a problem in wetlands and indeed in some cases wetlands  
372 receive sediments from flowing water from the adjoining upland areas, which add to  
373 the organic matter and nutrient pools.

374 Lowland soils conserve soil fertility and organic matter by net gains through  
375 various physical, chemical, and biological (including biological nitrogen fixation)  
376 processes. Wetlands also have relatively large capacity to sequester and store organic  
377 matter. Carbon sequestration under soil submergence is the foundation of  
378 sustainable fertility maintenance in wetland rice soils and is also a strategy to reduce  
379 atmospheric carbon dioxide concentration and mitigate climate change.

380 Growing rice under non-flooded moisture regime likely would influence soil  
381 fertility and nutrient availability and this understandably has implications for the  
382 growth and productivity of aerobic rice. Such soil quality and fertility effects could  
383 range from overall depletion of soil organic matter and nitrogen to the availability of  
384 nutrients such as P on acid soils and micronutrients such as Mn and Fe on calcareous  
385 soils (Ponnamperuma 1975; DeDatta 1981; Sahrawat 1998). With the land use change  
386 from paddy to aerobic rice is likely to have a reverse influence on the fertility  
387 benefits derived from soil submergence. This has implications for the growth and  
388 yield of aerobic rice through the depletion of soil organic matter and nitrogen  
389 (Sahrawat 2004b; Belder et al. 2005; Nie et al. 2009; Xiang et al. 2009); and the  
390 availability of some nutrients especially micronutrients such as Mn and Fe on  
391 calcareous soils (Ponnamperuma 1975; Yoshida 1981; Maruyama et al. 2005). While  
392 the long-term fertility benefits of wetland rice are established, the information on  
393 those under the aerobic rice needs to be generated by long-term studies for a

394 comparative evaluation of the long-term effects of growing aerobic rice on the  
395 dynamics of soil chemical and biological fertility.

396 To enhance rice productivity under non-flooded conditions, the role of  
397 cultivars that are adapted to aerobic soil is of critical importance, but there is genetic  
398 limit in the adaptation of the rice cultivars. The overall role of production (crop  
399 rotations or intercropping) systems and nutrient and water management assume  
400 greater importance in the sustainability of the system as a whole. The critical limits  
401 for deficiency and toxicity of nutrients vary in flooded and non-flooded rice ecology  
402 and there is need for research in this important area for optimizing nutrient  
403 management strategy in aerobic rice (Nie et al. 2009). Equally importantly, there is  
404 need to generate information on the long-term effects of growing aerobic rice on  
405 chemical and biological fertility of soils as such information will help in developing  
406 management strategies for a sustainable aerobic rice production system.

#### 407 **References**

- 408 Atlin GN, Lafitte HR, Tao D, Laza M, Amante M, Courtois B. 2006. Developing rice  
409 cultivars for high-fertility upland systems in the Asian tropics. *Field Crops*  
410 *Res.* 97 : 43-52.
- 411 Becker M, Asch F. 2005. Iron toxicity in rice – conditions and management concepts. *J*  
412 *Plant Nutr Soil Sci* 168 :558-573.
- 413 Belder P, Bouman BAM, Spiertz JHJ, Peng S, Castaneda AR, Visperas RM. 2005.  
414 Crop performance, nitrogen and water use in flooded and aerobic rice. *Plant*  
415 *Soil* 273 :167-182.
- 416 Bhandari AL, Ladha JK, Pathak H, Padre AT, Dawe D, Gupta RK. 2002. Yield and  
417 nutrient changes in a long-term rice-wheat rotation in India. *Soil Sci Soc Am J.*  
418 66: 162-170.

- 419 Bouman BAM, Tuong TP. 2001. Field water management to save water and increase  
420 its productivity in irrigated lowland rice. *Agric Water Manag.* 49: 11-30.
- 421 Bouman BAM, Peng S, Castaneda AR, Visperas RM. 2005. Yield and water use of  
422 irrigated tropical aerobic rice systems. *Agric Water Manag.* 74:87-105.
- 423 Bouman BAM, Lampayan RM, Tuong TP. 2007. *Water Management in Rice: Coping  
424 with Water Scarcity.* Los Baños, Philippines: International Rice Research  
425 Institute. 54 pp.
- 426 Bridge J, Plowright RA, Peng DL. 2005. Nematode parasites of rice. In: Luc M, Sikora  
427 RA, Bridge J, editors. *Plant parasitic nematodes in subtropical and tropical  
428 agriculture, second edition.* Cambridge, UK: CAB International Publishing.  
429 p. 87-130.
- 430 Cassman KG, Pingali PL. 1995. Intensification of irrigated rice systems: Learning  
431 from the past to meet future challenges. *Geojournal* 35: 299-305.
- 432 Cheng YQ, Yang LZ, Cao ZH, Ci E, Yin S. 2009. Chronosequential changes of  
433 selected pedogenic properties in paddy soils as compared with non-paddy  
434 soils. *Geoderma* 151:31-41.
- 435 Coyne DL, Sahrawat KL, Plowright RA. 2004. The influence of mineral fertilizer  
436 application and plant nutrition on plant-parasitic nematodes in upland and  
437 lowland rice in Cote d'Ivoire and its implications in long term agricultural  
438 Research trials. *Exp Agric.* 40:245-256.
- 439 De Datta SK. 1981. *Principles and practices of rice production.* New York, New York,  
440 USA: Wiley.
- 441 Fageria NK, Baligar VC. 2003. Upland rice and allelopathy. *Commun Soil Sci Plant  
442 Anal.* 34:1311-1329.

443 Fageria NK, Baligar VC. 2008. Ameliorating soil acidity of tropical Oxisols by liming  
444 for sustainable crop production. *Adv Agron* 99:345-399.

445 Fageria NK, Baligar VC, Clark RB. 2002. Micronutrients in crop production. *Adv*  
446 *Agron* 77:185-268.

447 Fiedler S, Vepraskas MJ, Richardson JL. 2007. Soil redox potential: importance, field  
448 measurements, and observations. *Adv Agron.* 94:1-54.

449 Foy CD. 2002. Soil chemical factors limiting plant root growth. *Adv Soil Sci.* 19:97-  
450 149.

451

452 Gambrell RP, Patrick WH Jr. 1978. Chemical and microbiological properties of  
453 anaerobic soils and sediments. In: Hook DD, Crawford RRM, editors. *Plant*  
454 *life in anaerobic environments.* Ann Arbor, Michigan, USA: Ann Arbor  
455 Science Publication. p. 375-423.

456 Gao X, Zou C, Fan X, Zhang F, Hoffland E. 2006. From flooded to aerobic conditions  
457 in rice cultivation: Consequences for zinc uptake. *Plant Soil* 280:41-47.

458 George T, Magbanua R, Garrity DP, Tubana BS, Quiton J. 2002. Rapid yield loss of  
459 rice cropped in aerobic soil. *Agron J.* 94:981-989.

460 Gupta RK, Abrol IP. 1990. Salt-affected soils: Their reclamation and management for  
461 crop production. *Adv Soil Sci.* 11:223-288.

462 Hiradate S, Ma JF, Matsumoto F. 2007. Strategies of plants to adapt to mineral  
463 stresses on problem soils. *Adv Agron.* 96:65-132.

464 Hu R, Hatano R, Kusa K, Sawamoto T. 2004. Soil respiration and net ecosystem  
465 production in an onion field in central Hokkaido, Japan. *Soil Sci Plant Nutr.*  
466 50:27-33.

467 Kamoshita A. 2007. Historical changes in urban rice production systems in Tokyo,  
468 Japan. *Plant Prod Sci.* 10:245-269.

469 Kato Y, Okami M, Katsura K. 2009. Yield potential and water use efficiency of  
470 aerobic rice (*Oryza sativa* L.) in Japan. *Field Crops Res.* 113:328-334.

471 Koizumi H, Usami Y, Satoh M. 1993. Carbon dynamics and budgets in three upland  
472 double-cropping agro-ecosystems in Japan. *Agric Ecosyst Environ.* 43:235-244.

473 Kreye C, Bouman BAM, Castaneda AR, Lampayan RM, Faronilo JE, Lactaoen AT,  
474 Fernandez L. 2009. Possible causes of yield decline in tropical aerobic rice.  
475 *Field Crops Res.* 111:197-206.

476 Kukal SS, Rehana-Rasool, Benbi DK. 2009. Soil organic carbon sequestration in  
477 relation to organic and inorganic fertilization in rice-wheat and maize-wheat  
478 systems. *Soil Till Res.* 102:87-92.

479 Kyuma K. 2004. *Paddy soil science.* Kyoto, Japan: Kyoto University Press.

480 Mahajan G, Chauhan BS, Johnson DE. 2009. Weed management in aerobic rice in  
481 northwestern Indo-Gangetic Plains. *J Crop Improv.* 23:366-382.

482 Maruyama T, Higuchi K, Yoshida M, Tadano T. 2005. Comparison of iron  
483 availability in leaves of barley and rice. *Soil Sci Plant Nutr.* 51:1035-1042.

484 Mitsuchi M. 1974. Characters of humus formed under rice cultivation. *Soil Sci Plant*  
485 *Nutr.* 20:249-259.

486 Narteh LT, Sahrawat KL. 1999. Influence of flooding on electrochemical and  
487 chemical properties of West African soils. *Geoderma* 87:179-207.

488 Nayak P, Patel D, Ramakrishnan B, Mishra AK, Samantaray RN. 2009. Long-term  
489 application effects of chemical fertilizer and compost on soil organic carbon  
490 under intensive rice-rice cultivation. *Nutr Cycl Agroecosyst.* 83:259-269.

491 Neue HU, Gaunt JL, Wang ZP, Becker-Heidmann P, Quijano C. 1997. Carbon in tropical  
492 wetlands. *Geoderma* 79:163-185.

493 Nie L, Peng S, Bouman BAM, Huang J, Cui K, Visperas RM, Xiang J. 2009. Alleviating soil  
494 sickness caused by aerobic monocropping: Responses of aerobic rice to various  
495 nitrogen sources. *Soil Sci Plant Nutr* 55:150-159.

496 Nishimura S, Yonemura S, Sawamoto T, Shirato Y, Akiyama H, Sudo S, Yagi K. 2008.  
497 Effect of land use change from paddy rice cultivation to upland crop  
498 cultivation on soil carbon budget of a cropland in Japan. *Agric Ecosyst*  
499 *Environ.* 125 :9-20.

500 Patrick WH Jr, Reddy CN. 1978. Chemical changes in rice soils. In : *Soils and rice.*  
501 *Manila, Philippines : International Rice Research Institute.* pp. 361-379.

502 Peng S, Bouman B, Visperas RM, Castaneda A. Nie L, Park H -K. 2006. Comparison  
503 between aerobic and flooded rice in the tropics : agronomic performance in an  
504 eight-season experiment. *Field Crops Res.* 96 :252-259.

505 Ponnampereuma FN. 1972. The chemistry of submerged soils. *Adv Agron.* 24:29-96.

506 Ponnampereuma FN. 1975. Growth-limiting factors of aerobic rice. In : *Major research*  
507 *in upland Rice. Los Baños, the Philippines : International Rice Research*  
508 *Institute.* p. 40-43.

509 Ponnampereuma FN. 1984. Effects of flooding on soils. In : Kozlowski T, editor.  
510 *Flooding and plant growth. New York : Academic Press.* P. 9-45.

511 Pampolino MF, Laureles EV, Gines HC, Buresh RJ. 2008. Soil carbon and nitrogen  
512 changes in long-term continuous lowland rice cropping. *Soil Sci Soc Am J.*  
513 2:798-807.

514 Qadir M, Oster JD, Schubert S, Noble AD, Sahrawat KL. 2007. Phytoremediation of  
515 sodic and saline-sodic soils. *Adv Agron* 96:97-247.

516 Rao DLN, Pathak H. 1996. Ameliorative influence of organic matter on biological  
517 activity of salt-affected soils. *Arid Soil Res Rehabil.* 10:311-319.

518 Rao AN, Johnson DE, Sivaprasad B, Ladha JK, Mortimer AM. 2007. Weed  
519 management in direct-seeded rice. *Adv Agron.* 93:153-255.

520 Reddy KR, De Laune RD. 2008. Biogeochemistry of wetlands: Science and  
521 applications. Boca Raton, Florida, USA: CRC Press, Taylor and Francis Group,  
522 774 pp.

523 Regmi A P, Ladha JK, Pathak H, Pasuquin E, Bueno C, Dawe D, Hobbs PR, Joshy D,  
524 Maskey SL, Pandey SP. 2002. Yield and soil fertility trends in a 20-year rice-  
525 rice-wheat experiment in Nepal. *Soil Sci Soc of Am J.* 66:857-867.

526 Sahrawat KL. 1983. Nitrogen availability indexes for submerged rice soils. *Adv Agron.* 36:  
527 415-451.

528 Sahrawat KL. 1998. Flooding soil: a great equalizer of diversity in soil chemical  
529 fertility. *Oryza* 35:300-305.

530 Sahrawat KL. 2004 a. Ammonium production in submerged soils and sediments: the  
531 role of reducible iron. *Commun Soil Sci Plant Anal.* 35:399-411.

532 Sahrawat KL. 2004 b. Organic matter accumulation in submerged soils. *Adv Agron.*  
533 81: 169-201.

534 Sahrawat KL. 2004 c. Iron toxicity in wetland rice and the role of other nutrients. *J Plant*  
535 *Nutr.* 27:1471-1504.

536 Sahrawat KL. 2009. The role of tolerant genotypes and plant nutrients in reducing acid-soil  
537 infertility in upland rice ecosystem: an appraisal. *Arch Agron Soil Sci.* 55:597-607.

538 Sahrawat KL. 2010. Nitrogen mineralization in lowland rice soils: The role of organic matter  
539 quantity and quality. *Arch Agron Soil Sci.* 56:337-353.

540 Sahrawat KL, Narteh LT. 2002. A fertility index for submerged rice soils. *Communic Soil*  
541 *Sci Plant Anal* 33:229-236.

542 Sahrawat KL, Bhattacharyya T, Wani SP, Chandran P, Ray SK, Pal DK, Padmaja  
543 KV. 2005. Long-term lowland rice and arable cropping effects on carbon and  
544 nitrogen status of some semi-arid tropical soils. *Current Sci.* 89:2159-2163.

545 Shrestha RK, Ladha JK, Gami SK. 2006. Total and organic soil carbon in cropping  
546 systems of Nepal. *Nutr Cycl Agroecosyst.* 75:257-269.

547 Takkar PN, 1996. Micronutrient research and sustainable agricultural production. *J*  
548 *Indian Soc Soil Sci.* 44:563-581.

549 Takkar PN, Chhibba IM, Mehta SK. 1989. Twenty Years of Coordinated Research on  
550 Micronutrients in Soils and Plants (1967-1987). Bulletin no. 1. Bhopal, India:  
551 Indian Institute of Soil Science.

552

553 Tao HB, Dittert K, Zhang LM, Lin S, Roemheld V, Sattelmacher B. 2007. Effects of  
554 soil water content on growth, tillering, and manganese uptake of lowland rice  
555 grown in the water-saving ground-cover rice-production system (GCRPS). *J*  
556 *Plant Nutr Soil Sci.* 170:7-13.

557 Tuong TP, Bouman BAM, Mortimer M. 2005. More rice, less water—integrated  
558 approaches for increasing water productivity in irrigated rice-based systems  
559 in Asia. *Plant Prod Sci.* 8:231-241.

560 van Asten PJA. 2003. Soil quality and rice productivity problems in Sahelian  
561 irrigation schemes. *Tropical Resource Management Papers* No. 46.  
562 Wageningen, the Netherlands: Department of Environmental Sciences,  
563 Wageningen University and Research Centre.

564 Ventura W, Watanabe I. 1978. Growth inhibition due to continuous cropping of  
565 dryland rice and other crops. *Soil Sci Plant Nutr.* 24:375-389.

566 Witt C, Cassman KG, Olk DC, Biker U, Liboon SP, Samson MI, Ottow JCG. 2000.  
567 Crop rotation and residue management effects on carbon sequestration,  
568 nitrogen cycling and productivity of irrigated rice systems. *Plant Soil.* 225:  
569 263-278.

570 Xiang J, Haden VR, Peng S, Bouman BAM, Visperas RM, Nie L, Huang J, Cui K.  
571 2009. Improvement in nitrogen availability, nitrogen uptake and growth of  
572 aerobic rice following soil acidification. *Soil Sci Plant Nutr* 55:705-714.

573 Xue C, Yang XG, Bouman BAM, Deng W, Zhang QP, Yan WX, Zhang TY, Rouzi A,  
574 Wang HQ. 2008. Optimizing yield, water requirements, and water  
575 productivity of aerobic rice for the North China Plain. *Irrig Sci.* 26:459-474.

576 Yoshida S. 1981. *Fundamentals of rice crop science.* 1981. Manila, the Philippines:  
577 International Rice Research Institute.

578 Zhang M, He Z. 2004. Long-term changes in organic carbon and nutrients of an Ultisol  
579 Under rice cropping in southeast China. *Geoderma.* 118:167-179.

580

581 Table1. Changes in soil organic matter, availability of plant nutrients and other factors affecting plant  
 582 growth under submerged and aerobic rice systems<sup>1</sup>

583

	Submerged rice	Aerobic rice
pH	Converges to neutral range	Ambient pH
Organic matter	Favors accumulation of C and N but reduction products can be toxic	Decomposition is rapid and the accumulation relatively slow
C:N ratio	Wider C:N ratio due to OC accumulation	Varies with soil and organic matter management
NH <sub>4</sub> -N	Production and accumulation favored	Oxidized to nitrate, which is liable to loss by leaching and denitrification
P	Improved P availability	Not applicable
K	Improved K availability	Not applicable
S	Reduced S availability likely due to sulfide formation	Normal availability due to sulfate
Fe	Improved Fe availability in alkali and calcareous soils, but Fe toxicity may occur in acidic soils high in reducible Fe	Iron deficiency is a serious problem in calcareous and high pH soils
Mn	Reduced solubility	Depends on pH, toxicity in acid soils
Cu, Zn and Mo	Improved availability of Cu and Mo but not of Zn	Depends on soil pH
Al	Not a problem except perhaps in acid sulfate soils	Serious problem on acidic soils
Reduction products	Sulfide and organic acids produced can be toxic	Not a problem
Root knot nematodes	Relatively less of a problem	Serious problem
Sustainability	Stability provided	Stability under question

584

585 <sup>1</sup>Gleaned from various sources in literature (for details see Ponnampereuma 1972, 1975,1984; DeDatta 1981;  
 586 Yoshida 1981; Bridge et al. 2005; Becker, Asch 2004; Coyne et al. 2004; Fageria et al. 2003; Kreye et al. 2009;  
 587 Sahrawat 1998, 2004a,b,c,2009,2010; Sahrawat et al. 2005).

588

589 Table2. The range of oxidation-reduction potential found in rice soils ranging from  
590 well drained to waterlogged conditions<sup>1</sup>

591

---

592 Soil water condition	Redox potential (mV)
594 Aerated or well-drained	+ 700 to + 500
595 Moderately reduced	+ 400 to + 200
596 Reduced	+ 100 to – 100
597 Highly reduced	- 100 to – 300

598

---

599 <sup>1</sup>Adapted from Patrick and Reddy (1978).

600

601

602

603 Table3. The range of redox potentials in which the main oxidized components in submerged  
604 soils become unstable<sup>1</sup>

---

606	Reaction	Redox potential (mV)
608	$O_2 - H_2O$	+ 380 to + 320
609	$NO_3^- - N_2, Mn^{4+} - Mn^{2+}$	+ 280 to + 220
610	$Fe^{3+} - Fe^{2+}$	+ 180 to + 150
611	$SO_4^{2-} - S^{2-}$	- 120 to - 180
612	$CO_2 - CH_4$	- 200 to - 280

---

613

614 <sup>1</sup>Adapted from Patrick and Reddy (1978).

615

616 Table 4. Estimated soil organic carbon and total soil nitrogen balance for the rice-upland  
 617 crop rotation experiment after five consecutive crops in 1993-1995. The data presented are  
 618 from treatments without any N fertilizer application<sup>1</sup>.  
 619

620 Cropping system	Rice-Rice	Maize-Rice
622 <b>Soil organic C, kg ha<sup>-1</sup></b>		
623 1993 wet season	19130 (827) <sup>2</sup>	19222 (791)
624 1995 wet season	20973 (494)	19105 (403)
625 Change	+ 1843 (440)	- 216 (502)
626 <b>Total soil N, kg ha<sup>-1</sup></b>		
627 1993 wet season	1811 (47)	1771 (56)
628 1995 wet season	1863 (49)	1720 (29)
629 Change	+ 52 (30)	- 51 (32)

630  
 631  
 632  
 633  
 634  
 635  
 636  
 637  
 638  
 639  
 640 <sup>1</sup>Adapted from Witt et al. (2000). Five consecutive crops under two rotations were grown in  
 641 wet and dry seasons under irrigated conditions. The crops received uniform application of P  
 642 (26 kg ha<sup>-1</sup>) and K (50 kg ha<sup>-1</sup>) each season. Zinc (10 kg Zn ha<sup>-1</sup>) was applied uniformly in  
 643 1993 wet season.

644  
 645 <sup>2</sup>Standard error.

646

647 .

648