

LETTER • OPEN ACCESS

## Impacts of European livestock production: nitrogen, sulphur, phosphorus and greenhouse gas emissions, land-use, water eutrophication and biodiversity

To cite this article: Adrian Leip *et al* 2015 *Environ. Res. Lett.* **10** 115004

View the [article online](#) for updates and enhancements.

You may also like

- [Impacts of supply-side climate change mitigation practices and trade policy regimes under dietary transition: the case of European agriculture](#)  
Francesco Clora, Wusheng Yu, Gino Baudry et al.
- [Global rangeland production systems and livelihoods at threat under climate change and variability](#)  
C M Godde, R B Boone, A J Ash et al.
- [Livestock in a changing climate: production system transitions as an adaptation strategy for agriculture](#)  
Isabelle Weindl, Hermann Lotze-Campen, Alexander Popp et al.



**The Breath Biopsy® Guide**  
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

## Environmental Research Letters



## LETTER

## OPEN ACCESS

RECEIVED  
27 June 2015

REVISED  
1 September 2015

ACCEPTED FOR PUBLICATION  
15 September 2015

PUBLISHED  
4 November 2015

Content from this work  
may be used under the  
terms of the [Creative  
Commons Attribution 3.0  
licence](#).

Any further distribution of  
this work must maintain  
attribution to the  
author(s) and the title of  
the work, journal citation  
and DOI.



# Impacts of European livestock production: nitrogen, sulphur, phosphorus and greenhouse gas emissions, land-use, water eutrophication and biodiversity

Adrian Leip<sup>1</sup>, Gilles Billen<sup>2</sup>, Josette Garnier<sup>2</sup>, Bruna Grizzetti<sup>1</sup>, Luis Lassalle<sup>2,3</sup>, Stefan Reis<sup>4,9</sup>, David Simpson<sup>5,6</sup>, Mark A Sutton<sup>4</sup>, Wim de Vries<sup>7,8</sup>, Franz Weiss<sup>1</sup> and Henk Westhoek<sup>3</sup>

<sup>1</sup> European Commission, Joint Research Centre (JRC), Institute for Environment and Sustainability, Ispra (VA), Italy

<sup>2</sup> CNRS-UPMC, UMR 7619 Metis, BP 105, 4 Place Jussieu, 75005 Paris, France

<sup>3</sup> PBL Netherlands Environmental Assessment Agency, PO Box 303, 3720 AH, Bilthoven, The Netherlands

<sup>4</sup> NERC Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

<sup>5</sup> EMEP MSC-W, Norwegian Meteorological Institute, NO-0313, Oslo, Norway

<sup>6</sup> Department Earth and Space Sciences, Chalmers University of Technology, SE41308 Gothenburg, Sweden

<sup>7</sup> Alterra, Wageningen University and Research Centre (WUR), PO Box 47, 6700 AA Wageningen, The Netherlands

<sup>8</sup> Environmental Systems Analysis Group, Wageningen University, PO Box 47, 6700 AA Wageningen, The Netherlands

<sup>9</sup> University of Exeter Medical School, Knowledge Spa, Truro, TR1 3HD, UK

E-mail: [adrian.leip@jrc.ec.europa.eu](mailto:adrian.leip@jrc.ec.europa.eu)

**Keywords:** livestock production, climate change, air quality, biodiversity loss, coastal eutrophication, soil acidification, European Union

Supplementary material for this article is available [online](#)

## Abstract


Livestock production systems currently occupy around 28% of the land surface of the European Union (equivalent to 65% of the agricultural land). In conjunction with other human activities, livestock production systems affect water, air and soil quality, global climate and biodiversity, altering the biogeochemical cycles of nitrogen, phosphorus and carbon. Here, we quantify the contribution of European livestock production to these major impacts. For each environmental effect, the contribution of livestock is expressed as shares of the emitted compounds and land used, as compared to the whole agricultural sector. The results show that the livestock sector contributes significantly to agricultural environmental impacts. This contribution is 78% for terrestrial biodiversity loss, 80% for soil acidification and air pollution (ammonia and nitrogen oxides emissions), 81% for global warming, and 73% for water pollution (both N and P). The agriculture sector itself is one of the major contributors to these environmental impacts, ranging between 12% for global warming and 59% for N water quality impact. Significant progress in mitigating these environmental impacts in Europe will only be possible through a combination of technological measures reducing livestock emissions, improved food choices and reduced food waste of European citizens.

## Introduction

Nowadays agricultural land occupies about 180 million hectares or 42% of the land area of the European Union<sup>10</sup> from which a great portion is used as grassland and for cultivating feed (FAO 2006). Historically, livestock helped to transform inedible materials (grass and waste) into high quality food. However today livestock production systems affect air quality, global climate, soil quality, biodiversity and

water quality (Sutton *et al* 2011b, 2011c), by altering the biogeochemical cycles of nitrogen, phosphorus and carbon. In particular, reactive nitrogen ( $N_r$ ) plays a key role in several environmental impacts ( $N_r$  represents all forms of nitrogen other than  $N_2$ , including ammonia ( $NH_3$ ), nitrogen oxides ( $NO_x$ ), nitrous oxide ( $N_2O$ ), and N losses to water bodies). Nitrogen cascades or recycles through crop and livestock production systems, in form of feed for livestock and of manure to grow crops, as illustrated in figure 1, leading to several un-intended flows that give rise to environmental concerns (Sutton *et al* 2011a, Leip *et al* 2011a).

<sup>10</sup> Data refer to the situation with 27 Member States, in this article we will use the term 'Europe' also to refer to the EU27.



The emissions from the livestock sector contribute to five major environmental impacts:

- (i) The emissions of ammonia (NH<sub>3</sub>) and nitrogen oxides (NO<sub>x</sub> = NO + NO<sub>2</sub>) contribute to the formation of secondary particulate matter (PM) and tropospheric ozone, both with serious impacts on air quality. Across Europe, ammonium in particles may account for 5–15% of total PM<sub>2.5</sub> (Putaud *et al* 2010). Loss of statistical life expectancy due to exposure to PM<sub>2.5</sub> is estimated at 6–12 months for large parts in Europe (Amann *et al* 2011).
- (ii) Emissions from the livestock sector affect radiative forcing in many ways; long-lived greenhouse gases (GHGs) occur as methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and carbon dioxide (CO<sub>2</sub>) from the use of fossil fuels or through land use and land use change. Emissions of NO<sub>x</sub> contribute to tropospheric ozone formation—an important greenhouse gas in its own right—which reduces carbon sequestration through damage to vegetation (Ainsworth *et al* 2012, Simpson *et al* 2014). Conversely, aerosols produced by NO<sub>x</sub>-driven photochemistry also affect climate, generally having a cooling effect (Simpson *et al* 2014, Shindell *et al* 2009). The interactions between

atmospheric chemistry, vegetation and aerosols are complex and only partially understood (Ainsworth *et al* 2012, Simpson *et al* 2014), but overall, it is estimated that the emissions of N<sub>r</sub> from Europe lead to a small, but uncertain, cooling (Butterbach-Bahl *et al* 2011). Deposition of N<sub>r</sub> contributes also to additional carbon sequestration in forests by stimulating plant growth and altering rates of ecosystem respiration, generally reducing CO<sub>2</sub> concentrations (de Vries *et al* 2011a, Zaehle *et al* 2011). Globally, livestock systems are estimated to contribute about 14.5% to total GHG emissions (Gerber *et al* 2013). Estimates of the share of GHG emissions from land use changes to total livestock emissions range between 9%–35% (FAO 2006, 2010, Lesschen *et al* 2011, Weiss and Leip 2012).

- (iii) Terrestrial biodiversity is affected by livestock production through land use (including historic land use changes), ammonia emissions and consequent deposition and climate change. Intensively managed grassland and arable land used to grow livestock feeds have a low biodiversity, while extensive grazing avoids shrub encroachment or reforestation and helps maintain landscapes of high biodiversity. Habitat changes and land fragmentation can lead to truncation of

**Table 1.** Overview of emissions caused by livestock rearing and feed that were quantified in the study.

	Livestock rearing	Feed	
	Direct and energy <sup>a</sup>	Cultivation excl. energy <sup>b</sup>	Energy incl. feed processing and transport <sup>c</sup>
Air quality	NH <sub>3</sub> , NO <sub>x</sub>	NH <sub>3</sub> , NO <sub>x</sub>	NO <sub>x</sub>
Climate change	CH <sub>4</sub> , N <sub>2</sub> O, CO <sub>2</sub> (NH <sub>3</sub> , NO <sub>x</sub> )	N <sub>2</sub> O, CO <sub>2</sub> (NH <sub>3</sub> , NO <sub>x</sub> )	CO <sub>2</sub>
Soil quality	NH <sub>3</sub> , NO <sub>x</sub> , SO <sub>2</sub>	NH <sub>3</sub> , NO <sub>x</sub>	NO <sub>x</sub> , SO <sub>2</sub>
Terrestrial biodiversity	CH <sub>4</sub> , N <sub>2</sub> O, CO <sub>2</sub> , NH <sub>3</sub> , NO <sub>x</sub>	N <sub>2</sub> O, CO <sub>2</sub> , NH <sub>3</sub> , NO <sub>x</sub> , land use	NO <sub>x</sub> , CO <sub>2</sub>
Quality of inland and coastal water	N and P losses	N and P losses	

Note: Direct livestock rearing includes livestock housing and manure management and storage; energy consumption from housing, milking, buildings etc. Cultivation of forages (grass, fodder maize, fodder beet etc) and other feed includes all direct and indirect emissions not linked with the consumption of energy.

<sup>a</sup> Place of emissions is EU27.

<sup>b</sup> Place of emissions is EU27 for forages and both EU27 and rest of the world for other feed.

<sup>c</sup> Place of emissions is both EU27 and rest of the world.

migratory routes or the replacement of native species with invasive ones (Reid *et al* 2010). N<sub>r</sub> deposition reduces species richness through eutrophication, acidification, direct foliar impacts, and exacerbation of other stresses (Dise *et al* 2011, Bobbink *et al* 2010).

- (iv) Finally, livestock production has a role in the deterioration of the quality of freshwater and coastal water, increasing losses of N and phosphorus (P) to the water system. There is evidence that concentrations of 25 mg L<sup>-1</sup> nitrate (NO<sub>3</sub><sup>-</sup>) in drinking water are related to an increase of incidences of colon cancer by about 3% (van Grinsven *et al* 2010). The stoichiometric excess of nitrogen and phosphorus in coastal water with respect to silica (Si), can enhance water eutrophication (Billen and Garnier 2007, Grizzetti *et al* 2011, Voß *et al* 2011).

In this study we quantified the contribution of livestock production systems to the above mentioned five impacts combining new data based on a cradle-to-gate Life Cycle Assessment (LCA) calculation with results of other studies on emissions at the European scale. The results are discussed with respect to total emissions in the EU27. Although several compounds are included in the analysis, this paper will give particular attention to reactive nitrogen flows in agricultural production systems.

## Methods

### Overall approach

We estimated emissions of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, NH<sub>3</sub>, NO<sub>x</sub>, SO<sub>2</sub>, and N and P losses to the hydrosphere related to livestock production in Europe and interpreted the results in the light of the five major environmental threats. This was done by (i) extending the LCA approach in the agro-economic Common

Agricultural Policy Regionalised Impact (CAPRI) modelling system (Britz and Witzke 2012)<sup>11</sup>, developed for the assessment of GHG emissions and N footprints, to provide cradle-to-farm gate LCA data for reactive nitrogen, in combination with (ii) model results for other compounds (P and SO<sub>2</sub>) to provide a comprehensive picture of the environmental impact of EU livestock production. Table 1 gives an overview of emission sources quantified indicating how they are linked to livestock production systems. As the table indicates, emissions caused by livestock rearing were calculated for EU27 only. Emissions from imported livestock products were not included in the analysis, while further emissions associated with cultivation, processing and transport of (non-forage) feed might occur within or outside the EU.

### The CAPRI N-LCA approach

The emissions of GHGs (N<sub>2</sub>O and CH<sub>4</sub>) and reactive nitrogen (N<sub>r</sub>) to air (NH<sub>3</sub>, NO<sub>x</sub>) and water (N leaching and runoff) and other N flows (such as trade of food and feed) were estimated with CAPRI, using the cradle-to-gate LCA approach implemented for the estimation of GHG emissions (Weiss and Leip 2012) and further extended to include the estimation of N footprints per product group (Leip *et al* 2014b). Here we present new data that have been calculated when further extending the model with flows of all N<sub>r</sub>. All new data presented here include emissions occurring within Europe and outside the EU territory for imported feed and/or land use change (LUC; LUC emissions were calculated using the land use transition probabilities of scenario II in Weiss and Leip (2012), table A1). Also included are credits for carbon sequestration in managed grassland as well as emissions due to foregone carbon calculation in croplands according to Weiss and Leip (2012). To analyze

<sup>11</sup> <http://www.capri-model.org/>

international N flows to and from EU27, the detailed trade data matrix of the trade module in the FAOSTAT database (FAOSTAT 2014) was used taking into account the N content to the 504 commodities involved (vegetal, animal and fiber products) (Lassalletta *et al* 2014). Internal flows among EU 27 countries were calculated and subtracted. The net N import to EU27 for each world country was estimated by the difference between total imports and exports.

A general description of the CAPRI model and its relevant modules can be found in Britz *et al* (2010), Jansson and Heckelei (2011), Leip *et al* (2011b, 2011d), Britz and Witzke (2012) and Perez-Dominguez *et al* (2012).

In the CAPRI LCA module (Weiss and Leip 2012, Leip *et al* 2014b, Westhoek *et al* 2015), total agricultural emissions  $E_{agri}$  were estimated as the sum of flows caused by agricultural production activities, plus emissions caused in earlier phases of the products life cycle, including energy use or land use change. Supplementary Information S1 gives detailed results of the N-LCA for the main six vegetable and six livestock product groups to which the data were aggregated. Differently from Weiss and Leip (2012) and in accordance to Leip *et al* (2014b) the allocation of flows from primary crop products to secondary products (e.g. soya to soybean oil and soybean cakes) is done by mass. The allocation of emissions from feed production to specific livestock products makes use of the animal budget module in CAPRI where energy and protein requirements are matched with domestic and imported feed supply, and data on farm expenditures for feed (Britz and Witzke 2012, Leip *et al* 2011d). In a first step, emissions from crop activities are converted into emission intensities and allocated to animal activities and in a second step to animal products (Weiss and Leip 2012).

For our purpose,  $E_{agri}$  was divided into emissions related to livestock production  $E_{lvst}$  and those related to production of crops for other purposes (food, fuel, fibre)  $E_{crop}$ .  $E_{lvst}$  includes emissions from livestock production systems  $E_{anim}$  (e.g. CH<sub>4</sub> emissions from enteric fermentation, emissions from energy use for milking etc) plus the emissions from feed production, transport and processing  $E_{feed}$ .

For each crop product, allocation to food or feed is done on the basis of the market balance which is available in CAPRI at the national level (see supplementary information S2). The share of total quantities of flows allocated to food crops was calculated on the basis of total domestic production (gross production) minus the quantity used for feed, while the share of the flow used to feed is allocated to livestock products based on feed intake quantities. Emission sources considered are given in table 2.1 by Westhoek *et al* (2015). While this study is restricted to quantifying emissions from EU27 agriculture,  $E_{feed}$  includes both emissions from domestic feed production  $E_{feedeu}$  and emissions from imported feed products  $E_{feedrow}$ :

$$\begin{aligned} E_{agri} &= E_{lvst} + E_{crop} \\ &= E_{feedrow} + E_{feedeu} + E_{anim} + E_{crop} \end{aligned}$$

### Global warming

The contribution to global warming was assessed as the sum of direct GHG emissions (CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub>), indirect N<sub>2</sub>O emissions and C sequestration. Direct N<sub>2</sub>O and CH<sub>4</sub> emissions were from CAPRI-LCA, using a global warming potential of 298 kg CO<sub>2</sub>eq (kg N<sub>2</sub>O)<sup>-1</sup> and 25 kg CO<sub>2</sub>eq (kg CH<sub>4</sub>)<sup>-1</sup> (IPCC 2007). Indirect N<sub>2</sub>O emissions were estimated as 1% of the emitted N (IPCC 2006). Indirect C sequestration in forests was calculated using an average carbon uptake (sequestration) factor of 35 kg C per kg N deposited for European boreal and temperate forests (de Vries *et al* 2014, 2009) and a fraction of 0.25 kg N deposited on forest per kg N emitted, based on EU27 total NH<sub>3</sub>-N emissions from agriculture and NH<sub>3</sub>-N deposition on forests (de Vries *et al* 2011b). Other 'cooling effects' (Butterbach-Bahl *et al* 2011) were not included in the quantification.

### Air quality

The contribution to air quality impacts was assessed on the basis of the sum of NH<sub>3</sub> and NO<sub>x</sub> emissions, as calculated with CAPRI N-LCA.

### Soil acidification


Contribution to soil acidification was assessed on the basis of the sum of SO<sub>2</sub>, NH<sub>3</sub> and NO<sub>x</sub> emissions. NH<sub>3</sub> and NO<sub>x</sub> emissions were calculated with CAPRI N-LCA. Total SO<sub>2</sub> emissions were obtained from the EDGAR data base (European Commission 2011). SO<sub>2</sub> emissions caused by agricultural activities were approximated by the ratio of CO<sub>2</sub> emissions from agriculture related to energy use (Weiss and Leip 2012) and total energy CO<sub>2</sub> emissions in EU27 (EEA 2011). This gives a share of about 6%, which is associated with livestock products (4%) and vegetable products (2%), in accordance with the global estimate of FAO (FAO 2006).

The contribution of agricultural sources to emissions of acidifying substances was estimated on the basis of acidity equivalents (Schöpp and Posch 2003). This method converts emissions of S, NO<sub>x</sub> and NH<sub>3</sub> to acidity equivalents on the basis of the molecular weight  $m$  (64, 46, and 17 for SO<sub>2</sub>, NO<sub>x</sub>, and NH<sub>3</sub>, respectively) and the charge per mole  $z$  (-2, -1, and +1 for SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup>, respectively) to get the conversion factors 0.03125 Geq (Gg SO<sub>2</sub>)<sup>-1</sup>, 0.02174 Geq (Gg NO<sub>2</sub>)<sup>-1</sup>, and 0.05882 Geq (Gg NH<sub>3</sub>)<sup>-1</sup>.

### Terrestrial biodiversity

The contribution of agriculture, livestock and feed to loss of relative mean species abundance (MSA) was estimated using shares of land use and emissions of





$\text{NH}_3$ ,  $\text{NO}_x$ , and net GHG exchange, accounting for C sequestration, as calculated with the CAPRI-LCA. The data are linked to the estimates of the effect of the main drivers for biodiversity loss as calculated with the GLOBIO-model (Alkemade *et al* 2009, Kram and Stehfest 2012, van Vuuren *et al* 2015). This model gives an absolute loss of 65% MSA caused by land conversion into arable, grazing and forestry (35%, 15% and 14%, respectively), and to pressures such as N deposition (2%), climate change (3%) and land fragmentation (30%) (for details see supplementary information S3).

### Water quality

The contribution of agriculture, livestock and feed to  $\text{N}_r$  losses to the hydrosphere was derived from the results of CAPRI N-LCA. Contribution of livestock and feed to dissolved inorganic phosphorus (DIP) losses has been calculated by applying the share of P in fertilizers (mineral fertilizer and manure) per crop from CAPRI LCA to Global NEWS results on total and agricultural flows of DIP (Mayorga *et al* 2010). We were unable to quantify the role of agriculture in the load of particulate phosphorus (PP).

A quantification of the impact of N and P losses was done by combining an analysis of potential risk of eutrophication, based on the Indicator for Coastal

Eutrophication Potential (ICEP, Garnier *et al* 2010, Billen *et al* 2011) with the estimation of livestock contribution to river nutrient loads provided by the model GREEN (Grizzetti *et al* 2012) in the different European coastal areas (see supplementary information S4).

## Results

### The role of trade in N emissions in EU27 and other world regions

N flows from EU27 to other world regions and vice versa for the year 2004 are illustrated in figure 2. Much of the proteins grown in Europe are used to feed livestock. From a total of 16.4 Tg N produced on agricultural land in the year 2004 only 2.4 Tg N  $\text{yr}^{-1}$  (about 15%) were supplied for direct human consumption or further processing. Most of it was used as animal feed (8.8 Tg N  $\text{yr}^{-1}$  or 54%) or returned to the soil as crop residue (5.1 Tg N  $\text{yr}^{-1}$  or 31%). Furthermore, we estimate that livestock received 4.2 Tg N  $\text{yr}^{-1}$  from imports or industry (Leip *et al* 2014b, see also details in supplementary information S2).

According to FAO trade statistics, EU27 was in 2004 a net importer of agricultural products with soy-bean products for animal feed produced in Argentina,

Brazil and USA representing 84% of the total net imports of EU27 (see figure 2; for a comparison of EU27 and global agricultural structure and emissions see supplementary information S5) that entails significant trade of embodied cropland surface (MacDonald *et al* 2015). According to calculations based on FAOSTAT (2014) data, in 2004 about 70% of the European livestock production was used for intra-national consumption and 18%–27% (respectively for chicken and cattle meat, expressed in N) was traded between EU27 countries with significant associated embodied GHG emissions (Caro *et al* 2014). The EU was thus close to self-sufficiency for meat and dairy products, but the share of pig meat production was much higher than in the rest of the world, while the share of ruminant meat was significantly lower (22% versus 29% globally).

### The environmental impact of agriculture and livestock production in EU27

Table 2 shows the results for total agricultural emissions from the EU27 agricultural sector and emissions related to livestock production, feed production and imported feed. Values are provided for  $\text{NH}_3$ ,  $\text{NO}_x$ ,  $\text{SO}_2$ , the combined effect of the three pollutants converted to acidity equivalents, GHG emissions ( $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , and  $\text{CO}_2$ ), C-sequestration, water pollution by emissions of N and P (as dissolved inorganic P, DIP), the land use and the contribution to the loss of the MSA. Values reported in table 2 refer to the year 2004. Detailed results of the LCA calculation ( $\text{N}_r$  and GHG emissions) are given in supplementary information S1.

Total agricultural emissions as compared with the total EU27 emissions from all sources (Leip *et al* 2011a) are given in table 3. Both total and agricultural emissions refer to emissions from EU27 territory. Therefore, agricultural emissions in table 3 do not include emissions associated with imported feed (see table 2).

#### Air quality

Agricultural sources of  $\text{NH}_3$  from manure management, and manure and mineral fertilizers on soils totalled  $2.8 \text{ Tg N yr}^{-1}$  in 2004. The contribution of livestock production to total agricultural emissions was particularly high for  $\text{NH}_3$  (82%) due to the importance of manure management. The share of  $\text{NH}_3$  emissions linked to livestock feed was 41% of agricultural emissions, of which about 8% occurred outside Europe.

Total agricultural  $\text{NO}_x$  emissions at  $0.46 \text{ Tg N yr}^{-1}$  were dominated by emissions from fossil fuel used for farm operations and during processing or transport of animal feed. The share of energy related emissions was higher for crop products (88%) than for livestock products (77%) with an overall contribution to emissions of 85% ( $0.39 \text{ Tg N yr}^{-1}$ ). As there were only small  $\text{NO}_x$

emissions from livestock production systems, most of the emissions were related to feed production, processing and transport ( $0.23 \text{ Tg N yr}^{-1}$ ) and we estimated that about 51% of those occurred outside the EU territory or were linked to feed transport.

For the sum of  $\text{NH}_3$  and  $\text{NO}_x$  emissions, the share of agricultural emissions from livestock was 80% due to the dominance of  $\text{NH}_3$  emissions. 42% of emissions were related to feed production, and 10% were associated with feed imports.

#### Global warming

The direct emissions of GHGs from the agriculture sector itself in 2003–2005 was  $483 \text{ Tg CO}_{2\text{eq}} \text{ yr}^{-1}$ , contributing about 10% of total anthropogenic GHG emissions in the European Union (EEA 2011). However, we estimated emissions of more than twice that amount when including associated emissions that agriculture causes in other sectors, such as energy, industry, or land use and land use change (Weiss and Leip 2012). Overall, 81% of total European agricultural emissions (including associated emissions and emissions from outside of the EU27) were caused by livestock production. As much as 39% of agricultural emissions were estimated to occur outside the EU territory or from associated emissions. This includes especially feed imports, feed transport and emissions from land use change. Carbon sequestration induced by N deposition on forests was found to reduce agricultural emissions by about  $100 \text{ Tg CO}_{2\text{eq}} \text{ yr}^{-1}$  (i.e., 10%). As agricultural N emissions are closely linked to manure management (see supplementary information S1), the N benefit for carbon sequestration was mainly located within the EU.

#### Soil quality

Emissions of acidity equivalents were dominated by  $\text{NH}_3$  which accounted for about 85% of the acidity equivalent emissions for livestock (including associated emissions).

#### Terrestrial biodiversity

Expressed in terms of MSA, we estimated that overall agriculture, through arable and grazing and emissions of N and GHG, caused a loss of 34% MSA, i.e., more than half of the overall loss of biodiversity (Alkemade *et al* 2009). Of this agriculture related loss, 76% was estimated to be caused by livestock, with most of this through feed production.

#### Quality of inland and coastal water

#### Nitrogen

Diffuse N losses from agricultural systems were estimated at  $6.0 \text{ Tg N yr}^{-1}$ . This represents percolation of nitrate and organic nitrogen below the rooting zone in agricultural soils, including both cropland and pasture land, and run-off from soils or barn yards.

**Table 2.** Share of the livestock sector, feed production and feed imports on the emissions of pollutants due to agriculture in EU27 with relevance for air quality, global warming, soil quality, biodiversity and water quality for the year 2004.

	Air and soil quality		Air and soil quality		Air quality		Global warming	
	NH <sub>3</sub>		NO <sub>x</sub>		NH <sub>3</sub> and NO <sub>x</sub>		GHG	
	Emissions [Tg N yr <sup>-1</sup> ]	Share	Emissions [Tg N yr <sup>-1</sup> ]	Share	Emissions [Tg N yr <sup>-1</sup> ]	Share	Emissions [Tg CO <sub>2eq</sub> yr <sup>-1</sup> ]	Share
Agriculture	2.8	100%	0.46	100%	3.2	100%	1062	100%
Livestock	2.3	82%	0.30	66%	2.6	80%	861	81%
Feed	1.1	41%	0.23	49%	1.4	42%	560	53%
Feed imports	0.2	8%	0.12	25%	0.3	10%	411	39%

	Global warming		Global warming		Soil quality		Soil quality	
	C-sequestration		GHG + sequestration		SO <sub>2</sub>		NH <sub>3</sub> and NO <sub>x</sub> and SO <sub>2</sub>	
	Emissions [Tg CO <sub>2eq</sub> yr <sup>-1</sup> ]	Share	Emissions [Tg CO <sub>2eq</sub> yr <sup>-1</sup> ]	Share total	Emissions [Tg yr <sup>-1</sup> ]	Share	Emissions [Tg yr <sup>-1</sup> ]	Share
Agriculture	−104	100%	958	100%	0.021	100% <sup>a,f</sup>	0.19	100%
Livestock	−82	80%	779	81%	0.014	67%	0.15	79%
Feed	−43	42%	516	54%	0.010	50%	0.08	42%
Feed imports	−10	10%	400	42%			0.01	8% <sup>d</sup>

	Biodiversity <sup>b</sup>		Biodiversity <sup>b</sup>		Water quality N		Water quality P <sup>b</sup>	
	Land Use		Loss of biodiversity		N		DIP <sup>c</sup>	
	Area [Mio km <sup>2</sup> ]	Share	Relative MSA [%]	Share	Emissions [Tg N yr <sup>-1</sup> ]	Share	Emissions [Tg P yr <sup>-1</sup> ]	Share
Agriculture	2.0	100%	−34%	100% <sup>a,c</sup>	6.0	100%	0.025	100%
Livestock	1.4	69%	−25%	76%	4.4	73%	0.018	73%
Feed	1.4	69%	−25%	74%	4.2	71%	0.018	73%
Feed imports	0.2	11%			0.6	10%		

## Notes

<sup>a</sup> Own calculation;<sup>b</sup> Emissions occurring outside Europe not included in these estimates;<sup>c</sup> DIP emissions represent about 50% of total P export to the coastal zones;<sup>d</sup> Not considering SO<sub>2</sub>;<sup>e</sup> Alkemade *et al* 2009;<sup>f</sup> EEA 2011.

73% of these emissions were associated with livestock, which was dominated by feed production. The share of leaching and runoff occurring outside of the EU territory was estimated at 10%.

### Phosphorus

Diffuse losses of DIP from agriculture were estimated at 0.025 Tg P yr<sup>-1</sup>, while weathering in agricultural systems contributes an estimated additional 0.003 Tg P yr<sup>-1</sup> (Mayorga *et al* 2010). By far the largest share of net P input (P input minus P crop removal) was retained in the soil, which is considered a benefit as long as this leads to increased soil fertility, i.e., with low erosion. We do not have an estimate of agricultural dissolved organic phosphorus (DOP) or PP, as it is very difficult to distinguish sources for DOP and particularly for PP export. However, most likely the contribution of agriculture is much higher for PP than for DIP while PP dominates P export. The data presented in table 2 relate to DIP only and are

therefore to be regarded as a conservative estimate for the total contribution of agriculture to P flows to coastal areas in Europe. Phosphorus losses from livestock were entirely attributed to feed production, with livestock DIP representing 73% of total agricultural losses, even though some additional losses from animal housing or manure storage systems might occur.

We estimate that in Europe the livestock sector accounts for 23%–47% of the nitrogen river load to coastal waters, and 17%–26% of the phosphorus river loads, where the lower limit is calculated considering the contribution of manure alone and the upper limit taking into account manure applications plus mineral fertilizer (see supplementary information S4).

## Discussion

To our knowledge, this is the first study to estimate the contribution of livestock production systems, feed and



**Table 3.** Comparison of estimated agricultural emissions in this study (from Table 2) and reported total EU27 emissions.

	<b>Total Agricultural LCA impact within EU27 territory</b>	<b>Total EU27 budget impact</b>	<b>Data source for total EU27 impact</b>
NH <sub>3</sub> emissions [Tg N yr <sup>-1</sup> ]	2.6	2.7	EDGAR data base (European Commission 2011) for non-agricultural emissions
NO <sub>x</sub> emissions [Tg N yr <sup>-1</sup> ]	0.3	2.6	EDGAR data base (European Commission 2011)
<b>GHG emissions</b> CH <sub>4</sub> +N <sub>2</sub> O+CO <sub>2</sub> emissions [Tg CO <sub>2eq</sub> yr <sup>-1</sup> ]	651	4889	EU GHG inventory (EEA 2011)
Carbon sequestration [Tg CO <sub>2eq</sub> yr <sup>-1</sup> ]	−93	−171	Same method as for agricultural C sequestration
SO <sub>2</sub> [Tg yr <sup>-1</sup> ]	0.021	0.3	EDGAR data base (European Commission 2011)
<b>Land Use</b> Area [Mio km <sup>2</sup> ]	1.8	4.2	FAOSTAT
<b>Air quality</b> NO <sub>x</sub> + NH <sub>3</sub> emissions [Tg N yr <sup>-1</sup> ]	2.9	5.3	
<b>Global warming</b> GHG + C sequestration [Tg CO <sub>2eq</sub> yr <sup>-1</sup> ]	558	4718	
<b>Soil quality</b> NO <sub>x</sub> +NH <sub>3</sub> +SO <sub>2</sub> [Tg yr <sup>-1</sup> ]	0.18	0.56	
<b>Loss of biodiversity</b> [relative MSA]	−25%	−65%	Alkemade <i>et al</i> (2009)
<b>Water quality N</b> N [Tg N yr <sup>-1</sup> ]	5.4	9.1	European Nitrogen Assessment (Leip <i>et al</i> 2011a) for non-agricultural sources (sewage, forest including background, deposition on water surfaces)
<b>Water quality P</b> DIP [Tg P yr <sup>-1</sup> ]	0.025	0.25	Mayorga <i>et al</i> (2010)

feed imports to total agricultural emissions and their related environmental impact at a comparable level of detail. Plausibility of results have been discussed in depth with regard to GHG emissions (Weiss and Leip 2012) and N-footprints (Leip *et al* 2014b). Estimates of the share of N<sub>r</sub> emissions however are different from those given in Leip *et al* (2014b), as the authors calculated the shares on the basis of domestic consumption (human consumption or processing) while in this study we calculated the share on the basis of total production. Below, we first discuss uncertainty aspects of our emission estimates and estimates of emission shares, followed by options to reduce the environmental impact.

### Uncertainty of emission estimates

For N, combining NH<sub>3</sub> + NO<sub>x</sub> emissions, our estimate of 2.6 Tg N yr<sup>-1</sup> from agricultural sources using the CAPRI model (excluding energy related NO<sub>x</sub>) is 19% and 7% smaller than official estimates of the European Union of 3.2 Tg N yr<sup>-1</sup> (EEA 2014) and the estimate of the MITERRA model of 2.8 Tg N yr<sup>-1</sup> (Westhoek *et al* 2014), respectively. The estimated total N excretion in CAPRI, at 8.9 Tg N yr<sup>-1</sup>, is 88% of the official estimate in EEA (2014, EEA 2014). This difference can be explained by the fact that CAPRI calculates N excretion on the basis of a consistent IPCC Tier 2 approach (animal budget, Leip *et al* 2011b, IPCC 2006) across all

countries, while national inventories in Europe are constructed with a large variety of methods and data quality (Leip 2010). National estimates would be of higher quality than CAPRI estimates from countries with good data (Leip *et al* 2014b), but some countries still need to improve their methodology (EEA 2014). Furthermore, CAPRI uses ammonia abatement measures from the GAINS model (Klimont and Winiwarter 2011) which may not have been considered in national inventories (Leip *et al* 2010).

In comparison with  $4.4 \text{ Tg N yr}^{-1}$  agricultural  $\text{NH}_3$  emissions in the EDGAR data base, our estimate of  $2.6 \text{ Tg N yr}^{-1}$  for EU27, excluding emissions from imported feed, is lower. The reason for this might be the lower excretion estimates, although the  $\text{NH}_3$  emissions are in line with estimates by the MITERRA model.

While 85% of  $\text{NO}_x$  emissions were related to energy use, only  $0.07 \text{ Tg N yr}^{-1}$  were from non-energy sources. A quality check of the total agricultural emission estimate for  $\text{NO}_x$  is difficult as no comparable study exists including both  $\text{NO}_x$  budget flows and  $\text{NO}_x$  emissions related to energy consumption in agricultural systems. Energy consumption in agriculture is calculated in CAPRI with a dedicated energy module which is also used for GHG emission estimates (Kempen and Kraenzlein 2008, Weiss and Leip 2012); this is also the basis of the estimated contribution of  $\text{SO}_2$  emissions. The share of agricultural  $\text{NO}_x$  and  $\text{NH}_3$  to total emissions is within  $-6\%$  to  $+16\%$  of earlier estimates (Leip *et al* 2011a).

Our estimate for the share of agricultural GHG emissions to total GHG emissions based on table 3 is about 13%. It ranges between the value of the official EU GHG inventory (10%, EEA 2014) and other estimates on the shares of agriculture or even livestock production on total GHG emissions (Gerber *et al* 2013, FAO 2006, Weiss and Leip 2012). The official GHG inventory considers only emissions reported in the agriculture sector, whereas LCA studies also include emissions from Land Use Change (LUC) and from imported feeds, which amounted to 39% of total agricultural emissions (see table 2).

Our estimate for  $\text{N}_2\text{O}$  emissions from agricultural soils is considerably lower than official estimates; a comparison of  $\text{N}_2\text{O}$  emissions between various models (de Vries *et al* 2011b) showed overall satisfying agreement. No methodology is able to capture the huge variability of  $\text{N}_2\text{O}$  emissions caused by changing soil and climate conditions. In view of the general lack of experimental observations, even process-based models are not able to achieve a closer match than independent calculations using inverse methods (Leip *et al* 2011c).

LUC is certainly one of the most difficult sources to quantify, as it requires data (or good assumptions) on *how much* LUC is occurring as a consequence of EU agricultural and livestock production, as well as *what*

*kind* of LUC is triggered. Indeed, the debate on the best method to estimate LUC emissions from agricultural products is still ongoing (European Commission 2013). The method developed in the CAPRI model (Leip *et al* 2010, Weiss and Leip 2012) was based on the assumptions that the agricultural market is very fluid and no differentiation between direct and indirect LUC is possible. The approach considers only LUC linked to an expansion of harvested area, very similar to the methods proposed by recent guidelines (Food SCP RT 2013, leap 2014). We used unique LUC factors for imports from a country outside the EU as weighted average for all importing countries accounting for globally connected and substitutable trade flows.

We are aware of the debate on the permanency of carbon sequestration in grassland (Smith 2014), however the approach by Weiss and Leip (2012) is based on the observation that enhanced carbon sequestration rates in grassland are observed also after the 20 years equilibrium time usually used by IPCC methodologies (IPCC 2006), which is also consistent with recent simulations with the CENTURY model (Lugato *et al* 2014).

Carbon sequestration in forests has been estimated earlier at the scale of the EU27 for the year 2000 by multiplying an estimated N deposition caused by agricultural  $\text{NH}_3$  emissions of  $0.61 \text{ Tg N yr}^{-1}$  with a C response of 50 kg C per kg N deposited leading to a C sequestration near  $30 \text{ Tg C yr}^{-1}$  or  $112 \text{ Tg CO}_2 \text{ yr}^{-1}$  (de Vries *et al* 2011a), being very close to our estimate of  $104 \text{ Tg CO}_2 \text{ yr}^{-1}$ . In our study the estimated N deposition was larger ( $0.82 \text{ Tg N yr}^{-1}$ ) while the C:N response was estimated at 35 kg C per kg N deposited.

Estimates of agricultural N-leaching range from 2.0 to  $5.7 \text{ Tg N yr}^{-1}$  (de Vries *et al* 2011b), the higher value being also found in the EU GHG inventory (EEA 2014). Possible reasons for these differences—in addition to those already discussed—are available calibration data for nitrate concentrations which might neglect flows of organic nitrogen to water, and the split of total N between the highly uncertain  $\text{N}_2$  emissions and N leaching/runoff (de Vries *et al* 2011b). Our estimate of nitrate leaching at  $5.4 \text{ Tg N yr}^{-1}$  is consistent with the estimate of the European Nitrogen Budget (Leip *et al* 2011a) which is used in table 3 for total N input to water. Although livestock dominated overall agriculture P flows (73%), agriculture is responsible for only 10% of the total riverine P export (table 3). This is because point sources from human wastewater dominate (accounting for  $0.21 \text{ Tg P yr}^{-1}$  out of a total riverine DIP export of  $0.25 \text{ Tg P yr}^{-1}$ ). The contribution of agriculture to the total P load (including DIP, dissolved organic P, DOP and particulate P, PP) may however be larger, specifically for PP, which is about 40% of the total P export to waters in Europe (50% is DIP and 10% is DOP). The share of agriculture in PP export is determined by (i) surface runoff of P in particles of fertilizer and manure, (ii) agricultural practices (e.g. tillage) that affect the erosion rate and (iii)

elevated P contents of soil material eroded from agricultural fields due to application of P fertilizers and animal manure. The effects of agriculture on PP export are likely to occur much faster than the strongly delayed effect of DIP export through the soil systems but these mixed contributions make it very hard to assess what the agricultural contribution to the PP load is.

Finally, we have used the MSA indicator as a measure for terrestrial biodiversity. MSA represents an index of the naturalness of an ecosystem. Compared with more traditional measures (e.g. monitoring species changes), this measure has two main advantages: it is possible to attribute biodiversity loss to certain sectors (PBL 2014), and the effect of alternative scenarios on biodiversity can be quantified (PBL 2012, 2014). The patterns of change indicated by the MSA are largely similar to those indicated by other measures as the living planet index as developed by WWF or red list indices (SCBD 2014). A limitation of the MSA indicator is that it does not yield comprehensive information on the actual distribution and abundance of species, such as the status of endangered or threatened endemic species.

#### Uncertainty of the estimated shares

While some uncertainty is associated with the individual emission estimates, other parameters might dominate the uncertainty of the estimated shares. For example, while the estimate of N<sub>2</sub>O emission factors might be associated with an uncertainty of up to 50% (at the European scale), the uncertainty around the estimated share of crops that is used as feed determines the uncertainty of livestock's contribution on total agricultural N<sub>2</sub>O emissions. A bias in the total feed translates directly into a bias of the estimated share of emissions from livestock production, as it not only determines the share of crops used as feed, but also the amount of CH<sub>4</sub> emissions from enteric fermentation in ruminants and manure excretion (which is calculated on the basis of animal retention data) and consequent emissions from manure management. This value is obtained from statistical sources (market balances) and is further constrained by energy and nutrient requirement calculations for major livestock types.

No data are available whether farmers prefer domestically produced crops or imported crops for feed; therefore this value is highly uncertain. Because of the lack of information, we considered equal preference. However, this uncertainty concerns only primary crops that are used for both food and feed, which make only 12% of the total feed, while the rest comes from non-marketable crops (82%), such as grass, fodder maize and beet, or secondary crops (6%). Non-marketable crops are all domestically produced; secondary feed stuff is dominated by imported soya bean.

#### Options for reducing the environmental impact of livestock production

There are two main routes to reduce the environmental impacts of livestock production:

- (i) technical measures (reduce emissions intensity/land use intensity and
- (ii) lower livestock production in the EU with demand side measures, i.e., a reduction of food losses and wastes and/or dietary shifts.

Our study presents a 'status quo' analysis (attributional LCA) and does not examine emissions without (or with less) livestock production (case ii). What would happen if livestock production is reduced in Europe has been discussed in depth in Westhoek *et al* (2014, 2015). Based on the observation that the intake of protein as well as saturated fats by European inhabitants is far above the maximum recommended level (WHO 2007, Westhoek *et al* 2011), the authors showed that reducing the consumption of meat, dairy and eggs in the EU27 by 50% would lead to a decrease of N<sub>r</sub> emissions by 40% and a reduction of GHG emissions by 25%–40% with expected substantial health benefits (Westhoek *et al* 2014, 2015). Those results hold for two contrasting scenarios on the use of the 'freed' land that would not anymore be required for feed production, i.e., a 'greening scenario' with enhanced production of bio-energy and a 'high price' scenario with increasing export of cereals. They can be regarded as conservative scenarios, as beneficial environmental effects outside Europe had not been quantified, such as the subsequent prevention of land conversion outside Europe (Stehfest *et al* 2013), or the reduction of GHG emissions from bio-energy production (or other options such as afforestation).

From the production side many technical, structural or policy mitigation options are being discussed, addressing feed production (e.g., precision agriculture and agronomic nitrogen use efficiencies), livestock production (e.g. grazing and feeding management and feed supplements, improved herd structures), or housing and manure management (Thornton and Herrero 2010, Gerber *et al* 2013, Golub *et al* 2013, USDA 2013, Cohn *et al* 2014, Havlik *et al* 2014, Hou *et al* 2014, Van Middelaar *et al* 2014, Winiwarter *et al* 2014, Van Doorslaer *et al* 2015). The benefits of sustainable extensification practices have also recently been explored for Europe (van Grinsven *et al* 2015). Bouraoui *et al* (2014) have shown that high reductions of nitrogen losses to water could be achieved in Europe by an optimized use of organic manure. Additional emission reductions could be achieved by decreasing the wastage of the supplied proteins (Westhoek *et al* 2011, Bellarby *et al* 2013, Grizzetti *et al* 2013).

## Conclusions

This analysis shows that, while agricultural activities are a major source of pollutants and land use change, livestock production systems dominate the environmental consequences. For the five threats considered here, livestock production contributed between 73% (water quality) to about 80% (biodiversity, air quality, soil acidification and global warming) of the overall agricultural impact.

The results point to the fact that in Europe serious efforts in mitigating the major environmental problems for Europe from agriculture need to address the livestock sector. While technical measures can clearly contribute significantly to emission reductions, they cannot alone be sufficient (Bellarby et al 2013, Bajželj et al 2014, Eshel et al 2014, Leip et al 2014a, Witzke et al 2014, Pierrehumbert and Eshel 2015, Vanham et al 2015). The issues of what European citizens eat and their food waste also need to be addressed. For example, recent scenarios showed that all these actions would be necessary to achieve a stabilization in global N<sub>2</sub>O emissions (UNEP 2013).

Moreover, while a shift of production from Europe to other world regions might make Europe 'cleaner', this would possibly come at the cost of higher emission intensities in other regions of the world where production systems might be less optimised (FAO 2010, Cederberg et al 2011, Gerber et al 2013); this could increase the environmental footprint of products consumed in Europe, unless additional actions were taken to address this.

Our study shows that there are intimate links between key environmental threats, emissions of N<sub>r</sub> to the environment, the production of animal products and our diet choices.

## Acknowledgments

The authors would like to thank the European Commission funded research projects NitroEurope-IP (FP6-017841), ANIMALCHANGE (FP7-266018), and ECLAIRE (FP7-282910) for supporting the research and collaboration underpinning the results presented in this paper. We also like to thank Dr Jean Philippe Putaud (EC-JRC) for helpful discussion on the aerosol phenomenology and Dr Faycal Bouraoui (EC-JRC) for discussion on the results of the model GREEN.

## References

- Ainsworth E A, Yendrek C R, Sitch S, Collins W J and Emberson L D 2012 The effects of tropospheric ozone on net primary productivity and implications for climate change *Annu. Rev. Plant Biol.* **63** 637–61
- Alkemade R, Oorschot M, Miles L, Nellemann C, Bakkenes M and ten Brink B 2009 GLOBIO3: a framework to investigate options for reducing global terrestrial biodiversity loss *Ecosystems* **12** 374–90
- Amann M et al 2011 Cost-effective control of air quality and greenhouse gases in Europe: modeling and policy applications *Environ. Modelling Softw.* **26** 1489–501
- Bajželj B, Richards K S, Allwood J M, Smith P, Dennis J S, Curmi E and Gilligan C A 2014 Importance of food-demand management for climate mitigation *Nat. Clim. Change* **4** 924–9
- Bellarby J, Tirado R, Leip A, Weiss F, Lesschen J P and Smith P 2013 Livestock greenhouse gas emissions and mitigation potential in Europe *Glob. Change Biol.* **19** 3–18
- Billen G et al 2011 Nitrogen flows from European regional watersheds to coastal marine waters *European Nitrogen Assessment* ed M Sutton et al (Cambridge: Cambridge University Press) pp 271–97
- Billen G and Garnier J 2007 River basin nutrient delivery to the coastal sea: assessing its potential to sustain new production of non-siliceous algae *Mar. Chem.* **106** 148–60
- Bobbink R et al 2010 Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis *Ecol. Appl.* **20** 30–59
- Bouraoui F, Thieu V, Grizzetti B, Britz W and Bidoglio G 2014 Scenario analysis for nutrient emission reduction in the European inland waters *Environ. Res. Lett.* **9** 125007
- Britz W, Domínguez I P and Heckelet T 2010 A comparison of CAPRI and SEAMLESS-IF as integrated modelling systems *Environmental and Agricultural Modelling: Integrated Approaches for Policy Impact Assessment* ed F M Brower and M K van Ittersum (Netherlands: Springer) pp 257–74
- Britz W and Witzke H-P 2012 CAPRI Model Documentation 2012 ed W Britz and H-P Witzke (Bonn: University Bonn) (<http://capri-model.org/dokuwiki/doku.php?id=start>)
- Butterbach-Bahl K et al 2011 Nitrogen as a threat to the European greenhouse balance *European Nitrogen Assessment* ed M Sutton et al (Cambridge: Cambridge University Press) pp 434–62
- Caro D, LoPresti A, Davis S J, Bastianoni S and Caldeira K 2014 CH<sub>4</sub> and N<sub>2</sub>O emissions embodied in international trade of meat *Environ. Res. Lett.* **9** 114005
- Cederberg C, Persson U M, Neovius K, Molander S and Clift R 2011 Including carbon emissions from deforestation in the carbon footprint of Brazilian beef *Environ. Sci. Technol.* **45** 1773–9
- Cohn A S, Mosnier A, Havlik P, Valin H, Herrero M, Schmid E, O'Hare M and Obersteiner M 2014 Cattle ranching intensification in Brazil can reduce global greenhouse gas emissions by sparing land from deforestation *Proc. Natl Acad. Sci. USA* **111** 7236–41
- De Vries W et al 2009 The impact of nitrogen deposition on carbon sequestration by European forests and heathlands *For. Ecol. Manage.* **258** 1814–23
- De Vries W, Du E and Butterbach-Bahl K 2014 Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems *Curr. Opin. Environ. Sustain.* **9–10** 90–104
- De Vries W, Kros J, Reinds G J and Butterbach-Bahl K 2011a Quantifying impacts of nitrogen use in European agriculture on global warming potential *Curr. Opin. Environ. Sustain.* **3** 291–302
- De Vries W, Leip A, Reinds G J, Kros J, Lesschen J P and Bouwman A F 2011b Comparison of land nitrogen budgets for European agriculture by various modeling approaches *Environ. Pollut.* **159** 3254–68
- Dise N B, Ashmore M, Belyazid S, Bleeker A, Bobbink R, de Vries W, Erismann J W, Spranger T, Stevensand C J and van den Berg L 2011 Nitrogen as a threat to European terrestrial biodiversity *European Nitrogen Assessment* vol 33 ed M Sutton et al (Cambridge: Cambridge University Press) pp 463–94
- EEA 2011 Annual European Union greenhouse gas inventory 1990–2009 and inventory report 2011. Submission to the UNFCCC Secretariat *Technical Report* No. 2/2011 (Copenhagen, Denmark: European Environment Agency)
- EEA 2014 Annual European Union greenhouse gas inventory 1990–2012 and inventory report 2014 Submission to the UNFCCC Secretariat *Technical report* No. 09/2014 (Copenhagen: European Environment Agency) ([www.eea.europa.eu](http://www.eea.europa.eu))



- [europa.eu/publications/european-union-greenhouse-gas-inventory-2014](http://europa.eu/publications/european-union-greenhouse-gas-inventory-2014))
- Eshel G, Shepon A, Makov T and Milo R 2014 Land, irrigation water, greenhouse gas, and reactive nitrogen burdens of meat, eggs, and dairy production in the United States *Proc. Natl Acad. Sci. USA* **111** 11996–2001
- European Commission 2011 *Emission Database for Global Atmospheric Research (EDGAR), release version 4.2* (European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL)) (<http://edgar.jrc.ec.europa.eu>)
- European Commission 2013 Commission Recommendations of 9 April 2013 on the use of common methods to measure and communicate the life cycle environmental performance of products and organisations (2013/179/EU) *Off. J. Eur. Union* **L124** 1–210 (<http://eurlex.europa.eu/legal-content/EN/TXT/?uri=CELEX:52013DC0196>)
- FAO 2010 Greenhouse gas emissions from the dairy sector A *Lifecycle Assessment* (Rome, Italy: FAO) (<http://fao.org/agriculture/lead/themes0/climate/emissions/en/>)
- FAO 2006 *Livestock's Long Shadow* ed H Steinfeld et al (Rome, Italy: Food and Agriculture Organization) (<http://fao.org/docrep/010/a0701e/a0701e00.HTM>)
- FAOSTAT 2014 FAOSTAT (<http://faostat.fao.org/>)
- Food SCP RT 2013 *ENVIFOOD Protocol, Environmental Assessment of Food and Drink Protocol, European Food Sustainable Consumption and Production Round table (SCP RT), Working Group 1* (Brussels, Belgium: Food SCP RT) (<http://food-scp.eu/node/25>)
- Garnier J, Beusen A, Thieu V, Billen G and Bouwman L 2010 N:P:Si nutrient export ratios and ecological consequences in coastal seas evaluated by the ICEP approach *Glob. Biogeochem. Cycles* **24** GB0A05
- Gerber P J, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, Falcucci A and Tempio G 2013 *Tackling Climate Change through Livestock. A Global Assessment of Emissions and Mitigation Opportunities* (Rome: Food and Agriculture Organization of the United Nations (FAO)) (<http://fao.org/docrep/018/i3437e/i3437e.pdf>)
- Golub A A, Henderson B B, Hertel T W, Gerber P J, Rose S K and Sohngen B 2013 Global climate policy impacts on livestock, land use, livelihoods, and food security *Proc. Natl Acad. Sci. USA* **110** 20894–9
- Grizzetti B, Bouraoui F and Aloe A 2012 Changes of nitrogen and phosphorus loads to European seas *Glob. Change Biol.* **18** 769–82
- Grizzetti B, Bouraoui F, Billen G, van Grinsven H, Cardoso A C, Thieu V, Garnier J, Curtis C, Howarth R W and Johnes P J 2011 Nitrogen as a threat to European water quality *European Nitrogen Assessment* ed M Sutton et al (Cambridge: Cambridge University Press) pp 379–404
- Grizzetti B, Pretato U, Lassaletta L, Billen G and Garnier J 2013 The contribution of food waste to global and European nitrogen pollution *Environ. Sci. Policy* **33** 186–95
- Havlik P et al 2014 Climate change mitigation through livestock system transitions *Proc. Natl Acad. Sci. USA* **111** 3709–14
- Hou Y, Velthof G L and Oenema O 2014 Mitigation of ammonia, nitrous oxide and methane emissions from manure management chains: a meta-analysis and integrated assessment *Glob. Change Biol.* **21** 1293–312
- IPCC 2007 *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* ed R Pachauri and R Reisinger (Geneva: IPCC)
- IPCC 2006 *Agriculture, forestry and other land use IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme* vol 4 ed H Eggleston et al (Japan: IGES)
- Jansson T and Heckelei T 2011 Estimating a primal model of regional crop supply in the European Union *J. Agric. Econ.* **62** 137–52
- Kempen M and Kraenzlein T 2008 Energy use in agriculture: a modeling approach to evaluate energy reduction policies key words: energy use in agriculture, energy cost, agricultural sector model
- Klimont Z and Winiwarter W 2011 *Integrated Ammonia Abatement—Modelling of Emission Control Potentials and Costs in GAINS vol IIASA Inte* (Laxenburg: International Institute for Applied System Analysis) ([www.iiasa.ac.at/publication/more\\_IR-11-027.php](http://www.iiasa.ac.at/publication/more_IR-11-027.php))
- Kram T and Stehfest E 2012 *The IMAGE Model Suite Used for the OECD Environmental Outlook to 2050* (Bilthoven, Netherlands: PBL Netherlands Environmental Assessment Agency)
- Lassaletta L, Billen G, Grizzetti B, Garnier J, Leach A M and Galloway J N 2014 Food and feed trade as a driver in the global nitrogen cycle: 50-year trends *Biogeochemistry* **118** 225–41
- Leap 2014 *Environmental Performance of Animal Feeds Supply Chains. Guidelines for Quantification—Draft for Public Review* Livestock environmental assessment and performance partnership (Rome, Italy: Food and Agriculture Organization of the United Nations (FAO)) (<http://fao.org/partnerships/leap/en/>)
- Leip A 2010 Quantitative quality assessment of the greenhouse gas inventory for agriculture in Europe *Clim. Change* **103** 245–61
- Leip A et al 2011a Integrating nitrogen fluxes at the European scale *European Nitrogen Assessment* ed M Sutton et al (Cambridge: Cambridge University Press) pp 345–76
- Leip A, Britz W, Weiss F and de Vries W 2011b Farm, land, and soil nitrogen budgets for agriculture in Europe calculated with CAPRI *Environ. Pollut.* **159** 3243–53
- Leip A, Busto M, Corazza M, Bergamaschi P, Koeble R, Dechow R, Monni S and de Vries W 2011c Estimation of N<sub>2</sub>O fluxes at the regional scale: data, models, challenges *Curr. Opin. Environ. Sustain.* **3** 328–38
- Leip A, Weiss F and Britz W 2011d Agri-environmental nitrogen indicators for EU27 *Bio-Economic Models Applied to Agricultural Systems* ed G Flichman (Dordrecht: Springer) pp 109–23
- Leip A et al 2014a Nitrogen-neutrality: a step towards sustainability *Environ. Res. Lett.* **9** 115001
- Leip A, Weiss F, Lesschen J P and Westhoek H 2014b The nitrogen footprint of food products in the European Union *J. Agric. Sci.* **152** 20–33
- Leip A, Weiss F, Wassenaar T, Perez-Dominguez I, Fellmann T, Loudjani P, Tubiello F, Grandgirard D, Monni S and Biala K 2010 Evaluation of the livestock sector's contribution to the EU greenhouse gas emissions (GGELS) *Final Report* European Commission, Joint Research Centre ([http://ec.europa.eu/agriculture/analysis/external/livestock-gas/full\\_text\\_en.pdf](http://ec.europa.eu/agriculture/analysis/external/livestock-gas/full_text_en.pdf))
- Lesschen J P, van den Berg M, Westhoek H J, Witzke H P and Oenema O 2011 Greenhouse gas emission profiles of European livestock sectors *Anim. Feed Sci. Technol.* **166–167** 16–28
- Lugato E, Bampa F, Panagos P, Montanarella L and Jones A 2014 Potential carbon sequestration of European arable soils estimated by modelling a comprehensive set of management practices *Glob. Change Biol.* **20** 3557–67
- MacDonald G K, Brauman K A, Sun S, Carlson K M, Cassidy E S, Gerber J S and West P C 2015 Rethinking agricultural trade relationships in an era of globalization *Bioscience* **65** 275–89
- Mayorga E, Seitzinger S P, Harrison J a., Dumont E, Beusen A H W, Bouwman A F, Fekete B M, Kroeze C and Van Drecht G 2010 Global nutrient export from watersheds 2 (news 2): model development and implementation *Environ. Modelling Softw.* **25** 837–53
- PBL 2012 *Roads from Rio + 20. Pathways to Achieve Sustainability Goals by 2050* (Hague: PBL, Netherlands Environmental Assessment Agency) (<http://roadsfromrio.pbl.nl/>)
- PBL 2014 How sectors can contribute to sustainable use and conservation of biodiversity *CBD Tech. Ser.* **79**
- Perez-Dominguez I, Fellmann T, Witzke H-P, Jansson T and Oudendag D 2012 *Agricultural GHG Emissions in the EU: An*



- Exploratory Economic Assessment of Mitigation Policy Options* (Luxembourg: Publication Office of the European Union) (doi:[10.2791/8124](https://doi.org/10.2791/8124))
- Pierrehumbert R T and Eshel G 2015 Climate impact of beef: an analysis considering multiple time scales and production methods without use of global warming potentials *Environ. Res. Lett.* **10** 085002
- Putaud J-P et al 2010 A European aerosol phenomenology: III. Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe *Atmos. Environ.* **44** 1308–20
- Reid R S, Bedelian C, Said M Y, Kruska R L, Mauricio R M, Castel V, Olson J and Thornton P K 2010 Global livestock impacts on biodiversity *Livestock in a Changing Landscape* vol 1 ed H Steinfeld et al (Washington, DC: Island Press) pp 111–37
- SCBD 2014 *Global Biodiversity Outlook 4* (Montreal: Secretariat of the Convention on Biological Diversity)
- Schöpp W and Posch M 2003 Long-term development of acid deposition (1880–2030) in sensitive freshwater regions in Europe *Hydrol. Earth Syst. Sci.* **7** 436–46
- Shindell D T, Faluvegi G, Koch D M, Schmidt G A, Unger N and Bauer S E 2009 Improved attribution of climate forcing to emissions *Science* **326** 716–8
- Simpson D, Arneth A, Mills G, Solberg S and Uddling J 2014 Ozone—the persistent menace: interactions with the N cycle and climate change *Curr. Opin. Environ. Sustain.* **9–10** 9–19
- Smith P 2014 Do grasslands act as a perpetual sink for carbon? *Glob. Change Biol.* **20** 2708–11
- Stehfest E, Van Den B M, Woltjer G, Msangi S and Westhoek H 2013 Options to reduce the environmental effects of livestock production—comparison of two economic models *Agric. Syst.* **114** 38–53
- Sutton M A, Billen G, Bleeker A, Erisman J W, Grennfelt P, Van Grinsven H, Grizzetti B, Howard C M and Leip A 2011a European nitrogen assessment—technical summary *European Nitrogen Assessment* ed M Sutton et al (Cambridge: Cambridge University Press) pp xxxv–lii
- Sutton M A, Howard C, Erisman J W, Billen G, Bleeker A, van Grinsven H, Grennfelt P and Grizzetti B 2011b *The European Nitrogen Assessment. Sources, Effects and Policy Perspectives* ed M Sutton et al (Cambridge: Cambridge University Press)
- Sutton M A, Oenema O, Erisman J W, Leip A, van Grinsven H and Winiwarter W 2011c Too much of a good thing *Nature* **472** 159–61
- Thornton P K and Herrero M 2010 Potential for reduced methane and carbon dioxide emissions from livestock and pasture management in the tropics *Proc. Natl Acad. Sci. USA* **107** 19667–72
- UNEP 2013 *Drawing Down N<sub>2</sub>O to Protect Climate and the Ozone Layer. A UNEP Synthesis Report* (Nairobi, Kenya: United Nations Environment Programme (UNEP)) (<http://unep.org/publications/ebooks/UNEPN2Oreport/>)
- USDA 2013 *Greenhouse Gas Mitigation Options and Costs for Agricultural Land and Animal Production within the United States* (Washington, DC: US Department of Agriculture Climate Change Program Office) ([http://usda.gov/oce/climate\\_change/mitigation\\_technologies/GHG\\_Mitigation\\_Options.pdf](http://usda.gov/oce/climate_change/mitigation_technologies/GHG_Mitigation_Options.pdf))
- Van Doorslaer B, Witzke P, Huck I, Weiss F, Fellmann T, Salputra G, Jansson T and Leip A 2015 An economic assessment of GHG mitigation policy options for EU agriculture *EcAMPA vol Report EUR* (Luxembourg: Publications Office of the European Union) (doi:[10.2791/180800](https://doi.org/10.2791/180800))
- Van Grinsven H J M, Erisman J W, de Vries W and Westhoek H 2015 Potential of extensification of European agriculture for a more sustainable food system, focusing on nitrogen *Environ. Res. Lett.* **10** 25002
- Van Grinsven H J M, Rabl A and de Kok T M 2010 Estimation of incidence and social cost of colon cancer due to nitrate in drinking water in the EU: a tentative cost-benefit assessment *Environ. Health* **9** 58
- Vanham D, Bouraoui F, Leip A, Grizzetti B and Bidoglio G 2015 Lost water and nitrogen resources due to EU consumer food waste *Environ. Res. Lett.* **10** 084008
- Van Middelaar C E, Dijkstra J, Berentsen P B M and De Boer I J M 2014 Cost-effectiveness of feeding strategies to reduce greenhouse gas emissions from dairy farming *J. Dairy Sci.* **97** 2427–39
- Van Vuuren D P et al 2015 Pathways to achieve a set of ambitious global sustainability objectives by 2050: explorations using the IMAGE integrated assessment model *Technol. Forecast. Soc. Change* **98** 303–23
- Voß M et al 2011 Nitrogen processes in coastal and marine ecosystems *European Nitrogen Assessment* ed M Sutton et al (Cambridge: Cambridge University Press) pp 147–76
- Weiss F and Leip A 2012 Greenhouse gas emissions from the EU livestock sector: a life cycle assessment carried out with the CAPRI model *Agric. Ecosyst. Environ.* **149** 124–34
- Westhoek H et al 2015 Nitrogen on the table: the influence of food choices on nitrogen emissions and the European environment *European Nitrogen Assessment Special Report on Nitrogen and Food* (Edinburgh: Centre for Ecology and Hydrology)
- Westhoek H, Lesschen J P, Rood T, Wagner S, De Marco A, Murphy-Bokern D, Leip A, van Grinsven H, Sutton M A and Oenema O 2014 Food choices, health and environment: effects of cutting Europe's meat and dairy intake *Glob. Environ. Change* **26** 196–205
- Westhoek H, Rood T, van den Berg M, Janse J, Nijdam D, Reudink M and Stehfest E 2011 The protein puzzle *The Consumption and Production of Meat, Dairy and Fish in the European Union* (The Hague: PBL Netherlands Environmental Assessment Agency)
- WHO 2007 Protein and amino acid requirements in human nutrition report of a joint FAO/WHO/UNU expert consultation *WHO Technical Report Series* 935 (Geneva: World Health Organization)
- Winiwarter W, Leip A, Tuomisto H L and Hastrup P 2014 A European perspective of innovations towards mitigation of nitrogen-related greenhouse gases *Curr. Opin. Environ. Sustain.* **9–10** 37–45
- Witzke P, van Doorslaer B, Huck I, Salputra G, Fellmann T, Drabik D, Weiss F and Leip A 2014 Assessing the importance of technological non-CO<sub>2</sub> GHG emission mitigation options in EU agriculture with the CAPRI model *EAAE Int. Congress (Ljubljana, Slovenia, 26–29 August)* pp 1–15
- Zaehle S, Ciais P, Friend A D and Prieur V 2011 Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions *Nat. Geosci.* **4** 601–5