

RESEARCH ARTICLE

Does a history of sexual and physical childhood abuse contribute to HIV infection risk in adulthood? A study among post-natal women in Harare, Zimbabwe

Simukai Shamu^{1,2,3*}, Patience Shamu⁴, Christina Zarowsky^{2,5}, Marleen Temmerman^{6,7}, Tamara Shefer⁸, Naeemah Abrahams^{2,9}

1 Health Systems Strengthening Division, Foundation for Professional Development, Pretoria, South Africa, **2** School of Public Health, University of the Western Cape, Belville, South Africa, **3** School of Public Health, University of the Witwatersrand, Johannesburg, South Africa, **4** Wits Reproductive Health Research Institute, University of the Witwatersrand, Johannesburg, South Africa, **5** University of Montreal Hospital Research Centre, Montreal, Canada, **6** International Centre for Reproductive Health, Ghent University, Ghent, Belgium, **7** Obstetrics and Gynaecology, Agha Khan University, Nairobi, Kenya, **8** Department of Women and Gender Studies, University of the Western Cape, Belville, South Africa, **9** Gender and Health Research Unit, South African Medical Research Council, Cape Town, South Africa

* shamuts@yahoo.com



OPEN ACCESS

Citation: Shamu S, Shamu P, Zarowsky C, Temmerman M, Shefer T, Abrahams N (2019) Does a history of sexual and physical childhood abuse contribute to HIV infection risk in adulthood? A study among post-natal women in Harare, Zimbabwe. PLoS ONE 14(1): e0198866. <https://doi.org/10.1371/journal.pone.0198866>

Editor: Soraya Seedat, Stellenbosch University, SOUTH AFRICA

Received: May 24, 2018

Accepted: November 14, 2018

Published: January 4, 2019

Copyright: © 2019 Shamu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The authors acknowledge funding and assistance from the following organisations: Flemish interuniversity cooperation (VLIR-UOS) for supporting SS in completing PhD studies on this topic in collaboration with Ghent University and the University of the Western Cape; African Population and Health Research Centre in partnership with the

Abstract

Background

Sexual and physical abuse in childhood creates a great health burden including on mental and reproductive health. A possible link between child abuse and HIV infection has increasingly attracted attention. This paper investigated whether a history of child physical and sexual abuse is associated with HIV infection among adult women.

Methods

A cross sectional survey was conducted among 2042 postnatal women (mean age = 26y) attending six public primary health care clinics in Harare, Zimbabwe within 6 weeks post-delivery. Clinic records were reviewed for mother's antenatal HIV status. Participants were interviewed about childhood abuse including physical or sexual abuse before 15 years of age, forced first sex before 16, HIV risk factors such as age difference at first sex before age 16. Multivariate analyses assessed the associations between mother's HIV status and child physical and sexual abuse while controlling for confounding variables.

Results

More than one in four (26.6%) reported abuse before the age of 15: 14.6% physical abuse and 9.1% sexual abuse, 14.3% reported forced first sex and 9.0% first sex before 16 with someone 5+ years older. Fifteen percent of women tested HIV positive during the recent antenatal care visit. In multivariate analysis, childhood physical abuse (aOR 3.30 95%CI 1.58–6.90), sexual abuse (3.18 95%CI: 1.64–6.19), forced first sex (aOR 1.42, 95%CI:

International Development Research Centre for funding data collection and analysis for the study; South African Medical Research Council for supporting data collection; and the University of Zimbabwe for providing office space during data collection.

Competing interests: The authors have declared that no competing interests exist.

1.00–2.02), and 5+ years age difference with first sex partner (aOR 1.66 95%CI 1.09–2.53) were independently associated with HIV infection.

Conclusion

This study highlights that child physical and/or sexual abuse may increase risk for HIV acquisition. Further research is needed to assess the pathways to HIV acquisition from childhood to adulthood. Prevention of child abuse must form part of the HIV prevention agenda in Sub-Saharan Africa.

Introduction

Child abuse is a violation of the fundamental rights of children. Although research on child abuse is growing globally, research on childhood exposure to violence is a neglected area in many parts of Africa. Globally, 180/1000 girls are abused sexually and 163/1000 are abused physically [1]. An extensive review of studies in Africa reported among the highest prevalence of child abuse globally [2]. The review revealed that both boys and girls are affected by abuse. The prevalence of physical or sexual violence against girls in Africa ranges between 2% and 64% [2] and the wide range is a result of the various definitions of violence adopted and settings where the study was conducted (see also <http://www.knowviolenceinchildhood.org/>). Most of the abuse of children happens at home or in school [3]. Drivers of child abuse include family disorganisation and dysfunction [2,4] often related to contexts of poverty and structural violence, cultural factors and community norms accepting violence such as bullying [5], legal frameworks that allow the use of violence to discipline children [6] and gender inequity and inequality [7,8]. For example, in Zimbabwe the use of corporal punishment in schools is permitted to discipline children under special circumstances—heads of schools exercise corporal punishment while teachers can obtain permission to use corporal punishment from the school head. In each instance where corporal punishment is used a register is kept (Secretary of Education Circular P 35, 1993; Statutory Instrument 65 of the Constitution of Zimbabwe, 1992). Nevertheless, corporal punishment is used outside of these conditions as it goes undocumented, goes beyond disciplining to set limits and having detrimental effects to children [6]. Research has also highlighted the absence of alternative ways of disciplining children leading to the use of violence both in schools and at home [8,9]. Shumba argues that the use of violence against children in Zimbabwean schools is an extension of violent ways of disciplining a child that teachers transfer from their roles as parents and adults in the community to the school [6].

The few studies on sexual violence in Zimbabwe show that sexual violence against girls is high [10] while more than one in five (21.6%) report forced first sexual intercourse. The malfunctioning of the family institution to protect or look after children, orphanhood, maternal challenges and non-attendance of school were found to be associated with sexual abuse in Zimbabwe [10]. Studies in high income countries have shown that abuse in childhood is likely to result in later life challenges including mental health problems (substance use, suicide behaviour, depression, stress) and sexually transmitted infections [11–14]. More literature is gathering evidence on the relationship with HIV [15] and these include a few studies being conducted in Sub-Saharan Africa [8,10,16]. Despite this, the evidence base is still thin and no studies have assessed multiple childhood adversities and their association with HIV in pregnancy in one study. Dedicated research on sexual violence is seldom a focus in Zimbabwe as

child sexual abuse is researched as part of other studies [17]. The intersection of age with gender and other inequalities facilitate HIV vulnerability. Women's sexual relationships with older men in Zimbabwe have been reported to increase HIV incidence [18]. Having sexual intercourse with a minor who is five years younger than the person having sex with them is also considered a legally abusive act as it indicates possible coercive or abusive experiences. Research shows an association between age and unwanted sexual practices [18] which increase the risk of HIV infection.

This study provides an opportunity to assess if child physical/sexual abuse is associated with HIV diagnosed in pregnancy. We used data in a women's health research project examining intimate partner violence and HIV diagnosed during pregnancy to estimate here the prevalence of childhood adversities (child sexual abuse, child physical abuse, forced first sex and first sex with older men) and to assess the relationship between these childhood adversities with HIV infection. We tested the hypothesis that a woman's history of childhood adversities contributes to her HIV positive status in antenatal care.

Methods

Design and setting

We conducted a cross sectional study in Harare, Zimbabwe in 2011 among 2042 women aged 18–49 years attending postnatal care at six public primary health care facilities in resource limited formal urban locations. We reported detailed methodology for the study elsewhere [19]. Face to face interviewer administered structured interviews were conducted by trained female fieldworkers (university entry level and graduates) to collect demographic and childhood abuse experiences data for this study. HIV status information were abstracted from clinic records. Women were recruited while queuing for postnatal services and baby immunisation sessions at the six clinics. Interviews were conducted in privacy. The study clinics had both antenatal and postnatal services and were owned and managed by the Harare City Council.

Measures

We measured child physical abuse by asking each participant whether anyone ever “excessively” beat or physically mistreated the woman in any way before age 15. We prefixed the word “excessive” to categorise abuse differently from culturally acceptable ways of enforcing discipline in Zimbabwe which were common in the study setting [6]. Any response in the affirmative was considered as childhood physical abuse. For child sexual abuse we followed the WHO Multi-country study on women's health and domestic violence [20] by asking whether anyone ever forced a woman to have sex or to perform a sexual act or ever touched her sexually when she did not want to before she turned age 15. Any response in the affirmative was recorded as childhood sexual abuse. To assess forced first sexual intercourse respondents were asked whether their first sexual intercourse occurred when they were willing, tricked, persuaded, forced or raped [20,21]. If a participant responded that they were forced or raped we defined it as forced first sexual intercourse. Participants who responded affirmatively and also reported first sex before age 16 were considered as having had forced first sex before age 16. We used this conceptualisation of forced first sex before age 16 in this analysis.

To assess first sex with someone 5+ years older while under 16 years we first asked women about their age and the age of their partners. The two ages were then compared to identify the age difference. Women were also asked the year in which they first had sexual intercourse. We constructed a variable that describes those who had sexual intercourse before age 16 and had first sex with someone five years or older than them. Although many studies have reported that age 18 was the median age of sexual debut of many women in Zimbabwe [22] we used age

16 as the cut off to define childhood sexual abuse following the Sexual Offences Act of Zimbabwe [23]. Intimate partner physical/sexual violence was measured using the WHO questionnaire on measuring domestic violence against women and girls [24]. The interviews also collected demographic information of the participants including education, employment status, marital status, and current or most recent partner's demographic characteristics. HIV was tested through rapid test kits [19]. Health facilities in the study used the Determine rapid test and positive results confirmed using Capillus, and the Western blot was used to resolve any conflicts. HIV results were abstracted from the patient records with permission from the City Health Department and the participant. This paper focuses on participants with available HIV test results.

Data analysis

We analysed data in Stata version 13 [25]. Variable frequencies were calculated and presented in tables. Sociodemographic and childhood abuse experiences were described by HIV status using chi-square tests to illustrate statistical differences between groups. We reported statistical differences at 95% confidence levels (CI). Univariate analyses were performed before building a multiple regression model of factors associated with testing HIV positive. We performed a multivariate logistic regression analysis with HIV as our outcome variable using a backward stepwise elimination process. We reported regression model output as adjusted Odds Ratios (aOR) with 95% CI and p values. The model controlled for known HIV covariates such as demographic variables (age, marital status and education), and history of intimate partner violence based on our knowledge of determinants of HIV in generalised epidemics such as in Zimbabwe.

Ethics

The study received ethics approval from the Medical Research Council of Zimbabwe and the University of the Western Cape Research Ethics Committees. The Harare City Health Department gave permission for the study to be conducted at the six clinics. All women were given contact details of organisations that provide counselling services for abused women for assistance as per the WHO guidelines for researching violence against women and girls [26]. We sought and received informed written consent for all participants while those mothers who were under 18 years of age provided informed written assent after parental/guardian consent was provided.

Findings

We approached 2101 women to participate in the study and 2042 (97%) agreed and completed the interviews. A total of 1951 (95.5%) of those interviewed had available HIV test results in the clinic records. Demographic characteristics of participants (age, education, partners' age and education, marital status, employment status, gravida and age differences with partners) with and without HIV test results did not differ ($p > 0.05$) (data not shown).

[Table 1](#) shows the participants' demographic characteristics. Women were on average 26.4 years old [95% Confidence interval (CI) 26.17–26.68], over 9 in 10 (92.7%) attended at least secondary education. Close to a third (30%) were employed, 88.2% were married. Mean gravidity was 2.2. Participant's partners were on average 31.3 years of age, that is, five years older than them. They were also more educated than them with 97% having completed at least secondary education (11 years of formal education).

[Table 2](#) shows the prevalence of participants' childhood sexual or physical abuse by HIV status. Fourteen percent (14.3%) of the women reported forced first sexual intercourse.

Table 1. Socio-demographic characteristics of the sample (N = 2042).

Variable	n	%	LCI	UCI
Age (mean)		26.4	26.17	26.68
Education: at least secondary	1889/2037	92.7		
Woman employed	606/2027	29.9		
Not married	240/2041	11.8		
Gravida (mean)		2.2	2.16	2.27
Partner has at least secondary education	1973/2030	97.2		
Partner's age (mean)		31.3	31.04	31.61

<https://doi.org/10.1371/journal.pone.0198866.t001>

Women in the sample generally partnered with men older than them. The mean age difference between women and their partners was 5.17 years and almost a tenth of the women (9.0%) reported having sex for the first time with someone five or more years older than them while they were below 16 years of age. Slightly more than a quarter (26.1%) of the women reported any form of childhood abuse before age 15. Almost one in ten (9.1%) women reported physical abuse before their 15th birthday while 14.6% reported sexual abuse before age 15. Significant differences were found among all the four experiences (forced first sex, child physical abuse, child sexual abuse and age difference between the woman and her partner at first sex) and HIV status. More women who reported these experiences tested HIV positive than women who did not report these childhood abusive experiences and HIV risk factors.

Fifteen percent (15.3%) of the women tested HIV positive. About 47% of the women reported experiencing physical or sexual intimate partner violence (IPV) in the 12 months preceding the most recent pregnancy.

Table 3 shows the results of multiple regression analysis that assessed factors associated with testing HIV positive in antenatal care. Childhood physical abuse (aOR 3.30 95%CI 1.58–6.90), sexual abuse (3.18 95%CI: 1.64–6.19), forced first sex (aOR 1.42, 95%CI: 1.00–2.02), and 5+ years age difference with first sex partner (aOR 1.66 95%CI 1.09–2.53) were associated with HIV infection. The strongest associations (at least aOR 3.18) were between HIV and child sexual abuse and physical abuse. Women who were older than 24, not married, being less educated and having a partner at least 30 years old had higher odds of being HIV infected.

Discussion

Our study highlights the prevalence of and association between childhood abuse experiences and subsequent HIV infection detected in antenatal care. With the exception of a few small scale studies profiling corporal punishment in schools [6,27] we are not aware of any study in Zimbabwe on child sexual/physical abuse since 2000 that presents the prevalence of child

Table 2. Prevalence of childhood abuse by HIV status (N = 1945).

Variables	Sample		HIV+ (n = 299)		HIV-(n = 1646)		p-value
	%	CI	n	%	n	%	
Forced first sexual intercourse	14.3	12.81–15.85	58/299	19.4	218/1647	13.2	0.005
5+ year age difference at 1 st intercourse before age 16	9.0	7.75–10.24	43/295	14.6	130/1649	7.8	<0.0001
Child sexual abuse	14.6	13.05–16.19	46/299	15.4	131/1643	8.0	<0.0001
Child physical abuse	9.1	7.83–10.39	57/299	19.3	227/1646	13.8	0.014
Adult experiences							
HIV positivity	15.3	13.72–16.92					
Last 12mo physical/sexual IPV	47.0	43.76–48.18					

<https://doi.org/10.1371/journal.pone.0198866.t002>

Table 3. Multivariate logistic regression showing factors associated with HIV after adjusting for demographic factors and partner violence.

Variable	aOR	95% CI	p-value
Child physical abuse	3.30	1.58–6.90	0.002
Child sexual abuse	3.18	1.64–6.19	0.001
5+ year age difference at 1 st intercourse before age 16	1.66	1.09–2.53	0.017
Forced first sexual intercourse	1.42	1.00–2.02	0.047
Woman's age: 25+ (ref:>25)	1.54	1.09–2.17	0.015
Not currently married (ref: married)	1.96	1.35–2.83	<0.001
Woman has at least 8 years of education	0.63	0.40–0.97	0.037
Current partner is at least 30 years old	1.43	1.02–2.01	0.037
Past year intimate partner violence	1.00	0.76–1.32	0.988

<https://doi.org/10.1371/journal.pone.0198866.t003>

sexual and physical abuse and association between childhood adversities and HIV positivity in antenatal care. Previous studies have found associations between more recent violent experiences or lifetime violent experiences against adult women [28] without specifically studying the abuse that faced the women in childhood. Our study investigated a wide range of childhood abuse and/or conditions that have been associated with physical and sexual abuse such as a significantly older male partner experiences including sexual abuse, physical abuse, high age differences with a partner at first sex, forced first sexual intercourse and all were associated with HIV positivity. We failed to reject the hypothesis that childhood experiences of abuse are associated with HIV diagnosed in adulthood.

The prevalence of sexual abuse in children reported in this study falls within the prevalence range found in a systematic review of literature [12,29] but is lower than found in African reviews [2] where in some countries such as Egypt in which corporal punishment is legal as many as 50–80% children reported physical punishment. The prevalence of physical abuse and sexual abuse is much lower than found in South Africa [9,30]. The reported prevalence may have been a methodological underestimation of abuse experiences experienced by participants in a clinic population especially as we defined sexual abuse as any sexual intercourse occurring before the 15th birthday when other studies define it until 16th birthday. Research has shown that experiences of childhood violence among women is seldom exaggerated but can be under reported and underestimated. For example, some previous psychological experiments have shown that abusive experiences are forgotten if a person recently experienced similar and less abusive experiences in their lives than before [31]. Studies that interview children may give accurate prevalence figures although interviewing young people before they reach 18 would miss an opportunity to capture future violence before age 18.

We found a rate of forced first sex (FFS) consistent with rates found in similar clinic based samples in Zimbabwe (11.8%) [22], South Africa (12.4%) [32] and Uganda (14%) [33] but lower than reported in the Zimbabwe demographic and health survey (ZDHS) (21.6%) [34] which collected data in the same year as in our study. While our study may have captured all childhood's experiences, it may have suffered from recall bias. This might be because of the differences in study designs since the ZDHS collected population level data while our study was a clinic based study. Forced first sexual intercourse is a risk factor for a number of HIV risk factors such as IPV during pregnancy or after disclosing HIV status and severe revictimisation [32,35] and as such prevention science must be concerned with such high prevalence to institute relevant behaviour change strategies.

We found that a significant proportion of women (14.3%) were sexually abused before age 15. Although this is lower than reported in South Africa it is of concern because early sexual

debut increases risk of HIV infection among girls [22]. Young women who have sex early have a greater lifetime number of sexual partners and increased chances of partnering high risk men, usually older than them, have less education and less social integration [36] than those who do not initiate sex early. Mathematical modelling with data in Zimbabwe has shown that a reduction in partnering with older partners reduces individual risk of HIV infection among women [37]. Therefore protecting children from abuse may be a viable entry point towards HIV prevention.

The analysis shows that both child sexual and physical abuse are major drivers of HIV risk as has been found in South Africa, also among low income women [30].

The link between abusive events and HIV cannot be overemphasized. Proponents of the child development model argue that exposure to abuse in childhood negatively impacts on the child's development leading to cognitive, psychological, and social impairment [13,38]. This may then lead to limited life skills consequently making them vulnerable to sexual abuse and HIV risk behaviours in relationships. Secondary victimisation in life is more likely in vulnerable people who have had the experience of violence and abuse normalised in their subjective experience [32]. Some researchers have suggested two pathways linking violence in childhood with HIV infection. The first is a direct pathway that includes abusive experiences that lead to HIV infection through rape and forced sex. There is established evidence that points to the high likelihood of being HIV positive for people who rape girls and women [15,39–43]. Secondly, violence against women reduces one's psychological wellbeing leading to risky behaviours and HIV infection as well as increasing fear and non-use of protective mechanisms that again lead to HIV infection [39]. Due to repeated exposure to violence one may lack the sense of urgency to resist unprotected sex which fuels HIV infection. Most of the discourse on child protection needs is contextualised more specifically in relation to orphans. However, our study suggests that child protection rights, including directed at non-orphans, must be discussed as part of HIV prevention strategies.

Despite significant findings, this study has several limitations that must be described to guide the interpretation of the results. Firstly, the study was cross sectional and involved a clinic population and as such does not allow making general conclusions that childhood experiences lead to HIV infection. Even if we controlled for co-variables there may be other factors that we did not include in our questionnaire that may have contributed to such positive relationships. Prospective studies are required to identify possible causal relationships. Nevertheless, since we controlled for covariates, our results highlight the presence of the relationship between childhood abusive relationships and HIV. Secondly, asking women whose average age is 26 years about what happened before they turned 15 years may be subject to recall bias thereby underreporting the violence that happened to them [44]. Thirdly, some women may not report experiences that they perceive as minor or unimportant to report [45,46] or simply do not recall abusive events that happened many years back in their childhood [47]. However, studies have shown that abusive experiences are difficult to forget [48] and can haunt a person for a long time in their lives [49] leading them to report them since a research setting presents an opportunity to vent their challenges and feel better [50] although new evidence from prospective and intergenerational studies is beginning to show different trends and dynamics [51]. Fourthly, a dedicated study on childhood exposure to violence using recognised tools such as the Child Trauma questionnaire [52] would have improved our understanding of the exposure to violence in childhood. Lastly, our study only presented HIV status results and did not assess when the HIV was acquired as it is possible that they may have acquired it at the same time or prior to the abuse experiences. It is also possible that perhaps women who are HIV-positive have suffered greater hardships in general and therefore perceive their childhood more negatively. Our methodology has its strengths. Privately asking women about violence in primary

health care settings offers a secure place to disclose painful experiences in one's life compared to disclosing in household surveys where partners are usually available and limit women's freedom to express themselves. A large sample size powered the study adequately to conduct multivariate analysis with a number of variables. Lastly, the analysis controlled for covariates and so helped to give a fuller picture of factors that are individually associated with HIV status.

Conclusion

Exposure to child abuse is commonplace in Africa and happens in several ways highlighted in this study from physical to sexual violence, although this subject is not finding adequate attention from academic, implementing professionals and policy makers. The study therefore highlights the importance of protecting children as a way of preventing HIV infections in women and more broadly in society. Interventions to prevent HIV in women must therefore consider protecting children while researchers need to give priority to child abuse research in African settings.

Supporting information

S1 File. Anonymised data set.
(DTA)

Author Contributions

Conceptualization: Simukai Shamu, Patience Shamu, Christina Zarowsky, Marleen Temmerman, Tamara Shefer, Naeemah Abrahams.

Data curation: Simukai Shamu.

Formal analysis: Simukai Shamu, Patience Shamu.

Funding acquisition: Simukai Shamu, Christina Zarowsky, Naeemah Abrahams.

Investigation: Simukai Shamu, Christina Zarowsky.

Methodology: Simukai Shamu, Patience Shamu, Christina Zarowsky, Marleen Temmerman, Naeemah Abrahams.

Project administration: Simukai Shamu.

Resources: Simukai Shamu.

Supervision: Simukai Shamu, Christina Zarowsky, Marleen Temmerman, Tamara Shefer, Naeemah Abrahams.

Validation: Simukai Shamu.

Writing – original draft: Simukai Shamu, Patience Shamu, Christina Zarowsky, Marleen Temmerman, Tamara Shefer, Naeemah Abrahams.

Writing – review & editing: Simukai Shamu, Patience Shamu, Christina Zarowsky, Marleen Temmerman, Tamara Shefer, Naeemah Abrahams.

References

1. Stoltenborgh M, Bakermans-Kranenburg MJ, Alink LRA, van IJzendoorn MH. The Prevalence of Child Maltreatment across the Globe: Review of a Series of Meta-Analyses. *Child Abuse Rev* 2015; 24:37–50. <https://doi.org/10.1002/car.2353>

2. Meinck F, Cluver LD, Boyes ME, Mhlongo EL. Risk and Protective Factors for Physical and Sexual Abuse of Children and Adolescents in Africa: A Review and Implications for Practice. *Trauma, Violence, Abus* 2015; 16:81–107. <https://doi.org/10.1177/1524838014523336> PMID: 24648489
3. Shamu S, Gevers A, Mahlangu BP, Jama Shai PN, Chirwa ED JR. Prevalence and risk factors for intimate partner violence among Grade 8 learners in urban South Africa: baseline analysis from the Skhokho Supporting Success cluster randomised controlled trial 2015; 8:18–26. <https://doi.org/10.1093/inthealth/ihv068>
4. Meinck F, Cluver LD, Orkin FM, Kuo C, Sharma AD, Hensels IS, et al. Pathways From Family Disadvantage via Abusive Parenting and Caregiver Mental Health to Adolescent Health Risks in South Africa. *J Adolesc Heal* 2017; 60:57–64. <https://doi.org/10.1016/j.jadohealth.2016.08.016> PMID: 27793729
5. Meinck F., Cluver L.D., Boyes M.E. Ndhlovu LD. Risk and protective factors for physical and emotional abuse victimisation amongst vulnerable children in South Africa. *Child Abus Rev* 2015; 24:182–97.
6. Shumba A. Epidemiology and etiology of reported cases of child physical abuse in Zimbabwean primary schools. *Child Abuse Negl* 2001; 25:265–77. PMID: 11330924
7. Carey, Paul D., Jennifer L. Walker, Rossouw Wendy, Soraya Seedat DJS. Risk indicators and psychopathology in traumatised children and adolescents with a history of sexual abuse. *Eur Child Adolesc Psychiatry* 2008; 17:93–8. <https://doi.org/10.1007/s00787-007-0641-0> PMID: 17876504
8. Jewkes RK, Dunkle K, Nduna M, Jama PN, Puren A. Associations between childhood adversity and depression, substance abuse and HIV and HSV2 incident infections in rural South African youth. *Child Abuse Negl* 2010; 34:833–41. <https://doi.org/10.1016/j.chiabu.2010.05.002> PMID: 20943270
9. Shamu S, Gevers A, Mahlangu BP, Jama PN, Chirwa ED. Prevalence and risk factors for intimate partner violence among Grade 8 learners in urban South Africa: baseline analysis from the Skhokho Supporting Success cluster randomised controlled trial 2015; 8:18–26. <https://doi.org/10.1093/inthealth/ihv068>
10. Birdthistle IJ, Floyd S, Mwanasa S, Nyagadza A, Gwiza E, Glynn J. Child sexual abuse and links to HIV and orphanhood in urban Zimbabwe. *J Epidemiol Community Health* 2011; 65:1075–82. <https://doi.org/10.1136/jech.2009.094359> PMID: 20628080
11. Bensley LS, Van Eenwyk J, Simmons KW. Self-reported childhood sexual and physical abuse and adult HIV-risk behaviors and heavy drinking. *Am J Prev Med* 2000; 18:151–8. [https://doi.org/10.1016/S0749-3797\(99\)00084-7](https://doi.org/10.1016/S0749-3797(99)00084-7) PMID: 10698246
12. Brown DW, Riley L, Butchart A, Meddings DR, Kann L, Harvey AP. Exposure to physical and sexual violence and adverse health behaviours in African children: Results from the Global School-based Student Health Survey. *Bull World Health Organ* 2009; 87:447–55. <https://doi.org/10.2471/BLT.07.047423> PMID: 19565123
13. Norman RE, Byambaa M, De R, Butchart A, Scott J, Vos T. The Long-Term Health Consequences of Child Physical Abuse, Emotional Abuse, and Neglect: A Systematic Review and Meta-Analysis. *PLoS Med* 2012; 9. <https://doi.org/10.1371/journal.pmed.1001349> PMID: 23209385
14. Lindert J, Von Ehrenstein OS, Grashow R, Gal G, Braehler E, Weisskopf MG. Sexual and physical abuse in childhood is associated with depression and anxiety over the life course: Systematic review and meta-analysis. *Int J Public Health* 2014; 59:359–72. <https://doi.org/10.1007/s00038-013-0519-5> PMID: 24122075
15. Jewkes R, Dunkle K, Nduna M, Levin J, Jama N, Khuzwayo N, et al. Factors associated with HIV serostatus in young rural South African women: Connections between intimate partner violence and HIV. *Int J Epidemiol* 2006; 35:1461–8. <https://doi.org/10.1093/ije/dyl218> PMID: 17008362
16. Cohen M, Deamant C, Barkan S, Richardson J, Young M, Holman S, et al. Domestic violence and childhood sexual abuse in HIV-infected women and women at risk for HIV. *Am J Public Health* 2000; 90:560–5. <https://doi.org/10.2105/AJPH.90.4.560> PMID: 10754970
17. Gavin L, Galavotti C, Dube H, McNaghten AD, Murwirwa M, Khan R, et al. Factors Associated with HIV Infection in Adolescent Females in Zimbabwe. *J Adolesc Heal* 2006; 39. <https://doi.org/10.1016/j.jadohealth.2006.03.002> PMID: 16982397
18. Schaefer R, Gregson S, Eaton JW, Mugurungi O, Rhead R, Takaruzza A, et al. Age-disparate relationships and HIV incidence in adolescent girls and young women: Evidence from Zimbabwe. *AIDS* 2017; 31:1461–70. <https://doi.org/10.1097/QAD.0000000000001506> PMID: 28426534
19. Shamu S, Abrahams N, Zarowsky C, Shefer T, Temmerman M. Intimate partner violence during pregnancy in Zimbabwe: A cross-sectional study of prevalence, predictors and associations with HIV. *Trop Med Int Heal* 2013; 18:696–711. <https://doi.org/10.1111/tmi.12078> PMID: 23414103
20. Garcia-Moreno C, Jansen H a FM, Ellsberg M, Heise L, Watts CH. WHO Multi-country Study on Women ' s Health and Domestic Initial results on prevalence. *Genetics* 2005; 151:277–83. [https://doi.org/10.1016/S0140-6736\(06\)69523-8](https://doi.org/10.1016/S0140-6736(06)69523-8)

21. Dunkle KL, Jewkes RK, Brown HC, Gray GE, McIntyre JA, Harlow SD. Gender-based violence, relationship power, and risk of HIV infection in women attending antenatal clinics in South Africa. *Lancet* (London, England) 2004; 363:1415–21. [https://doi.org/10.1016/S0140-6736\(04\)16098-4](https://doi.org/10.1016/S0140-6736(04)16098-4) PMID: 15121402
22. Pettifor AE, Van Der Straten A, Dunbar MS, Shiboski SC, Padian NS. Early age of first sex: A risk factor for HIV infection among women in Zimbabwe. *AIDS* 2004; 18:1435–42. <https://doi.org/10.1097/01.aids.0000131338.61042.b8> PMID: 15199320
23. Government of Zimbabwe. Sexual offences act. 2001.
24. Garcia-Moreno C, Jansen H a FM, Ellsberg M, Heise L, Watts CH. Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *Lancet* 2006; 368:1260–9. [https://doi.org/10.1016/S0140-6736\(06\)69523-8](https://doi.org/10.1016/S0140-6736(06)69523-8) PMID: 17027732
25. StataCorp. Stata Statistical Software: Release 13. 2013 2013. <https://doi.org/10.2307/2234838>
26. World Health Organisation. Putting Women First: Ethical and safety recommendations for research on domestic violence against women. *World Heal Organ* 2001:33.
27. Shumba A, Mpofu E, Chireshe R, Mapfumo J. Corporal Punishment in Zimbabwean Schools: Aetiology and Challenges 1995:1–10.
28. Widom CS, Czaja S, Dutton MA. Child abuse and neglect and intimate partner violence victimization and perpetration: A prospective investigation. *Child Abuse Negl* 2014; 38:650–63. <https://doi.org/10.1016/j.chiabu.2013.11.004> PMID: 24325940
29. Barth J, Bermetz L, Heim E, Trelle S, Tonia T. The current prevalence of child sexual abuse worldwide: A systematic review and meta-analysis. *Int J Public Health* 2013; 58:469–83. <https://doi.org/10.1007/s00038-012-0426-1> PMID: 23178922
30. Gibbs A, Dunkle K, Washington L, Willan S, Shai N, Jewkes R. Childhood traumas as a risk factor for HIV-risk behaviours amongst young women and men living in urban informal settlements in South Africa: A cross-sectional study. *PLoS One* 2018; 13:e0195369. <https://doi.org/10.1371/journal.pone.0195369> PMID: 29624612
31. Weems CF, Russell JD, Banks DM, Graham R a, Neill EL, Scott BG. Memories of Traumatic Events in Childhood Fade After Experiencing Similar Less Stressful Events: Results From Two Natural Experiments. *J Exp Psychol Gen* 2014; 143:No Pagination Specified. <https://doi.org/10.1037/xge0000016> PMID: 25068536
32. Dunkle KL, Jewkes RK, Brown HC, Yoshihama M, Glenda E, Mcintyre JA, et al. Prevalence and Patterns of Gender-based Violence and Revictimization among Women Attending Antenatal Clinics in Soweto, South Africa 2004; 160:230–9. <https://doi.org/10.1093/aje/kwh194>
33. Koenig M a, Zablotska I, Lutalo T, Nalugoda F, Wagman J, Gray R. Coerced first intercourse and reproductive health among adolescent women in Rakai, Uganda. *Int Fam Plan Perspect* 2004; 30:156–63. <https://doi.org/10.1363/iffp.30.156.04> PMID: 15590381
34. Zimbabwe National Statistics Agency ICFII. Zimbabwe Demographic and Health Survey 2010–11 2012:1–470.
35. Shamu S. The dynamics of intimate partner violence during pregnancy and linkages with hiv infection and disclosure in zimbabwe. Ghent: International Centre for Reproductive Health; 2013.
36. Hallett TB, Lewis JJC, Lopman BA, Nyamukapa CA, Mushati P, Wambe M, et al. Age at first sex and HIV infection in rural Zimbabwe. *Stud Fam Plann* 2007; 38:1–10. <https://doi.org/10.1111/j.1728-4465.2007.00111.x> PMID: 17385378
37. Hallett TB, Gregson S, Lewis JJC, Lopman BA, Garnett GP. Behaviour change in generalised HIV epidemics: Impact of reducing cross-generational sex and delaying age at sexual debut. *Sex Transm Infect* 2007; 83. <https://doi.org/10.1136/sti.2006.023606> PMID: 17314125
38. Perry BD. The Neurodevelopmental Impact of Violence in Childhood. *Textb Child Adolesc Forensic Psychiatry* 2001:221–38.
39. Dunkle KL, Decker MR. Gender-Based Violence and HIV: Reviewing the Evidence for Links and Causal Pathways in the General Population and High-risk Groups. *Am J Reprod Immunol* 2013; 69:20–6. <https://doi.org/10.1111/aji.12039> PMID: 23216606
40. Jewkes R, Nduna M, Levin J, Jama N, Dunkle K, Khuzwayo N, et al. A cluster randomized-controlled trial to determine the effectiveness of Stepping Stones in preventing HIV infections and promoting safer sexual behaviour amongst youth in the rural Eastern Cape, South Africa: Trial design, methods and baseline findings. *Trop Med Int Heal* 2006; 11:3–16. <https://doi.org/10.1111/j.1365-3156.2005.01530.x> PMID: 16398750
41. Khanna A, Goyal RS, Bhawsar R. Menstrual Practices and Reproductive Problems: A Study of Adolescent Girls in Rajasthan. *J Health Manag* 2005; 7:91–107. <https://doi.org/10.1177/097206340400700103>

42. Raj A, Santana MC, Marche A La, Amaro H, Cranston K, Silverman JG. Perpetration of Intimate Partner Violence Associated With Sexual Risk Behaviors Among Young Adult Men 2006; 96:1873–8. <https://doi.org/10.2105/AJPH.2005.081554>
43. Silverman JG, Decker MR, Kapur NA, Gupta J, Raj A. Violence against wives, sexual risk and sexually transmitted infection among Bangladeshi men 2007:211–5. <https://doi.org/10.1136/sti.2006.023366>
44. Williams LW. Recall of childhood trauma: A prospective study of women's memories of child sexual abuse. *J Consult Clin Psychol* 1994; 62:1167–76. <https://doi.org/10.1037/0022-006X.62.6.1167> PMID: 7860814
45. Pallitto CC, García-Moreno C, Jansen HAFM, Heise L, Ellsberg M, Watts C. Intimate partner violence, abortion, and unintended pregnancy: results from the WHO Multi-country Study on Women's Health and Domestic Violence. *Int J Gynaecol Obstet* 2013; 120:3–9. <https://doi.org/10.1016/j.ijgo.2012.07.003> PMID: 22959631
46. Gil-González D, Vives-Cases C, Ruiz MT, Carrasco-Portiño M, Álvarez-Dardet C. Childhood experiences of violence in perpetrators as a risk factor of intimate partner violence: A systematic review. *J Public Health (Bangkok)* 2008; 30:14–22. <https://doi.org/10.1093/pubmed/fdm071> PMID: 17986717
47. Williams LM. Recall of childhood trauma: A prospective study of women's memories of child sexual abuse. *J Consult Clin Psychol* 1994; 62:1167–76. <https://doi.org/10.1037/0022-006X.62.6.1167> PMID: 7860814
48. Bernet C.Z.; Stein MB. Relationship of childhood maltreatment to the onset and course of major depression in adulthood. *Depress Anxiety* 1999; 9:169–74. PMID: 10431682
49. Schwab G. *Haunting legacies: Violent histories and transgenerational trauma.*, Columbia University Press; 2010.
50. Dyregrov KM, Dieserud G, Hjelmeland HM, Straiton M, Rasmussen ML, Knizek BL, et al. Meaning-making through psychological autopsy interviews: The value of participating in qualitative research for those bereaved by suicide. *Death Stud* 2011; 35:685–710. <https://doi.org/10.1080/07481187.2011.553310> PMID: 24501818
51. Widom CS, Czaja SJ, DuMont KA. Intergenerational transmission of child abuse and neglect: Real or detection bias? *Science (80-)* 2015; 347:1480–5. <https://doi.org/10.1126/science.1259917> PMID: 25814584
52. Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abus Negl* 2003; 27:169–90. [https://doi.org/10.1016/S0145-2134\(02\)00541-0](https://doi.org/10.1016/S0145-2134(02)00541-0)