

Test of the Conserved Vector Current Hypothesis by β -ray Angular Distribution Measurement in the Mass-8 System

T. Sumikama,^{1,2} K. Matsuta,¹ T. Nagatomo,³ M. Ogura,¹
 T. Iwakoshi,¹ Y. Nakashima,¹ H. Fujiwara,¹ M. Fukuda,¹
 M. Mihara,¹ K. Minamisono,⁴ T. Yamaguchi,⁵ and T. Minamisono⁶

¹*Department of Physics, Osaka University,*

1-1 Machikaneyama, Toyonaka, Osaka 560-0043, Japan

²*Department of Physics, Faculty of Science and Technology,
 Tokyo University of Science, Noda, Chiba 278-8510, Japan*

³*Department of Chemistry, International Christian
 University, Mitaka, Tokyo 181-8585, Japan*

⁴*National Superconducting Cyclotron Laboratory,
 Michigan State University, East Lansing, Michigan 48824, USA*

⁵*Department of Physics, Saitama University, Saitama 338-8570, Japan*

⁶*Fukui University of Technology, 3-6-1 Gakuen, Fukui 910-8505, Japan*

(Dated: April 22, 2022)

Abstract

The β -ray angular correlations for the spin alignments of ${}^8\text{Li}$ and ${}^8\text{B}$ have been observed in order to test the conserved vector current (CVC) hypothesis. The alignment correlation terms were combined with the known β - α -angular correlation terms to determine all the matrix elements contributing to the correlation terms. The weak magnetism term, 7.5 ± 0.2 , deduced from the β -ray correlation terms was consistent with the CVC prediction 7.3 ± 0.2 , deduced from the analog- γ -decay measurement based on the CVC hypothesis. However, there was no consistent CVC prediction for the second-forbidden term associated with the weak vector current. The experimental value for the second-forbidden term was 1.0 ± 0.3 , while the CVC prediction was 0.1 ± 0.4 or 2.1 ± 0.5 .

PACS numbers: 23.20.En, 23.40.Bw

I. INTRODUCTION

In the standard electroweak model, the weak vector current of nucleons and the isovector part of the electromagnetic current form a single isovector electroweak current [1]. The conserved vector current (CVC) hypothesis is analogous to the electromagnetic current conservation law. The weak vector current is conserved despite the influence of strong interactions such as the contribution from the pion cloud around a nucleon. The minimum conditions required for the CVC hypothesis are the universality of the vector coupling constant g_V and the absence of the induced scalar term g_S in the weak nucleon current. The universality of g_V is confirmed at the level of 1.2×10^{-4} and g_S is limited to $m_e g_S / 2M_n g_V = -(0.0011 \pm 0.0013)$ from 20 superallowed $0^+ \rightarrow 0^+$ β decays [2], where M_n and m_e are the nucleon and electron masses, respectively. In this article, the coupling constant and the induced term in the weak nucleon current, including g_V and g_S , are written following Holstein's expression [3].

In addition, the so-called strong CVC hypothesis demands that the weak vector current is paired with the isovector electromagnetic current. For the isospin triplet state, the strong CVC requires that a matrix element from the weak vector current for β decay is identical to one from the isovector electromagnetic current for analog- γ decay. To test the strong CVC hypothesis, the weak magnetism term a_{WM}^β has been compared with the CVC prediction deduced from the isovector M1 component of the analog- γ transition strength [4]. The term a_{WM}^β was detected using one of the following: a spectral shape factor, a β -ray angular correlation with a spin orientation, or a correlation with a delayed α or γ ray [4–7]. The experimentally determined a_{WM}^β may include a possible G -parity irregular term, g_{II} , in the weak axial-vector current; therefore, the strong CVC has been tested under the assumption of G -parity conservation. Among those studies, Minamisono *et al.* [6] determined the most accurate a_{WM}^β in the mass $A = 12$ system. In their paper, the g_{II} was determined using the CVC prediction, $a_{\text{WM}}^{\text{CVC}}$, for the weak magnetism term. When G -parity conservation was assumed, the strong CVC was confirmed as $a_{\text{WM}}^\beta / a_{\text{WM}}^{\text{CVC}} = 1.04 \pm 0.03$ [8].

Earlier, the strong CVC in the $A = 8$ system was tested using the β - α angular correlation terms of ${}^8\text{Li}$ and ${}^8\text{B}$ [9–11]. Among the previous measurements, those by Tribble and McKewen [10, 11] were performed for a wide energy range of β rays, and the mirror difference $\delta_{\beta\alpha}^-$ of the β - α angular correlation term was determined. $\delta_{\beta\alpha}^-$ has a contribution of a_{WM}^β and a second-forbidden term a_{WE2}^β associated with the weak vector current. While the Tribble's

data did not reproduce the kinematic shift term for the angular correlation, the McKeown's data reproduced this term properly, and the result was consistent with the CVC prediction value δ_{CVC}^- [7] as $\delta_{\beta\alpha}^-/\delta_{\text{CVC}}^- = 0.93 \pm 0.03 \pm 0.05$, where the first uncertainty was from the β - α measurement and the second one was from the CVC prediction.

The β -ray angular distributions of ^8Li and ^8B are given by a combination of several matrix elements; not only a_{WM}^β and a_{WE2}^β but also the Gamow-Teller, axial charge, and second forbidden terms from the axial-vector currents. In spite of this complexity, we previously showed [12] that a_{WM}^β and a_{WE2}^β could be determined separately by combining the alignment correlation term and the β - α angular correlation term. Thus, the strong CVC can be tested for the second-forbidden transition for the first time.

In our previous letter [12], we reported the measurement of the β -ray angular correlation term from the spin aligned ^8Li and ^8B ($J^\pi = 2^+$) and the limitation of g_{II} under the assumption of CVC. In the present study, we reanalyzed the data with the assumption of G -parity conservation in order to test the strong CVC hypothesis for the weak magnetism and for the second-forbidden transition separately.

II. BETA-RAY ANGULAR CORRELATION TERMS AND ANALOG GAMMA DECAY

The two kinds of β -ray angular correlation term, i.e., the alignment correlation term and the β - α angular correlation term, are similar to each other. The alignment correlation term is associated with the spin alignment of parent nucleus. Because β -delayed α particles are emitted in the direction perpendicular to the angular momentum of the daughter nucleus ^8Be ($J^\pi = 2^+$), the β - α angular correlation term is associated with the spin alignment of the daughter nucleus. As a result, the alignment correlation term and the β - α angular correlation term have the same formula except for the signs of several second-forbidden terms. This complementary relationship allows all the matrix elements to be separately determined, as follows.

The β -ray angular distribution from purely spin-aligned nuclei is given by $W(E, \theta_{I\beta}) \propto pE(E_0 - E)^2\{B_0(E) + \mathcal{A}B_2(E)P_2(\cos \theta_{I\beta})\}$, where p , E , E_0 , and $\theta_{I\beta}$ are the β -ray momentum, energy, end-point energy, and ejection angle with respect to the spin-orientation axis, respectively. P_2 is the Legendre polynomial. The ^8Li and ^8B nuclei decay to the broad first

excited state of ${}^8\text{Be}$, thus the end-point energy E_0 is given as $E_0 = E_{\text{max}} - E_x$. E_{max} is the energy release during the β decay to the ${}^8\text{Be}$ ground state, while E_x is the excitation energy of ${}^8\text{Be}$. The nuclear-spin alignment $\mathcal{A} = (2a_{+2} - a_{+1} - 2a_0 - a_{-1} + 2a_{-2})/2$ is given by the population a_m of the magnetic substate m , with $\sum a_m = 1$. The alignment correlation terms $B_2(E)/B_0(E)$ for ${}^8\text{Li}$ and ${}^8\text{B}$ are given by $K(E, 0)$ in [3] as

$$K(E, s) = -\frac{E}{3M_n} \left[\frac{1}{A} \pm \frac{b}{Ac} - \frac{d_1}{Ac} \mp \frac{g_{\text{II}}}{g_A} \right. \\ \left. + \frac{(-)^s}{\sqrt{14}} \left\{ \pm \frac{f}{Ac} \frac{E_0 + 2E}{E_0} + \frac{3}{2} \frac{j_2}{A^2c} \frac{E_0 - 2E}{M_n} \right\} \right. \\ \left. - \frac{3}{\sqrt{35}} \frac{j_3}{A^2c} \frac{E}{M_n} \right], \quad (1)$$

where g_A is the axial-vector coupling constant, c is the Gamow-Teller matrix element, b is the weak magnetism matrix element, d_1 is the axial charge, f is the second-forbidden term from the vector current, j_2 and j_3 are the second-forbidden terms from the axial-vector current, and A is the mass number of the nucleus. a_{WM} and a_{WE2} are given by the ratios $a_{\text{WM}} = b/Ac$ and $a_{\text{WE2}} = f/Ac$. The β - α angular correlation term, on the other hand, is given by $W(E, \theta_{\beta\alpha}) \propto pE(E_0 - E)^2 \{1 + a_{\mp}(E) \cos \theta_{\beta\alpha} + p_{\mp}(E) \cos^2 \theta_{\beta\alpha}\}$, where $\theta_{\beta\alpha}$ is the angle between the momenta of β and α rays. $a_{\mp}(E)$ is the kinematic shift term associated with the recoil of the daughter nucleus. The β - α angular correlation term $p_{\mp}(E)$ is given as $-\frac{2}{3}p_{\mp}(E) = K(E, 1)$, which is also defined by Eq. (1). The difference in the correlation terms between the mirror pair, $\delta_{\text{align.}}^- = (B_2/B_0)_{\text{sLi}} - (B_2/B_0)_{\text{sB}}$ and $\delta_{\beta\alpha}^- = (-2/3)(p_- - p_+)$, consists of only three terms, b/Ac , g_{II}/g_A , and f/Ac . The b/Ac term is determined under the assumption that $g_{\text{II}} = 0$; f/Ac is completely separated from the others as follows:

$$\frac{\delta_{\text{align.}}^- + \delta_{\beta\alpha}^-}{2} = -\frac{2E}{3M_n} \frac{b}{Ac} \quad (2)$$

$$\frac{\delta_{\text{align.}}^- - \delta_{\beta\alpha}^-}{2} = -\frac{2E}{3M_n} \frac{f}{Ac} \frac{E_0 + 2E}{\sqrt{14}E_0}. \quad (3)$$

The c , b , and f terms described by reduced matrix elements as follows: $c = g_A \langle f || \tau^{\pm} \boldsymbol{\sigma} || i \rangle$, $b = A (g_{\text{M}} \langle f || \tau^{\pm} \boldsymbol{\sigma} || i \rangle + g_{\text{V}} \langle f || \tau^{\pm} \mathbf{L} || i \rangle)$, and $f = 2(2\pi/15)^{1/2} AM_n E_0 g_{\text{V}} \langle f || \tau^{\pm} r^2 Y_2(\hat{r}) || i \rangle$, where g_{M} is the weak magnetism coupling constant in the weak vector current. c is determined from the β -decay half-lives of ${}^8\text{Li}$ and ${}^8\text{B}$.

The matrix elements depend on the final state energy which is broadly distributed. The E_x dependence of c and b is taken into account by using R -matrix theory as described

in Secs. VB and VD. The E_x dependence of the others is considered as a systematic uncertainty as described in Sec. VD.

The requirement by strong CVC is that b and f contribute also to the electromagnetic transition from the isobaric analog state in ${}^8\text{Be}$. b and f are related to the isovector components of the M1 and E2 transition strengths, $\Gamma_{\text{M1}}^{T=1}$ and $\Gamma_{\text{E2}}^{T=1}$, i.e. $b = AM_n \{6\Gamma_{\text{M1}}^{T=1}/(\alpha E_\gamma^3)\}^{1/2}$ and $f/b = \sqrt{10/3}\delta_1$ [7]. Here E_γ is the γ -ray energy, the fine structure constant $\alpha = 1/137$, and the M1/E2 ratio $\delta_1 = (\Gamma_{\text{E2}}^{T=1}/\Gamma_{\text{M1}}^{T=1})^{1/2}$.

The initial state of the analog- γ decay splits into two isospin mixing states with $T = 0$ and 1. In addition, the electromagnetic transitions from these states include the isoscalar and isovector components. Two strengths $\Gamma_{\text{M1}}^{T=1}$ and $\Gamma_{\text{E2}}^{T=1}$ are the isovector component from the state with $T = 1$. The measurement of the γ decay from these states and the extraction of $\Gamma_{\text{M1}}^{T=1}$ and $\Gamma_{\text{E2}}^{T=1}$ were performed in the previous work by De Braeckeleer *et al.* [7].

III. EXPERIMENTAL

In this section, the experimental details for the alignment correlation term measurement is described. Figure 1 shows the experimental setup, which is essentially similar to the previous experiment for the alignment correlation terms of ${}^{12}\text{B}$ and ${}^{12}\text{N}$ [6].

A. Production of unstable ${}^8\text{Li}$ and ${}^8\text{B}$

The ${}^8\text{Li}$ [${}^8\text{B}$] nuclei were produced through the nuclear reactions ${}^7\text{Li}(d,p){}^8\text{Li}$ [${}^6\text{Li}({}^3\text{He},n){}^8\text{B}$]. Hereafter, information with the parentheses represent the conditions for ${}^8\text{B}$. A Li_2O [enriched metal ${}^6\text{Li}$] target was bombarded by a deuteron [${}^3\text{He}$] beam at 3.5 MeV [4.7 MeV] with a typical intensity of 9 μA [40 μA]. A rotating target, which occupied one third of the circumference of the target rotor, was cooled from inside the holder by a compressed air jet in order to withstand the high-intensity ${}^3\text{He}$ beam, which operated at 4.7 MeV up to 40 μA . The pulsed beam was synchronized to the rotational period of 2.4 s. The beam-on and beam-off times were 0.8 s and 1.6 s, respectively. The target material was vacuum evaporated on a backing ribbon made of molybdenum [phosphor bronze]. Phosphor bronze was used to reduce a Rutherford scattering of ${}^3\text{He}$, which could otherwise have bombarded the recoil catcher and have been an origin of disturbing background activities. A new ion-source




FIG. 1. Schematic view of the experimental setup. The rotational target with an air cooling system was used to reduce the background from the target. The catcher and the rf coil were placed at the center of two telescopes. Each plastic scintillation detector telescope consisted of two thin ΔE detectors (A and B), one veto detector (\bar{C}), and one energy detector (E).

bottle, made of glass, was used for the ^3He beam to prevent a very weak HD^+ molecular ion beam from mixing with the ^3He beam. The HD^+ ion beam was formed by H_2 and D_2 gases oozing out from the inner wall of the ion-source bottle, which were in turn used for the production of p^+ or d^+ beams.

B. Recoil implantation of polarized nuclei

The recoil angle of the nuclear-reaction products was selected in the range 14° - 40° [7° - 18°] to optimize the obtained polarization. The polarized ^8Li [^8B] nuclei were implanted in Zn [TiO_2 (rutile structure)] single crystals by using a recoil energy of 1.7 MeV [2.3 MeV] obtained by the nuclear reaction. The crystals were placed in a static magnetic field B_0 to maintain the polarization and to manipulate the spin orientation using the β -NMR technique. The c axis of the single crystals was set parallel to B_0 , which was 60 mT [230 mT]. An asymmetry of β -rays emitted from polarized nuclei was detected by two sets of the counter telescope placed in the opposite direction. The obtained polarization was determined to be 7.2% [5.4%] from the β -ray asymmetry by using the β -NMR technique.




FIG. 2. Focused view of NMR equipment. An rf oscillating magnetic field B_1 , which was applied by the rf coil, is perpendicular to the external magnetic field B_0 .

The recoil catcher consisted of a pair of crystals, which were tilted 45° with respect to the magnetic field but in opposite directions in order to form a dog-leg shape (similar to a half-opened book) as seen from the side, as shown in Fig. 2. The implantation depth was uniformly distributed at $2.4 \mu\text{m}$ [$3.1 \mu\text{m}$] from the surface. The recoil nuclei were implanted from the inner side of the two crystals, making the path length and the energy loss of the β rays in the catcher less sensitive to the β -emitter position. The thickness of crystals was $360 \pm 20 \mu\text{m}$ for the guph Zn crystal and $250 \pm 20 \mu\text{m}$ for the gdownh crystal, and $100 \pm 10 \mu\text{m}$ for both TiO_2 crystals. The systematic uncertainty due to the ambiguity in the thickness was considered as discussed in Sec. IV D.

C. Spin manipulation

In order to convert the initial polarization into positive and negative alignments with, ideally, zero polarization, the nuclear spin was manipulated using the NMR technique. The Larmor frequency for spin $J = 2$ nucleus splits into four resonance frequencies because of hyperfine interaction between the electric quadrupole moment Q of the implanted nucleus and the electric field gradient (EFG) at an implantation site in the crystal. EFG is defined by $V_{ii} = d^2V/di^2$, where i is the principal axes of EFG, i.e., X, Y , and Z , $V_{XX} + V_{YY} + V_{ZZ} = 0$, and $|V_{XX}| \leq |V_{YY}| \leq |V_{ZZ}|$. Therefore, once principal axes are chosen, EFG is given by

two parameters $q = V_{ZZ}$ and $\eta = (V_{XX} - V_{YY})/V_{ZZ}$. The resonance frequency between two neighboring magnetic substates, $(m - 1) \leftrightarrow m$, is given in [13] as

$$\nu_{m-1 \leftrightarrow m} = \nu_L - \frac{\nu_Q}{4}(3 \cos^2 \theta - 1 + \eta \sin^2 \theta \cos 2\phi)(2m - 1), \quad (4)$$

where ν_L is the Larmor frequency, $\nu_Q = eQq/4h$, and θ and ϕ are the Euler angle between the principal axes of EFG and the external magnetic field, respectively.

Populations of two neighboring magnetic substates can be manipulated independently by applying an rf oscillating magnetic field at each frequency. EFG at implantation sites in crystals has been studied by the β -NMR technique [14–16]. The number of possible implantation sites is one for ${}^8\text{Li}$ in Zn and two for ${}^8\text{B}$ in TiO_2 . The relative populations are 90% for ${}^8\text{B}$ implanted in the major site of TiO_2 and 10% for that in the minor site. ν_Q and η have been determined as $\nu_Q = +8.4 \pm 0.5$ kHz and $\eta = 0$ for the implantation site of ${}^8\text{Li}$ in Zn [14] and as $\nu_Q = +144.5 \pm 0.6$ kHz and $\eta < 0.03$ for the major implantation site of ${}^8\text{B}$ in TiO_2 [16]. Because of a small population for the minor site, it was difficult to detect a β -NQR signal for ${}^8\text{B}$ in the minor site. ν_Q and η at the minor site of ${}^8\text{B}$ in TiO_2 was evaluated as $\nu_Q = +1185 \pm 8$ kHz and $\eta = 0.020 \pm 0.006$ from ν_Q at the minor site of ${}^{12}\text{B}$ [15] and the ratio of the Q moments of ${}^8\text{B}$ and ${}^{12}\text{B}$ [16]. The directions of q at an implantation site of ${}^8\text{Li}$ in Zn and at the major site of ${}^8\text{B}$ in TiO_2 [14, 16] were parallel to the c axis of the crystals, i.e., $\theta = 0$, thus giving four frequencies split at regular intervals. For the minor site of ${}^8\text{B}$ in TiO_2 , the direction of q was inclined at 106° relative to the $\langle 100 \rangle$ axis on the (001) plane and the direction of V_{YY} was parallel to the c axis of the crystal [15], i.e., $\theta = \phi = 90^\circ$, as the c axis was parallel to B_0 . Frequencies for ${}^8\text{B}$ at the major and the minor sites are shown as a function of B_0 in Fig. 3. Frequencies for the major site were isolated from those of the minor site only for the experimental condition of $B_0 = 230$ mT. Under this condition, only the nuclear spin of ${}^8\text{B}$ implanted in the major site can be manipulated. The β -ray angular distribution from the unmanipulated ${}^8\text{B}$ in the minor site was stable. Because the alignment correlation term was derived from the dependence of the β -ray angular distribution on the degree of the alignment, the effect of ${}^8\text{B}$ in the minor site was canceled.

The spin-aligning procedure for spin $J = 2$ was newly developed as part of our study. Figure 4 shows the schematic aligning procedure using ${}^8\text{Li}$ as an example. Immediately after the pulsed beam was stopped, the nuclear spin was manipulated by applying two kinds of




FIG. 3. External magnetic field dependence of the resonance frequencies of ^8B in TiO_2 . The solid and dashed lines denote the frequencies of major and minor sites, respectively.

β -NMR technique, the adiabatic fast passage (AFP) and the depolarization methods. The populations between the two neighboring magnetic substates were interchanged by the AFP method and equalized by the depolarization method. To convert a positive polarization to a positive alignment \mathcal{A}^+ , the populations in $m = +2$ and $+1$, as well as in $m = -1$ and 0 were first of all equalized using the depolarization method. Following this, the positive alignment was produced by sequentially applying the AFP method four times, by which the populations between $m = +1$ and 0 , $m = -1$ and 0 , $m = -2$ and -1 , as well as $m = -1$ and 0 were interchanged. A negative alignment was produced immediately after the beam was stopped in the next beam-count cycle following a similar procedure applied to the magnetic substates as shown in the \mathcal{A}^- part of Fig. 4. For ^8B , an opposite sign of alignment was produced using the same procedure as for ^8Li , because the polarization initially obtained for ^8B was negative while the other parameters, i.e., the direction of the holding magnetic field and the field gradient, were similar. The alignment was converted back to a polarization to check the spin manipulation and to measure the relaxation time of the alignment. Subsequently, in the same beam-count cycle, the polarization was converted to an alignment with the opposite sign, as shown in Fig. 5. This method of data acquisition using the present timing program removed the systematic uncertainty due to beam fluctuation as described in Sec. IV B.




FIG. 4. Spin-aligning procedure for ${}^8\text{Li}$. The change in the populations, a_m , of the magnetic substate are shown. The spin manipulations with the AFP and depolarization methods of the NMR technique are denoted by the solid and dashed arrows, respectively. The two open bars in each orientation show the manipulated populations. The upper and lower parts show the production procedure for the positive and negative alignments, respectively. The polarizations of the three orientation patterns framed by the separate squares were measured to determine the alignment. The timing program for the measurement is shown in Fig. 5.

D. Beta-ray energy spectra

The β rays were detected by two sets of plastic scintillation counter telescopes placed above ($\theta_{I\beta} = 0^\circ$) and below (180°) the crystal as shown in Fig. 1. Each telescope consisted of two thin ΔE (A and B) detectors of $12\text{ mm}\phi \times 0.5\text{ mm}$ and $55\text{ mm}\phi \times 1\text{ mm}$, one β -ray energy (E) detector of $160\text{ mm}\phi \times 120\text{ mm}$, and one cone-shaped veto (\bar{C}) detector. The \bar{C} detector was used to reject the β rays scattered at the magnet. A typical counting rate of β rays from β emitters stopped in the catcher was 4 kcps (1.5 kcps).

The energy spectra of β -rays emitted from purely aligned ${}^8\text{Li}$ and ${}^8\text{B}$ are shown in Fig. 6. The gain in the analog signal was stabilized using the standard light pulse from a light-emitting-diode (LED) pulser whose the circuit was maintained at a constant temperature.

The energy deposit in the E detector for a monoenergetic β ray was obtained by a Monte Carlo simulation with the EGS4 code [17]. The detector telescopes, the catcher of the reaction products ${}^8\text{Li}$ and ${}^8\text{B}$, the catcher holder, and the vacuum chamber near the β -ray window were arranged in the simulation. The distribution of the reaction products on the catcher was given using the reaction kinematics. The response function was obtained by convoluting the deposit function with a detector resolution as shown in Fig. 7. The




FIG. 5. Timing program for the spin-aligning process. In each cycle, the positive and negative alignments were produced as shown in (1). The timing program for the spin manipulation and the β -ray-angular-distribution measurement are shown in (2). The ellipses and squares with two numbers show the spin manipulation with the AFP and depolarization methods, respectively, used for the relevant transition between the magnetic substate nominally shown by the two numbers. The β -ray angular distribution was observed at the half-height squares. The alignment was converted back to the polarized form to check the spin manipulation and to measure alignment relaxation.

resolution of the Gaussian function was determined by the χ^2 fitting of the β -ray energy spectra of ^8Li and ^8B with $\sigma = \sigma_0 \sqrt{E_{dep}}$, where $\sigma_0 = 0.10 \pm 0.02 \text{ (MeV)}^{1/2}$. Here, E_{dep} is the energy deposit in the E detector, which was observed, whereas the alignment correlation term needed to be extracted as a function of the β -ray energy just as it was emitted from the nucleus. The peak position of the energy deposit for monoenergetic β rays was scaled to the incident energy of the β -ray. The β -ray energy spectrum for the χ^2 fitting was obtained by convoluting the β -ray continuous energy spectrum with the response function of the monoenergetic β ray.




FIG. 6. Typical β -ray energy spectra for ${}^8\text{Li}$ (upper) and ${}^8\text{B}$ (lower). The dots are the experimental data and the solid curves are the best-fit lines. The energy region lying between the two vertical lines in each energy spectrum shows the region used for the line fitting. The background β rays in the low energy region of the ${}^8\text{B}$ spectrum were from ${}^{15}\text{O}$.

The β -ray energy was scaled by determining the end-point energies of several β -emitters, which were ${}^8\text{Li}$ itself, ${}^{28}\text{Al}$ ($E_0 = 2.86$ MeV), ${}^{20}\text{F}$ (5.39 MeV), and ${}^{12}\text{B}$ (13.37 MeV) for the ${}^8\text{Li}$ experiment, and ${}^8\text{B}$ itself, ${}^{15}\text{O}$ (1.73 MeV), ${}^{20}\text{F}$ (5.39 MeV), and ${}^{12}\text{N}$ (16.32 MeV) for the ${}^8\text{B}$ experiment.

IV. ANALYSIS

First, the determination of the degree of polarization and alignment, and then that of the alignment correlation terms are described. The evaluation of corrections and systematic uncertainties follows.

A. Degree of polarization and alignment

The polarization was determined from the β -ray asymmetry where β rays from 5 to 13 MeV were used. The β -ray angular distribution from the polarized nuclei is given by




FIG. 7. Response function of the E detector for β^- ray with 8 MeV. The horizontal axis has been rescaled from the simulated energy deposit in the E detector to the β -ray energy just after the emission from the nucleus.

$W(\theta_{I\beta}) \propto B_0(E) + B_1(E)\mathcal{P} \cos \theta_{I\beta} \propto 1 + A_s\mathcal{P} \cos \theta_{I\beta}$, where \mathcal{P} is the degree of the polarization. The asymmetry parameter, $A_s = B_1(E)/B_0(E)$, has an energy independent-main term and an energy-dependent higher order term. For the determination of the degree of the polarization and alignment, A_s was approximated as $-1/3$ for ${}^8\text{Li}$ and $+1/3$ for ${}^8\text{B}$. The effect on the alignment correlation term by the higher order term of A_s was corrected as described in Sec. IV C 2. The counting ratio of the top and bottom telescopes, i.e. $\theta_{I\beta} = 0^\circ$ and 180° , was caused by the β -ray asymmetry from the polarization \mathcal{P} as well as the possible geometrical asymmetry g resulting from the geometrical misalignment between two telescopes. This ratio is expressed as $R_\beta = W(0^\circ)/W(180^\circ) = g(1 + A_s\mathcal{P})/(1 - A_s\mathcal{P})$. To determine g , the polarization was inverted by applying a series of 10 AFPs. The measured counting ratios for the initial polarization $R_{\mathcal{P}_+}$, the inverted polarization $R_{\mathcal{P}_-}$, and the twice inverted polarization $R_{\mathcal{P}_{++}}$ are given by

$$R_{\mathcal{P}_+} = g(1 + A_s\mathcal{P}_0)/(1 - A_s\mathcal{P}_0) \quad (5)$$

$$R_{\mathcal{P}_-} = g(1 + \alpha A_s\mathcal{P}_0)/(1 - \alpha A_s\mathcal{P}_0) \quad (6)$$

$$R_{\mathcal{P}_{++}} = g(1 + \alpha^2 A_s\mathcal{P}_0)/(1 - \alpha^2 A_s\mathcal{P}_0). \quad (7)$$

From these equations, the initial polarization, \mathcal{P}_0 , g and the polarization inversion efficiency α were deduced, as shown in Table I. The inversion efficiency η for the populations between the two magnetic substates by one AFP were determined from the relationship between α

and η , $\alpha \approx 4 - 5\eta$. The relaxation time of the polarization T_1 was determined from the time spectrum of polarization. These parameters are given in Table I.

The extraction of the degree of alignment from the negative alignment section shown in Fig. 4 was performed as follows. The β -ray asymmetry of the three orientations shown in Fig. 4 was observed during the aligning process. The polarization was determined from the measured asymmetry and g , as shown in Fig. 8. The population of the magnetic substate at the first orientation is given by $[a_{-2}, a_{-1}, a_0, a_{+1}, a_{+2}] = [r(1 - \epsilon_1), r(1 + \epsilon_1), s(1 - \epsilon_2), s(1 + \epsilon_2), t]$. The parameters r , s , and t satisfy the relation $2r + 2s + t = 1$. ϵ_1 and ϵ_2 are the parameters describing incompleteness in the depolarization method for the two different frequencies. These two parameters yielded a small residual polarization at the pure alignment section. The polarization of the first orientation is given by $\mathcal{P}_{1st} = \frac{1}{2}\{r(\epsilon_1 - 3) + s(\epsilon_2 + 1) + 2t\}$. The population after the spin manipulation using the AFP method, for example, between $m = +2$ and $m = +1$ is given as a matrix:

$$\begin{pmatrix} 1 - \eta & \eta & 0 & 0 & 0 \\ \eta & 1 - \eta & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a_{+2} \\ a_{+1} \\ a_0 \\ a_{-1} \\ a_{-2} \end{pmatrix}, \quad (8)$$

where $\eta \approx 1$. The spin manipulation shown in Fig. 4-(2) can be described as the product of the matrices. Therefore, the population at each orientation is given by r , s , t , η , ϵ_1 , and ϵ_2 . The polarization at the second orientation is given by $\mathcal{P}_{2nd} \approx \frac{1}{2}\{4r\epsilon_1 + s(\epsilon_2 - 1) + t + (1 - \eta)(-6r(\epsilon_1 + 1) - s(\epsilon_2 - 5) + t)\}$ under the approximation, up to the first order, that $(1 - \eta) \ll 1$. The pure alignment is produced at the third orientation. The residual polarization is given by $\mathcal{P}_{3rd} \approx \frac{1}{2}\{4r\epsilon_1 + 2s\epsilon_2 + (1 - \eta)(-6r(\epsilon_1 + 1) - 3s(\epsilon_2 - 1) + 3t)\}$. η was determined from the measurement of $R_{\mathcal{P}^+}$, $R_{\mathcal{P}^-}$, and $R_{\mathcal{P}^{++}}$. Therefore, the number of free parameters is three by assuming $\epsilon_1 = \epsilon_2$ and giving the relation $2r + 2s + t = 1$. All the population parameters were determined from the polarization change of the three orientations.

The alignment in the third orientation can be calculated from the population parameters using $\mathcal{A} \approx \frac{1}{2}\{-2 + 8r + 2s + (1 - \eta)(-10r(\epsilon_1 + 1) - s(\epsilon_2 - 5) + 5t)\}$. This equation gives the alignment prior to the alignment section. In order to consider the alignment relaxation in the crystal, the alignments prior to and after the alignment section were determined




FIG. 8. Polarization change in the timing program for the spin-aligning process. The filled circles and the open squares are for the cycles A and B in Fig. 5, respectively. The beam was chopped and was stopped for the cycle at the time 0.

from the polarization change before and after the alignment section, respectively. Then the effective alignment and the relaxation time of the alignment were deduced. Using a different assumption that $\epsilon_1 = 10\epsilon_2$ or $\epsilon_1 = 0.1\epsilon_2$, the systematic uncertainty was estimated. The change in the alignment was less than the statistical uncertainty. The results for the spin manipulation are summarized in Table I.

B. Alignment correlation term

The alignment correlation term was obtained from the ratio of counts, $R(E) = N(E, d\mathcal{P}^+, \mathcal{A}^+)/N(E, d\mathcal{P}^-)$ at the positive and negative alignment sections. \mathcal{A} and $d\mathcal{P}$ are the alignment and the residual polarization at the alignment section, respectively. The signs given by the superscript in \mathcal{A}^\pm and $d\mathcal{P}^\pm$ are the alignment signs. The counts are proportional to the β -ray angular distribution as expressed by $N(E, d\mathcal{P}, \mathcal{A}) \propto B_0(E)[1 \pm (B_1(E)/B_0(E))d\mathcal{P} + (B_2(E)/B_0(E))\mathcal{A}]$,

TABLE I. Results of the spin manipulation. \mathcal{P}_0 is the initial polarization. $\mathcal{A}_{1/2}^\mp$ is the alignment, where subscripts 1 and 2 indicate the first and second halves of the timing program, respectively, and the superscript is the sign of the alignment. $\Delta\mathcal{A}_{1+2}$ is the sum of the absolute value of the alignments. T_1 and $T_{\mathcal{A}}$ are the relaxation times of the polarization and the alignment, respectively. α and η are the efficiencies of the polarization inversion and the population inversion between the two neighboring magnetic substates. ϵ is a parameter of the incompleteness of the depolarization.

	^8Li	^8B
\mathcal{P}_0 (%)	7.18 ± 0.10	5.42 ± 0.19
\mathcal{A}_1^+ (%)	$+3.96 \pm 0.20$	$+4.9 \pm 0.4$
\mathcal{A}_1^- (%)	-4.93 ± 0.20	-5.6 ± 0.4
\mathcal{A}_2^+ (%)	$+2.29 \pm 0.19$	$+3.9 \pm 0.4$
\mathcal{A}_2^- (%)	-1.91 ± 0.19	-3.2 ± 0.4
$\Delta\mathcal{A}_{1+2}$ (%)	13.1 ± 0.4	17.7 ± 0.8
T_1 (s)	13.0 ± 1.6	13 ± 4
$T_{\mathcal{A}}$ (s)	2.0 ± 0.7	2.2 ± 1.2
α (%)	-85.5 ± 0.3	-94.8 ± 0.9
η (%)	97.09 ± 0.07	98.95 ± 0.18
ϵ (10^{-3})	4.4 ± 0.2	-0.9 ± 0.4

where the upper and lower signs are for the top and bottom telescopes, respectively.

For the first half of the cycles A and B shown in Fig. 5, the counting ratio $R_1(E)$ is given by

$$R_1(E) = \frac{T^+ N(E, d\mathcal{P}_1^+, \mathcal{A}_1^+)}{T^- N(E, d\mathcal{P}_1^-, \mathcal{A}_1^-)}, \quad (9)$$

where the values with subscript 1, such as \mathcal{A}_1^+ , are for the first half. T^- and T^+ are the beam-current integral for the cycles A and B, respectively. The alignment correlation term was derived using the well-approximated formula as

$$R_1(E) \approx \frac{T^+}{T^-} \left\{ 1 \pm \frac{B_1(E)}{B_0(E)} d\mathcal{P}_1 + \frac{B_2(E)}{B_0(E)} \Delta\mathcal{A}_1 \right\}, \quad (10)$$

where the upper and lower signs are for the top and bottom telescopes, respectively. $d\mathcal{P}_1 = d\mathcal{P}_1^+ - d\mathcal{P}_1^-$ and $\Delta\mathcal{A}_1 = \mathcal{A}_1^+ - \mathcal{A}_1^-$. The ratio of T^+ and T^- caused a spurious β -ray

asymmetry in $R_1(E)$. The counting ratio $R_2(E)$ at the second half of the cycles A and B is given by

$$\begin{aligned} R_2(E) &= \frac{T^- N(E, d\mathcal{P}_2^+, \mathcal{A}_2^+)}{T^+ N(E, d\mathcal{P}_2^-, \mathcal{A}_2^-)} \\ &= \frac{T^-}{T^+} \left\{ 1 \pm \frac{B_1(E)}{B_0(E)} d\mathcal{P}_2 + \frac{B_2(E)}{B_0(E)} \Delta\mathcal{A}_2 \right\}, \end{aligned} \quad (11)$$

where the values with subscript 2, such as \mathcal{A}_2^\pm , are for the second half. In the double ratio $R_1(E)R_2(E)$, T^+ and T^- are canceled as

$$\begin{aligned} R_1(E)R_2(E) &= \frac{N(E, d\mathcal{P}_1^+, \mathcal{A}_1^+) N(E, d\mathcal{P}_2^+, \mathcal{A}_2^+)}{N(E, d\mathcal{P}_2^-, \mathcal{A}_2^-) N(E, d\mathcal{P}_1^-, \mathcal{A}_1^-)} \\ &\approx 1 \pm \frac{B_1(E)}{B_0(E)} d\mathcal{P}_{1+2} \\ &\quad + \frac{B_2(E)}{B_0(E)} \Delta\mathcal{A}_{1+2}, \end{aligned} \quad (12)$$

where $d\mathcal{P}_{1+2} = d\mathcal{P}_1 + d\mathcal{P}_2$ and $\Delta\mathcal{A}_{1+2} = \Delta\mathcal{A}_1 + \Delta\mathcal{A}_2$. The alignment correlation terms were extracted from the simple average of the double ratios $R_1(E)R_2(E)$ for the top and bottom telescopes so that the influence of the residual polarization was canceled.

C. Corrections

In the extraction procedure for the alignment correlation terms described above, the β -ray angular distribution for ${}^8\text{Li}$ is given by

$$W(E) \propto pE(E_0 - E) \left\{ 1 \mp \frac{1}{3} \mathcal{P} + \frac{B_2(E)}{B_0(E)} \mathcal{A} \right\}, \quad (13)$$

where the upper and lower signs are for telescopes with $\theta_{I\beta} = 0^\circ$ and 180° , respectively, instead of the following:

$$\begin{aligned} W(E, \theta_{I\beta}) &\propto pE(E_0 - E) \\ &\quad \left\{ 1 + \frac{B_1(E)}{B_0(E)} \frac{p}{E} \mathcal{P} P_1(\cos(\theta_{I\beta})) \right. \\ &\quad \left. + \frac{B_2(E)}{B_0(E)} \left(\frac{p}{E} \right)^2 \mathcal{A} P_2(\cos(\theta_{I\beta})) \right\}. \end{aligned} \quad (14)$$

The correction for the $P_1(\cos(\theta_{I\beta}))$, and $P_2(\cos(\theta_{I\beta}))$ is given in Sec. IV C 1. The corrections for (p/E) , $(p/E)^2$, and $B_1(E)/B_0(E)$ are given in Sec. IV C 2.

1. *Solid angle of β -ray telescope*

The polarization and alignment correlation terms in the β -ray angular distribution are proportional to the Legendre polynomials $P_1(\cos(\theta_{I\beta}))$ and $P_2(\cos(\theta_{I\beta}))$, i.e., the $\cos\theta_{I\beta}$ and $\frac{3}{2}(\cos^2\theta_{I\beta} - 1/3)$ terms, respectively. $R_1(E)R_2(E)$ in Eq. (12) includes $B_2(E)/B_0(E)$, so the $\frac{3}{2}(\cos^2\theta_{I\beta} - 1/3)$ contribution should be corrected. $\Delta\mathcal{A}_{1+2}$ was determined from the degree of polarization, so the $\cos\theta_{I\beta}$ contribution should be corrected. In order to take the finite solid angle of the detector into account, the detection efficiency as a function of E and $\theta_{I\beta}$ was simulated using the EGS4 code. The correction for the solid angle was evaluated by convoluting the simulated efficiency, as shown in Fig. 9.

2. *Higher order term in the polarization and alignment correlation terms*

The p/E term and the $B_1(E)/B_0(E)$ term in the polarization correlation term were assumed to be 1 and $\mp 1/3$, where the upper and lower signs are for ${}^8\text{Li}$ and ${}^8\text{B}$, respectively, when the polarization was determined from the β -ray asymmetry. The correction for the polarization is independent of energy, because the polarization was determined from the total count from 5 to 13 MeV. The correction for the p/E term was 0.9972 and 0.9973 for ${}^8\text{Li}$ and ${}^8\text{B}$, respectively. The $B_1(E)/B_0(E)$ term is given in [3] as

$$\begin{aligned} \frac{B_1(E)}{B_0(E)} = \mp \frac{1}{3} \left[1 + \frac{E}{3M_n} \left(\frac{1}{A} \pm \frac{b}{Ac} - \frac{d_1}{Ac} \right) \right. \\ \left. - \frac{\sqrt{21}}{4} \left\{ \pm \frac{f}{Ac} \frac{4E + E_0 + 4E^2/E_0}{3M_n} \right. \right. \\ \left. \left. - \frac{j_2}{A^2c} \frac{8E^2 - 5EE_0}{2M_n^2} \right\} \right]. \end{aligned} \quad (15)$$

To avoid the large systematic uncertainty from the j_2/A^2c term, the correction factor was evaluated using the product of the correction factor at $\frac{5}{8}E_0$ and the ratio of the value at $\frac{5}{8}E_0$ to the averaged value from 5 to 13 MeV. The ratio was determined, from the observed energy dependence of the polarization correlation term, to be 0.983 ± 0.007 for ${}^8\text{Li}$ and 1.013 ± 0.014 for ${}^8\text{B}$. The correction factor at $\frac{5}{8}E_0$ was self-consistently evaluated using iteration to be 0.98 ± 0.03 for ${}^8\text{Li}$ and 0.99 ± 0.03 for ${}^8\text{B}$ from the matrix elements b/Ac , d_1/Ac , f/Ac , and j_2/A^2c , which in the present study were determined from the alignment correlation terms and the β - α angular correlation terms. The uncertainty of this correction

included the uncertainty of the matrix elements and a 100% uncertainty of the higher order contribution from f/Ac , thus implying a severe evaluation. Accordingly, the correction factor for $B_1(E)/B_0(E)$ was 0.96 ± 0.03 for ${}^8\text{Li}$ and 1.00 ± 0.03 for ${}^8\text{B}$.

The $(p/E)^2$ term in the alignment correlation term is assumed to be 1 for the first-order analysis. The evaluated correction factor for the $(p/E)^2$ term is shown in Fig. 9.

3. *Detector response*

The observed alignment correlation term includes the contribution from the neighboring energy region to some extent because of the finite detector resolution and the low-energy tail component of the detector response, as shown in Fig. 7. The correction factor was evaluated self-consistently using the known detector response and the alignment correlation term, as shown in Fig. 9. Here, the alignment correlation term was approximated by a quadratic curve, $c_1E + c_2E^2$, with two parameters c_1 and c_2 . The correction factor for ${}^8\text{B}$ from 6 to 12 MeV was close to 1.0 because the alignment correlation term was almost constant and the influence of the different energy was small.

4. *Background*

The main backgrounds for ${}^8\text{Li}$ and ${}^8\text{B}$ below 4 MeV were ${}^{17}\text{F}(T_{1/2} = 64.5 \text{ s}, Q_{EC} = 2.76 \text{ MeV})$ and ${}^{15}\text{O}(T_{1/2} = 122 \text{ s}, Q_{EC} = 2.75 \text{ MeV})$, respectively. The correction for the background is also shown in Fig. 9. The systematic uncertainty in the alignment correlation term was estimated by assuming 20% ambiguity in the background fraction.

D. **Systematic uncertainties**

In this subsection, the systematic uncertainties of the alignment correlation term are described. They are also summarized in Table II.

1. *Polarization relaxation and rank-three spin orientation*

The polarization and alignment were relaxed as a function of time. While the alignment relaxation during the alignment correlation term measurement was taken into account in the




FIG. 9. Energy dependent correction factors and total correction for ${}^8\text{Li}$ (upper) and ${}^8\text{B}$ (lower). The correction factors for the solid angle, C_{solid} , the detector response, C_{res} , the background, C_{BG} , and the $(p/E)^2$ term in the β -ray angular distribution, $C_{(p/E)^2}$, are shown. The total correction factor, C_{total} also includes the energy independent correction factors for the (p/E) and B_1/B_0 terms in the β -ray angular distribution.

procedure of the alignment extraction, the relaxation during conversion from the polarization to alignment was evaluated as a systematic uncertainty.

The β -ray asymmetry was caused by the odd rank of the spin orientation. The degree of the rank-three spin orientation was determined from the population parameters similar to the case of the degree of alignment, as described in Sec. IV A. The polarization was evaluated by taking the degree of the rank-three orientation into account. The effect on the polarization was considered as a systematic uncertainty.

2. *Uncertainty in the correction factor resulting from self-consistent evaluation*

The correction factor for the detector response was self-consistently evaluated using the results of the alignment correlation term. The energy dependence of the alignment correlation term was estimated as a quadratic curve without a constant term. The statistical uncertainty of the quadratic curve was propagated to the systematic uncertainty.

The correction factor for $B_1(E)/B_0(E)$ was evaluated using the matrix elements, such as d_1/Ac . The systematic uncertainty for this correction factor has been discussed in Sec. IV C.

3. *Relative positions of the beam, recoil catcher, and telescope*

The implanted recoil nuclei distributed widely on the catcher except for the part in the shadow part of the collimator. The relative positions of the beam spot and the catcher were able to change the distribution of the reaction products on the catcher. The beam spot was tuned using a fluorescent target with accuracy 0.5 mm and 1 mm in the horizontal and vertical directions, respectively. The relative position of the catcher and the ΔE detector had the potential to change the solid angle of the telescope and β -ray efficiency. The uncertainty of this relative position was 2 mm. The correction factors were evaluated by using the detector response simulated for different conditions using the EGS4 code. The change in the alignment correlation terms because of these two uncertainties was considered to be the systematic uncertainty.

4. *Detector response function*

The reliability of the low-energy tail in the simulated response function of a monoenergetic β -ray was evaluated. The low-energy tail was mainly caused by the energy loss straggling in the material between the positions of the β -ray emitter and E detector. The largest uncertainty was due to the catcher thickness. The relative uncertainty of the thickness was 10% for all the crystals.

The reliability of the simulated low-energy tail has been studied experimentally [18]. ^{12}B and ^{12}N were produced as emitters. The β -ray energy was selected via a dipole magnet. The shape and amount of low-energy tail were confirmed to within 20% statistical uncertainty.

The catcher thickness uncertainty of 10% and the simulation reliability of 20% for the low-energy tail were simulated simultaneously by varying the crystal thickness by 30% in the EGS4 simulation. The correction factors were evaluated by simulating the detector response at a crystal thickness varied by 30%. The systematic uncertainty in the alignment correlation terms was evaluated using these correction factors.

The detector resolution was determined from that reproducing the most complete experimental β -ray spectra of ^8Li and ^8B . The uncertainty of the counter resolution was 20%. The correction factors were evaluated using counter resolutions both the 20% larger and 20% smaller than the most probable resolution. The systematic uncertainty in the alignment correlation terms was evaluated using these correction factors.

5. *Energy scaling, gain fluctuation, and pileup*

The systematic uncertainty due to the energy-scale uncertainty δE was evaluated using $\frac{d}{dE}(B_2(E)/B_0(E))\delta E$. $B_2(E)/B_0(E)$ was given by the polynomial for E and E^2 , where the coefficients were determined by the χ^2 fit analysis.

The gain fluctuation was typically within 40 keV. The systematic uncertainty due to the gain fluctuation of the E detector was evaluated using the same procedure as that for the energy-scale uncertainty.

For a pileup event caused by two β rays, the obtained alignment correlation term is determined on the basis of the contributions of the two β rays at their respective energies. This effect was evaluated as a systematic uncertainty by integrating its contribution over the energy of two β rays.

V. RESULTS AND DISCUSSION

After the obtained alignment correlation terms are shown, the results given by Eq. (2) are compared with the CVC prediction. And then the weighted mean value of end-point energy over final-state distribution is described, which is used when the matrix elements are determined from the alignment correlation terms and the β - α correlation terms. Finally, the extraction of the weak magnetism and the second-forbidden term is described and these terms are compared with the CVC prediction.

TABLE II. Systematic uncertainties of the alignment correlation term at 9MeV.

	${}^8\text{Li}$	${}^8\text{B}$
	$\times 10^{-2}$	$\times 10^{-2}$
Polarization relaxation	0.002	0.003
Third-order orientation	0.015	0.026
Uncertainty of B_2/B_0 in detector-response correction	0.010	0.007
Uncertainty of matrix elements in B_1/B_0 correction	0.107	0.133
Position of beam spot and catcher	0.011	0.027
Position of ΔE detector and catcher	0.015	0.039
Low-energy tail of detector response	0.050	0.034
Detector resolution	0.021	0.017
Energy scaling	0.053	0.001
Gain fluctuation	0.028	< 0.001
Pileup	0.001	< 0.001
Background	< 0.001	< 0.001
Total	0.137	0.148

A. Alignment correlation terms and β - α correlation terms

The alignment correlation terms that were obtained are shown in Fig. 10. The statistical uncertainty of the alignment $\Delta\mathcal{A}_{1+2}$ in Eq. (12) could shift all data points of the alignment correlation term in the same direction. The statistical uncertainty of $\Delta\mathcal{A}_{1+2}$ is not included in each data point of Fig. 10 in order to retain a statistical fluctuation among the different points; however, the statistical uncertainties of the final results, such as a_{WM}^β , include the statistical uncertainty of $\Delta\mathcal{A}_{1+2}$.

The β - α correlation terms $-\frac{2}{3}p_\pm(E)$ [11] are also shown as crosses. The weak magnetism term, $-(3M_n/4E)(\delta_{\text{align.}}^- + \delta_{\beta\alpha}^-) = b/Ac$, was derived combining the two types of correlation term shown in Fig. 11. Figure 11 also reflects a reanalysis using the same energy bin as the β - α correlation terms [11].




FIG. 10. Alignment correlation terms and β - α correlation terms. The circles are the alignment correlation terms and the crosses are the β - α angular correlation terms. The open circles are not used for the derivation of the matrix elements. The lines are the best-fit curves.




FIG. 11. Weak magnetism term derived from the β -ray correlation terms. The CVC predictions are shown by the 1σ error band. The solid, dashed and dotted bands are the present, De Braeckeleer's [7], and Winter's [19, 20] predictions, respectively. Winter's prediction was re-evaluated using the mirror-averaged end-point energy.

B. CVC prediction

The experimental b/Ac results shown in Fig. 11 indicates a slight E dependence. The CVC prediction of an energy dependent b/Ac has in previous studies been indicated and described by introducing the dependence into the matrix elements, $b(E_x)$ and $c(E_x)$, of the final-state energy E_x in ${}^8\text{Be}$ [7, 21–23]. The final-state energy distributes widely because several states

with spin and parity of 2^+ are mixed because of the wide decay width. This final-state distribution can be formulated using the R -matrix theory with four final states [24, 25]. We re-evaluated the CVC prediction in Ref. [12] by using the analog- γ -decay measurement by De Braekeleer *et al.* [7], and the recent measurement of the β -delayed- α energy spectra from ^8Li and ^8B by Bhattacharya *et al.* [25]. The procedure for this re-evaluation was same as for the previous work [7] except for the number of final states; three final states were used in it, while four final states were used in the present evaluation, similar to that for the Gamow-Teller matrix element $c(E_x)$ in Ref. [25]. The procedure is summarized below.

The E_x dependence of $c(E_x)$ gives the final-state distribution for the β decay, i.e., the delayed α energy spectrum. The mirror-averaged $c(E_x)$ was determined from the delayed α energy spectra of ^8Li and ^8B based on the R -matrix formalism by Bhattacharya *et al.* [25].

$b(E_x)$ is given by the isovector M1 transition strength of the analog- γ decay, based on the strong CVC. The isobaric analog state in ^8Be was produced using the $^4\text{He}(\alpha, \gamma)$ reaction and the de-excited γ ray was measured [7]. The E_x dependence of $b(E_x)$ gives the final-state distribution in the analog- γ decay, which has been measured through the γ -ray energy spectra shown in Fig. 4 of Ref. [7]. The matrix elements, \mathcal{M}_1^γ and R_γ in $b(E_x)$, gives the E_x dependence of $b(E_x)$ and were determined using three final states in Ref. [7]. The \mathcal{M}_1^γ is the weak magnetism matrix element for the transition to the first excited state, and R_γ is the ratio, $\mathcal{M}_{16}^\gamma/\mathcal{M}_1^\gamma$, where \mathcal{M}_{16}^γ represents the transition to an isospin doublet at 16 MeV. These matrix elements were re-determined for the four final states so as to reproduce the γ -ray energy spectra, which were $\mathcal{M}_1^\gamma = 8.71 \pm 0.28$ and $R_\gamma = 1.5 \pm 1.4$.

The E dependent b/Ac is given in [7] by the weighted average as

$$\frac{b}{Ac} \rightarrow \frac{\int b(E_x)c(E_x)(E_{\max} - E_x - E)dE_x}{A \int c^2(E_x)(E_{\max} - E_x - E)dE_x}. \quad (16)$$

Here, $c(E_x)$ and E_{\max} averaged between the mirror pair were used, because b/Ac was derived from the β -ray angular correlations of both ^8Li and ^8B . The CVC prediction of b/Ac determined from Eq. (16) is shown in Fig. 11.

The CVC prediction determined by De Braekeleer *et al.* [7] and Winter *et al.* [19, 20] are also shown in Fig. 11. The previous predictions have a problem in regards to final-state treatment. The delayed- α spectra have been reproduced very well using four final states [24, 25]. De Braekeleer *et al.*, however, used three states for both $b(E_x)$ and $c(E_x)$, and Winter *et al.* used three states only for $b(E_x)$. The present CVC prediction was slightly

smaller than the previous predictions at a higher energy region.

The transition to the first excited state of ${}^8\text{Be}$ was predominant for the analog- γ transition [7]. Therefore, for a comparison between the b/Ac extracted from the β decay and its CVC prediction, the weak magnetism for the first excited state, a_{WM} , was used. This was given by the matrix elements of the transition to the first excited state, $a_{\text{WM}}^\beta = \mathcal{M}_1^\beta/A\mathcal{M}_{\text{GT1}}^\beta$ for β decay and $a_{\text{WM}}^{\text{CVC}} = \mathcal{M}_1^\gamma/A\mathcal{M}_{\text{GT1}}^\beta$ for the CVC prediction, where \mathcal{M}_1^β and $\mathcal{M}_{\text{GT1}}^\beta$ are the weak magnetism, b , and Gamow-Teller, c , matrix elements for the weak transition to the first excited state, respectively. The expression for b/Ac using \mathcal{M}_1 and \mathcal{M}_{GT1} was given in Ref. [7]. The CVC prediction was determined to be $a_{\text{WM}}^{\text{CVC}} = 7.3 \pm 0.2$ based on $\mathcal{M}_1^\gamma = 8.71 \pm 0.28$ and the mirror-averaged $\mathcal{M}_{\text{GT1}}^\beta = 0.1496 \pm 0.0005$ [25]. The CVC prediction of f was determined by the isovector M1/E2 ratio $\delta_1 = 0.01 \pm 0.03$ [7] as $a_{\text{WE2}}^{\text{CVC}} = \sqrt{10/3}\delta_1 a_{\text{WM}}^{\text{CVC}} = 0.1 \pm 0.4$. The values are summarized in Tables III and IV.

TABLE III. Decay widths and matrix elements for the γ decay from the isobaric analog state in ${}^8\text{Be}$. $\Gamma_{M1}^{T=1}$ is the decay width for the isovector component of the M1 transition from the isobaric analog state ($T = 1$). δ_1 is the isovector M1/E2 ratio from the isobaric analog state. Definition of \mathcal{M}_1^γ and R_1^γ is described in the text. Average value $\mathcal{M}_{\text{GT1}}^\beta$ of the Gamow-Teller matrix elements of ${}^8\text{Li}$ and ${}^8\text{B}$ is also shown.

Analog γ decay	Value	Matrix Element	Value
$\Gamma_{M1}^{T=1}$ [7]	2.80 ± 0.18 eV	\mathcal{M}_1^γ [7] ^a	8.7 ± 0.3
$\Gamma_{M1}^{T=1}$ [23] ^b	3.6 ± 0.3 eV	\mathcal{M}_1^γ [23] ^c	9.9 ± 0.6
$\Gamma_{M1}^{T=1}$ [22] ^b	4.1 ± 0.6 eV	\mathcal{M}_1^γ [22] ^c	10.5 ± 0.9
δ_1 [7]	0.01 ± 0.03	R_1^γ [7] ^a	1.5 ± 1.4
δ_1 [23] ^b	0.14 ± 0.03	$\mathcal{M}_{\text{GT1}}^\beta$ [25]	0.1496 ± 0.0005

^a Reanalyzed in the present work using the four final states in the R -matrix formalism.

^b Reanalyzed in Ref. [7].

^c Calculated from \mathcal{M}_1^γ of Refs. [7] and $\Gamma_{M1}^{T=1}$ of Ref. [7, 23] or Refs. [7, 22] using the relation $\mathcal{M}_1^\gamma \propto \sqrt{\Gamma_{M1}^{T=1}}$.




FIG. 12. Weighted mean value of the end-point energy over the broad-final-state distribution as a function of β -ray energy.

C. End-point energy

The end-point energy of the β ray is not a constant because of the broad final state. The alignment correlation terms and the β - α correlation terms given in Eq. (1) were measured as a function of β -ray energy without measurement of the end-point energy. Therefore, the end-point energy was averaged over the final-state-energy distribution. The weight is the product of $pE(E_0(E_x) - E)^2$ and the final-state distribution $c^2(E_x)$ of the β decay. When a certain β -ray energy is chosen, the weighted mean value of the end-point energy is given by

$$\begin{aligned} \bar{E}_0(E) &= \frac{\int pE(E_0(E_x) - E)^2 c^2(E_x) E_0(E_x) dE_x}{\int pE(E_0(E_x) - E)^2 c^2(E_x) dE_x} \\ &= \frac{\int (E_0(E_x) - E)^2 c^2(E_x) E_0(E_x) dE_x}{\int (E_0(E_x) - E)^2 c^2(E_x) dE_x}, \end{aligned} \quad (17)$$

where $E_0(E_x) = E_{\max} - E_x$ and the integral range is from 0 to $(E_{\max} - E)$. Figure 12 shows $\bar{E}_0(E)$ calculated using $c(E_x)$ determined in Ref. [25]. This $\bar{E}_0(E)$ was used in the analysis to determine the matrix elements.

D. Weak magnetism and second-forbidden terms from the weak vector current

The mirror difference δ^- consists of b/Ac and a small contribution of j_2/A^2c due to the mirror asymmetry of E_0 . To avoid the influence of this mirror asymmetry, the χ^2 fit analysis was performed simultaneously on the four correlation terms, i.e., both alignment correlation terms and β - α angular correlation terms of ${}^8\text{Li}$ and ${}^8\text{B}$. The E_x dependent b/Ac in the

β -ray angular correlation terms was given by the same formula as the CVC prediction of b/Ac . $a_{\text{WM}}^\beta = \mathcal{M}_1^\beta/A\mathcal{M}_{\text{GT1}}^\beta$ was used as a free parameter for the χ^2 fit analysis, where R_β was assumed to be the same as R_γ . The E dependences of a_{WE2}^β , d_1/Ac , j_2/A^2c , and j_3/A^2c were not clearly seen in the β -ray correlation terms because of the relatively large statistical uncertainties. These terms were assumed to be constant and were chosen as free parameters for the χ^2 fit analysis. The obtained terms were considered as the value averaged over the analyzed energy region. The best-fit curves are shown in Fig. 10 and the results are summarized in Table IV. The weak magnetism and the second-forbidden terms were $a_{\text{WM}}^\beta = 7.54 \pm 0.12(\text{stat.}) \pm 0.15(\text{syst.})$ and $a_{\text{WE2}}^\beta = 1.0 \pm 0.2(\text{stat.}) \pm 0.2(\text{syst.})$, respectively. The systematic uncertainty because of the E dependence of a_{WE2}^β was estimated to be 0.05 for a_{WE2}^β by assuming that the E_x dependence of $f(E_x)$ was the same as $b(E_x)$. The other systematic uncertainties in the alignment correlation terms and the β - α correlation terms were independently propagated to those in a_{WM}^β and a_{WE2}^β by performing the χ^2 fit analysis for the data applied to the different correction factors. a_{WM}^β was consistent with the CVC prediction from De Braeckeleer's data, i.e., $a_{\text{WM}}^\beta/a_{\text{WM}}^{\text{CVC}} = 1.03 \pm 0.04$. However, the present a_{WE2}^β is inconsistent with the De Braeckeleer's data, $a_{\text{WE2}}^{\text{CVC}} = 0.1 \pm 0.4$. The deviation of a_{WE2}^β was 1.8σ as $a_{\text{WE2}}^\beta - a_{\text{WE2}}^{\text{CVC}} = 0.9 \pm 0.5$.

TABLE IV. Ratio of matrix elements contributing to the β -ray angular correlations. The CVC predictions are also shown. $a_{\text{WM}}^{\text{CVC}} = \mathcal{M}_1^\gamma/A\mathcal{M}_{\text{GT1}}^\beta$ and $a_{\text{WE2}}^{\text{CVC}} = \sqrt{10/3}\delta_1 a_{\text{WM}}^{\text{CVC}}$.

Matrix Element	Value	Matrix Element	Value
a_{WM}^β	7.5 ± 0.2	$a_{\text{WM}}^{\text{CVC}}$ [7]	7.3 ± 0.2
a_{WE2}^β	1.0 ± 0.3	$a_{\text{WM}}^{\text{CVC}}$ [23]	8.3 ± 0.5
d_1/Ac	5.5 ± 1.7	$a_{\text{WM}}^{\text{CVC}}$ [22]	8.8 ± 0.7
j_2/A^2c	-490 ± 70	$a_{\text{WE2}}^{\text{CVC}}$ [7]	0.1 ± 0.4
j_3/A^2c	-980 ± 280	$a_{\text{WE2}}^{\text{CVC}}$ [23]	2.1 ± 0.5

We compared these results with the other analog- γ -decay measurements by Bowles and Garvey [23] and Paul *et al.* [22]. The CVC prediction was $a_{\text{WM}}^{\text{CVC}} = 8.3 \pm 0.5$ and $a_{\text{WE2}}^{\text{CVC}} = 2.1 \pm 0.5$ for Bowles' data, and $a_{\text{WM}}^{\text{CVC}} = 8.8 \pm 0.7$ for Paul's data. These predictions were inconsistent with the De Braeckeleer's data and also with the present β -decay results; that is, both $a_{\text{WM}}^{\text{CVC}}$ were larger than a_{WM}^β , and the deviation of a_{WE2}^β was 1.8σ , as -1.1 ± 0.6 . It

was pointed out by De Braeckeleer *et al.* [7] that there were problems in these measurements in regards to the absolute cross section, the photon angular distribution and the neutron background. The difference between the two a_{WE2}^{CVC} was due to deviation of δ_1 , i.e., 0.01 ± 0.03 [7] compared to 0.14 ± 0.03 [7, 23]. This deviation was determined via the relatively difficult measurement of the photon angular distribution. The inconsistency might be due to an underestimated background for the photon angular distribution. Although the CVC prediction by De Braeckeleer *et al.* was adopted in the present work, De Braeckeleer's data need to be confirmed with more accurate measurements.

VI. SUMMARY

The nuclear-spin-aligned nuclei ${}^8\text{Li}$ and ${}^8\text{B}$ were produced from spin-polarized nuclei using the β -NMR technique to test the strong CVC at a zero momentum transfer limit. The strong CVC could be tested for the second-forbidden transition for the first time. The alignment correlation terms for the β -ray angular distribution were determined using both positively and negatively aligned nuclei. The weak magnetism and the second-forbidden terms originating from the weak vector current were determined by combining the present alignment correlation terms and the previously known β - α angular correlation terms. The CVC predictions of the weak magnetism and the second-forbidden terms were re-evaluated using the most precise data set of the analog- γ decay in ${}^8\text{Be}$. Although the weak magnetism term was consistent with the CVC prediction obtained from the isovector-M1-transition strength, the second-forbidden term was inconsistent with that from the isovector-E2-transition strength. For more reliable tests for the second-forbidden transition, the CVC predictions need to be confirmed by more accurate measurements especially with regard to the isovector M1/E2 ratio δ_1 .

ACKNOWLEDGMENTS

This work was supported by KAKENHI (21740189).

[1] See e.g., I.S. Towner and J.C. Hardy, in *Symmetries and Fundamental Interactions in Nuclei*,

- edited by W.C. Haxton and E.M. Henley (World Scientific, Singapore, 1995), nucl-th/9504015.
- [2] J.C. Hardy and I.S. Towner, Phys. Rev. C **79**, 055502 (2009).
 - [3] B.R. Holstein, Rev. Mod. Phys. **46**, 789 (1974).
 - [4] L. Grenacs, Ann. Rev. Nucl. Part. Sci. **35**, 455 (1985).
 - [5] Y. Lee, L. Mo and C. Wu, Phys. Rev. Lett. **10**, 253 (1963).
 - [6] K. Minamisono, K. Matsuta, T. Minamisono, T. Yamaguchi, T. Sumikama, T. Nagatomo, M. Ogura, T. Iwakoshi, M. Fukuda, M. Mihara, K. Koshigiri, and M. Morita, Phys. Rev. C **65**, 015501 (2002).
 - [7] L. De Braekeleer, E. G. Adelberger, J. H. Gundlach, M. Kaplan, D. Markoff, A. M. Nathan, W. Schieff, K. A. Snover, D. W. Storm, K. B. Swartz, D. Wright, and B. A. Brown, Phys. Rev. C **51**, 2778 (1995).
 - [8] K. Minamisono, et al., Phys. Rev. C (to be submitted).
 - [9] Y.G. Abov *et al.*, Nucl. Phys. **34**, 505 (1962); H. Eichner *et al.*, Z. Naturforsch. **21a**, 908 (1966).
 - [10] R.E. Tribble and G.T. Garvey, Phys. Rev. C **12**, 967 (1975).
 - [11] R.D. McKeown, G.T. Garvey, and C.A. Gagliardi, Phys. Rev. C **22**, 738 (1980).
 - [12] T. Sumikama, K. Matsuta, T. Nagatomo, M. Ogura, T. Iwakoshi, Y. Nakashima, H. Fujiwara, M. Fukuda, M. Mihara, K. Minamisono, T. Yamaguchi, and T. Minamisono, Phys. Lett. B **664**, 235 (2008).
 - [13] A. Abragam, *Principles of Nuclear Magnetism* (Oxford, New York, 1983).
 - [14] T. Ohtsubo, Y. Nakayama, I. Minami, M. Tanigaki, S. Fukuda, A. Kitagawa, M. Fukuda, K. Matsuta, Y. Nojiri, H. Akai, and T. Minamisono, Hyperfine Interact. **80**, 1051 (1993).
 - [15] T. Sumikama, M. Ogura, Y. Nakashima, T. Iwakoshi, M. Mihara, M. Fukuda, K. Matsuta, T. Minamisono, and H. Akai, Hyperfine Interact. **158**, 413 (2004).
 - [16] T. Sumikama, T. Nagatomo, M. Ogura, T. Iwakoshi, Y. Nakashima, H. Fujiwara, K. Matsuta, T. Minamisono, M. Fukuda, and M. Mihara, Phys. Rev. C **74**, 024327 (2006).
 - [17] W.R. Nelson and H. Hirayama and D.W.O. Rogers, Stanford University Report, SLAC-265; D.W.O. Rogers, Nucl. Instr. and Meth. **A227**, 535 (1984).
 - [18] K. Tanaka, M. Fukuda, M. Mihara, K. Kidera, A. Takemura, H. Kenzaki, K. Matsuta, and T. Minamisono, OULNS Ann. Rep. , p. 38 (1998).
 - [19] W. T. Winter, S. J. Freedman, K. E. Rehm, I. Ahmad, J. P. Greene, A. Heinz, D. Henderson, R. V. F. Janssens, C. L. Jiang, E. F. Moore, G. Mukherjee, R. C. Pardo, T. Pennington,

- G. Savard, J. P. Schiffer, D. Seweryniak, G. Zinkann, and M. Paul, Phys. Rev. Lett. **91**, 252501 (2003).
- [20] W. T. Winter, S. J. Freedman, K. E. Rehm, and J. P. Schiffer, Phys. Rev. C **73**, 025503 (2006), nucl-ex/0406019.
- [21] A. M. Nathan, G. T. Garvey, P. Paul, and E. K. Warburton, Phys. Rev. Lett. **35**, 1137 (1975).
- [22] P. Paul, M. Suffert, and P. Gorodetzky, Phys. Lett. B **71**, 71 (1975).
- [23] T. J. Bowles and G. T. Garvey, Phys. Rev. C **18**, 1447 (1978).
- [24] E. K. Warburton, Phys. Rev. C **33**, 303 (1986).
- [25] M. Bhattacharya, E. G. Adelberger, and H. E. Swanson, Phys. Rev. C **73**, 055802 (2006).