

Ab-initio theory of NMR chemical shifts in solids and liquids

Francesco Mauri, Bernd G. Pfrommer, and Steven G. Louie

*Department of Physics, University of California at Berkeley, Berkeley, CA 94720, USA, and
Materials Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.*

(August 14, 2018)

Abstract

We present a theory for the *ab-initio* computation of NMR chemical shifts (σ) in condensed matter systems, using periodic boundary conditions. Our approach can be applied to periodic systems such as crystals, surfaces, or polymers and, with a super-cell technique, to non-periodic systems such as amorphous materials, liquids, or solids with defects. We have computed the hydrogen σ for a set of free molecules, for an ionic crystal, LiH, and for a H-bonded crystal, HF, using density functional theory in the local density approximation. The results are in excellent agreement with experimental data.

76.60.Cq, 71.15.Mb

arXiv:cond-mat/9611055v1 [cond-mat.mtrl-sci] 7 Nov 1996

Nuclear magnetic resonance (NMR) is one of the most widely used experimental techniques in structural chemistry. In particular, the chemical shift (σ) spectra are a fingerprint of the molecular geometry and the chemical structure of the material under study. Although the interpretation of these spectra generally relies on empirical rules, *ab-initio* calculations of σ for molecules have led in many cases to an unambiguous determination of the microscopic structure [1]. So far, *ab-initio* calculations of σ have been restricted to finite systems such as isolated molecules or clusters [2,3]. This is a serious limitation, because most of the NMR experiments are performed on liquid samples. Moreover, it is nowadays possible to measure σ also in solids with the resolution required for structural determinations [4,5]. E.g., σ -spectra have been used for the characterization of amorphous carbon [6]. In this letter, we present a formalism to compute, from first principles, σ in extended systems using periodic boundary conditions. Our approach can be applied to periodic systems like crystals, surfaces, or polymers and, using a super-cell technique, to non-periodic systems such as amorphous materials, liquids, or solids with defects. In the case of the amorphous solid or liquid, the atomic coordinates may be generated by *ab-initio* molecular dynamic simulations [7,8].

The chemical shift measures the local magnetic field in a sample induced by a uniform applied magnetic field. The computation of σ in an extended system is not straightforward, since the expectation values of the individual terms of the perturbative Hamiltonian for extended eigenstates are not well-defined quantities [9]. To overcome this problem, we follow Ref. [10], in which a theory for the computation of the macroscopic magnetic susceptibility is presented. In particular: i) we obtain the magnetic response to a uniform field as the long-wave limit of a periodic field; and ii) we use a generalized f-sum rule to remove the numerical instability which occurs in this limit. We apply the resulting equations to real systems, describing the electronic structure within density functional theory (DFT) in the local density approximation (LDA). We compute the hydrogen σ for a set of small molecules, for an ionic crystal, and for a H-bonded crystal. Our results are in excellent agreement with experimental data.

A uniform, external magnetic field \mathbf{B}_{ext} , applied to a sample induces an electronic current density $\mathbf{J}_{in}(\mathbf{r})$. This current produces an induced magnetic field $\mathbf{B}_{in}(\mathbf{r})$. If \mathbf{B}_{ext} is small enough, a condition realized in NMR experiments, then:

$$\mathbf{B}_{in}(\mathbf{r}) = -\overset{\leftrightarrow}{\sigma}(\mathbf{r})\mathbf{B}_{ext}. \quad (1)$$

Here $\overset{\leftrightarrow}{\sigma}(\mathbf{r})$ is the chemical shift tensor. With NMR spectroscopy, it is possible to measure the symmetric part of $\overset{\leftrightarrow}{\sigma}(\mathbf{r})$, or more often its trace, $\sigma(\mathbf{r}) = (1/3)\text{Tr}[\overset{\leftrightarrow}{\sigma}(\mathbf{r})]$, at the position of the non-zero spin nuclei.

In the bulk of a periodic system, $\overset{\leftrightarrow}{\sigma}(\mathbf{r})$ is also periodic. We may write:

$$\overset{\leftrightarrow}{\sigma}(\mathbf{r}) = \sum_{\mathbf{G}} \overset{\leftrightarrow}{\sigma}(\mathbf{G})e^{i\mathbf{G}\cdot\mathbf{r}}, \quad (2)$$

where \mathbf{G} are the reciprocal lattice vectors. For $\mathbf{G} \neq \mathbf{0}$, $\overset{\leftrightarrow}{\sigma}(\mathbf{G})$ is a bulk property:

$$\overset{\leftrightarrow}{\sigma}(\mathbf{G}) = -4\pi\overset{\leftrightarrow}{\chi}(\mathbf{G}, \mathbf{0}), \quad (3)$$

where $\overset{\leftrightarrow}{\chi}(\mathbf{G}, \mathbf{G}')$ is the magnetic susceptibility matrix. However, for $\mathbf{G} = \mathbf{0}$, $\overset{\leftrightarrow}{\sigma}(\mathbf{0})$ is *not* a bulk property. Its value depends on the shape of the sample, and is determined by macroscopic magnetostatic. In our calculations we assume a spherical sample, for which:

$$\overset{\leftrightarrow}{\sigma}(\mathbf{0}) = -\frac{8\pi}{3}\overset{\leftrightarrow}{\chi}(\mathbf{0}, \mathbf{0}), \quad (4)$$

where $\overset{\leftrightarrow}{\chi}(\mathbf{0}, \mathbf{0})$ is the macroscopic susceptibility [11]. Thus the calculation of σ in a periodic system requires the knowledge of $\overset{\leftrightarrow}{\chi}(\mathbf{G}, \mathbf{0})$. We compute the macroscopic susceptibility $\overset{\leftrightarrow}{\chi}(\mathbf{0}, \mathbf{0})$ following Ref. [10]. The other elements of $\overset{\leftrightarrow}{\chi}(\mathbf{G}, \mathbf{0})$ are computed as described below.

The susceptibility matrix is defined as the second derivative of the total energy with respect to the external magnetic field. In particular,

$$\hat{\mathbf{b}}_{-\mathbf{G}} \cdot \overset{\leftrightarrow}{\chi}(\mathbf{G}, \mathbf{0})\hat{\mathbf{b}}_{\mathbf{0}} = - \left. \frac{\partial^2 E[\mathbf{B}]}{\partial B_{\mathbf{0}} \partial B_{-\mathbf{G}}} \right|_{\mathbf{B}=\mathbf{0}} = -E''_{\mathbf{0}, -\mathbf{G}}, \quad (5)$$

where $E[\mathbf{B}]$ is the total energy of the system per unit volume in the external magnetic field $\mathbf{B}(\mathbf{r})$, $\mathbf{B}(\mathbf{r}) = [B_{\mathbf{0}}\hat{\mathbf{b}}_{\mathbf{0}} + B_{-\mathbf{G}}\hat{\mathbf{b}}_{-\mathbf{G}} \exp(-i\mathbf{G} \cdot \mathbf{r})]$, and $\hat{\mathbf{b}}_{-\mathbf{G}}$ and $\hat{\mathbf{b}}_{\mathbf{0}}$ are vectors of unit length.

Thus $\hat{\chi}(\mathbf{G}, \mathbf{0})$ can be evaluated using perturbation theory. However the expectation values of the perturbative Hamiltonians for a uniform field between extended eigenstates are ill defined. To avoid this problem, we modulate the external periodic field with a finite wave-vector \mathbf{q} , $\mathbf{B}(\mathbf{r}) = [B_{\mathbf{q}}\hat{\mathbf{b}}_{\mathbf{q}} \exp(i\mathbf{q} \cdot \mathbf{r}) + B_{-\mathbf{G}-\mathbf{q}}\hat{\mathbf{b}}_{-\mathbf{G}-\mathbf{q}} \exp(-i(\mathbf{G} + \mathbf{q}) \cdot \mathbf{r})]$ [12], and we recover the results for the uniform field by considering the limit [10]:

$$\hat{\mathbf{b}}_{-\mathbf{G}} \cdot \hat{\chi}(\mathbf{G}, \mathbf{0}) \hat{\mathbf{b}}_{\mathbf{0}} = - \lim_{q \rightarrow 0} E''_{\mathbf{q}, -\mathbf{G}-\mathbf{q}}. \quad (6)$$

We now consider a spin compensated system described by a single particle Hamiltonian. The derivatives of the Hamiltonian required to compute $E''_{\mathbf{q}, -\mathbf{G}-\mathbf{q}}$ are:

$$\begin{aligned} H'_{\mathbf{q}} &= \frac{-i}{2} \left(e^{i\mathbf{q} \cdot \mathbf{r}} \mathbf{a}_{\mathbf{q}} \cdot \nabla + \mathbf{a}_{\mathbf{q}} \cdot \nabla e^{i\mathbf{q} \cdot \mathbf{r}} \right), \\ H'_{-\mathbf{G}-\mathbf{q}} &= \frac{-i}{2} \left(e^{-i(\mathbf{G}+\mathbf{q}) \cdot \mathbf{r}} \mathbf{a}_{-\mathbf{G}-\mathbf{q}} \cdot \nabla \right. \\ &\quad \left. + \mathbf{a}_{-\mathbf{G}-\mathbf{q}} \cdot \nabla e^{-i(\mathbf{G}+\mathbf{q}) \cdot \mathbf{r}} \right), \\ H''_{\mathbf{q}, -\mathbf{G}-\mathbf{q}} &= \mathbf{a}_{\mathbf{q}} \cdot \mathbf{a}_{-\mathbf{G}-\mathbf{q}} e^{-i\mathbf{G} \cdot \mathbf{r}}, \end{aligned} \quad (7)$$

where $\mathbf{a}_{\mathbf{s}} = (i\mathbf{s} \times \hat{\mathbf{b}}_{\mathbf{s}})/(cs^2)$, and c is the speed of light. Using perturbation theory we obtain:

$$\begin{aligned} E''_{\mathbf{q}, -\mathbf{G}-\mathbf{q}} &= f(\mathbf{a}_{\mathbf{q}}, \mathbf{a}_{-\mathbf{G}-\mathbf{q}}, \mathbf{G}, \mathbf{q}) \\ &\quad + 2\mathbf{a}_{\mathbf{q}} \cdot \mathbf{a}_{-\mathbf{G}-\mathbf{q}} \int \frac{d^3k}{(2\pi)^3} \sum_{i \in \mathcal{O}} \langle u_{\mathbf{k}, i} | e^{-i\mathbf{G} \cdot \mathbf{r}} | u_{\mathbf{k}, i} \rangle, \end{aligned} \quad (8)$$

where $|u_{\mathbf{k}, i}\rangle$ is the periodic part of the Bloch eigenstate of the unperturbed Hamiltonian $H_{\mathbf{k}}$ with eigenvalue $\epsilon_{\mathbf{k}, i}$, \mathcal{O} are the sets of occupied bands, and

$$\begin{aligned} f(\mathbf{a}_1, \mathbf{a}_2, \mathbf{G}, \mathbf{q}) &= \int d^3r [h(\mathbf{a}_1, \mathbf{a}_2, \mathbf{r}, \mathbf{q}) \\ &\quad + h^*(\mathbf{a}_1^*, \mathbf{a}_2^*, \mathbf{r}, -\mathbf{q})] e^{-i\mathbf{G} \cdot \mathbf{r}}, \end{aligned} \quad (9)$$

where the integral is performed in the periodic cell, with

$$\begin{aligned} h(\mathbf{a}_1, \mathbf{a}_2, \mathbf{r}, \mathbf{q}) &= \int \frac{d^3k}{(2\pi)^3} \sum_{i \in \mathcal{O}} \\ &\quad [\langle u_{\mathbf{k}, i} | \mathbf{r} \rangle \langle \mathbf{r} | \mathbf{a}_2 \cdot (-i\nabla + \mathbf{k} + \mathbf{q}) | u_{\mathbf{k}, i}^{\mathbf{q}, \mathbf{a}_1} \rangle \\ &\quad + \langle u_{\mathbf{k}, i} | \mathbf{a}_2 \cdot (-i\nabla + \mathbf{k}) | \mathbf{r} \rangle \langle \mathbf{r} | u_{\mathbf{k}, i}^{\mathbf{q}, \mathbf{a}_1} \rangle]. \end{aligned} \quad (10)$$

$|u_{\mathbf{k},i}^{\mathbf{q},\mathbf{a}_1}\rangle$ is the first order change of the eigenstate $|u_{\mathbf{k},i}\rangle$ due to a field with wavevector \mathbf{q} . It can be obtained by solving the linear system:

$$(\epsilon_{\mathbf{k},i} - H_{\mathbf{k}+\mathbf{q}})|u_{\mathbf{k},i}^{\mathbf{q},\mathbf{a}_1}\rangle = Q_{\mathbf{k}+\mathbf{q}}\mathbf{a}_1 \cdot \left(-i\nabla + \mathbf{k} + \frac{\mathbf{q}}{2}\right)|u_{\mathbf{k},i}\rangle, \quad (11)$$

where $Q_{\mathbf{k}+\mathbf{q}} = (\mathbf{1} - \sum_{i \in \mathcal{O}} |u_{\mathbf{k}+\mathbf{q},i}\rangle\langle u_{\mathbf{k}+\mathbf{q},i}|)$ is the projector onto the empty subspace.

The first term on the r.h.s. of Eq. (8) is obtained as second order perturbation with the first order derivatives of the Hamiltonian H'_q and $H'_{-\mathbf{G}-\mathbf{q}}$. The second term in the r.h.s. of Eq. (8) is obtained as first order perturbation with the second order derivative of the Hamiltonian $H''_{\mathbf{q},-\mathbf{G}-\mathbf{q}}$. Since $\mathbf{a}_{\mathbf{q}}$ diverges as $1/q$ for $q \rightarrow 0$, the two terms on the r.h.s. of Eq. (8) individually diverge as $1/q$. To remove this divergence, which would produce a numerical instability, we use the generalized f-sum rule:

$$2\mathbf{a}_1 \cdot \mathbf{a}_2 \int \frac{d^3k}{(2\pi)^3} \sum_{i \in \mathcal{O}} \langle u_{\mathbf{k},i} | e^{-i\mathbf{G}\cdot\mathbf{r}} | u_{\mathbf{k},i} \rangle = -f(\mathbf{a}_1, \mathbf{a}_2, \mathbf{G}, \mathbf{0}). \quad (12)$$

Substituting the f-sum rule into Eq. (8) we obtain: $E''_{\mathbf{q},-\mathbf{G}-\mathbf{q}} = [f(\mathbf{a}_{\mathbf{q}}, \mathbf{a}_{-\mathbf{G}-\mathbf{q}}, \mathbf{G}, \mathbf{q}) - f(\mathbf{a}_{\mathbf{q}}, \mathbf{a}_{-\mathbf{G}-\mathbf{q}}, \mathbf{G}, \mathbf{0})]$. Then, for $\mathbf{G} \neq \mathbf{0}$,

$$\begin{aligned} \hat{\mathbf{b}}_{-\mathbf{G}} \cdot \overleftrightarrow{\chi}(\mathbf{G}, \mathbf{0}) \hat{\mathbf{b}}_0 &= -\lim_{q \rightarrow 0} E''_{\mathbf{q},-\mathbf{G}-\mathbf{q}} \\ &= -\frac{\partial}{c^2 G^2 \partial q} f(\hat{\mathbf{q}} \times \hat{\mathbf{b}}_0, \mathbf{G} \times \hat{\mathbf{b}}_{-\mathbf{G}}, \mathbf{G}, q\hat{\mathbf{q}}) \Big|_{q=0}, \end{aligned} \quad (13)$$

where $\hat{\mathbf{q}}$ is the unit vector in the direction of \mathbf{q} . Finally, the derivative with respect to q in Eq. (13) can also be evaluated using the following limit:

$$\begin{aligned} \hat{\mathbf{b}}_{-\mathbf{G}} \cdot \overleftrightarrow{\chi}(\mathbf{G}, \mathbf{0}) \hat{\mathbf{b}}_0 &= -\lim_{q \rightarrow 0} [f(\hat{\mathbf{q}} \times \hat{\mathbf{b}}_0, \mathbf{G} \times \hat{\mathbf{b}}_{-\mathbf{G}}, \mathbf{G}, q\hat{\mathbf{q}}) \\ &\quad - f(\hat{\mathbf{q}} \times \hat{\mathbf{b}}_0, \mathbf{G} \times \hat{\mathbf{b}}_{-\mathbf{G}}, \mathbf{G}, -q\hat{\mathbf{q}})] / (2qc^2G^2). \end{aligned} \quad (14)$$

Note that, for $\mathbf{G} \neq \mathbf{0}$, $\overleftrightarrow{\chi}(\mathbf{G}, \mathbf{0})$ is proportional to the first derivative of f with respect to q , whereas the macroscopic susceptibility $\overleftrightarrow{\chi}(\mathbf{0}, \mathbf{0})$ is proportional to the second derivative of f with respect to q [10].

In practice, we evaluate numerically $\overleftrightarrow{\chi}(\mathbf{G}, \mathbf{0})$ using Eq. (14) with a small, but finite q , and the \mathbf{k} -integral in Eq. (10) with a finite summation in the irreducible wedge of the Brillouin zone.

We describe the electrons using DFT-LDA; i.e., we neglect any explicit dependence of the exchange-correlation functional (E_{xc}) on the current density. The current dependence of E_{xc} could be taken into account using the approximate functional proposed in Ref. [13], but in practice, this produces only negligible corrections to σ in real systems [3]. An *ad-hoc* procedure to include many-body effects beyond DFT in the calculation of σ has been proposed in [14]. While this approach improves over DFT in small molecules, the corrections to DFT vanish for periodic systems, where the eigenstates are always extended. In general, to compute the second order variation in the DFT total energy with respect to an external perturbation, one should take into account the linear variation of the Hamiltonian induced by the linear variation of the charge $\delta\rho$. However, if the perturbation is a magnetic field, $\delta\rho$ is zero by time reversal symmetry. Thus Eq.s (7-14) are correct within DFT.

In the present calculation, we will consider the magnetic response of valence electrons only. We describe the ionic cores by norm conserving pseudopotentials [15] in the Kleinman-Bylander form [16]. This approximation does not affect σ of the nuclei without core electrons, such as H [2,17], but those containing core electrons. In the latter case σ computed with pseudopotentials differs from the one computed with an all-electron scheme by three different terms: i) the diamagnetic core contribution, which is independent of the chemical environment; ii) a contribution due to the transitions from valence states to core states [10]; iii) a contribution due to the difference between the all-electron valence wavefunctions and the pseudo wavefunctions in the core region. We found that, for first row atoms like carbon, the error due to the pseudopotential is minor, since the terms ii) and iii) are usually much smaller than the range of variation of σ with the chemical environment. In the present paper, however, we present only the results for σ of H, which is not affected by the use of pseudopotentials. Finally, in our pseudopotential calculation, we replace $(-i\nabla + \mathbf{k} + \mathbf{q}/2)$ in Eq. (11) with $(\mathbf{v}_{\mathbf{k}}^p + \mathbf{v}_{\mathbf{k}+\mathbf{q}}^p)/2$, where $\mathbf{v}_{\mathbf{k}}^p = (d/d\mathbf{k})H_{\mathbf{k}}^p$ is the velocity operator, and $H_{\mathbf{k}}^p$ is the pseudo-Hamiltonian [18].

We expand the wave-functions using a plane wave (PW) basis set. We obtain $|u_{\mathbf{k},i}^{\mathbf{q},\mathbf{a}_1}\rangle$ by solving Eq. (11) with a conjugate gradient minimization [19]; i.e., we never use the

unoccupied eigenstates of $H_{\mathbf{k}}^p$. We obtain $f(\mathbf{a}_1, \mathbf{a}_2, \mathbf{G}, \mathbf{q})$ from $|u_{\mathbf{k},i}^{\mathbf{q},\mathbf{a}_1}\rangle$ using fast Fourier transforms. Thus, the computational effort to evaluate σ for all nuclei in the sample is comparable to that required for a computation of the total energy [20]. We compute σ in systems containing atoms of H, C, Li, and F. For H we use a local pseudopotential to speed up the convergence with respect to the PW basis [21]. For C, Li and F atoms we use a pseudopotential with a non-local s-projector, to describe the interaction of valence states with the 1s core states. We use a PW energy cut-off of 100 Ry for the F atom, and of 70 Ry for the other atoms. Molecular and crystalline structures are taken from experiments [22]. The results for the free molecules are obtained using super-cells, such that the error on σ due to interaction between a molecule and its periodic replica is smaller than 0.1 parts per million (ppm).

The results for a set of molecules are reported in Table I. All the computed values are in excellent agreement with experimental data. The discrepancies are larger for C_2H_4 and C_2H_2 , in which a double and a triple C-C bond exist, respectively. This could indicate a relatively larger inaccuracy of LDA in describing these types of bonds. Similar agreement with experiments for the hydrogen σ in molecules is found in previous calculations [2,3].

The results for molecular and crystalline LiH are reported in Table II. LiH is an ionic system, which crystallizes in the rock-salt structure. The Li-H equilibrium distance in the crystal is significantly larger than that of the molecule. Our results show that the difference between σ in the molecule and in the solid is very small. To verify if σ is sensitive to the geometry, we consider a crystal at a pressure of 65 GPa, in which the Li-H distance is equal to that of the molecule. In this case we find a much larger value of σ . Thus, σ of the crystal and of the free molecule are only similar at the equilibrium geometries.

The results for molecular and crystalline HF are reported in Table III. The intra-molecular bond of HF is covalent. In the liquid and solid phases, the HF molecules are bonded together in zigzag chains via H-bonds. HF has the strongest H-bonding found in nature. This strong inter-molecular bond is reflected in the large variation of σ observed in the transition from the gas phase to the liquid phase, and in the large temperature depen-

dence of σ in the liquid phase [24]. The HF crystal is constituted of a 2D stacking of parallel 1D zigzag chains. We compute σ for the crystal at the experimental geometries measured for a DF crystal at two temperatures, 4.2K and 85K [25]. The computed σ in the free HF molecule is in outstanding agreement with experimental data. In the crystal, the computed σ decreases dramatically to a value close to the one measured in the liquid phase (we are not aware of any experimental measurement of σ for solid HF). Also the theoretical variation of σ with temperature, $\delta\sigma/\delta T = 0.013$ ppm/K, is very close to the one measured in the liquid, which is $d\sigma/dT = 0.0135$ ppm/K [24]. Thus, with respect to σ , the behavior of the solid is very similar to that of the liquid. Interestingly, a single H-bond does not account for the large decrease of σ observed in the condensed phases. Indeed the computed σ for the central H atom in a HF-HF dimer, is closer to the σ of the free molecule than to that of the liquid [2].

In conclusion, we have presented an *ab-initio* theory for the evaluation of NMR chemical shifts (σ) in extended systems. We have computed, for the first time, σ in real solids. Our results show that DFT-LDA predicts hydrogen σ which are in excellent agreement with experiment. The evaluation of σ requires the same numerical effort as the computation of the total energy [20]. Thus, our approach can be applied to super-cells containing up to a hundred atoms, which are sufficient to model liquids or amorphous materials [6–8].

We thank Prof. U. Haeberlen and Prof. M. Mehring for useful discussions. This work was supported by the National Science Foundation under Grant No. DMR-9520554, by the Office of Energy Research, Office of Basic Energy Sciences, Materials Sciences Division of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098, and by the Miller Institute for Basic Research in Science. Computer time was provided by the NSF at the NCSA in Illinois and by NERSC at LBNL.

REFERENCES

- [1] See, e. g., M. Schindler, J. Am. Chem. Soc. **109**, 1020 (1987).
- [2] W. Kutzelnigg, U. Fleischer and M. Schindler, in *NMR basic principles and progress* vol. 23, ed. by P. Diehl, *at al* pag. 165 (Springer-Verlag, Berlin ,1990).
- [3] A. M. Lee, N. C. Handy, and S. M. Colwell, J. Chem. Phys. **103**, 10095 (1995).
- [4] U. Haeberlen, in *Encyclopedia of NMR*, ed. by D. M. Grant and R. K. Harris, vol. 1, pag. 368 (John Wiley & Sons, England, 1996).
- [5] R. A. Wind, in *Modern NMR technique and their application in chemistry*, ed. by A. I. Popov and K. Hallenga, pag.125 (Marcel Dekker, New York, 1991).
- [6] C. Jäger, J. Gottwald, H. W. Spiess, and R. J. Newport Phys. Rev. B **50**, 846 (1994).
- [7] K. Laasonen, M. Sprik, M. Parrinello, R. Car, J. of Chem. Phys. **99**, 9080 (1993).
- [8] N. A. Marks, D. R. McKenzie, B. A. Pailthorpe, M. Bernasconi, M. Parrinello, Phys. Rev. Lett. **76**, 768 (1996).
- [9] K. Kobayashi, and M. Tsukada, Phys. Rev. B **38**, 8566 (1988); and ref. therein.
- [10] F. Mauri and S. G. Louie, Phys. Rev. Lett. **76**, 4246 (1996).
- [11] This macroscopic term is important for the determination of the hydrogen σ in condensed matter. E.g., in water, $4\pi\chi(\mathbf{0}, \mathbf{0}) = -9.2$ ppm, which is comparable to variation of σ with the chemical environment. We consider a spherical geometry for two reasons. Firstly, it is the most suitable to identify inter-molecular interactions, because σ measured in a spherical sample of liquid is equal to that of the isolated molecule, if the molecules in the liquid do not interact. Secondly, if the magic angle spinning (MAS) technique is used [5] for a solid with isotropic $\vec{\chi}(\mathbf{0}, \mathbf{0})$, σ is independent of the particular shape of the sample, and is equal to the one obtained without MAS but with a spherical shape.

- [12] Since $\nabla \cdot \mathbf{B}(\mathbf{r}) = 0$, $\mathbf{q} \cdot \hat{\mathbf{b}}_{\mathbf{q}} = 0$ and $(\mathbf{q} + \mathbf{G}) \cdot \hat{\mathbf{b}}_{-\mathbf{G}-\mathbf{q}} = 0$.
- [13] G. Vignale, M. Rasolt, Phys. Rev. Lett. **59**, 2360 (1987).
- [14] V. G. Malkin, O. L. Malkina, M. E. Casida, and D. R. Salahub, J. Am. Chem. Soc. **116**, 5898 (1995).
- [15] N. Troullier and J. L. Martins, Phys. Rev. B **43**, 1993 (1991).
- [16] L. Kleinman, and D. M. Bylander, Phys. Rev. Lett. **48**, 1425 (1982).
- [17] M. Kaupp, V. G. Malkin, O. L. Malkina, D. R. Salahub, Chem. Phys. Lett. **235**, 382 (1995).
- [18] Note that $\mathbf{v}_{\mathbf{k}}^p = (-i\nabla + \mathbf{k})$ if $H_{\mathbf{k}}^p$ is a local Hamiltonian. For a more detailed discussion see: F. Mauri, B. Pfommer, S. G. Louie, unpublished.
- [19] W. H. Press, A. A. Teukolsky, W. T. Vetterling and B. P. Flannery *Numerical Recipes*, pag. 77 and pag. 413 (Cambridge University Press, USA, 1992).
- [20] The computational time required for the evaluation of all σ in a unit cell scales with the system size as $O(N_{\mathcal{O}}^2 N_{pw})$, where $N_{\mathcal{O}}$ is the number of occupied states and N_{pw} is the number of PWs.
- [21] We verified that the values for H σ obtained with the H pseudopotential are rigidly shifted by -0.3 ppm with respect to the values obtained without the H pseudopotential. The results reported in the present paper have been corrected for this effect.
- [22] H₂: $r_{\text{HH}} = 0.767 \text{ \AA}$; CH₄: $r_{\text{CH}} = 1.094 \text{ \AA}$; C₂H₆: $r_{\text{CH}} = 1.096 \text{ \AA}$, $r_{\text{CC}} = 1.532 \text{ \AA}$, $\theta_{\text{CCH}} = 111.5^\circ$; C₂H₄: $r_{\text{CH}} = 1.103 \text{ \AA}$, $r_{\text{CC}} = 1.337 \text{ \AA}$, $\theta_{\text{CCH}} = 121.4^\circ$; C₂H₂: $r_{\text{CH}} = 1.061 \text{ \AA}$, $r_{\text{CC}} = 1.206 \text{ \AA}$; HF (molecule) $r_{\text{HF}} = 0.926 \text{ \AA}$.
- [23] W. T. Raynes, Nuclear Magnetic Resonance **7**, 1 (The Chemical Society, London, 1977).
- [24] σ relative to CH₄ are reported in W. G. Schneider, H. G. Bernstein, J. A. Pople, J.

Chem. Phys. **28**, 601 (1958). We use Ref. [23] to convert to the absolute scale.

[25] M. W. Johnson, E. Sándor, and E. Arzi, Acta Cryst. **B31**, 1998 (1975).

TABLES

	σ -theory (ppm)	σ -experiment (ppm)
H ₂	25.9	26.2 ^a
CH ₄	30.7	30.61 ^a
C ₂ H ₆	29.7	29.86 ^b
C ₂ H ₄	24.5	25.43 ^b
C ₂ H ₂	28.6	29.26 ^b

TABLE I. Computed and measured H chemical shifts (σ) for a set of free molecules.^a Ref. [23],^b Ref. [24].

		σ -theory (ppm)
LiH molecule	$d_0 = 1.595$	26.6
LiH crystal	$a = a_0 = 4.08$	26.3
LiH crystal	$a = 2d_0 = 3.19$	31.2

TABLE II. Computed H chemical shifts (σ) in LiH for a free molecule and a spherical crystalline sample (in the rock-salt structure). d_0 is the equilibrium bond-length of the molecule, a is the lattice constant, and a_0 is the experimental equilibrium lattice constant. For $a = 2d_0$, the shortest Li-H distance in the crystal is equal to d_0 .

	T (K)	σ -theory (ppm)	σ -experiment (ppm)
HF molecule		28.4	28.5 ^a
HF crystal	4.2	20.71	
HF crystal	85	21.76	
HF liquid	214		21.45 ^b

TABLE III. Computed and measured hydrogen chemical shifts σ in HF for a free molecule, a spherical crystalline sample, and a spherical liquid sample. T is the temperature. ^a Ref. [23], ^b Ref. [24].