

PDF issue: 2024-05-24

The IGF-1/Akt/S6 pathway and expressions of glycolytic myosin heavy chain isoforms are upregulated in chicken skeletal muscle during the first week after hatching

Saneyasu, Takaoki ; Tsuchihashi, Tatsuya ; Kitashiro, Ayana ; Tsuchii, Nami ; Kimura, Sayaka ; Honda, Kazuhisa ; Kamisoyama, Hiroshi

(Citation)

Animal Science Journal, 88(11):1779-1787

(Issue Date) 2017-11

(Resource Type) journal article

(Version) Accepted Manuscript

(Rights)

© 2017 Japanese Society of Animal Science. This is the peer reviewed version of the following article: [Animal Science Journal, 88(11):1779-1787, 2017], which has been published in final form at http://dx.doi.org/10.1111/asj.12847. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for...

(URL)

https://hdl.handle.net/20.500.14094/90004617



The IGF-1/Akt/S6 pathway and expressions of glycolytic myosin heavy chain isoforms are upregulated in chicken skeletal muscle during the first week after hatching.

Takaoki SANEYASU, Tatsuya TSUCHIHASHI, Ayana KITASHIRO, Nami TSUCHII, Sayaka KIMURA, Kazuhisa HONDA and Hiroshi KAMISOYAMA

Graduate School of Agricultural Science, Kobe University, Kobe, Japan

Running title: Protein synthesis in skeletal muscle of newly hatched chicks

Correspondence: Takaoki Saneyasu, PhD.,

Tel: +81 78 803 5808; Fax: +81 78 803 5809

Abstract

Skeletal muscle mass is an important trait in animal industry. We previously reported an age-dependent downregulation of the IGF-1/Akt/S6 pathway, major protein synthesis pathway, in chicken breast muscle after 1 week of age, despite a continuous increase of breast muscle weight. Myosin heavy chain (HC), a major protein in muscle fiber, has several isoforms depending on chicken skeletal muscle types. HC I (fast-twitch glycolytic type) is known to be expressed in adult chicken breast muscle. However, little is known about the changes in the expression levels of protein synthesis-related factors and HC isoforms in perihatching chicken muscle. In the present study, protein synthesis-related factors, such as IGF-1 mRNA levels, phosphorylation of Akt, and phosphorylated S6 content, increased in an age-dependent manner after post-hatch day (D) 0. The mRNA levels of HC I, III, and V (fast-twitch glycolytic type) dramatically increased after D0. The increase ratio of breast muscle weight was approximately 1,100% from D0 to D7. To our knowledge, these findings provide the first evidence that upregulation of protein synthesis pathway and transcription of fast twitch glycolytic HC isoforms play critical roles in the increase of chicken breast muscle weight during the first week after hatching.

Keywords: myosin heavy chain, perihatching chick, protein synthesis, skeletal muscle

1. Introduction

Broiler chickens have been genetically selected for growth rate and skeletal (especially breast) muscle weight (Arthur and Alber, 2003), resulting in larger number and diameter of muscle fiber than in layer chickens (Scheuermann et al., 2004). Since the total number of muscle fibers is set at hatching (Smith, 1963; Velleman, 2007), the posthatch growth of muscle is mainly attributable to muscle fiber enlargement (Velleman, 2007). For that to occur there must be an increase in protein synthesis (Velleman, 2007). Therefore, to clarify how the protein synthesis are regulated in posthatch skeletal muscle is important in poultry meat production industry.

The increase in muscle fiber size requires more nuclei for upregulated protein synthesis through the promotion of transcription and translation (Velleman, 2007). The additional nuclei are provided from satellite cells located underneath the basement membrane of the myofiber; satellite cells proliferate and then fuse with the existing myofibers (Mauro, 1961; Moss and LeBlond, 1971). Post-hatch day (D) 2 and 3 show the greatest number of satellite cells after which there is a subsequent decrease (Sobolewska et al., 2011), suggesting that the additional nuclei are actively provided to the existing myofibers and consequently, protein synthesis may be upregulated after D3 in chicken skeletal muscle. Lines of evidence suggest that the IGF-1-induced phosphorylation of Akt and ribosomal protein S6 (S6) are critical events in protein synthesis in mammals (Glass, 2005; Sandri, 2008; Schiaffino *et al.*, 2013; Shah *et al.*, 2000). Phosphorylated S6 (p-S6) can be used as a marker of translational efficiency in muscle fibers (Goodman *et al.*, 2012). Although IGF-1 is known to be synthesized in both the skeletal muscle and liver, previous studies using transgenic mice indicated that skeletal IGF-1 plays a major role in skeletal muscle growth (Sjögren et al., 1999; Yakar et al., 1999).

Previous studies in suckling rats (between 1 and 28 days postpartum) and chickens (between 1 and 6 weeks) showed that fractional protein synthesis rates decreased agedependently (Davis et al., 1989; Kang et al., 1985), although skeletal muscle weight continuously increased. In addition, we previously reported that IGF-1 expression, phosphorylation rate of Akt, and pS6 content decreased coordinately in chicken breast muscle from 1 to 7 weeks of age (Saneyasu *et al.*, 2016). These findings likely suggest the age-dependent downregulation of IGF-1/Akt/S6 pathway in chicken skeletal muscle form 1 to 7 week after hatching. However, it remains to be elucidated how the IGF-1/Akt/S6 pathway is regulated in neonatal chicken skeletal muscle.

Myosin heavy chain (HC) is a major component of muscle fiber, and several isoforms have been identified in chickens (Rushbrook *et al.*, 1997). The adult breast muscle (*pectoralis major*) consists of HC I only (Rushbrook *et al.*, 1997, 1998). In contrast, several HC isoforms were shown to be expressed during the embryonic period; HC VI, VII, and IV were the major isoforms on embryonic day (E) 12, E14, and E19, respectively (Rushbrook *et al.*, 1998). Furthermore, the post-hatch isoform HC V was reported to be expressed during D3 to D12 (Rushbrook *et al.*, 1998). However, it remains unclear how the expression levels of HC isoforms (especially HC I, the major isoform in adults) change in chicken breast muscle during the perihatching period.

In an effort to understand the changes in protein synthesis and expression of HC isoforms in neonatal chicken muscle, we investigated the mRNA and protein levels of the protein synthesis-related factors and HC isoforms in the breast muscle of perihatch chickens. Our results indicate that IGF-1-stimulating protein synthesis pathway and transcription of HC I and V are markedly upregulated after hatching.

2. Materials and Methods

Animals and sampling

Commercial broiler (Ross308) eggs were purchased from a local hatchery (Ishii Poultry Farming Cooperative Association, Tokushima, Japan) and incubated at 37.5°C. After hatching, the chicks were given free access to water and a commercial starter diet (23.5% crude protein and 3,050 kcal/kg metabolizable energy; Nippon Formula Feed Mfg. Co. Ltd., Kanagawa, Japan). Six embryos or chicks of each sex were sampled at E18, E20, D0, D3, D5, and D7. D0 (post-hatch) chicks were euthanized before feeding. Body weights were measured, and embryos/chicks were euthanized by decapitation. The yolk sac and *pectoralis major* muscle were excised and measured. The dissected muscles were immediately frozen in liquid nitrogen and stored at -80°C until further analyses. The experimental procedures involving animals were approved by the Institutional Animal Care and Use Committee and carried out according to the Kobe University Animal Experimental Regulation.

Sexing

The sex of chicks was determined as described by Fridolfsson and Ellegren (1999). Briefly, DNA was extracted from approximately 50 mg of *pectoralis major* muscle using Kaneka Easy DNA Extraction Kit for animal tissue (Kaneka Co., Osaka, Japan). The gene encoding chromo-helicase-DNA binding protein (CHD) was amplified using Kaneka High-Speed DNA polymerase (Kaneka Co., Osaka, Japan). PCR products were separated in 2% agarose gel including SYBR[®] safe Gel stain (Invitrogen, Carsbad, CA) and visualized in a transilluminator. Samples taken from males showed a single band (putatively 600 bp), whereas two different bands (putatively 450 bp and 600 bp) were observed in those taken from females.

Real-time PCR analysis

Total RNA was extracted from the muscles using Sepazol-RNA I (Nacalai Tesque, Inc., Kyoto, Japan). First-strand cDNA was synthesized from total RNA using ReverTra Ace[®] qPCR RT Master Mix with gDNA Remover (Toyobo Co. Ltd, Osaka, Japan). mRNA levels were analyzed using Thermal Cycler Dice[®] Real Time System (Takara Bio Inc., Otsu, Japan) and SYBR Premix Ex Taq II (Tli RNaseH Plus; Takara Bio Inc., Otsu, Japan) according to the manufacturer's recommendations; sequences of the gene-specific primers used during the study are shown in Table 1. The expression levels of target genes were normalized to those of ribosomal protein S17 (RPS17).

Western blot analysis

Frozen muscles were homogenized by polytron in lysis buffer described previously by Duchêne *et al.* (2008). The homogenate were centrifuged at 17,900 g for 15 min at 4°C and the supernatants were stored at -80°C. Protein concentrations were determined by the method of Lowry *et al.* (1951). Muscle lysates were subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting using the HorizeBlot (ATTO Co., Tokyo, Japan) according to the manufacturer's recommendations. Bands were detected using Chemi-Lumi one Super (Nacalai Tesque, Inc., Kyoto, Japan), visualized with the Lumicube (Liponics Inc., Tokyo, Japan) and quantified using CS Analyzer (ATTO Co., Tokyo, Japan). Anti-Akt (#9272), anti-p-Akt (S473) (#9271), anti-S6 (#2217), anti-p-S6 (S240/244) (#5364), anti- α/β -tubulin (#2148), and horseradish peroxidase-conjugated anti-rabbit IgG (#7074) were purchased from Cell Signaling Technology (Beverly, MA, USA). Band intensity of α/β -tubulin was used as a loading control.

Statistical analysis

Data were analyzed using the Tukey-Kramer method. All statistical analyses were performed using Excel 2013 (Microsoft, USA) with Statcel 3 add-in software (OMS, Tokyo, Japan).

3. Results

Figure 1 shows the changes in the weights of body without yolk sac, yolk sac, and *pectoralis major* muscle in male chicks. Body weight and weight of *pectoralis major* muscle increased in an age-dependent manner during the experimental period, whereas weight of yolk sac decreased in an age-dependent manner. A marked increase was observed in breast muscle weight from D3 to D5. Thus, we examined the changes in the expression levels of myogenic regulatory factors (MRFs), protein synthesis-related factors and HC isoforms in the breast muscle.

The MRF MyoD is expressed in the activated and proliferating satellite cells (Bentzinger *et al.*, 2012; Legerlotz and Smith, 2008; Sobolewska *et al.*, 2011), and activates the transcription of muscle-specific genes such as myogenin (Berkes et al., 2004) and HC (Beylkin *et al.*, 2006; Wheeler *et al.*, 1999). Myogenin is also categorized as a member of the MRFs and induces the fusion of myocytes (Nabeshima *et al.*, 1993; Bentzinger *et al.*, 2012). In the present study, MyoD mRNA levels increased in an age-dependent manner from E18 to D3 and maintained the highest level until D7 (Fig. 2A). In agreement with the results of previous studies (Halevy *et al.*, 2004; Sobolewska *et al.*, 2011),

myogenin expression peaked on D3 (Fig. 2B). These results likely suggest an increasing number of activated satellite cells (E18-D3) and their fusion with myofibers (at approximately D3).

The expression of breast muscle IGF-1 increased in an age-dependent manner after hatching (Fig. 3A). Phosphorylation of Akt and S6, and p-S6 contents increased in an age-dependent manner (Fig. 3C, D and E) in association with the changes in IGF-1 expression, although the total S6 contents, which is used as translational capacity (Goodman et al., 2012), did not change age-dependently (Fig. 3E). These results suggest that the IGF-1/Akt/S6 signaling pathway is upregulated in chicken breast muscle during the first week post-hatch.

Myostatin was reported to inhibit the IGF-1/Akt signaling pathway in mammals (Morissette *et al.*, 2009; Gumucio and Mendias, 2013). In chicken, the myostatin mRNA level decreased in an age-dependent manner before hatching and gradually increased after hatching in a manner similar to the IGF-1 mRNA levels (Fig. 2B), indicating the possibility that IGF-1 regulates myostatin expression in the skeletal muscle of newly hatched chicks.

Figure 4 shows the mRNA levels of HC and troponin I (Tn I) isoforms. The expression levels of the embryonic/perihatch isoform HC IV and the Tn I type 1 (slow type) decreased significantly in chicken breast muscle after D0 and E20, respectively (Fig. 3C and E). In contrast, significant increases were observed in the mRNA levels of the post-hatch isoform HC V, adult isoform HC I, Tn I type 2 (fast type), and ratio of Tn I type 2 to type 1 (Fig. 3A, D, F, and G). Unexpectedly, HC III mRNA level peaked on D3 (Fig. 3B); previous studies report that this isoform does not appear in the developing and adult *pectoralis major* muscle (Rushbrook *et al.*, 1997, 1998). To our knowledge, these findings

provide the first evidence that the expression of isoforms of HC and Tn I changes drastically in chicken breast muscle during the perihatching period.

The weights of body and tissue, and the levels of mRNA and proteins were examined in females as well as in males. They changed in females in a manner similar to those in males (Supplementary Figure S1, S2, S3, and S4).

4. Discussion

Early post-hatch fasting is shown to depress skeletal muscle growth in chickens, resulting in the lower body weight at marketing compared to those exposed to feed earlier (Gonzales *et al.*, 2003; Halevy *et al.*, 2000; Noy and Sklan, 1997). Previous studies in chickens and turkeys reported that mRNA/protein levels of myogenin, IGF-1, and p-Akt were lower in the breast muscle of unfed birds from D0 to D2 than in those of fed birds (Guernec *et al.*, 2004; Halevy *et al.*, 2003). In the present study, the upregulations of myogenin expression and IGF-1/Akt/S6 pathway were observed after feeding. In addition, oral injection of carboxymethylcellulose to neonatal chick failed to increase the IGF-1 mRNA level in the breast muscle (unpublished data). All these findings suggest that initial intake of exogenous nutrients except fiber is one of the triggers to stimulate post-hatch growth via upregulation of the fusion of activated satellite cells and IGF-1/Akt/S6 pathway in the skeletal muscle of newly hatched chicks.

In contrast to the results of a previous study (Saneyasu et al., 2016), IGF-1/Akt/S6 pathway is upregulated in chicken breast muscle during the first week after hatching. The reason for the different pattern of changes in the basal level of IGF-1/Akt/S6 pathway before and after 1 week of age in chicken skeletal muscle remains unclear. However, our present and previous studies shows that the increase ratio of the *pectoralis major* muscle

weight is the highest during the first week after hatching: 80% (E18 to D0), 1,100% (D0 to D7), 495% (D7 to D14), 404% (D14 to D28), and 171% (D28 to D49) (Fig. 1, 3, and Saneyasu et al., 2016). It is therefore reasonable to consider that the IGF-1/Akt/S6 is upregulated in chicken skeletal muscle from hatching to 1 week of age.

Myostatin has been demonstrated to be a specific target of MyoD in *in vitro* study using C2C12 cells (Spiller *et al.*, 2002). However, our data did not show any correlating changes in mRNA levels of MyoD and myostatin. Previously, an *in vitro* study on the differentiation of primary chicken myoblasts into the myotube showed that myostatin expression was enhanced by IGF-1 (Kurokawa *et al.*, 2009). Our data shows that changes in mRNA levels of myostatin and IGF-1 were coordinated after D0, in agreement with previous study (Guernec *et al.* 2004). It is therefore likely that IGF-1 regulates myostatin transcription in chicken muscle post-hatch.

The postnatal development of muscle fiber type has been studied histochemically in chicken pectoral muscle (Ashmore and Doerr, 1971). At hatching, a majority of these fibers have many large diformazan granules. Neutral lipid droplets are large and numerous inside fibers that have higher mitochondrial density (Ashmore and Doerr, 1971). By 4 days of age, mitochondria and lipid-rich muscle fibers are clearly in a minority, these fibers being transformed rapidly into mitochondria-poor fibers in which histologically visible lipid is absent (Ashmore and Doerr, 1971). However, the mechanism underlying this histochemical change in the muscle fiber after hatching remains unclear. Previous studies reported that HC I, III, and V appeared to be associated with fast-twitch glycolytic fiber, and that HC IV appeared to be associated with fast-twitch glycolytic fiber, and that HC IV appeared to be associated with fast-twitch fiber (Barnard *et al.*, 1982; Rushbrook *et al.*, 1997, 1998). Therefore, we provide the first evidence on the drastic change in gene expressions of HC and Tn I

isoforms from oxidative to glycolytic type in the breast muscle of perihatching chicks, and suggest that this change may underlie the histochemical changes in the muscle fiber during the first week after hatching.

It remains unclear how expression of HC isoforms is regulated during the perihatching period in chicken muscle. The present results show a coordinated increase in the mRNA levels of IGF-1 and glycolytic type HC post-hatch and a preceding increase in MyoD mRNA level during perihatching period. *In vitro* studies using the C2C12 myotubes have shown that IGF-1 and MyoD activate the type IIB (fast-twitch glycolytic type) HC promotor (Shanely et al., 2009; Wheeler et al., 1999). Thus, our findings raise the possibility that IGF-1 and/or MyoD promote transcription of HC I, III, and V in the skeletal muscle of newly hatched chicks.

In conclusion, we show that protein synthesis pathway and transcription of fast-twitch glycolytic type HC are upregulated in an age-dependent manner in chicken breast muscle during the first week after hatching. These findings can provide new insights for further research on skeletal muscle growth in newly-hatched chicks, e.g., how protein synthesis can be enhanced in skeletal muscles after hatching, thereby contributing to improved industrial poultry meat productivity.

Acknowledgment

This work was supported by JSPS KAKENHI Grant Number 15K18770.

References

Arthur, J., Albers, G. 2003. Industrial perspective on problems and issues associated with

poultry breeding. In: Muir WM, Aggrey SE. (eds), *Poultry Genetics, breeding and biotechnology*, pp. 1-12. CABI Publishing.

- Ashmore, C.R., Doerr, L. 1971. Postnatal development of fiber types in normal and dystrophic skeletal muscle of the chick. *Experimental neurology* **30**, 431-446.
- Barnard EA, Lyles JM, Pizzey JA. 1982. Fibre types in chicken skeletal muscles and their changes in muscular dystrophy. *The Journal of Physiology* **331**, 333-354.
- Bentzinger, C.F., Wang Y.X., Rudnicki M.A. 2012 building Muscle: Molecular regulation of myogenesis. *Cold Spring Harbor Perspectives in Biology* **4**, a008342.
- Berkes, C.A., Bergstrom D.A., Penn B.H., Seaver K.J., Knoepfler P.S., Tapscott S.J. 2004.
 Pbx marks genes for activation by MyoD indicating a role for a homeodomain protein in establishing myogenic potential. *Molecular cell* 14, 465-477.
- Beylkin, D.H., Allen D.L., Leinwand L.A. 2006. MyoD, Myf5, and the calcineurin pathway activate the developmental myosin heavy chain genes. *Developmental Biology* 294, 541-553.
- Davis, T.A., Fiorotto, M.A., Nguyen, H.V., Reeds, P.J. 1989. Protein turnover in skeletal muscle of suckling rats. *The American Journal of Physiology* 257, R1141-R1146.
- Duchêne, S., Metayer, S., Audouin, E., Bigot, K., Dupont, J., Tesseraud, S. 2008.
 Refeeding and insulin activate the Akt/p70S6 kinase pathway without affecting IRS1 tyrosine phosphorylation in chicken muscle. *Domestic Animal Endocrinology* 3, 1-13.
- Fridolfsson, A.K. and Ellegren, H. 1999. A simple and universal method for molecular sexing of non-ratite birds. *Journal of Avian Biology* **30**, 116-121.
- Glass, D.J. 2005. Skeletal muscle hypertrophy and atrophy signaling pathways. *The International Journal of Biochemistry & Cell Biology* **37**, 1974-1984.

- Gonzales, E., Kondo, N., Saldanha, E.S., Loddy, M.M., Careghi, C., Decuypere, E. 2003. Performance and physiological parameters of broiler chickens subjected to fasting on the neonatal period. *Poultry Science* 82, 1250-1256.
- Goodman, C.A., Kotecki, J.A., Jacobs, B.L., Hornberger, T.A. 2012. Muscle fiber typedependent differences in the regulation of protein synthesis. *PLos ONE* **7**, e37890
- Guernec, A., Chevalier, B., Duclos, M.J. 2004. Nutrient supply enhances both IGF-1 and MSTN mRNA levels in chicken skeletal muscle. *Domestic Animal Endocrinology* 26, 143-154.
- Gumucio, J.P., Mendias, C.L. 2013. Atrogin-1, MuRF-1, and sarcopenia. *Endocrine* **43**, 12-21.
- Halevy, O., Geyra, A., Barak, M., Uni, Z., Sklan, D. 2000. Early posthatch starvation decreases satellite cell proliferation and skeletal muscle growth in chicks. *The Journal of Nutrition* 130, 858-864.
- Halevy, O., Nadel, Y., Barak, M., Rozenboim, I., Sklan, D. 2003. Early posthatch feeding stimulates satellite cell proliferation and skeletal muscle growth in turkey poults. *The Journal of Nutrition* 133, 1376-1382.
- Halevy, O., Piestun, Y., Allouh, M.Z., Rosser, B.W.C., Rinkevich, Y., Reshef, R., Rozenboim, I., Wleklinski-Lee, M., Yablonka-Reuveni, Z. 2004. Pattern of Pax7 expression during myogenesis in the posthatch chicken establishes a model for satellite cell differentiation and renewal. *Developmental Dynamics* 231, 489-502.
- Kimball, S.R., Farrell, P.A., Nguyen, H.V., Jefferson, L.S., Davis, T.A. 2002. Developmental decline in components of signal transduction pathways regulating protein synthesis in pig muscle. *American journal of physiology. Endocrinology and metabolism* 282, E585-E592.

- Kurokawa, M., Sato, F., Aramaki, S., Soh, T., Yamauchi, N., Hattori, M. 2009. Monitor of the myostatin autocrine action during differentiation of embryonic chicken myoblasts into myotubes: effect of IGF-I. *Molecular Cell Biochemistry* 331, 193-199.
- Legerlotz, K., Smith, H.K. 2008. Role of MyoD in denervated, disused, ad exercised muscle. *Muscle Nerve* **38**, 1087-1100.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J. 1951. Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry* **193**, 265-275.
- Mauro, A. 1961. Satellite cells of skeletal muscle fibers. *The Journal of biophysical and biochemical cytology* **9**, 493-495.
- Morissette M.R., Cook S.A., Buranasombati C., Rosenberg M.A., Rosenzweig A. 2009. Myostatin inhibits IGF-1-induced myotube hypertrophy through Akt. American journal of physiology. Cell Physiology 297, C1124-C1132.
- Moss, F.P., LeBlond, C.P. 1971. Satellite cells are source of nuclei in muscles of growing rats. *The Anatomical record* **170**, 421-436.
- Nabeshima Y., Hanaoka K., Hayasaka M., Esumi E., Li S., Nonaka I., Nabeshima Y. 1993. Myogenin gene disruption results in perinatal lethality because of severe muscle defect. *Nature* 364, 532-535.
- Noy, Y., Sklan, D. 1997. Posthatch development in poultry. *The Journal of Applied Poultry Research* **6**, 344-354.
- Rushbrook, J.I., Huang, J., Weiss, C., Siconolfi-Baez, L., Yao, T.T., Becker, E., Feuerman,
 M. 1997. Characterization of the myosin heavy chains of avian adult fast muscles at
 the protein and mRNA levels. *Journal of Muscle Research and Cell Motility* 18, 449-463.

Rushbrook, J.I., Huang, J., Weiss, C., Yao, T.T., Siconolfi-Baez, L., Becker, E. 1998. Protein and mRNA analysis of myosin heavy chains in the developing avian pectoralis major muscle. *Journal of Muscle Research and Cell Motility* **19**, 157-168.

Sandri M. 2008. Signaling in muscle atrophy and hypertrophy. *Physiology* 23, 160-170.

- Saneyasu T., Inui M., Kimura S., Yoshimoto Y., Tsuchii N., Shindo H., Honda K., Kamisoyama H. 2016. The IGF-1/Akt/S6 signaling pathway is age-dependently downregulated in chicken breast muscle. *The Journal of Poultry Science* 53, 213-219.
- Saneyasu, T., Kimura, S., Inui, M., Yoshimoto, Y., Honda, K., Kamisoyama, H. 2015. Differences in the gene expression involved in skeletal muscle proteolysis during food deprivation between broiler and layer chicks. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **186**, 36-42.
- Schiaffino S, Dyer KA, Cicilot S, Blaauw B, Sandi M. 2013. Mechanisms regulating skeletal muscle growth and atrophy. *FEBS Journal* 280, 4294-4314.
- Scheuermann, G.N., Bilgili, S.F., Tuzun, S., Mulvaney, D.R. 2004, Comparison of chicken genotypes: myofiber number in pectoralis muscle and myostatin ontogeny. *Poultry Science* 83, 1404-1412.
- Shah, O.J., Anthony, J.C., Kimbakk, S.R., Jefferson, L.S. 2000. 4E-BP1 and S6K1: translational integration sites for nutritional and hormonal information in muscle. *American Journal of Physiology. Endocrinology and Metabolism* 279, E715-E729.
- Shanely, R.A., Zwetsloot, K.A., Childs, T.E., Lees, S.J., Tsika, R.W., Booth, F.W. 2009. IGF-I activates the mouse type IIb myosin heavy chain gene. *American journal of physiology. Cell physiology* 297, C1019-C1027.

Sjögren, K., Liu, J.L., Blad, K., Skrtic, S., Vidal, O., Wallenius, V., LeRoith, D., Törnell,

J., Isaksson, O.G., Jansson, J.O., Ohlsson, C. 1999. Liver-derived insulin-like growth factor I (IGF-I) is the principal source of IGF-I in blood but is not required for postnatal body growth in mice. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 7088-7092.

- Smith, J.H. 1963. Relation of body size to muscle cell size and number in the chicken. *Poultry Science* **42**, 283-290.
- Sobolewska, A., Elminowska-Wenda, G., Bogucka, J., Szpinda, M., Walasik, K., Bednarczyk, M., Paruszewska-Achtel, M. 2011. Myogenesis - possibilities of its stimulation in chickens. *Folia biologica* 59, 85-90.
- Spiller, M.P., Kambadur, R., Jeanplong, F., Thomas, M., Martyn, J.K., Bass, J.J., Sharma,
 M. 2002. The myostatin gene is a downstream target gene of basic helix-loop-helix
 transcription factor MyoD. *Molecular and cellular biology* 22, 7066-7082.
- Velleman, S.G. 2007. Muscle development in the embryo and hatchling. *Poultry Science* 86, 1050-1054.
- Wheeler, M.T., Snyder, E.C., Patterson, M.N., Swoap, S.J. 1999. An E-box within the MHC IIB gene is bound by MyoD and is required for gene expression in fast muscle. *American Journal of Physiology. Cell Physiology* 276, C1069-C1078.
- Yakar, S., Liu, J.L., Stannard, B., Butler, A., Accili, D., Sauer, B., LeRoith, D. 1999.
 Normal growth and development in the absence of hepatic insulin-like growth factor
 I. *Proceedings of the National Academy of Sciences of the United States of America* 96, 7324-7329.

Figure legends

Figure 1. Changes in the weights of body, yolk sac, and skeletal muscle in perinatal

male chicks. Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (*P* < 0.05).

- Figure 2. Changes in mRNA levels of MyoD and myogenin in breast muscles of male chicks. Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (P < 0.05).
- Figure 3. Changes in mRNA levels of IGF-1 and myostatin, phosphorylation of Akt and S6, and contents of phosphorylated and total S6 in breast muscles of male chicks. Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (P < 0.05).
- Figure 4. Changes in mRNA levels of myosin heavy chain and troponin I isoforms in breast muscles of male chicks. Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (P < 0.05). HC, myosin heavy chain

孵化後1週間におけるニワトリ骨格筋の IGF-1/Akt/S6 経路及び解糖型ミオシン 重鎖アイソフォームの遺伝子発現量は加齢依存的に上向き調節される

實安隆興・土橋竜也・来代紋菜・槌井七海・木村沙也加・本田和久・上曽山博

神戸大学大学院農学研究科 神戸市 657-8501

骨格筋量は畜産業において重要な形質である。これまでに我々は、ニワトリのム ネ肉重量は増加し続けるにもかかわらず、ムネ肉における IGF-1/Akt/S6 経路は 1週齢以降加齢依存的に下向き調節されることを報告している。骨格筋線維中の 主要なタンパク質であるミオシン重鎖(HC)には複数のアイソフォームが存在 し、骨格筋の種類によって発現するアイソフォームが異なる。また、成鶏のムネ 肉には HCI のみが発現していることが知られている。しかしながら、孵化前後 のニワトリ骨格筋におけるタンパク質合成関連因子や HC アイソフォームの発 現変動については殆ど知られていない。本研究では、タンパク質合成因子である、 IGF-1 mRNA 量、Akt のリン酸化率、及びリン酸化 S6 タンパク質量は0日齢以 降、加齢依存的に増加した。解糖型速筋線維型である HC I, III 及び V の mRNA 量は孵化後増加した。また、ムネ肉重量の増加率は0日齢から7日齢にかけて 約1,100%であった。これらの結果から、ニワトリのムネ肉では、タンパク質合 成経路と解糖型速筋線維型 HC アイソフォームの転写は孵化後1週間において 上向き調節されることが示唆された。

| Gene name | Forward primer | Reverse primer | Accession number |
|---------------------|---|---|------------------|
| MyoD | 5'-gga atc acc aaa tga ccc aaa g-3' | 5'-ggc agt cga ggc tgg aaa c-3' | NM_204214 |
| Myogenin | 5'-ggg aga agc gga ggc tga ag-3' | 5'-cag agt gct gcg ttt cag agc c-3' | NM_204184 |
| IGF-1 | 5'-gct gcc ggc cca gaa-3' | 5'-acg aac tga aga gca tca acc a-3' | NM_001004384 |
| Myostatin | 5'-atg cag atc gcg gtt gat c-3' | 5'-gcg ttc tct gtg ggc tga ct-3' | NM_001001461 |
| HC I [#] | 5'-aga gaa gat gtg ccg cac tct gga gga c-3' | 5'-gta aat ccc tgc ttg cct cta gag-3' | |
| HC III [#] | 5'-gcg gat gat caa tga cct caa tgc t-3' | 5'-ctc ttg gcc ttt atc tcc tcc tct aaa tgc-3' | |
| HC IV [#] | 5'-cct gga gaa gat tgc cgc aca ct-3' | 5'-gag taa atg cct gtt tgc ctc ttg ata ac-3' | |
| HC V [#] | 5'-acg gat gat caa tga tgt taa tgc t-3' | 5'-agc aga ctg caa acc atg ggc t-3' | |
| Tn I type 1 | 5'-tga gcc aac tcc agg atc tgt-3' | 5'-tet teg tee aca ate tea ace tt-3' | XM_004934839 |
| Tn I type 2 | 5'-gtg ggc atg tct gct gat g-3' | 5'-cat gtt gac ctt gtg ctt gga-3' | NM_205417 |
| RPS17 | 5'-gcg ggt gat cat cga gaa gt-3' | 5'-gcg ctt gtt ggt gtg aag t-3' | NM_204217 |

Table 1. Primer sequences used for real-time PCR analysis

IGF-1, insulin like growth factor-1; HC, myosin heavy chain; RPS17, ribosomal protein S17; Tn I, troponin I. The primers of IGF-1, myostatin, RPS17 were used in previous studies (Saneyasu *et al.*, 2015, 2016). [#], Refer to corresponding nucleotide sequences reported by Rushbrook *et al.* (1997, 1998).















Figure 4













Supplementary Figure 1. Changes in the weight of body, yolk sac, and skeletal muscle in perinatal female chicks.

Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (*P* < 0.05).



Supplementary Figure 2. Changes in mRNA levels of MyoD and myogenin in breast muscles of female chicks.

Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (*P* < 0.05).





Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (*P* < 0.05).



Supplementary Figure 4. Changes in mRNA levels of myosin heavy chain isoforms in breast muscles of chicks. Data are expressed as mean \pm SEM (n = 6). The Tukey-Kramer method was used to analyze differences. Groups with different letters were significantly different (P < 0.05). HC, myosin heavy chain.