

Review Article Clinical Development of Immune Checkpoint Inhibitors

Ayumu Ito,¹ Shunsuke Kondo,^{2,3} Kohei Tada,⁴ and Shigehisa Kitano^{2,4}

¹Department of Hematopoietic Stem Cell Transplantation, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

²Department of Experimental Therapeutics, Exploratory Oncology Research and Clinical Trial Center (EPOC), National Cancer Center, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

³Department of Hepatobiliary and Pancreatic Oncology, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

⁴Division of Cancer Immunotherapy, Exploratory Oncology Research and Clinical Trial Center (EPOC), National Cancer Center, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

Correspondence should be addressed to Shigehisa Kitano; skitano@ncc.go.jp

Received 25 July 2014; Accepted 7 December 2014

Academic Editor: Swaleha Zubair

Copyright © 2015 Ayumu Ito et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recent progress in cancer immunotherapy has been remarkable. Most striking are the clinical development and approval of immunomodulators, also known as immune checkpoint inhibitors. These monoclonal antibodies (mAb) are directed to immune checkpoint molecules, which are expressed on immune cells and mediate signals to attenuate excessive immune reactions. Although mAbs targeting tumor associated antigens, such as anti-CD20 mAb and anti-Her2 mAb, directly recognize tumor cells and induce cell death, immune checkpoint inhibitors restore and augment the antitumor immune activities of cytotoxic T cells by blocking immune checkpoint molecules on T cells or their ligands on antigen presenting and tumor cells. Based on preclinical data, many clinical trials have demonstrated the acceptable safety profiles and efficacies of immune checkpoint inhibitors in a variety of cancers. The first in class approved immune checkpoint inhibitor is ipilimumab, an anti-CTLA-4 (cytotoxic T lymphocyte antigen-4) mAb. Two pivotal phase III randomized controlled trials demonstrated a survival benefit in patients with metastatic melanoma. In 2011, the US Food and Drug Administration (FDA) approved ipilimumab for metastatic melanoma. Several clinical trials have since investigated new agents, alone and in combination, for various cancers. In this review, we discuss the current development status of and future challenges in utilizing immune checkpoint inhibitors.

1. Introduction

In this decade, remarkable progress has been made in the clinical application of cancer immunotherapies. Most notable is the emergence of immune checkpoint inhibitors. Largescale clinical trials have shown their feasibility and efficacy for patients with advanced malignancies. The therapeutic targets, or "immune checkpoints," are also known as coinhibitory molecules or costimulatory molecules expressed on T cells.

As the name implies, costimulatory/inhibitory molecules mediate positive/negative signals that modify MHC-TCR (major histocompatibility complex-T-cell receptor) signaling pathways. These signals each regulate T-cell survival, proliferation, differentiation, or responsiveness to cognate antigens. The net effect depends on the balance among signals [1]. Tcell activation requires costimulatory signals. If they contact antigens without costimulatory ligands on antigen presenting cells (APCs), T cells remain inactivated in a state of anergy.

Coinhibitory molecules induce T-cell dysfunction (so called "T-cell exhaustion") or apoptosis. Employing this inhibitory pathway, the immune system can attenuate excessive immune reactions and ensure self-tolerance, which is important for maintaining immune homeostasis. These functions involve programmed cell death protein-1 (PD-1), programmed cell death-1 ligand-1/2 (PD-L1/2), cytotoxic T lymphocyte antigen-4 (CTLA-4), lymphocyte-activation gene 3 (LAG-3), T-cell immunoglobulin mucin-3 (TIM-3), and B and T lymphocyte attenuator (BTLA). Tumor cells harness

these suppressive effects as one of their "immunoediting" mechanisms [2]. As shown in recent clinical trials, immune checkpoint blockade with monoclonal antibody promotes endogenous antitumor activities of immune cells and achieves clinically significant benefits for cancer patients [3, 4].

In this review, we focus on the current development status of and future challenges in utilizing immune checkpoint inhibitors, especially CTLA-4, PD-1, and PD-L1.

2. Anti-CTLA-4 Antibody

CTLA-4 (also known as CD152) is a member of the CD28 family of receptors [21, 22]. CTLA-4 is inducibly expressed on the surfaces of activated conventional CD4⁺ and CD8⁺ T cells. CTLA-4 binds to ligands B7.1 (CD80) and B7.2 (CD86) on APCs, where it competes with costimulatory receptor CD28 to bind with shared ligands. As CTLA-4 binds with higher affinity than CD28, it reduces CD28-dependent costimulation. CTLA-4 also mediates direct inhibitory effects on the MHC-TCR pathway [23]. CTLA-4 recruits 2 phosphatases, SHP-2 and PP2A, to its intracellular YVKM domain. SHP-2 dephosphorylates the CD3 ζ chain, attenuating the TCR signal. PP2A inhibits downstream Akt phosphorylation, further impairing TCR signaling. Furthermore, CTLA-4 is constitutively and highly expressed on CD4⁺CD25⁺FOXP3⁺ regulatory T cells (T regs) and plays a role in their suppressive functions [24-26]. CTLA-4 knockout mice have a lethal autoimmune-like syndrome. Prominent infiltration of CD4⁺ T cells is detected in multiple organs. Thus, CTLA-4 is considered to be indispensable for maintaining immune homeostasis.

In the tumor microenvironment, CTLA-4 suppresses antitumor immune activities. In animal models, it has been shown that CTLA-4 blockade leads to reactivation of the antitumor immune response and tumor shrinkage [27–29]. The mechanism of action has not yet been fully elucidated. Observations made to date suggest that anti-CTLA-4 antibodies function not only by blocking inhibitory signals from reaching effector T cells but also by depleting regulatory T cells in the tumor microenvironment [30, 31]. For use in humans, based on preclinical studies, two anti-CTLA-4 antibodies have been developed: ipilimumab (Bristol-Myers Squibb) and tremelimumab (Pfizer).

2.1. Ipilimumab. Ipilimumab is a fully humanized IgG1 monoclonal antibody that inhibits CTLA-4 [32, 33].

Early clinical trials evaluated ipilimumab in patients with a variety of malignancies, including melanoma, prostate cancer, renal cell carcinoma, and non-Hodgkin lymphoma [34–45]. Some of these studies combined ipilimumab with a peptide vaccine, chemotherapy, or IL-2. Based on preclinical data, ipilimumab was administered at a dose range of 0.1– 20 mg/kg, employing single or multiple dosing schedules (every 3-4 weeks).

A phase I study evaluated a single 3 mg/kg dose of ipilimumab for patients with metastatic hormone-refractory prostate cancer. Two (14%) of 14 patients showed $\geq 50\%$

decline in prostate specific antigen. One (7%) patient developed grade 3 rash/pruritus requiring systemic corticosteroid administration [36]. Another phase I trial combined ipilimumab (administered at 3 mg/kg every 3 weeks) with a glycoprotein (gp) 100 peptide vaccine for patients with metastatic melanoma. Three (21%) of 14 patients responded to this treatment, including 2 showing complete responses (CRs). Grade 3 to 4 immune-related adverse events (irAEs) occurred in 6 (43%) patients. These irAEs included dermatitis, enterocolitis, hepatitis, and hypophysitis [34]. On the whole, irAEs were mild and manageable with therapy discontinuation and/or appropriate treatments, including corticosteroids.

A phase II trial compared 3 doses (0.3, 3, or 10 mg/kg) administered every 3 weeks for a total of 4 doses. Eligible patients were permitted to receive reinduction therapy (at a dose of 10 mg/kg) or maintenance therapy (administered at the previously assigned dose level every 12 weeks). The overall response rate (ORR) in the 10 mg/kg arm was superior to those in the other arms (11.1% versus 4.2% versus 0.0%), but irAEs were also higher in the 10 mg/kg arm [43]. The optimal dosing and scheduling are as yet unknown. A phase III randomized trial (NCT01515189) is currently comparing 2 doses (3 mg/kg versus 10 mg/kg). No consensus has yet been reached on the relative significance of reinduction versus maintenance therapy [46, 47]. A prospective study comparing reinduction therapy versus the physician's choice of chemotherapy (NCT00495066) is currently underway.

Based on pivotal phase III randomized controlled trials (RCTs) showing survival benefit, ipilimumab was approved by the US Food and Drug Administration (FDA) for metastatic melanoma [5, 6]. In the landmark phase III trial for patients with previously treated metastatic melanoma, ipilimumab (administered at 3 mg/kg every 3 weeks for a total of 4 doses) with or without the gp 100 peptide vaccination was compared with the gp 100 peptide vaccine alone. Eligible patients were permitted to receive reinduction therapy. The median OSs in the ipilimumab-containing arms were significantly superior to that in the gp 100 alone arm (10.1 months in ipilimumab/gp 100, 10.0 months in ipilimumab alone, and 6.4 months in gp 100 alone, hazard ratio (HR) 0.68; P < 0.001). Grade 3 to 4 irAEs were seen in 10-15% of patients in the ipilimumab-containing arms, while 3% in the gp 100 alone arm experienced irAEs. There were 14 treatment-related deaths (2.1%), including 7 patients with irAEs [5]. Long-term follow-up analysis confirmed an approximately 20% survival rate for patients in the ipilimumab-containing arms. Safety profiles in long-term survivors were comparable among the 3 groups, and new onset irAEs after the last dose of ipilimumab were infrequent (8%; all grades) [48]. The other phase III trial compared ipilimumab (at 10 mg/kg every 3 weeks for 4 doses)/dacarbazine with dacarbazine/placebo, followed by maintenance therapy with ipilimumab or placebo administered every 12 weeks for eligible patients. Overall survival (OS) was significantly longer in the ipilimumab/dacarbazine arm (11.2 versus 9.1 months), and the higher survival rates were durable (47.3% versus 36.3% at 1 year, 28.5% versus 17.9% at 2 years, 20.8% versus 12.2% at 3 years, HR for death 0.72; P < 0.001). Grade 3 to 4 AEs were seen in more patients in the ipilimumab/dacarbazine arm (56.3% versus 27.5%;

P < 0.001). No drug-related deaths occurred among those in the ipilimumab/dacarbazine arm [6].

The analysis of the collected data from 12 previous clinical trials, which include 1861 ipilimumab-treated patients with advanced melanoma, demonstrated a median OS of 11.4 months and 3-year OS rate of 22%. The OS curve started to show plateau around year 3, which was independent of the dose of ipilimumab (3 or 10 mg/kg), therapy line (treatment-naïve or not), or use of maintenance therapy [49].

2.2. Tremelimumab. Tremelimumab is a human IgG2 monoclonal antibody that blocks CTLA-4 [50].

Early clinical trials on tremelimumab monotherapy showed response rates of 2–17%, and these responses were durable (>150 days) [51–57]. Based on preclinical and clinical data, the standard regimen is 15 mg/kg every 90 days. Most adverse events were mild and manageable. These adverse events included skin rash, diarrhea, and endocrine abnormalities.

A phase III study compared tremelimumab (15 mg/kg every 3 months) with chemotherapy (physician's choice) in patients with untreated advanced melanoma [7]. This study demonstrated no benefits in either ORR (10.7% versus 9.8%) or OS (12.6 mo versus 10.7 mo), but a superior response duration was seen (35.8 versus 13.7 months). This observation might be explained by patient selection bias (exclusion of patients with lactate dehydrogenase (LDH) >2x upper limit of normal), drug crossover (to ipilimumab) in the control arm, and even a potentially suboptimal dosing regimen. Tremelimumab is still being investigated for other tumors, both alone and as combination therapy (Table 1).

3. Anti-PD-1 Antibodies

Programmed cell death protein-1 (PD-1; also known as CD279), like CTLA-4, is a coinhibitory CD28-family molecule [22]. While CTLA-4 works in the early phase of naïve-T-cell activation, PD-1 functions mainly in the late phase, in which PD-1 induces exhaustion or anergy in effector T cells. Thus, PD-1 is considered to play an important role in chronic inflammation such as that associated with viral infection or tumor exposure [58]. PD-1 is expressed on activated T cells, T regs [59], activated B cells, NK cells, and monocytes. It binds to the B7-family ligands PD-L1 (programmed death ligand-1, B7-H1) and PD-L2 (programmed death ligand-2, B7-DC) on APCs. PD-1 has cytoplasmic domain motifs known as ITIM (immunoreceptor tyrosine-based inhibitory motif) and ITSM (immunoreceptor tyrosine-based switch motif) [23]. When these motifs are phosphorylated, they recruit two inhibitory phosphatases, SHP-1 and SHP-2 (SHP: SH2containing-phosphatase). These phosphatases dephosphorylate the CD3 ζ chain, decreasing TCR signaling. Although the inhibitory mechanisms of CTLA-4 and PD-1 have some similarity in terms of inhibiting Akt activation, CTLA-4 can also interfere with Akt independently via PP2A [23]. PD-1 knockout mice show a milder lupus-like syndrome than CTLA-4 knockout mice [60].

Tumor cells utilize the PD-1-PD-L1/2 pathway to evade immune-cell attack [61]. Blockade of this pathway was shown to restore and augment antitumor immune activities [62].

3.1. Nivolumab (BMS-936558/ONO-4538). Nivolumab is a fully humanized IgG4 monoclonal antibody that blocks PD-1 [62].

Phase I studies tested nivolumab in such cancers as melanoma, non-small cell carcinoma of the lung (NSCLC), ovarian cancer, and renal cell carcinoma. These studies showed response rates of approximately 20-30%, durable tumor regression (>1 year), and an acceptable safety profile, with Grade 3 to 4 irAEs developing in about 20% of patients [8, 9, 63–65]. In long-term follow-up of the phase I trial for advanced melanoma, median OS was 16.8 months and survival rates were 62% at 1 year and 43% at 2 years. The patients requiring discontinuation of treatment maintained their tumor responses for at least 16 months (16–56 months). Long-term safety profiles were acceptable and similar to those described in a previous report [8]. The preliminary results of a phase I study evaluating nivolumab (at 3 mg/kg q2w) for untreated advanced NSCLC were recently reported. The ORR was 30% with 2 complete remissions (CRs), as measured by RECIST. ORR and progression-free survival (PFS) correlated with PD-L1 positivity (67% versus 0% for ORR, 45.6 mo versus 36.1 mo for median PFS). AEs were generally manageable and grade 3 to 4 AEs occurred in 3 patients, including rash, increased transaminase, and hyperglycemia [66].

Recently the interim analysis report of a phase III study (NCT01721746), comparing nivolumab monotherapy (at 3 mg/kg q2w) with investigator's choice chemotherapy in ipilimumab-refractory advanced melanoma, was shown. The ORRs were 32% in the nivolumab arm and 11% in the control arm, with the median duration of response in the nivolumab arm not reached. Grade 3 to 4 drug-related AEs were less frequent in the nivolumab arm (9% versus 31%) [10]. Another phase III study (NCT01721772) compared nivolumab monotherapy (at 3 mg/kg q2w) with dacarbazine in 418 patients with previously untreated stage III or IV melanoma. This study was stopped ahead of schedule and unblinded after independent data monitoring committee found significant survival superiority in nivolumab over dacarbazine. The results from the double-blind part of the study before the stoppage showed that the OS rate at 1 year was significantly higher in the nivolumab arm (72.9% versus 42.1%, HR for death 0.42; P < 0.001), and the median PFS was also significantly longer in the nivolumab arm (5.1 versus 2.2 months, HR for death or progression 0.43; P < 0.001). Grade 3 to 4 drug-related AEs occurred in more patients in the dacarbazine arm (11.7% versus 17.6%). No drug-related deaths occurred in both arms [11]. A phase II study (NCT01927419) of nivolumab in combination with ipilimumab compared with ipilimumab alone for advanced melanoma is currently ongoing (recruitment has been completed).

In 2013, nivolumab received Fast Track designation for the treatment of NSCLC, melanoma, and renal cell carcinoma (RCC) from the FDA. In April 2014, a rolling submission to the FDA for nivolumab in third-line pretreated NSCLC was started. In May 2014, nivolumab received a Breakthrough

TABLE 1	Reference	[5]	[6]	[2]	
	Treatment-related adverse events (≧Gr3)	lpi + gp100: drug-related 17.4%; irAEs 10.2%; diarrhea 4.5%; fatigue 5.0%; dyspnea 3.7%; anemia 2.9%; endocrine abnl. 11%; AST↑ 0.5%; ALT↑ 0.3% lpi + DTIC:	immune-related 41.7%; pruritus 2.0%; rash 1.2%; diarrhea 4.0%; colitis 6.1%; AST† 17.4%; ALT† 20.7%	52%; diarrhea/colitis 18%; fatigue 6%; rash 2%; pruritus 1%; dyspnea 3%; hypothalamus and pituitary disorders 1%; hepatitis 1%	
	Survival	Ipi + gp100 versus gp100: 10.1 versus 6.4 mos	lpi + DTIC versus PBO + DTIC: 11.2 versus 9.1 mos	Treme. versus chemo.: 12.6 versus 10.7 mos (NS)	
	Response	Ipi + gp100: ORR 5.7%; SD 14.4%	Ipi + DTIC: ORR 15.2%; SD 18.0%	ORR 10.7%	
TABLE 1	Study design	Endpoint: safety/efficacy Ipi + gp100 versus Ipi versus gp100	Endpoint: efficacy Ipi + DTIC versus PBO + DITC	Endpoint: efficacy treme. versus chemo.	
	Number of patients	676	502	655	
	Disease	Melanoma	Melanoma	Melanoma	
	Status/NCT number	Completed (NCT00094653)	Completed (NCT00324155)	Completed (NCT00257205)	
	Phase	III	III	Ш	
	Drug name	Ipilimumab		Tremelimumab	
	Target molecule		CTLA-4		

Γ
Щ
H
В
.≺
H

name Phase Status/NCT Disease Number of Study design Response Sindy Respo	PhaseStatus/NCTDiseaseNumber of patientsStudy designResponseStudy design	Status/NCT Disease Number of Study design Response Si number	Disease Number of Study design Response S	Number of Study design Response S	f Study design Response S	Response S	N N	urvival	Treatment-related adverse events (≧Gr3)	Reference
Ongoing (notEndpoint:ORR 30.8%;Irecruiting)Melanoma107safety/efficacy 5median durat(NCT00730639)(NCT00730639)dosing regimensSD (≥24 wks)	ICongoing (not recruiting)Endpoint: MelanomaORR 30.8%; Median duratIrecruiting)Melanoma107safety/efficacy 5median durat(NCT00730639)dosing regimensSD (≥24 wks)	Ongoing (notEndpoint:ORR 30.8%;recruiting)Melanoma107safety/efficacy 5median durat(NCT00730639)dosing regimensSD (≥24 wks)	Endpoint:ORR 30.8%;Melanoma107safety/efficacy 5response 104 and duratdosing regimensSD (≥24 wks)	Endpoint: ORR 30.8%; 107 safety/efficacy 5 response 104 durat dosing regimens SD (≥24 wks)	Endpoint: ORR 30.8%; safety/efficacy 5 response 104 ¹ dosing regimens SD (≥24 wks)	ORR 30.8%; median durat response 104 ¹ SD (≥24 wks)	ion of wks; 6.5%	OS 16.8 mos; PFS 3.7 mos	22.4%; fatigue 1.9%; diarrhea 1.9%; abdominal pain 1.9%; lymphopenia 2.8%	[8]
S- Ongoing (not Endpoint: ORR 25%; SE 58/ONO- I recruiting) Melanoma 90 safety/efficacy 3 (≥24 wks) 219 dosing regimens (≥24 wks) 219	IEndpoint:ORR 25%; SEIrecruiting)Melanoma90safety/efficacy 3(≥24 wks) 219(NCT01176461)dosing regimens(≥24 wks) 219	Ongoing (notEndpoint:ORR 25%; SErecruiting)Melanoma90safety/efficacy 3(≥24 wks) 219(NCT01176461)dosing regimens(≥24 wks) 219	Endpoint:ORR 25%; SEMelanoma90safety/efficacy 3(≥24 wks) 219dosing regimens(≥24 wks) 219	Endpoint: ORR 25%; SE 90 safety/efficacy 3 (≧24 wks) 219 dosing regimens	Endpoint: ORR 25%; SE safety/efficacy 3 (≧24 wks) 219 dosing regimens	ORR 25%; SI (≧24 wks) 219	○ [∞]	PFS (at 24 wks) 46%	5.6%; rash 2.2%; interstitial pneumonitis 2.2%	[6]
Ongoing (not III recruiting) Melanoma 370 Endpoint: efficacy ORR 32% ver (NCT 01721772) Nivo. versus ICC 11%	Ongoing (not Endpoint: efficacy ORR 32% vei III recruiting) Melanoma 370 Nivo. versus ICC 11% (NCT 01721772) Nivo. versus ICC 11% 11% 11%	Ongoing (not recruiting) Melanoma 370 Endpoint: efficacy ORR 32% ver (NCT 01721772) Nivo. versus ICC 11%	Melanoma 370 Endpoint: efficacy ORR 32% ver Nivo. versus ICC 11%	370 Endpoint: efficacy ORR 32% ver Nivo. versus ICC 11%	Endpoint: efficacy ORR 32% ver Nivo. versus ICC 11%	ORR 32% vei 11%	sns	NA	9% versus 31%	[10]
III Completed Endpoint: efficacy ORR 40.0% Nivo. versus 13.9% dacarbazine 13.9%	III Completed Endpoint: efficacy ORR 40.0% (NCT01721772) Melanoma 418 Nivo. versus 13.9% dacarbazine	CompletedEndpoint: efficacyORR 40.0%(NCT01721772)Melanoma418Nivo. versus13.9%dacarbazine13.9%	Endpoint: efficacyORR 40.0%Melanoma418Nivo. versusdacarbazine13.9%	Endpoint: efficacyORR 40.0%418Nivo. versus13.9%dacarbazine13.9%	Endpoint: efficacy ORR 40.0% Nivo. versus 13.9% dacarbazine	ORR 40.0% 13.9%	versus	OS (at 1 yr) 72.9% versus 42.1%, median PFS 5.1 versus 2.2 mo	11.7% versus 17.6%; fatigue 0.5%; diarrhea 1.0%; rash 0.5%; vomiting 0.5%	[11]
izumab II Completed Endpoint: 311) II (NCT01435369) Melanoma 103 safety/efficacy 2 ORR 5.9% dosing regimens	II Completed Endpoint: (NCT01435369) Melanoma 103 safety/efficacy 2 ORR 5.9% dosing regimens	Completed Endpoint: (NCT01435369) Melanoma 103 safety/efficacy 2 ORR 5.9% dosing regimens	Endpoint: Melanoma 103 safety/efficacy 2 ORR 5.9% dosing regimens	Endpoint: 103 safety/efficacy 2 ORR 5.9% dosing regimens	Endpoint: safety/efficacy 2 ORR 5.9% dosing regimens	ORR 5.9%		OS (at 1 yr): 64.5%	NA	[12]
Ongoing (notEndpoint:ORR 38% bIrecruiting)Melanoma135safety/efficacy 3RECIST and (NCT01295827)(NCT01295827)dosing regimensby irRC	IOngoing (notEndpoint:ORR 38% bIrecruiting)Melanoma135safety/efficacy 3RECIST and(NCT01295827)dosing regimensby irRC	Ongoing (notEndpoint:ORR 38% brecruiting)Melanoma135safety/efficacy 3RECIST and(NCT01295827)dosing regimensby irRC	Endpoint:ORR 38% bMelanoma135safety/efficacy 3RECIST and dosing regimensby irRC	Endpoint: ORR 38% b safety/efficacy 3 RECIST and dosing regimens by irRC	Endpoint: ORR 38% b safety/efficacy 3 RECIST and dosing regimens by irRC	ORR 38% b RECIST and by irRC	y d 37%	Median PFS >7 mos	13%; hypothyroidism 1%; diarrhea 1%; fatigue 1%; AST7 1%; renal failure 1%; rash 2%; pruritus 1%	[13]
Ongoing (not rrolizumabUntreated safety/efficacy 3Endpoint: SAF6 bORR 26% b.3475)(NCT01295827)NSCLC57safety/efficacy 3RECIST an dosing regimens	IOngoing (not recruiting)Untreated 57Endpoint: safety/efficacy 3ORR 26% b RECIST an 0 kirRCIrecruiting)NSCLC57safety/efficacy 3RECIST an by irRC	Ongoing (not recruiting)Untreated S57Endpoint: safety/efficacy 3ORR 26% bNSCLC57safety/efficacy 3RECIST and dosing regimens(NCT01295827)NSCLCdosing regimensby irRC	Untreated 57 Endpoint: ORR 26% b NSCLC 57 safety/efficacy 3 RECIST an dosing regimens by irRC	Endpoint: ORR 26% b 57 safety/efficacy 3 RECIST an dosing regimens by irRC	Endpoint: ORR 26% b safety/efficacy 3 RECIST an dosing regimens by irRC	ORR 26% b RECIST an by irRC	y d 47%	Median OS NR; OS at 1 yr 80%; median PFS 45.6%; PFS at 24 wks 70%	CK↑ 2%; pericardial effusion 2%; pneumonitis 2%; acute kidney injury 2%	[14]
Ongoing (notHead andEndpoint:ORR 19.6%Irecruiting)Head and60safety/efficacy20.0% in HI(NCT01848834)neck cancer60single armHPV-;	IOngoing (notHead andEndpoint:ORR 19.6%Irecruiting)Head and60safety/efficacy20.0% in HI(NCT01848834)neck cancer60single armHPV-;	Ongoing (notHead andEndpoint:ORR 19.6%recruiting)Head and60safety/efficacy20.0% in HI(NCT01848834)neck cancer60single armHPV-;	Head andEndpoint:ORR 19.6%Head and60safety/efficacy20.0% in HIneck cancer60safety/efficacyand 19.4% inhead andsingle armHPV-;	Endpoint: ORR 19.6% Endpoint: 20.0% in HI safety/efficacy and 19.4% in single arm HPV-;	Endpoint: ORR 19.6% safety/efficacy 20.0% in HI and 19.4% ir single arm HPV-;	ORR 19.6% 20.0% in HI and 19.4% ir HPV-;	in total, V+, 1	NA	Gr3–5 16.7%; Rash 3.3%	[15]
Ongoing (not recruiting)Gastric GastricBrdpoint: Safety/efficacyORR 30.2% ORR 30.2%Irecruiting)cancer39safety/efficacy single armRECIST	IOngoing (not recruiting)Gastric Safety/efficacyEndpoint: ORR 30.2%Irecruiting)39safety/efficacy single armRECIST	Ongoing (not recruiting)Gastric astericBndpoint: Safety/efficacyORR 30.2% RECIST(NCT01848834)cancer39safety/efficacyRECIST	GastricEndpoint:ORR 30.2%39safety/efficacyRECISTcancersingle arm	Endpoint: ORR 30.2% safety/efficacy RECIST single arm	Endpoint: ORR 30.2% safety/efficacy RECIST single arm	ORR 30.2% RECIST	by	NA	7.7%; hypoxia 2.6%; peripheral neuropathy 2.6%; pneumonia 2.6%	[16]

TABLE 1: Continued.

	Reference			[17]		[18]	[61]	[20]	se; SD, stable not reached; sferase; CPK,
	Treatment-related adverse events (≧Gr3)		9%; fatigue 1%; infusion	reaction 1%; lymphopenia 1%		4%; no irAE	Any Gr 34%; Gr3/4 0%; no DLT; no MTD	Treatment discontinuation 52.2% (8.7% for AEs); drug-related AEs 11.1%; DLT 3.7% (CPK↑, myositis, and myocarditis)	e response rate; PR, partial respon. t available; NS, not significant; NR, ansferase; ALT, alanine aminotran.
	Survival	PFS (at 24 wks) 42%	PFS (at 24 wks) 31%	PFS (at 24 wks) 22%	PFS (at 24 wks) 53%	NA	NA	NA	tcebo; ORR, objective pillomavirus; NA, noi .ST, aspartate aminotr
	Response	ORR 17%; SD (≧24 wks) 27%	ORR 10%; SD (≥24 wks) 12%	ORR 6%; SD (≥24 wks) 18%	ORR 12%; SD (>24 wks) 41%	ORR: PD-L1 + 43% (at 6 wks) and 52% (at 12 wks); PD-L1	 – 11% (at 6 wks); PR 15.4%; disease control rate (≥12 wks) 46% 	NA	, dacarbazine; PBO, pli riteria; HPV, human pa s; abnl., abnormality; A therapy.
ABLE I. COULUINCO.	Study design		Padanint, mentatry 1	dose levels		Endpoint: safety/efficacy/ biomarker single	arm Endpoint: safety/efficacy single arm	Endpoint: safety single arm	peptide vaccine; DITC, nune-related response c ddverse event; Gr, Grad, stigator's choise chemo
-	Number of patients	52	49	17	17	68	26 (as of Jan 2014)	27 (as of Jan 2014)	coprotein 100 J ors; irRC, imm mune-related a dose; ICC, inve
	Disease	Melanoma	NSCLC	Ovarian cancer	Renal cell carcinoma	Urothelial bladder cancer	Advanced solid tumors	Refractory malignancies	umab; gp100, gly teria in solid tum e event; irAE, im imum tolerance
	Status/NCT number		Ongoing (not	recruiting) (NCT00729664)		Recruiting (NCT01375842)	Recruiting (NCT01693562)	Recruiting (NCT01772004)	g cancer; Ipi, Ipilim ponse evaluation cri survival; AE, advers ţ toxicity; MTD, max
	Phase			Ι		П	Ι	Г	nall cell lun RECIST, res ession-free lose limiting
	Drug name			BMS-936559		MPDL3280A	MEDI4736	MSB0019718C	ıs: NSCLC, non-sr month; wk, week; urvival; PFS, progi sphokinase; DLT, d
	Target molecule					PD-L1			Abbreviation disease; mo, OS, overall <i>s</i> creatine phos

TABLE 1: Continued.

BioMed Research International

Therapy designation for non-Hodgkin lymphoma from the FDA. In Japan, in July 2014, nivolumab received manufacturing and marketing approval for unresectable melanoma from the domestic regulator, the Ministry of Health Labor and Welfare, which made nivolumab the first in anti-PD-1 antibody to receive regulatory approval in the world.

3.2. Pidilizumab (CT-011). Pidilizumab (CT-011) is a humanized IgG-1 κ monoclonal antibody that blocks PD-1. In animal models, an antitumor effect was achieved with BAT monoclonal antibody (a murine mAb developed against a membrane preparation of a Burkitt lymphoma cell line), from which pidilizumab is derived [67, 68].

In humans, the safety and tolerability of the single dose regimen were shown in a phase I study of patients with advanced hematologic malignancies [69]. No treatment-related toxicities occurred and the maximum tolerated dose was not identified in this trial (0.2–6 mg/kg).

Pidilizumab has been tested in phase II trials, as monotherapy for patients with diffuse large B-cell lymphoma after autologous hematopoietic stem-cell transplantation [70] and as combined therapy with rituximab for relapsed follicular lymphoma [71]. Both trials showed promising efficacies even in high-risk patients.

The results of a phase II trial in patients with pretreated advanced melanoma were recently reported. ORR was 5.9%, measured by immune-related response criteria (irRC), and the OS rate at 1 year was 64.5%. The patients who had been pretreated with ipilimumab (51% of patients) tended to experience a higher rate of immune-related stable disease (irSD) and longer PFS (2.8 mo versus 1.9 mo) [12].

3.3. Pembrolizumab (MK-3475, Formally Known as Lambrolizumab). Pembrolizumab (MK-3475) is a humanized monoclonal IgG- 4κ antibody that blocks PD-1.

A phase I dose-escalation study evaluated three dose levels, 1 mg/kg, 3 mg/kg, and 10 mg/kg, administered every 2 weeks, in patients with multiple solid tumors [72]. All dose levels were found to be safe, and the maximum tolerated dose was not identified. Clinical responses were observed at all dose levels. Another phase I study tested 3 regimens (2 mg/kg every 3 weeks and 10 mg/kg every 2 or 3 weeks) in patients with advanced melanoma [13]. AEs were generally mild and grade 3 to 4 AEs were seen in 13% of patients. The ORRs ranged from 38% to 52%, in the biweekly 10 mg/kg cohort (measured by RECIST), showing no significant differences. These responses were durable, with the median PFS exceeding 7 months for all three regimens.

An ongoing phase II trial is now comparing 2 dose levels of pembrolizumab with investigator-choice chemotherapy in patients with previously treated advanced melanoma (NCT01704287). Another ongoing phase II trial is also evaluating 2 dose schedules of pembrolizumab (10 mg/kg q2w or q3w) compared with ipilimumab (3 mg/kg q3w) for advanced melanoma (NCT01866319).

In April 2013, pembrolizumab received the Breakthrough Therapy designation for advanced melanoma from the FDA. After being reviewed under the FDA's Accelerated Approval program, in September 2014, pembrolizumab received 7

approval for treatment of patients with advanced melanoma by the FDA.

Besides melanoma, several early trials have showed the tolerability and antitumor effects of pembrolizumab in other tumors. The preliminary results of another phase I study evaluating pembrolizumab in untreated PD-L1-positive NSCLC were recently reported. The overall objective response rate was 25% (33% in the 2 mg/kg q3w, 20% in the 10 mg/kg q3w, and 31% in the 10 mg/kg q2w group), as measured by RECIST. AEs were generally mild and grade 3 to 4 AEs occurred in 3 patients, including pneumonitis requiring treatment discontinuation [14]. Another preliminary result was reported for the phase I trial of pembrolizumab as monotherapy, administered at 2 mg/kg every 2 weeks, to 60 patients with recurrent/metastatic head and neck cancers. Grade 3 to 4 drugrelated AEs were reported in 16.7% of patients. The best ORR was 20% in all patients (assessed by RECIST 1.1). Efficacies were comparable between human papilloma virus- (HPV-) positive and HPV-negative patients (20.0% versus 19.4%) [15]. Another phase I study (NCT01848834) assessed pembrolizumab in the patients with previously treated advanced gastric cancer that expressed PD-L1. The enrolled 39 patients were treated with pembrolizumab at 10 mg/kg q2w. Median follow-up period was 6 months. Treatment-related AEs occurred in 24 patients (61.5%), and those of grade 3 to 5 occurred in 3 patients (pneumonitis, peripheral neuropathy, and hypoxia). ORR was 30.8% and disease control rate was 43.6%. Responses were mostly ongoing and the median response duration was not reached [16].

4. Anti-PD-L1 Antibodies

PD-L1 (also known as B7-H1 or CD274) and PD-L2 (also known as B7-DC or CD273) are inhibitory B7-family molecules that bind the PD-1 receptor. PD-L1 is inducibly expressed on a variety of hematopoietic and nonhematopoietic cells, including most human tumor cells and cells within the tumor microenvironment [61]. PD-L1 expression has been shown to correlate inversely with the clinical outcomes of some malignancies. PD-L2 is expressed on hematopoietic cells. PD-L1 knockout mice show infiltration of lymphocytes into nonlymphoid organs and exacerbation of preexisting autoimmune diseases [73, 74].

As mentioned above, the PD-1-PD-L1 axis is one of the main mechanisms by which cancer cells evade immune-cell attack [61]. Blockade of this pathway was shown to reinforce antitumor immune activities [62]. Because PD-L1 also interacts with CD80 [75, 76], anti-PD-L1 antibody might have optimal clinical potency against PD-1.

4.1. *BMS*-936559. BMS-936559 is a fully humanized IgG4 monoclonal anti-PD-L1 antibody. It inhibits the binding of PD-L1 to PD-1 and CD80. A phase I dose-escalation study evaluated BMS-936559 in 207 patients with selected cancers, including melanoma, NSCLC, ovarian cancer, and renal cell carcinoma. The study drug was administered at 4 dose levels (0.3–10 mg/kg) every 14 days, 3 times in each 6-week course for up to 16 cycles, when either CR or disease progression was confirmed. The ORRs were 6–17% and efficacy was durable

(>1 year in 8 of 16 patients who responded). Grade 3 to 4 irAEs, seen in 9% of the patients, were treatment-related in 5% [17].

4.2. MPDL3280A. MPDL3280A is a humanized IgG-1 κ monoclonal anti-PD-L1 antibody. It is genetically engineered to modify the Fc domain, thereby impairing the antibody-dependent cellular cytotoxicity of PD-L1 expressing cells [77, 78].

A phase I trial of MPDL3280A as monotherapy for advanced melanoma achieved a response rate of 26% and PFS of 35% at 24 weeks. Grade 3 to 4 AEs were seen in 33% of patients [79]. The results of another phase I trial were recently reported. MPDL3280A was tested in patients with pretreated metastatic urothelial bladder cancer. ORR in PD-L1-positive patients was superior to that in PD-L1-negative patients (43% versus 11% at 6 weeks). ORR at 12 weeks was 52% in PD-L1positive patients. Grade 3 to 4 AEs were seen in 4% of patients, with no irAEs [18]. The FDA has granted the Breakthrough Therapy designation to MPDL3280A.

4.3. *MEDI4736*. MEDI4736 is a humanized IgG- 1κ monoclonal antibody that blocks PD-L1. MEDI4736 demonstrated tumor regression and improved survival in a mouse model.

A "first-time-in-human" phase I study evaluating the safety, tolerability, and pharmacokinetics of this agent in patients with advanced solid tumors is currently underway (NCT01693562). The interim report was recently presented. As of January 2014, 26 patients were receiving dose-escalation treatments and had been given a median of 5 (1-25) q2w and 4.5 (1-7) q3w doses of MEDI4736 across 6 cohorts (0.1-10 mg/kg q2w; 15 mg/kg q3w). No dose limiting toxicities (DLTs) or maximum tolerated dose was identified. Treatment-related AEs occurred in 34% of patients, but all were grade 1 to 2 and did not lead totreatment discontinuation. Four of the 26 patients showed partial responses (PRs). The rate (PR + stable disease \geq 12 weeks) was 46%. Clinical responses were durable, with 11 patients remaining in the study (2+ to 14.9+ months) [19]. Another phase I trial is now testing the combination of MEDI4736 plus tremelimumab (NCT01975831).

4.4. MSB0010718C. MSB0010718 is a fully humanized IgG1 monoclonal antibody directed to PD-L1. A phase I trial is currently testing MSB0010718 to assess its safety, tolerability, and pharmacokinetics in patients with refractory malignancies (NCT01772004). As of January 2014, 27 patients had been enrolled and were participating in a dose-escalation study (3 + 3 design; 1, 3, 10, and 20 mg/kg, q2w). Twenty-three patients had been followed for at least 4 weeks. Discontinuation of the treatment had been necessary in 12 patients (52.2%): 9 (39.1%) due to progression of disease, 2 (8.7%) for AEs, and 1 (4.3%) because the patient died. Grade 3 to 4 drug-related toxicities included laboratory abnormalities in 3 patients. One DLT was observed in 1 patient at dose level 4 (20 mg/kg): an irAE with creatine kinase elevation, myositis, and myocarditis [20].

5. Combination Therapy

Recent clinical trials have actively investigated the potential for synergistic effects by combining immune checkpoint inhibitors with other agents. The partner agents/therapies include other checkpoint agents, cytotoxic agents, anticancer vaccines, cytokines, and radiotherapy.

A phase I study evaluated combined therapy with ipilimumab plus nivolumab in patients with advanced melanoma [80]. The patients received ipilimumab once every 3 weeks for 4 doses and nivolumab once every 3 weeks for 8 doses concurrently. Then, eligible patients were permitted to receive both once every 12 weeks up to 8 doses. Grade 3 to 4 treatmentrelated AEs were seen in 53% of the concurrent-cohort patients but were mild and manageable. The maximum tolerated dose was 3 mg/kg of ipilimumab and 1 mg/kg of nivolumab, a dosing regimen at which 53% of patients showed responses. Recent follow-up surveys confirmed OS to be 94% at 1 year and 88% at 2 years in this cohort. An expansion cohort, with the patients receiving 3 mg/kg of ipilimumab and 1 mg/kg of nivolumab every 3 weeks for 4 doses and 1 mg/kg of nivolumab every 2 weeks until disease progression, is currently being evaluated in a phase II/III study [81]. A phase III trial (NCT01844505) evaluating this combination is currently ongoing (recruitment has been completed).

6. Biomarkers for Predicting Clinical Benefits and Adverse Reactions

Although immune checkpoint inhibitors have shown promising safety and efficacy, to date only a small proportion of patients have achieved long-term survival, with severe irAEs occurring on occasion. Biomarkers predicting clinical benefit may enable physicians to select individualized treatments for their patents and thereby maximize clinical benefits. Thus, there is an urgent need to identify "baseline (pretreatment)" biomarkers predicting responses or toxicities. Several biomarkers for examining T-cell proliferation or activation and other forms of antigen-specific immunity have been assessed in the context of immune checkpoint inhibitors.

Immunohistochemical PD-L1 expression in a tumor specimen is among the potential markers for PD-1-PD-L1directed therapies. In a phase I study of nivolumab, though the data obtained are preliminary, an objective response was seen only in the patients who showed immunohistochemical PD-L1 expression in pretreatment tumor specimens [63]. These observations may support the strategy of selecting PD-L1-positive patients for therapy. However, PD-L1 expression on tumor cells is inducible and is susceptible to influences of the tumor microenvironment. Furthermore, technical advances in PD-L1 immunostaining are still needed. Also, the value of PD-L1 IHC staining as a predictive biomarker for combination therapy with nivolumab plus ipilimumab has yet to be validated [80]. As yet, the applicability and significance of PD-L1 expression as a baseline biomarker must be interpreted with caution and further prospective evaluations are needed, including the results of ongoing randomized

clinical trials that are prospectively evaluating PD-L1 IHC as a companion diagnostic platform (NCT01721746).

Another potential biomarker is pretreatment levels of monocytic myeloid-derived suppressor cells (m-MDSCs) [82, 83]. A recent retrospective study suggested higher pretreatment quantities of Lin⁻CD14⁺HLA-DR^{low/-} m-MDSC to be associated with inferior OS in patients with metastatic melanoma treated with ipilimumab [83].

Recent genetic analysis using whole-exome sequencing showed the significance of somatic mutational load as predictive biomarker of clinical benefit in melanoma patients treated with CTLA-4 blockade. The neopeptide signature associated with clinical response was identified and predicted mutant peptides were verified to activate patient T cell *in vitro* [84].

Other potential predictive/prognostic biomarkers include the gene expression profiles obtained employing tumor biopsies [85, 86], CRP level [87], absolute lymphocyte and eosinophil counts [88], and LDH levels [89]. These possibilities await further research.

7. Conclusion

Immune checkpoint inhibitors have opened a new era of cancer immunotherapy. Since the FDA approval was obtained for the anti-CTLA-4 monoclonal antibody ipilimumab, several large-scale clinical trials have evaluated new agents both alone and in combinations with other conventional or new therapies. Future challenges include exploring new target molecules and immune cells, optimizing dosing regimens and combination therapies, validating the safety and efficacy of these novel treatment strategies in many other malignancies, establishing an immunomonitoring system to be applied during therapy, and identifying biomarkers predicting clinical responses and toxicities. Active, ongoing investigations are anticipated to provide further clinical benefits for patients with cancers that are currently refractory to treatment.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- L. Chen and D. B. Flies, "Molecular mechanisms of T cell costimulation and co-inhibition," *Nature Reviews Immunology*, vol. 13, no. 4, pp. 227–242, 2013.
- [2] R. D. Schreiber, L. J. Old, and M. J. Smyth, "Cancer immunoediting: integrating immunity's roles in cancer suppression and promotion," *Science*, vol. 331, no. 6024, pp. 1565–1570, 2011.
- [3] D. B. Page, M. A. Postow, M. K. Callahan, J. P. Allison, and J. D. Wolchok, "Immune modulation in cancer with antibodies," *Annual Review of Medicine*, vol. 65, pp. 185–202, 2014.
- [4] D. M. Pardoll, "The blockade of immune checkpoints in cancer immunotherapy," *Nature Reviews Cancer*, vol. 12, no. 4, pp. 252– 264, 2012.
- [5] F. S. Hodi, S. J. O'Day, D. F. McDermott et al., "Improved survival with ipilimumab in patients with metastatic melanoma,"

- [6] C. Robert, L. Thomas, I. Bondarenko et al., "Ipilimumab plus dacarbazine for previously untreated metastatic melanoma," *The New England Journal of Medicine*, vol. 364, no. 26, pp. 2517– 2526, 2011.
- [7] A. Ribas, R. Kefford, M. A. Marshall et al., "Phase III randomized clinical trial comparing tremelimumab with standardof-care chemotherapy in patients with advanced melanoma," *Journal of Clinical Oncology*, vol. 31, no. 5, pp. 616–622, 2013.
- [8] S. L. Topalian, M. Sznol, D. F. McDermott et al., "Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab," *Journal of Clinical Oncology*, vol. 32, no. 10, pp. 1020–1030, 2014.
- [9] J. S. Weber, R. R. Kudchadkar, B. Yu et al., "Safety, efficacy, and biomarkers of nivolumab with vaccine in ipilimumab-refractory or -naive melanoma," *Journal of Clinical Oncology*, vol. 31, no. 34, pp. 4311–4318, 2013.
- [10] J. S. Weber, S. D'Angelo, F. S. Hodi et al., "A phase 3 randomized, open-label study of nivolumab (anti-PD-1; BMS-936558; ONO-4538) versus investigator's choice chemotherapy (ICC) in patients with advanced melanoma after prior anti-CTLA-4 therapy," ESMO Meeting Abstracts, 2014.
- [11] C. Robert, G. V. Long, B. Brady et al., "Nivolumab in previously untreated melanoma without BRAF mutation," *The New England Journal of Medicine*, 2014.
- [12] M. B. Atkins, R. R. Kudchadkar, M. Sznol et al., "Phase 2, multicenter, safety and efficacy study of pidilizumab in patients with metastatic melanoma," in *Proceedings of the ASCO Annual Meeting*, vol. 32, p. 9001, 2014.
- [13] O. Hamid, C. Robert, A. Daud et al., "Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma," *The New England Journal of Medicine*, vol. 369, no. 2, pp. 134–144, 2013.
- [14] N. A. Rizvi, E. B. Garon, A. Patnaik et al., "Safety and clinical activity of MK-3475 as initial therapy in patients with advanced non-small cell lung cancer (NSCLC)," *Proceedings of the ASCO Meeting Abstracts*, vol. 32, p. 8007, 2014.
- [15] T. Y. Seiwert, B. Burtness, J. Weiss et al., "A phase Ib study of MK-3475 in patients with human papillomavirus (HPV)-associated and non-HPV-associated head and neck (H/N) cancer," ASCO Meeting Abstracts, vol. 32, article 6011, 2014.
- [16] K. Muro, Y. Bang, V. Shankaran et al., "A phase 1b study of pembrolizumab (Pembro; MK-3475) in patients (Pts) with advanced gastric cancer," ESMO Meeting Abstracts, 2014.
- [17] J. R. Brahmer, S. S. Tykodi, L. Q. M. Chow et al., "Safety and activity of anti-PD-L1 antibody in patients with advanced cancer," *The New England Journal of Medicine*, vol. 366, no. 26, pp. 2455–2465, 2012.
- [18] T. Powles, N. J. Vogelzang, G. D. Fine et al., "Inhibition of PD-L1 by MPDL3280A and clinical activity in pts with metastatic urothelial bladder cancer (UBC)," ASCO Meeting Abstracts, vol. 32, article 5011, 2014.
- [19] J. Lutzky, S. J. Antonia, A. Blake-Haskins et al., "A phase 1 study of MEDI4736, an anti–PD-L1 antibody, in patients with advanced solid tumors," *ASCO Meeting Abstracts*, vol. 32, abstract 3001, 2014.
- [20] C. R. Heery, G. H. O'Sullivan Coyne, R. A. Madan et al., "Phase I open-label, multiple ascending dose trial of MSB0010718C, an anti-PD-L1 monoclonal antibody, in advanced solid malignancies," in *Proceedings of the ASCO Meeting Abstracts*, vol. 32, p. 3064, 2014.

- [21] C. E. Rudd, A. Taylor, and H. Schneider, "CD28 and CTLA-4 coreceptor expression and signal transduction," *Immunological Reviews*, vol. 229, no. 1, pp. 12–26, 2009.
- [22] L. Chen, "Co-inhibitory molecules of the B7-CD28 family in the control of T-cell immunity," *Nature Reviews Immunology*, vol. 4, no. 5, pp. 336–347, 2004.
- [23] C. J. Nirschl and C. G. Drake, "Molecular pathways: coexpression of immune checkpoint molecules: Signaling pathways and implications for cancer immunotherapy," *Clinical Cancer Research*, vol. 19, no. 18, pp. 4917–4924, 2013.
- [24] D. M. Sansom and L. S. K. Walker, "The role of CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4) in regulatory T-cell biology," *Immunological Reviews*, vol. 212, pp. 131–148, 2006.
- [25] K. Wing, Y. Onishi, P. Prieto-Martin et al., "CTLA-4 control over Foxp3⁺ regulatory T cell function," *Science*, vol. 322, no. 5899, pp. 271–275, 2008.
- [26] L. S. K. Walker, "Treg and CTLA-4: two intertwining pathways to immune tolerance," *Journal of Autoimmunity*, vol. 45, pp. 49– 57, 2013.
- [27] D. R. Leach, M. F. Krummel, and J. P. Allison, "Enhancement of antitumor immunity by CTLA-4 blockade," *Science*, vol. 271, no. 5256, pp. 1734–1736, 1996.
- [28] A. van Elsas, A. A. Hurwitz, and J. P. Allison, "Combination immunotherapy of B16 melanoma using anti-cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) and granulocyte/ macrophage colony- stimulating factor (GM-CSF)-producing vaccines induces rejection of subcutaneous and metastatic tumors accompanied by autoimmune depigmentation," *Journal* of Experimental Medicine, vol. 190, no. 3, pp. 355–366, 1999.
- [29] S. A. Quezada, T. R. Simpson, K. S. Peggs et al., "Tumor-reactive CD4⁺ T cells develop cytotoxic activity and eradicate large established melanoma after transfer into lymphopenic hosts," *The Journal of Experimental Medicine*, vol. 207, no. 3, pp. 637– 650, 2010.
- [30] K. S. Peggs, S. A. Quezada, C. A. Chambers, A. J. Korman, and J. P. Allison, "Blockade of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of anti-CTLA-4 antibodies," *The Journal of Experimental Medicine*, vol. 206, no. 8, pp. 1717–1725, 2009.
- [31] T. R. Simpson, F. Li, W. Montalvo-Ortiz et al., "Fc-dependent depletion of tumor-infiltrating regulatory t cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma," *The Journal* of *Experimental Medicine*, vol. 210, no. 9, pp. 1695–1710, 2013.
- [32] A. Hoos, R. Ibrahim, A. Korman et al., "Development of ipilimumab: contribution to a new paradigm for cancer immunotherapy," *Seminars in Oncology*, vol. 37, no. 5, pp. 533–546, 2010.
- [33] E. J. Lipson and C. G. Drake, "Ipilimumab: an anti-CTLA-4 antibody for metastatic melanoma," *Clinical Cancer Research*, vol. 17, no. 22, pp. 6958–6962, 2011.
- [34] G. Q. Phan, J. C. Yang, R. M. Sherry et al., "Cancer regression and autoimmunity induced by cytotoxic T lymphocyteassociated antigen 4 blockade in patients with metastatic melanoma," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 14, pp. 8372–8377, 2003.
- [35] A. V. Maker, G. Q. Phan, P. Attia et al., "Tumor regression and autoimmunity in patients treated with cytotoxic T lymphocyteassociated antigen 4 blockade and interleukin 2: a phase I/II study," *Annals of Surgical Oncology*, vol. 12, no. 12, pp. 1005–1016, 2005.

- [36] E. J. Small, N. S. Tchekmedyian, B. I. Rini, L. Fong, I. Lowy, and J. P. Allison, "A pilot trial of CTLA-4 blockade with human anti-CTLA-4 in patients with hormone-refractory prostate cancer," *Clinical Cancer Research*, vol. 13, no. 6, pp. 1810–1815, 2007.
- [37] S. M. Ansell, S. A. Hurvitz, P. A. Koenig et al., "Phase I study of ipilimumab, an anti-CTLA-4 monoclonal antibody, in patients with relapsed and refractory B-cell non-Hodgkin lymphoma," *Clinical Cancer Research*, vol. 15, no. 20, pp. 6446–6453, 2009.
- [38] L. Fong, S. S. Kwek, S. O'Brien et al., "Potentiating endogenous antitumor immunity to prostate cancer through combination immunotherapy with CTLA4 blockade and GM-CSF," *Cancer Research*, vol. 69, no. 2, pp. 609–615, 2009.
- [39] F. S. Hodi, M. Butler, D. A. Oble et al., "Immunologic and clinical effects of antibody blockade of cytotoxic T lymphocyteassociated antigen 4 in previously vaccinated cancer patients," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 8, pp. 3005–3010, 2008.
- [40] J. C. Yang, M. Hughes, U. Kammula et al., "Ipilimumab (anti-CTLA4 antibody) causes regression of metastatic renal cell cancer associated with enteritis and hypophysitis," *Journal of Immunotherapy*, vol. 30, no. 8, pp. 825–830, 2007.
- [41] J. Weber, J. A. Thompson, O. Hamid et al., "A randomized, double-blind, placebo-controlled, phase II study comparing the tolerability and efficacy of ipilimumab administered with or without prophylactic budesonide in patients with unresectable stage III or IV melanoma," *Clinical Cancer Research*, vol. 15, no. 17, pp. 5591–5598, 2009.
- [42] S. J. O'Day, M. Maio, V. Chiarion-Sileni et al., "Efficacy and safety of ipilimumab monotherapy in patients with pretreated advanced melanoma: a multicenter single-arm phase II study," *Annals of Oncology*, vol. 21, no. 8, pp. 1712–1717, 2010.
- [43] J. D. Wolchok, B. Neyns, G. Linette et al., "Ipilimumab monotherapy in patients with pretreated advanced melanoma: a randomised, double-blind, multicentre, phase 2, dose-ranging study," *The Lancet Oncology*, vol. 11, no. 2, pp. 155–164, 2010.
- [44] E. M. Hersh, S. J. O'Day, J. Powderly et al., "A phase II multicenter study of ipilimumab with or without dacarbazine in chemotherapy-naïve patients with advanced melanoma," *Investigational New Drugs*, vol. 29, no. 3, pp. 489–498, 2011.
- [45] K. Margolin, M. S. Ernstoff, O. Hamid et al., "Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial," *The Lancet Oncology*, vol. 13, no. 5, pp. 459–465, 2012.
- [46] C. Robert, D. Schadendorf, M. Messina, F. S. Hodi, and S. O'Day, "Efficacy and safety of retreatment with ipilimumab in patients with pretreated advanced melanoma who progressed after initially achieving disease control," *Clinical Cancer Research*, vol. 19, no. 8, pp. 2232–2239, 2013.
- [47] V. Chiarion-Sileni, J. Pigozzo, P. A. Ascierto et al., "Ipilimumab retreatment in patients with pretreated advanced melanoma: the expanded access programme in Italy," *British Journal of Cancer*, vol. 110, no. 7, pp. 1721–1726, 2014.
- [48] D. McDermott, J. Haanen, T.-T. Chen, P. Lorigan, and S. O'Day, "Efficacy and safety of ipilimumab in metastatic melanoma patients surviving more than 2 years following treatment in a phase III trial (MDX010-20)," *Annals of Oncology*, vol. 24, no. 10, pp. 2694–2698, 2013.
- [49] D. Schadendorf, C. Robert, J. S. Weber et al., "Pooled analysis of long-term survival data from phase II and phase III trials of ipilimumab in metastatic or locally advanced, unresectable melanoma," *European Journal of Cancer*, vol. 49, article S11, 2013.

- [50] A. Ribas, D. C. Hanson, D. A. Noe et al., "Tremelimumab (CP-675,206), a cytotoxic T lymphocyte—associated antigen 4 blocking monoclonal antibody in clinical development for patients with cancer," *The Oncologist*, vol. 12, no. 7, pp. 873–883, 2007.
- [51] A. Ribas, L. H. Camacho, G. Lopez-Berestein et al., "Antitumor activity in melanoma and anti-self responses in a phase I trial with the anti-cytotoxic T lymphocyte-associated antigen 4 monoclonal antibody CP-675,206," *Journal of Clinical Oncology*, vol. 23, no. 35, pp. 8968–8977, 2005.
- [52] L. H. Camacho, S. Antonia, J. Sosman et al., "Phase I/II trial of tremelimumab in patients with metastatic melanoma," *Journal* of Clinical Oncology, vol. 27, no. 7, pp. 1075–1081, 2009.
- [53] A. Ribas, B. Comin-Anduix, B. Chmielowski et al., "Dendritic cell vaccination combined with CTLA4 blockade in patients with metastatic melanoma," *Clinical Cancer Research*, vol. 15, no. 19, pp. 6267–6276, 2009.
- [54] K. Y. Chung, I. Gore, L. Fong et al., "Phase II study of the anti-cytotoxic T-lymphocyte-associated antigen 4 monoclonal antibody, tremelimumab, in patients with refractory metastatic colorectal cancer," *Journal of Clinical Oncology*, vol. 28, no. 21, pp. 3485–3490, 2010.
- [55] J. M. Kirkwood, P. Lorigan, P. Hersey et al., "Phase II trial of tremelimumab (CP-675,206) in patients with advanced refractory or relapsed melanoma," *Clinical Cancer Research*, vol. 16, no. 3, pp. 1042–1048, 2010.
- [56] B. Sangro, C. Gomez-Martin, M. de la Mata et al., "A clinical trial of CTLA-4 blockade with tremelimumab in patients with hepatocellular carcinoma and chronic hepatitis C," *Journal of Hepatology*, vol. 59, no. 1, pp. 81–88, 2013.
- [57] C. Ralph, E. Elkord, D. J. Burt et al., "Modulation of lymphocyte regulation for cancer therapy: a phase II trial of tremelimumab in advanced gastric and esophageal adenocarcinoma," *Clinical Cancer Research*, vol. 16, no. 5, pp. 1662–1672, 2010.
- [58] D. L. Barber, E. J. Wherry, D. Masopust et al., "Restoring function in exhausted CD8 T cells during chronic viral infection," *Nature*, vol. 439, no. 7077, pp. 682–687, 2006.
- [59] L. M. Francisco, V. H. Salinas, K. E. Brown et al., "PD-L1 regulates the development, maintenance, and function of induced regulatory T cells," *The Journal of Experimental Medicine*, vol. 206, no. 13, pp. 3015–3029, 2009.
- [60] T. Okazaki, Y. Tanaka, R. Nishio et al., "Autoantibodies against cardiac troponin I are responsible for dilated cardiomyopathy in PD-1-deficient mice," *Nature Medicine*, vol. 9, no. 12, pp. 1477– 1483, 2003.
- [61] W. Zou and L. Chen, "Inhibitory B7-family molecules in the tumour microenvironment," *Nature Reviews Immunology*, vol. 8, no. 6, pp. 467–477, 2008.
- [62] S. L. Topalian, C. G. Drake, and D. M. Pardoll, "Targeting the PD-1/B7-H1(PD-L1) pathway to activate anti-tumor immunity," *Current Opinion in Immunology*, vol. 24, no. 2, pp. 207–212, 2012.
- [63] S. L. Topalian, F. S. Hodi, J. R. Brahmer et al., "Safety, activity, and immune correlates of anti-PD-1 antibody in cancer," *The New England Journal of Medicine*, vol. 366, no. 26, pp. 2443– 2454, 2012.
- [64] J. R. Brahmer, C. G. Drake, I. Wollner et al., "Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates," *Journal of Clinical Oncology*, vol. 28, no. 19, pp. 3167–3175, 2010.
- [65] E. J. Lipson, W. H. Sharfman, C. G. Drake et al., "Durable cancer regression off-treatment and effective reinduction therapy with

an anti-PD-1 antibody," *Clinical Cancer Research*, vol. 19, no. 2, pp. 462–468, 2013.

- [66] S. N. Gettinger, F. A. Shepherd, S. J. Antonia et al., "First-line nivolumab (anti-PD-1; BMS-936558, ONO-4538) monotherapy in advanced NSCLC: Safety, efficacy, and correlation of outcomes with PD-L1 status," in *Proceedings of the ASCO Annual Meeting*, ASCO Meeting Abstracts 32:8024, 2014.
- [67] B. Hardy, L. Indjiia, G. Rodionov, A. Raiter, and A. Inbal, "Treatment with BAT monoclonal antibody decreases tumor burden in a murine model of leukemia/lymphoma," *International Journal of Oncology*, vol. 19, no. 5, pp. 897–902, 2001.
- [68] B. Hardy, Y. Niv, L. Fadaeev, and A. Raiter, "BAT mAb induces lymphopoiesis in nude mice," *International Immunology*, vol. 17, no. 5, pp. 615–619, 2005.
- [69] R. Berger, R. Rotem-Yehudar, G. Slama et al., "Phase i safety and pharmacokinetic study of CT-011, a humanized antibody interacting with PD-1, in patients with advanced hematologic malignancies," *Clinical Cancer Research*, vol. 14, no. 10, pp. 3044–3051, 2008.
- [70] P. Armand, A. Nagler, E. A. Weller et al., "Disabling immune tolerance by programmed death-1 blockade with pidilizumab after autologous hematopoietic stem-cell transplantation for diffuse large B-cell lymphoma: results of an international phase II trial," *Journal of Clinical Oncology*, vol. 31, no. 33, pp. 4199– 4206, 2013.
- [71] J. R. Westin, F. Chu, M. Zhang et al., "Safety and activity of PD1 blockade by pidilizumab in combination with rituximab in patients with relapsed follicular lymphoma: a single group, open-label, phase 2 trial," *The Lancet Oncology*, vol. 15, no. 1, pp. 69–77, 2014.
- [72] A. Patnaik, S. P. Kang, A. W. Tolcher et al., "Phase I study of MK-3475 (anti-PD-1 monoclonal antibody) in patients with advanced solid tumors," in *Proceedings of the ASCO Annual Meeting*, abstract 2512, May 2012.
- [73] H. Dong, G. Zhu, K. Tamada, D. B. Flies, J. M. A. van Deursen, and L. Chen, "B7-H1 determines accumulation and deletion of intrahepatic CD8⁺ T lymphocytes," *Immunity*, vol. 20, no. 3, pp. 327–336, 2004.
- [74] Y. E. Latchman, S. C. Liang, Y. Wu et al., "PD-L1-deficient mice show that PD-L1 on T cells, antigen-presenting cells, and host tissues negatively regulates T cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 29, pp. 10691–10696, 2004.
- [75] M. J. Butte, M. E. Keir, T. B. Phamduy, A. H. Sharpe, and G. J. Freeman, "Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses," *Immunity*, vol. 27, no. 1, pp. 111–122, 2007.
- [76] J.-J. Park, R. Omiya, Y. Matsumura et al., "B7-H1/CD80 interaction is required for the induction and maintenance of peripheral T-cell tolerance," *Blood*, vol. 116, no. 8, pp. 1291–1298, 2010.
- [77] D. S. Chen, B. A. Irving, and F. S. Hodi, "Molecular pathways: next-generation immunotherapy-inhibiting programmed death-ligand 1 and programmed death-1," *Clinical Cancer Research*, vol. 18, no. 24, pp. 6580–6587, 2012.
- [78] A. J. Furness, F. A. Vargas, K. S. Peggs, and S. A. Quezada, "Impact of tumour microenvironment and Fc receptors on the activity of immunomodulatory antibodies," *Trends in Immunol*ogy, vol. 35, pp. 290–298, 2014.
- [79] R. S. Herbst, M. S. Gordon, G. D. Fine et al., "A study of MPDL3280A, an engineered PD-L1 antibody in patients with locally advanced or metastatic tumors," ASCO Meeting Abstracts, vol. 31, p. 3000, 2013.

- [80] J. D. Wolchok, H. Kluger, M. K. Callahan et al., "Nivolumab plus Ipilimumab in advanced melanoma," *The New England Journal* of *Medicine*, vol. 369, no. 2, pp. 122–133, 2013.
- [81] M. Sznol, H. M. Kluger, M. K. Callahan et al., "Survival, response duration, and activity by BRAF mutation (MT) status of nivolumab (NIVO, anti-PD-1, BMS-936558, ONO-4538) and ipilimumab (IPI) concurrent therapy in advanced melanoma (MEL)," in *Proceedings of the ASCO Annual Meeting*, ASCO Meeting Abstracts 32:LBA9003, 2014.
- [82] C. Meyer, L. Cagnon, C. M. Costa-Nunes et al., "Frequencies of circulating MDSC correlate with clinical outcome of melanoma patients treated with ipilimumab," *Cancer Immunology, Immun*otherapy, vol. 63, no. 3, pp. 247–257, 2014.
- [83] S. Kitano, M. A. Postow, C. G. Ziegler et al., "Computational algorithm-driven evaluation of monocytic myeloid-derived suppressor cell frequency for prediction of clinical outcomes," *Cancer Immunology Research*, vol. 2, no. 8, pp. 812–821, 2014.
- [84] A. Snyder, V. Makarov, T. Merghoub et al., "Genetic basis for clinical response to CTLA-4 blockade in melanoma," *The New England Journal of Medicine*, vol. 371, pp. 2189–2199, 2014.
- [85] O. Hamid, H. Schmidt, A. Nissan et al., "A prospective phase II trial exploring the association between tumor microenvironment biomarkers and clinical activity of ipilimumab in advanced melanoma," *Journal of Translational Medicine*, vol. 9, no. 1, article 204, 2011.
- [86] R.-R. Ji, S. D. Chasalow, L. Wang et al., "An immune-active tumor microenvironment favors clinical response to ipilimumab," *Cancer Immunology, Immunotherapy*, vol. 61, no. 7, pp. 1019–1031, 2012.
- [87] S. Wilgenhof, S. D. Four, F. Vandenbroucke et al., "Single-center experience with ipilimumab in an expanded access program for patients with pretreated advanced melanoma," *Journal of Immunotherapy*, vol. 36, no. 3, pp. 215–222, 2013.
- [88] J. Delyon, C. Mateus, D. Lefeuvre et al., "Experience in daily practice with ipilimumab for thetreatment of patients with metastatic melanoma: anearly increase in lymphocyte and eosinophil countsis associated with improved survival," *Annals* of Oncology, vol. 24, no. 6, pp. 1697–1703, 2013.
- [89] S. Kelderman, B. Heemskerk, H. van Tinteren et al., "Lactate dehydrogenase as a selection criterion for ipilimumab treatment in metastatic melanoma," *Cancer Immunology, Immunotherapy*, vol. 63, no. 5, pp. 449–458, 2014.



The Scientific World Journal



Gastroenterology Research and Practice





Journal of Diabetes Research



Disease Markers



Immunology Research





International Journal of Endocrinology



BioMed **Research International**





Computational and Mathematical Methods in Medicine





Behavioural Neurology



Complementary and Alternative Medicine













Oxidative Medicine and Cellular Longevity