

# Biphasic Effects of Prostaglandin E<sub>2</sub> on the Human Fat Cell Adenylate Cyclase

HORST KATHER and BERND SIMON, *Klinisches Institut für Herzinfarktforschung  
an der Medizinischen Universitätsklinik Heidelberg, D-69  
Heidelberg, West Germany*

**ABSTRACT** Adenylate cyclase of human fat cell ghosts shows a biphasic response towards prostaglandin E<sub>2</sub> with inhibition occurring at nanomolar concentrations of the hormone and stimulation at concentrations beyond 10<sup>-6</sup> mol/liter. The expression of the inhibitory effect is critically dependent on GTP. Under the conditions employed (1 mmol/liter ATP, 5 mmol/liter Mg<sup>2+</sup>, 30°C) the inhibitory component of prostaglandin E<sub>2</sub> became apparent at GTP concentrations exceeding 10<sup>-6</sup> mol/liter. The prostaglandin E<sub>2</sub>-induced inhibition displayed characteristic features of prostaglandin action in intact fat cells with respect to the effective concentrations and degree of inhibition. It is concluded that prostaglandin E<sub>2</sub> is capable of inducing antagonistic effects upon lipolysis via interaction with the membrane-bound adenylate cyclase.

## INTRODUCTION

Prostaglandins are among the most potent inhibitors of hormone-activated lipolysis in adipocytes of various species including man (1). Their inhibitory effect is associated with a decrease of intracellular 3',5'-cyclic AMP (cAMP).<sup>1</sup> Because prostaglandins are synthesized in response to lipolytic hormones (2, 3), it has been proposed that these C-20 fatty acids might act as feedback inhibitors of hormone-activated lipolysis at the level of the membrane-bound adenylate cyclase (4).

One of the major obstacles in elucidating the site of action of prostaglandins in fat cells is that their inhibitory effects have not yet been elicited in cell-free systems. In the rat, various types of prostaglandins had no direct effects on the adenylate cyclase of membrane preparations (1, 5). In other species, the direct effects of the prostaglandins were opposite to those to be expected from the negative feedback con-

cept. We have recently shown that prostaglandins of the E and F type are stimulators of the human fat cell adenylate cyclase when tested over a concentration range of 10<sup>-6</sup>–10<sup>-3</sup> mol/liter (6). Gorman et al. (5) demonstrated that prostaglandin endoperoxide intermediates (0.28–28 μmol/liter) inhibited basal and noradrenaline stimulated adenylate cyclase from rat adipocyte ghosts. In intact cell preparations, however, these intermediates were rapidly converted to prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), which was physiologically more active, but had no direct effects on adenylate cyclase activity (7).

In this report, it is shown that the human fat cell adenylate cyclase exhibits biphasic responses to PGE<sub>2</sub>, with inhibition occurring at submicromolar concentrations and stimulation at higher concentrations of the hormone.

## METHODS

Biopsies of subcutaneous adipose tissue (10–15 g) were obtained from surgical patients not selected on the basis of age, sex, weight, or disease. The subjects were operated after an overnight fast. Anesthesia was induced with a short-acting barbiturate and maintained with halothane, nitrous oxide, and oxygen. The biopsies were usually obtained after the skin incision.

Experimental procedures were the same as described, (8). Fat cells and fat cell ghosts were prepared according to Rodbell (9). The adenylate cyclase activity was determined by the method of Salomon et al. (10) at 30°C. The assay mixture contained 25 mmol/liter Tris-HCl, pH 8.0, 5 mmol/liter MgCl<sub>2</sub>, 20 mmol/liter creatine phosphate, 100 U/ml creatine phosphokinase, 1 mmol/liter cAMP, [ $\alpha$ -<sup>32</sup>P]ATP (1 mmol/liter; 40–50 cpm/pmol), and GTP as stated in the legends to figures and tables. The protein content of the samples was determined by the Lowry et al. (11) method. Statistical analysis was by the Wilcoxon test.

[ $\alpha$ -<sup>32</sup>P]ATP (9–11 Ci/mmol) and [<sup>3</sup>H]cAMP (27 Ci/mmol) were purchased from the Radiochemical Centre Amersham, Bucks, England. Epinephrine-bitartrate and isoproterenol were from Merck AG, Darmstadt, West Germany. Enzymes, coenzymes and nucleotides were from Boehringer GmbH Mannheim, West Germany. PGE<sub>2</sub> was a gift of Dr. Brunnberg, Upjohn GmbH, Heppenheim, West Germany.

Received for publication 2 March 1979.

<sup>1</sup>Abbreviations used in this paper: cAMP, 3',5'-cyclic AMP; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>.

## RESULTS

Fig. 1 shows a dose-response curve for  $\text{PGE}_2$  over a wide range of concentrations ( $10^{-10}$  mol/liter– $10^{-3}$  mol/liter) in the presence of 0.1 mmol/liter GTP. Under these conditions, the hormone displayed biphasic effects with inhibition occurring at  $10^{-9}$ – $10^{-7}$  mol/liter and stimulation at higher concentrations. The inhibitory effect was maximal at  $2.8 \times 10^{-8}$  mol/liter of  $\text{PGE}_2$ . At this concentration the hormone depressed basal enzyme activity by about 40%. The stimulatory effects of  $\text{PGE}_2$  were detectable at concentrations beyond  $2.8 \times 10^{-7}$  mol/liter. The  $\text{PGE}_2$ -induced activation was much more impressive than the inhibitory effect. Activation was maximal at  $2.8 \times 10^{-4}$  mol/liter (five-fold increase of enzyme activity). Half-maximal effects occurred at a  $\text{PGE}_2$ -concentration of about  $2.8 \times 10^{-5}$  mol/liter.

Fig. 2 illustrates the inhibitory effects of two concentrations of  $\text{PGE}_2$  on the dose-response curve for epinephrine. The GTP concentration was 0.1 mmol/liter. At concentrations of  $10^{-9}$  mol/liter and  $10^{-8}$  mol/liter the prostaglandin caused a dose-dependent and parallel rightward shift of the epinephrine dose-response curve, which has also been observed in metabolic experiments designed to investigate the antilipolytic effects of the E-type prostaglandins (12).




FIGURE 1 Dose-response curve for  $\text{PGE}_2$ . A representative experiment out of four is shown. Values are means of triplicate determinations.




FIGURE 2 Effects of nanomolar concentrations of  $\text{PGE}_2$  on the dose-response relationships of epinephrine. Values are means of triplicate determinations.

The  $\text{PGE}_2$ -induced inhibition was critically dependent on the presence of GTP in the assay medium. Fig. 3 shows the effect of the guanine nucleotide on the human fat cell adenylate cyclase activated by  $10^{-5}$  mol/liter of isoproterenol in the absence and presence of 28 nmol/liter of  $\text{PGE}_2$ . Increasing concentrations of GTP induced a biphasic response of the isoproterenol-activated enzyme. Peak activity occurred at a GTP concentration of  $\approx 10^{-4}$  mol/liter, whereas higher concentrations of the nucleotide depressed the rate of




FIGURE 3 GTP-dependence of the  $\text{PGE}_2$ -induced inhibition of the human fat cell adenylate cyclase. A representative experiment out of four is shown. Values are means of triplicate determinations.

cAMP formation relative to peak activity. PGE<sub>2</sub> had no effect at GTP concentrations below 10<sup>-6</sup> mol/liter. Inhibition was most pronounced at GTP concentrations beyond 10<sup>-5</sup> mol/liter.

The inhibitory effects of nanomolar concentrations of PGE<sub>2</sub> were confirmed in six separate experiments carried out with membranes from different subjects. The catecholamine concentrations were chosen such as to elicit submaximal activation. PGE<sub>2</sub> caused a significant inhibition ( $P \leq 0.05$ ) by about 25% in these experiments. Quantitatively and qualitatively similar results were obtained with either isoproterenol or the naturally occurring catecholamine epinephrine (Table I).

## DISCUSSION

The inhibition of the human fat cell adenylate cyclase demonstrated in this report reflects the characteristic features of the antilipolytic effects of the E-prostaglandins in intact fat cells in many aspects. The effects of micromolar concentrations of lipolytic hormones can be inhibited by nanomolar concentrations of the E-type prostaglandins, making this class of hormone's orders of magnitude more potent than agents like nicotinic acid or propranolol (1). The PGE<sub>2</sub> concentrations inducing an inhibition of the human fat cell enzyme are in the physiological range. In addition, the maximal inhibition of hormone-stimulated lipolysis because of E-prostaglandins is only about 50% (12). This applies to the depression of the activity of the human fat cell adenylate cyclase by physiological concentrations of PGE<sub>2</sub> too. The prostaglandins share this latter property, i.e., incomplete inhibition of lipolysis, in common with adenosine (1). Interestingly, adenosine analogues, in a very recent report, have been shown to induce an inhibition of isoproterenol-stimulated rat fat cell adenylate cyclase that was also GTP-dependent (13).

TABLE I  
Inhibition of the Catecholamine-Activated Adenylate Cyclase by Prostaglandin E<sub>2</sub> ( $2.8 \times 10^{-8}$  mol/liter)

Additions	Adenylate cyclase activity*	
	Control†	PGE <sub>2</sub> (28 nmol/liter)‡
	nmol cAMP/mg protein/15 min	
Epinephrine ( $5 \times 10^{-5}$ mol/liter)	1.7±0.2	1.3±0.2§
Isoproterenol ( $1 \times 10^{-5}$ mol/liter)	2.4±0.3	1.9±0.2§

\* The assay mixture contained 0.1 mmol/liter GTP.

† Values are mean±SEM of six separate experiments carried out in triplicate.

§ Significantly lower ( $P \leq 0.05$ ) than the corresponding controls.

The main effect of the prostaglandins appears to be an inhibitory one on the mobilization of free fatty acids and glycerol from adipose tissue. However, these C-20 fatty acids can also stimulate lipolysis under certain conditions suggesting that not only inhibition but also stimulation of the human fat cell adenylate cyclase is of physiological importance in human fat cell function (14, 15). Our results are thus compatible with the concept that the antagonistic effects of the E-type prostaglandins on lipolysis are mediated via interaction with the membrane-bound adenylate cyclase thereby suggesting that the role of these hormones is more complex than assumed by the negative feedback concept. This hypothesis considers only an inhibition of adenylate cyclase but does not take into account the biphasic prostaglandin effects shown in this report. In contrast to the negative feedback concept our results, therefore, support the contention that the actual effects of prostaglandins are the result of a delicately regulated balance between two opposing prostaglandin effects, which is not only dependent on the concentration of these C-20 fatty acids but is also extremely sensitive to modulation by guanine nucleotides.

## ACKNOWLEDGMENTS

The authors wish to thank Mrs. I. Ring for her excellent technical assistance.

This work was supported by grants of the Deutsche Forschungsgemeinschaft, Bonn-Bad Godesberg, West Germany.

## REFERENCES

1. Fain, J. N. 1973. Biochemical aspects of drug and hormone action on adipose tissue. *Pharmacol. Rev.* **25**: 67-118.
2. Dalton, C., and W. C. Hope. 1974. Cyclic AMP regulation of prostaglandin biosynthesis in fat cells. *Prostaglandins*. **6**: 227-242.
3. Chang, J., G. P. Lewis, and P. J. Piper. 1977. The effect of anti-inflammatory drugs on the prostaglandin system in human subcutaneous adipose tissue. *Br. J. Pharmacol.* **61**: 446P-447P.
4. Bergström, S. 1967. Prostaglandins: members of a new hormonal series. *Science (Wash. D. C.)*. **157**: 595-606.
5. Gorman, R. R., M. Hamberg, and B. Samuelsson, 1975. Inhibition of basal and hormone-stimulated adenylate cyclase in adipocyte ghosts by the prostaglandin endoperoxide prostaglandin H<sub>2</sub>. *J. Biol. Chem.* **250**: 6460-6463.
6. Kather, H., and B. Simon. 1977. Adenylate cyclase of human fat cell ghosts: stimulation of enzyme activity by prostaglandins. *J. Cyclic Nucleotide Res.* **3**: 199-206.
7. Fredholm, B. B., and M. Hamberg. 1976. Metabolism and effect of prostaglandin H<sub>2</sub> in adipose tissue. *Prostaglandins*. **11**: 507-518.
8. Kather, H., and B. Simon, 1977. Adenylate cyclase of human fat cell ghosts: stimulation of enzyme activity by parathyroid hormone. *J. Clin. Invest.* **59**: 730-733.
9. Rodbell, M. 1972. Methods for the isolation of rat liver plasma membranes and fat cell "Ghosts": an assay method for adenylate cyclase. In *Methods of Cyclic Nucleotide Research*. M. Chasin, editor. Marcel Dekker, Inc., New York. 101-124.

10. Salomon, Y., C. Londos, and M. Rodbell. 1974. A highly sensitive adenylate cyclase assay. *Anal. Biochem.* **58**: 541–548.
11. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265–275.
12. Stock, K., A. Aulich, and E. Westermann, 1968. Studies on the mechanism of antilipolytic action of prostaglandin E<sub>1</sub>. *Life Sci.* **7**: 113–124.
13. Londos, C., D. M. F. Cooper, W. Schlegel, and M. Rodbell. 1978. Adenosine analogs inhibit adipocyte adenylate cyclase by a GTP-dependent process: Basis for actions of adenosine and methylxanthines on cyclic AMP production and lipolysis. *Proc. Natl. Acad. Sci. U. S. A.* **75**: 5362–5366.
14. Carlson, L. A. 1965. Inhibition of the mobilization of free fatty acids from adipose tissue. *Ann. N. Y. Acad. Sci.* **131**: 119–142.
15. Rosenquist, U., and S. Efendic. 1971. Stimulatory effect of prostaglandin E<sub>1</sub> on noradrenaline-induced lipolysis in subcutaneous adipose tissue from hypothyroid subjects. *Acta Med. Scand.* **190**: 341–345.