

RESEARCH ARTICLE

Open Access

Characterization of resistance to pine wood nematode infection in *Pinus thunbergii* using suppression subtractive hybridization

Tomonori Hirao^{1†}, Eitaro Fukatsu^{2†} and Atsushi Watanabe^{3*†}

Abstract

Background: Pine wilt disease is caused by the pine wood nematode, *Bursaphelenchus xylophilus*, which threatens pine forests and forest ecosystems worldwide and causes serious economic losses. In the 40 years since the pathogen was identified, the physiological changes occurring as the disease progresses have been characterized using anatomical and biochemical methods, and resistant trees have been selected via breeding programs. However, no studies have assessed the molecular genetics, e.g. transcriptional changes, associated with infection-induced physiological changes in resistant or susceptible trees.

Results: We constructed seven subtractive suppression hybridization (SSH) cDNA libraries using time-course sampling of trees inoculated with pine wood nematode at 1, 3, or 7 days post-inoculation (dpi) in susceptible trees and at 1, 3, 7, or 14 dpi in resistant trees. A total of 3,299 sequences was obtained from these cDNA libraries, including from 138 to 315 non-redundant sequences in susceptible SSH libraries and from 351 to 435 in resistant SSH libraries. Using Gene Ontology hierarchy, those non-redundant sequences were classified into 15 subcategories of the biological process Gene Ontology category and 17 subcategories of the molecular function category. The transcriptional components revealed by the Gene Ontology classification clearly differed between resistant and susceptible libraries. Some transcripts were discriminative: expression of antimicrobial peptide and putative pathogenesis-related genes (e.g., PR-1b, 2, 3, 4, 5, 6) was much higher in susceptible trees than in resistant trees at every time point, whereas expression of PR-9, PR-10, and cell wall-related genes (e.g., for hydroxyproline-rich glycoprotein precursor and extensin) was higher in resistant trees than in susceptible trees at 7 and 14 dpi.

Conclusions: Following inoculation with pine wood nematode, there were marked differences between resistant and susceptible trees in transcript diversity and the timing and level of transcripts expressed in common; in particular, expression of stress response and defense genes differed. This study provided new insight into the differences in the physiological changes between resistant and susceptible trees that have been observed in anatomical and biochemical studies.

Background

Pine wilt disease is caused by the pine wood nematode (PWN), *Bursaphelenchus xylophilus*, and was first reported by Tokushige and Kiyohara [1]; this disease threatens pine forests and forest ecosystems around the world and causes significant economic losses [2]. Pine wilt disease is a chronic problem in pine forests (*Pinus*

thunbergii and *Pinus densiflora*) in Japan, where approximately 40,000,000 m³ of pine forests have been blighted by the PWN [3]. Over the past 40 years, public administration and central, prefectural, and city governments have attempted to stem the spread of PWN and pine wilt disease by controlling the pine sawyer beetle (*Monochamus* sp.), the vector of PWN, with insecticides and cutting down infected trees. Additionally, national and prefectural forestry institutes have established breeding programs to develop resistant pine varieties. A breeding project to develop pine varieties resistant to pine wilt disease was started in 1978 in western Japan,

* Correspondence: nabeatsu@affrc.go.jp

† Contributed equally

³Forest Tree Breeding Center, Forestry and Forest Products Research Institute, 3809-1 Ishi, Juo, Hitachi, Ibaraki 319-1301, Japan
Full list of author information is available at the end of the article

and related projects were promoted throughout Japan, excluding Hokkaido Island, as the damage spread. In this breeding project, trees are screened for resistance using an artificial inoculation test that follows a strict protocol; during the first breeding program, which ran from 1978 to 1984, 92 resistant *P. densiflora* individuals were selected from 11,000 candidate trees, and only 16 resistant *P. thunbergii* individuals were selected from 14,000 candidate trees. The breeding projects continued, and 204 resistant *P. densiflora* and 90 resistant *P. thunbergii* individuals were generated. Resistant *P. densiflora* (n = 92) and resistant *P. thunbergii* (n = 16) were ranked with regard to resistance (levels 1-5) based on the survival rate of openly pollinated progeny following inoculation; higher survival rates are thought to indicate greater resistance. Average rates of survival of openly pollinated progeny from the selected pines (i.e., resistant trees) were 64% for *P. densiflora* and 53% for *P. thunbergii*, which was respectively 16% and 40% higher than for unselected populations [4].

Since the causative pathogen was identified [1], many researchers have characterized the physiological changes associated with progression of pine wilt disease, and by the mid-1990s, more than two thousand papers on the disease had been published [5]. Many symptoms associated with PWN infection, including decreased photosynthesis, denaturation of xylem and cortex parenchyma cells, traumatic resin canal formation, cambium destruction, production of phytotoxic substances and ethylene, reduced water potential and transpiration in leaves and heat pulse velocity have been studied (for review, see reference [6]). Based on the anatomical and biochemical evidence gathered during these 40 years, development of symptoms is thought to occur in two stages, early and advanced stages [6]. In the early stage, a small number of nematodes migrate and symptoms such as necrosis and destruction of cortex and phloem tissue and cambium, destruction of cortex resin canals, formation of wound periderm in cortex parenchyma around resin canals, and ethylene release all occur near the inoculation site. In the advanced stage, ethylene production is enhanced and coincides with the broad destruction of cortex parenchyma, cambial destruction, and cavitation-induced embolism of the tracheids in the xylem. The cavitation-induced embolism causes a decrease in leaf water potential and cessation of photosynthesis. After cessation of photosynthesis, severe symptoms develop rapidly with a burst in the nematode population; this population increase results in wilting and eventual tree death. Resistance against PWN infection depends on stopping the symptoms at the early stage or preventing the progression of infection to the advanced stage. While the physiological changes that occur as pine wilt disease progresses have been characterized anatomically

and biochemically, molecular genetic events such as changes in transcript profiles that may be associated with the physiological changes in either resistant or susceptible trees remain poorly understood.

Recently, the gene expression profiles of resistant (resistant variety of *P. thunbergii* 'Sendai-290'; resistant rank 1) and non-resistant Japanese black pine (*P. thunbergii*) trees were assessed using the LongSAGE technique on stems collected 3 days after PWN inoculation [7]. The researchers found that catalase and proteins in the dienelactone hydrolase family were highly expressed in resistant trees, but not non-resistant trees, whereas pathogenesis-related (PR)-1, 2, 3, leucoanthocyanidin dioxygenase and cell wall-related genes were expressed at higher levels in the non-resistant trees. Although the study assessed a difference of one time point in the defense responses of resistant and non-resistant *P. thunbergii* following PWN inoculation, the defense response is continuous, and the differences in resistance and susceptibility do not depend only upon qualitative differences in the activated defense genes, but also on differences in the timing and magnitude of their expression [8]. To characterize the differences in transcript profiles of resistant and susceptible trees as pine wilt symptoms develop, it is necessary to sample resistant and susceptible individuals over a defined time course.

Subtractive suppression hybridization (SSH) is a powerful tool for gene expression profiling that effectively identifies genes differentially expressed under different conditions or in different tissues [9]. This method is relatively simple and easy, it can be used with species for which there is little or no genomic information, and the cDNAs isolated are typically longer than 100 bp and can be effectively annotated using comparative genomics (e.g., BLAST analysis). This method is often used to isolate plant genes specifically expressed in response to pathogen infection and to identify differences in the transcript profiles of infected resistant and susceptible plants [10-14]. SSH selection reduces the cloning of abundantly expressed housekeeping genes or genes commonly expressed in both "tester" and "driver" samples, and therefore normalizes the expressed cDNA profiles during library construction. As a result, SSH selection significantly enhances the chances of cloning differentially expressed genes.

The goal of this study was to identify differences in the transcript profiles of PWN-inoculated *P. thunbergii* to understand the difference in the defense responses of resistant and susceptible individuals. Three important experimental design elements enhanced the clarity and relevance of our findings: 1) We used the most resistant variety of *P. thunbergii*, 'Namikata-73'; resistant rank 5. 2) We sampled inoculated trees (both resistant and susceptible) 1, 3, and 7 dpi before any macroscopic changes

usually occur; resistant trees were also sampled 14 dpi, when macroscopic changes are usually evident. 3) We used SSH, a powerful approach for identifying differentially expressed genes regardless of their abundance.

Results

Analysis of sequences in SSH libraries

Six subtractive libraries constructed from samples taken at three time points, 1, 3, and 7 days dpi, were used to identify genes that are differentially expressed in resistant and susceptible trees as disease symptoms develop following PWN inoculation. In addition, a seventh library contained genes expressed predominantly in resistant trees at 14 dpi compared to susceptible trees at 7 dpi (see Methods). We sequenced nucleotides from the 5'-end of 3,299 cDNA inserts that were recovered in the seven libraries (Table 1). Insert length varied from 100 to 780 bp; the median length ranged from 366 to 429 bp, depending on the library. The redundancy within each library varied from 3.9% to 27.2% in libraries from resistant trees and from 28.4% to 72.8% from susceptible trees. The obtained non-redundant expressed sequence tags (ESTs) ranged from 351 to 435 in libraries from resistant trees and from 138 to 315 in libraries from susceptible trees. Importantly, the overlap between the two libraries at each time point was extremely low; for example, 0.4% of the cDNA inserts were shared between libraries from resistant and susceptible trees sampled at 1 dpi. Similarly, only 3.7% of the cDNA inserts were shared by the library from resistant trees sampled at 14 dpi and from susceptible trees sampled at 7 dpi. This high level of specificity is expected from efficient SSH procedures.

Functional classification of the expressed genes in each SSH library and identification of differentially accumulated genes

In order to annotate the putative gene and functional Gene Ontology (GO) categories for the transcripts of each SSH library, we compared non-redundant ESTs of each SSH library with the GenBank non-redundant and EST databases using the search programs blastx, tblastx, and blastn using the Blast2GO program [15] (Additional file 1). The transitional hit rate of the BLAST analysis and the annotation rate per library are indicated in Table 1. For each library, 98% or more of the clones were matched in a BLAST search using the nr and EST databases at the National Center for Biotechnology Information (NCBI). Similarly, 97% or more of the clones in each library were matched in a BLAST search (blastn) using the Dana-Farber Cancer Institute *Pinus taeda* Gene Index (DFCI PGI, release 8.0). The annotation rates for GO terms resulting from automated and manual annotation varied from 75.45% to 91.00% per library.

Non-redundant ESTs recovered in the seven libraries were classified by function into the three principal Level 1 GO categories: biological process, molecular function, and cellular component. Furthermore, the biological process category was split into 24 subcategories at Level 2, molecular function was split into 61 subcategories at Level 3, and cellular component was split into 39 subcategories at Level 7. The level presented herein corresponds to the depth of hierarchical categories within each principal GO category, and higher levels represent more general classifications. The classification for biological process, divided into 15 major subcategories and 24 lower-level subcategories, is shown in Figure 1a. Of the 15 subcategories under biological process, the subcategory of response to stimulus was present at a higher percentage in the susceptible libraries at each time point, and three categories—cell wall organization or biogenesis, immune system process, multi-organism process—were present at a higher percentage in the susceptible libraries than in the resistant libraries at each time point. In contrast, transcripts related to cellular process and metabolic process were present at a higher percentage in the resistant libraries. In addition, the category of response to stimulus was present at a higher percentage even in the resistant libraries at 7 and 14 dpi. The classification for molecular function, divided into 17 major subcategories and 61 lower-level subcategories, is shown in Figure 1b. At each time point, susceptible trees expressed more transcripts in the four subcategories of carbohydrate binding, enzyme inhibitor activity, hydrolase activity, and pattern binding than did resistant trees, whereas transcripts related to several molecular binding categories such as ion binding and protein binding and to several molecular activity categories such as oxidoreductase activity and transferase activity were present at a higher percentage in the resistant libraries. The classification for cellular component, divided into 18 major subcategories and 39 lower-level subcategories, is shown in Figure 1c. At each time point, susceptible trees expressed more transcripts in the cytoplasmic membrane-bounded vesicle category than did resistant trees. In contrast, the transcripts related to organelle categories such as chloroplast, plastid and mitochondrion were present at a higher percentage in the resistant libraries at each time point.

In order to evaluate the specificity and diversity of the transcripts that were specific in a time course from infected resistant and susceptible trees, ESTs were selected from each SSH library (Table 2, Additional file 2). The libraries from susceptible infected trees showed very limited transcript diversity and were chiefly composed of PR proteins, such as basic PR-1, PR-2 (beta-1,3-glucanase), PR-3 (class I, IV chitinase), PR-4 (chitinase type I & II), PR-5 (thaumatin-like protein), PR-6

Table 1 Characterization of seven SSH libraries

SSH libraries ^a	Total ESTs sequences	Length of sequence (bp)	Contigs	Singletons	Non-redundant ESTs	Redundant (%)	Unique (%)	Hit rate using the BLAST program in Blast2GO tool (%) ^b			Hit rate using the local BLAST program (%)	Automatic annotation rate (%)	Manual annotation rate (%) ^d	Total annotation rate (%)		
								blastx to nrDB ¹⁾	tblastx to nrDB ²⁾	tblastx to estDB ³⁾					blastn to hit estDB ⁴⁾	
Tester	Drivers															
R_1dpi	S_1dpi	437	102-775	14	406	3.9	96.1	71.0	17.1	7.0	3.8	1.1	97.1	65.8	9.7	75.5
S_1dpi	R_1dpi	449	108-777	52	214	40.8	59.2	79.7	10.8	7.5	1.6	0.4	96.9	73.3	6.9	80.2
R_3dpi	S_3dpi	455	102-775	43	344	14.9	85.1	72.4	15.7	7.5	3.5	0.9	97.6	65.1	10.6	75.7
S_3dpi	R_3dpi	440	101-760	43	272	28.4	71.6	81.1	12.2	4.1	1.4	1.2	96.4	77.0	7.4	84.4
R_7dpi	S_7dpi	528	100-767	41	394	17.6	82.4	79.5	10.3	6.9	3.0	0.3	98.0	74.1	7.8	81.9
S_7dpi	R_7dpi	508	104-780	48	90	72.8	27.2	91.8	5.9	1.7	0.6	0.0	99.4	84.7	6.3	91.0
R_14dpi	S_7dpi	482	106-764	52	299	27.2	72.8	73.8	15.3	7.2	3.3	0.4	98.0	66.7	11.0	77.7

^a R = resistant, S = susceptible, dpi = days post-inoculation

^b The edited sequences in each library were compared to the NCBI database using the following algorithms: 1) blastx program against the non-redundant protein database of NCBI with a threshold value of e-6, 2) tblastx program against the nucleotide database of NCBI with a threshold value of e-6, 3) tblastx program against the EST database of NCBI with a threshold value of e-6, 4) blastn program against the EST database of NCBI with a threshold value of e-10

^c The edited sequences in each library were compared to the PGI database (DFCI Pine Gene Index, release 8.0) using the blastn program with a threshold value of e-10

^d Manual annotation based on the annotated gene information in the PGI database




Figure 1 Gene Ontology mapping for SSH libraries from resistant and susceptible trees. EST distribution in the categories (a) Biological process, (b) Molecular function, and (c) Cellular component.

(type II proteinase inhibitor family protein), and antimicrobial peptide. In contrast, the ESTs from resistant trees were more diverse. Transcripts encoding metallothionein-like protein, heat shock protein 70 (HSP70),

xyloglucan endotransglycosylase (XET), and cytochrome P450 (CYP450) were discernible in the resistant library at 1 dpi and 3 dpi. The libraries from resistant trees sampled at 7 dpi and 14 dpi had transcripts encoding

Table 2 Characteristic ESTs within each SSH library

Putative gene categories based on BLAST annotations	Number of ESTs from each SSH library*						
	R_1dpi	R_3dpi	R_7dpi	R_14dpi	S_1dpi	S_3dpi	S_7dpi
PR-1 family	0	0	1	0	2	6	1
PR-2 family (beta-1, 3-glucanase)	1	0	1	2	7	12	21
PR-3 family (class i chitinase)	2	1	4	1	6	2	1
PR-3 family (class iv chitinase)	1	0	2	5	9	9	58
PR-4 family	1	0	1	1	14	5	56
PR-5 family (thaumatin-like)	0	0	12	62	55	52	129
PR-6 family (proteinase-inhibitor)	0	0	2	10	56	52	130
PR-9 family (peroxidase)	1	0	5	1	3	5	0
PR-10 family (ribonuclease-like)	0	8	57	1	24	11	20
Antimicrobial peptide	0	0	0	1	27	13	8
Cytochrome P450	4	4	1	0	0	0	0
Extensin	0	0	0	27	10	5	4
Heat Shock Proteins	10	4	3	22	1	0	0
Hydroxyprolinerich glycoprotein precursor	0	0	6	16	1	1	1
metallothionein-like protein	7	26	4	0	1	0	0
Xyloglucan endotransglycosylase	5	4	2	0	0	1	0
Rate of occupying a library (%)	6.8	10.1	18.8	30.5	47.7	39.2	84.0

* R = resistant, S = susceptible, dpi = days post-inoculation

PR-5, PR-9 (peroxidase), PR-10 (ribonuclease-like), hydroxyproline-rich glycoprotein (HRGP) precursor, and extensin.

Validation of differential expression using selected SSH clones and quantitative real-time PCR (qRT-PCR)

To validate the results of the SSH procedures, expression of 16 ESTs recovered in one or more of the seven libraries was assayed using qRT-PCR; samples for the qRT-PCR analysis were collected from resistant and susceptible trees at 0, 1, 3, 7, and 14 dpi. (Figure 2). Furthermore, expression of 16 ESTs of mock samples at each time points was also assayed using qRT-PCR to monitor expression changes induced either by PWN infection or by cutting (Additional file 3). The results of the qRT-PCR analysis and SSH were consistent.

Expression of 12 of 16 ESTs was clearly upregulated following inoculation with PWN. Among the upregulated ESTs, expression of putative PR-1b, PR-2, PR-3 (class I & IV chitinase), PR-4, PR-5, PR-6, and antimicrobial peptide was much higher in the susceptible trees than in the resistant trees at each time point. Expression of these ESTs was much higher in susceptible trees at 1 dpi and expression had increased further by 7 dpi. In the resistant trees, expression of these ESTs was at a relatively low level at 1 dpi and 3 dpi, but was abundant at 7 dpi and 14 dpi. ESTs encoding PR-9, PR-10, HRSP, and extensin were discernible in resistant SSH libraries. Expression of PR-9 and PR-10 was much higher in the

susceptible trees at 1 dpi and 3 dpi; however, levels of these two transcripts were much higher in resistant trees at 7 dpi, though it was lower at 14 dpi than 7 dpi. HRGP and extensin transcript accumulation in resistant trees was high at 14 dpi, although expression of both ESTs was higher in susceptible trees than in resistant trees at 1 dpi, 3 dpi, and 7 dpi.

Based on qRT-PCR analysis, three ESTs—CYP450, metallothionein-like protein, and XET—were downregulated in both resistant and susceptible trees after PWN inoculation, with greater downregulation in susceptible trees. Transcript levels of these three ESTs decreased in susceptible trees within 1 dpi, and were even lower at 3 and 7 dpi. The levels of these ESTs in resistant trees at 7 and 14 dpi were similar to those in susceptible trees at 7 dpi. The transcript level of HSP70 was lower than the other downregulated genes, and its decrease was more moderate in resistant trees, with an expression pattern similar to that of mock samples.

Discussion

We sequenced cDNAs in seven SSH libraries to characterize transcriptional differences between resistant and susceptible *P. thunbergii* trees in response to inoculation with PWN. In susceptible trees, transcript diversity was statistically lower than in resistant trees at the three time points jointly tested, 1, 3, and 7 dpi. In susceptible trees, several transcripts encoding pathogenesis related proteins were present at a higher constitutive level than




Figure 2 Quantitative real-time PCR of transcripts differentially expressed in resistant and susceptible trees following PWN inoculation. The putative functional genes from (a) to (h) were clearly discernible ESTs in susceptible SSH libraries. The putative functional genes from (i) to (p) were clearly discernible ESTs in resistant SSH libraries. Elongation factor 1-alpha (EF1a) was used as the reference gene, and the data were calibrated relative to the transcript levels in resistant trees prior to nematode inoculation (at 0 days). The data are presented as the mean \pm S.D. of three replicates. Means designed by the same letter did not significantly differ at $P < 0.05$ according to Tukey's HSD test.

in resistant trees. In resistant trees at 14 dpi, several transcripts encoding cell wall proteins were identified. The results of the SSH approach were validated by qRT-PCR. We clearly demonstrated that transcript composition, temporal changes, and levels of gene expression involved in the stress/defense response to PWN inoculation in resistant trees differs from susceptible trees.

There was a significant difference in transcript diversity between resistant and susceptible trees after infection with PWN. The percentage of unique sequences in susceptible libraries ranged from 71.59 to 27.17%, whereas the percentage of unique sequences in resistant libraries ranged from 72.82 to 96.11%. Although transcript diversity of the susceptible library at 3 dpi was higher than the susceptible libraries at 1 and 7 dpi, transcript diversity in resistant libraries was higher than susceptible libraries at all time points after PWN infection. According to the GO classification of the differentially expressed transcripts, a large percentage in susceptible trees was involved in stress/defense response categories such as the response to stimulus, multi-organism process, and immune system process within the biological process category and the enzyme inhibitor activity, carbohydrate binding, pattern binding, and hydrolase activity within the molecular function category. Furthermore, a large percentage of transcripts in the cellular component category were in the cytoplasmic membrane-bounded vesicle subcategory. On the other hand, a large percentage of transcripts in resistant SSH libraries was categorized into the cellular process, metabolic process and response to stimulus subcategories of the biological process category and in the ligand binding and transferase activity subcategories of the molecular function category. Many transcripts recovered in libraries from resistant trees were assigned to the cellular component subcategory and further classified into organelle subcategories of plastid, chloroplast, and mitochondrion. The difference in transcript diversity between libraries from resistant and susceptible trees revealed by the GO classification indicated a qualitative difference in the stress/defense response of resistant and susceptible trees to PWN infection.

Resistance and susceptibility do not depend only upon qualitative differences in the activated defense genes, but also on differences in the timing and magnitude of their expression [8]. The gene regulation patterns of pathogenesis related defense proteins such as PR-1b, PR-2, PR-3, PR-4, PR-5, PR-6 and antimicrobial peptide indicated temporal and quantitative differences between resistant and susceptible trees in response to PWN infection. In regulating the plant defense response, most pathogenesis related proteins are induced through the action of the signaling compounds salicylic acid (SA),

jasmonic acid (JA), or ethylene [16]. PR-1b, PR-2 and PR-5 genes are SA-responsive genes and also indicators of systemic acquired resistance; additionally, the PR-6 gene is a JA and ethylene responsive gene [17]. Although the relationship between phytohormones and the defense response in the PWN-nematode interaction is not clear from this study, it is interesting that expression of pathogenesis related genes associated with phytohormones such as SA and JA/ethylene and with antimicrobial activities were induced more quickly and to a higher level in susceptible than in resistant trees.

The same three phytohormones, SA, JA, and ethylene, are important for both basal and R-gene mediated defense responses to foliar pathogens and insects. The JA and ethylene signaling pathways seem to work synergistically, whereas the SA and JA/ethylene signaling pathways inhibit each other, and negative cross-talk exists between SA and JA/ethylene signaling pathways [18]. In interactions between *Hero A*-resistant tomatoes and cyst nematode, expression of SA-responsive genes PR-1 and PR-5 is a hallmark of the resistant cultivar, and expression of JA-dependent PR-6 is higher in the susceptible cultivar, indicating that SA plays some role in the resistance to the nematode and that JA and ethylene in susceptible tomatoes are likely to interfere with the SA-inducible resistance pathway [19,20]. The involvement of SA in resistance and expression of SA-responsive genes in resistant plants has been observed in interactions between other plant species and nematodes (e.g., an *Arabidopsis thaliana* mutant [21] and a root knot nematode resistant peanut [13]). In our study, the higher induction of both SA-responsive genes such as PR-1b, PR-2, PR-5 and JA/ethylene-responsive genes such as PR-6 in susceptible trees indicates that the defense response mediated by these phytohormones was not very effective in controlling PWN infestations.

We detected notable levels of putative HRGPs, extensin and peroxidase (PR-9) in resistant trees at 7 dpi and 14 dpi, though expression in susceptible trees was higher than in resistant trees at 1 and 3 dpi. Cell wall-mediated resistance is the first line of plant defense against pathogens, and the components of plant cell wall are modified by production of reactive oxygen species (ROS) such as H₂O₂ during attack by pathogens [22]. The structural cell wall proteins extensin and HRGPs play an essential role in biotic and abiotic stress responses due to their oxidative cross-linking, which contributes to the strength of cell walls and is catalyzed by an oxidizing system based on peroxidase and H₂O₂ [23-25]. The cross-linking of HRGPs and extensin is involved in cell wall-mediated resistance and has a major role in arresting pathogens at the site of entry [23]; these proteins accumulate in the walls of a number of plant species during interactions with microbes [26].

A number of reports describe the response of plant cell wall HRGPs and extensin to pathogens such as viruses, bacteria, and fungi (for review, see reference [27]). In plant-nematode interactions, high extensin gene expression was observed in the cortical region of tobacco at 7 and 14 days after inoculation with root knot nematodes [28]. In *rhg1* resistant soybean, extensin was identified as one of the genes characteristically expressed in syncytia after inoculation with soybean cyst nematode, indicating that altered cell wall composition is important in the defense response [29]. Also, in *Mi* resistant tomato, extensin and glycosyltransferase may play a role in cell wall synthesis, which is an essential defense against root knot nematode [30,31]. In anatomical studies of PWN infection, Ishida et al. [32] inoculated virulent nematode (*B. xylophilus*) isolate S6-1 and avirulent nematode (*B. mucronatus*) isolate B. m to Japanese black pine, and observed accumulation of lignin- and suberin-like substances around the resin canals in the cortex with both nematode isolates at 7 dpi. Kusumoto et al. [33] also inoculated a virulent nematode (*B. xylophilus*) isolate, Ka-4, to Japanese black pine and found more frequent accumulation of phenolic compounds around the cortex resin canals in resistant trees at 7 dpi and 14 dpi after inoculation with PWN (*B. xylophilus*), and suggested that this accumulation was a very effective defense against infection due to restricting PWN migration. Although the relationship between HRGPs or extensin and other substances in the cell wall was not examined in this study, it is possible that upregulation of expression of cell wall-related genes such as those for HRGPs or extensin and oxidative cross-linking of these proteins by peroxidases plays a role in the effective defense response of trees resistant to PWN infection at 7 dpi and 14 dpi.

PR-10 was also one of the characteristically significantly upregulated genes in libraries in the 7 dpi subtraction library from resistant trees. Although the biological function of PR-10 remains unclear, many PR-10 genes are upregulated when plants are exposed to abiotic stresses, such as SA, CuCl₂, H₂O₂, cold, darkness and wounding [34], and biotic stresses, such as viruses [35], bacteria [34,36], fungi [37-40] and insects [41,42]. We observed synchronized expression of PR10 with peroxidase in resistant trees, which indicates this gene may be induced by ROS such as H₂O₂. However, PR10 (CpPRI) acts against a digestive proteinase from the root knot nematode *Meloidogyne incognita* [43]. Therefore, PR10 may act as a proteinase against cellulases, beta-1,3-glucanase, and pectate lyases secreted from PWN [44-46].

Heat shock protein (HSP) ESTs were characteristically recovered in libraries from resistant trees. In particular, stable HSP70 expression in infected resistant trees was

validated by qRT-PCR. HSP family members, which consist of HSP70, HSP60 and HSP90, are required for folding of nascent proteins and intracellular transportation in addition to stress responses, and are collectively called molecular chaperones [47]. In the interaction between soybean and soybean cyst nematode, Klink et al. [48] observed the induction of HSP70 and ROS responsive genes such as lipoxygenase and superoxidase dismutase isolated from 3 dpi syncytial cells showing an incompatible response to soybean cyst nematode infection, and suggested that HSP70 may be involved in maintaining a properly functioning environment for other defense responses. In the present study, it is unclear how HSP70 is involved in the defense response to PWN infection.

Three ESTs—putatively encoding CYP450, metallothionein-like protein, and XET—were detected in resistant SSH libraries at 1 dpi and 3 dpi, which depend on the genes significantly downregulated in susceptible trees at 1 and 3 dpi. In plants, CYP450 monooxygenases play paramount roles in the synthesis of lignin intermediates, sterols, terpenes, flavonoids, isoflavonoids, furanocoumarins, and a variety of other secondary plant products [49]. In conifers, diterpene resin acids are prominent defense compounds against insect pests and pathogens in conifers [50-52], and CYP450s are involved in the formation of a suite of diterpene resin acids of conifer oleoresin; they oxidize abietadienol and abietadienal to abietic acid in vitro and in vivo [53,54]. Keeling and Bohlmann [50,51] indicated that oleoresin may contain specific components that are toxic or deterrent to insect herbivores or may affect adults or broods physiologically and thus prevent successful colonization or reproduction. The downregulation of CYP450 observed in the present study may cause a reduction in diterpene resin acids in pine trees infested with PWN. Consequently, PWN may expand its invasion and habitat, and the rapid reduction in CYP450 expression in susceptible trees may trigger PWN expansion.

Metallothioneins are involved in ROS scavenging, and in rice, downregulation of metallothionein expression is observed during the oxidative burst phase in elicitor-treated cells, and metallothionein expression is important for defense signaling [55,56]. The metallothionein expression we observed indicated that ROS accumulation and defense signaling may have been induced by 1 dpi in susceptible trees infected with PWN, whereas it may not have been induced much in resistant trees 3 dpi later; this provides evidence for rapid induction of defense response genes such as those encoding pathogenesis related proteins in susceptible trees.

XET action is thought to regulate wall loosening during turgor-driven expansion by rearranging load-bearing xyloglucan cross-links between cellulose microfibrils,

and its activity and expression have been detected in growing tissues [57-64]. We found that XET was down-regulated in both resistant and susceptible trees following PWN infection, and its regulation was induced relatively early, by 1 dpi. These findings suggest that the expansion of cell walls in xylem, phloem or both is inhibited by the downregulated XET during PWN infection; alternatively, the cell wall may be immobilized by the cross-linking of HRGPs or extensin.

In pine-nematode interactions, Myers [65] and Futai [66] suggested that invasion and rapid migration of a few mobile parasites through tree tissues invokes an innate hypersensitive reaction such as death of the parenchyma, production of toxins, and leakage of oleoresins and other material into tracheids. Furthermore, the population of PWN spreads throughout the whole body, and a series of hypersensitive reactions eventually leads to tree death in susceptible pine species. In this study, defense response genes, antimicrobial peptide, SA-responsive genes such as PR-1b, PR-2, PR-5 and JA/ET-responsive genes such as PR-6 were induced more quickly and to a higher level in susceptible than in resistant trees. These defense responses in susceptible trees would not be effective in controlling PWN nematode infestations, and defense signaling induced within the tree may then induce a series of hypersensitive reactions through cellular interactions that subsequently lead to death, as Myers [65] and Futai [66] suggested. In contrast, a moderate hypersensitive reaction along with upregulation of pathogenesis related genes followed by upregulation of cell wall-related genes contributing to the strength of cell walls would be a very effective defense against PWN infection, because these events might restrict PWN migration.

Conclusions

This is the first study to assess the difference in EST transcript diversity of activated defense genes and differences in the timing and magnitude of expression of these genes between resistant and susceptible *P. thunbergii* trees following PWN inoculation. In susceptible trees after PWN inoculation, pathogenesis related genes and antimicrobial-related genes were rapidly induced to high levels within 1 dpi; this finding indicated that a hypersensitive reaction-like response was induced in susceptible trees and supported the hypothesis of Myers [63] and Futai [64]. In contrast, a moderate defense response mediated by pathogenesis related protein expression followed by significant upregulation of cell wall-related genes induced by ROS was a very effective defense against PWN infection. The ESTs generated in our study will certainly advance understanding of defense mechanisms against PWN at the transcriptional level in other varieties or other *Pinus* species.

Methods

Plant materials and nematode inoculation

A resistant tree of 'Namikata 73,' which is the most highly resistant variety of the 16 resistant varieties selected out of 15,000 individuals from 1990 to 1998 [4], was planted in the Forest Products Research Institute, Forest Tree Breeding Center (FFPRI-FTBC) in Ibaraki, Japan. A susceptible tree of the variety 'Kataura 1,' selected as a plus-tree for growth traits, was also in the FFPRI-FTBC. Both clones were grafts obtained from the original trees at the FFPRI-FTBC in 2005. The PWN used in this study was the Ka-4 isolate, which has been used in pine wilt disease resistance breeding projects since 2003.

Inoculation with PWN was conducted on July 1, 2007. In four susceptible clones and four resistant clones, 2 cm at the tip of the main stem was cut off, the cut edge was quickly crushed with pliers, and 10,000 nematodes that had been suspended in 100 μ l sterile water were injected into the cut edge. As a mock sample, sterile water (without nematodes) was injected into the cut edge of the main stem of another four susceptible clones and four resistant clones. Stem tissue of inoculated samples and mock samples was collected 5 cm below the inoculated stem apex at 1, 3, 7, and 14 dpi. A 2 cm segment of stem was cut, frozen immediately in liquid nitrogen, and stored at -80°C .

RNA isolation

Total RNA was isolated from 1.5 g of stem that included bark and wood tissue using the RNeasy plant mini kit (QIAGEN) following the protocol supplied by the manufacturer. RNA concentration was determined using a GeneQuant pro spectrophotometer (Amersham Biosciences). Integrity of the RNA was tested using the Agilent 2100 bioanalyzer (Agilent Technologies).

SSH library construction, DNA sequencing, data analysis and dbEST submission

Six subtractive libraries were constructed from samples taken at three time points; specifically, two libraries—one from resistant trees and one from susceptible trees—were constructed from samples taken 1, 3, or 7 days dpi. Forward libraries containing genes expressed predominantly in resistant trees were constructed by subtracting driver RNA sampled from susceptible trees from tester RNA sampled from resistant trees, and reverse libraries containing genes expressed predominantly in susceptible trees were constructed by subtracting driver RNA sampled from resistant trees from tester RNA sampled from susceptible trees. Additionally, a seventh library was constructed by subtracting driver RNA sampled from susceptible trees 7 dpi from tester RNA sampled from resistant trees 14 dpi; the driver RNA

Table 3 Primers used in this study

Putative gene function	Forward primer	Reverse primer	Fragment length (bp)	GenBank Acc.
PR-1b family	5' -TGCCCTTCAGGTAAATCGT-3'	5' -GCGGGTCGTAGTTGCAGATAA-3'	125	FY841927
PR-2 family (Beta-1,3-glucanase)	5' -CGACAACATTGCCCCCTTCT-3'	5' -CTGCAGCGCGGTTTGAATAT-3'	130	FY843702
PR-3 family (class I chitinase)	5' -ACCTACAGCGCCTTCATTGC-3'	5' -TGTGTTTCATGCGACGTTT-3'	120	FY841849
PR-3 family (class Iv chitinase)	5' -CCATCGAAGCCCAGGTAATTT-3'	5' -AGCCGGGAAGCAATATTATGGT-3'	90	FY843470
PR-4 family	5' -CCCCGTTACTGTCAATTGCAT-3'	5' -AAAGCGTGACGGTGCAT-3'	90	FY841704
PR-5 family (thaumatin-like)	5' -GAACCAGTGCCCATACACAGTCT-3'	5' -CCTGCGGCAACGTTAAAAGTC-3'	96	FY842709
PR-6 family (proteinase inhibitor)	5' -TGCTGGCGCATCTATTTTA-3'	5' -TAACACCTGCGCAAATGCA-3'	90	FY843534
PR-9 family (peroxidase)	5' -ACACCACCGTGCTGGACATT-3'	5' -GTGCGGGAGTCGGTGTAGAG-3'	118	FY842918
PR-10 family (ribonuclease-like)	5' -TGCTCAAGTGAGGCAAGGA-3'	5' -AAGCGACAATTCAGGCAAAAC-3'	90	FY842956
Antimicrobial peptide	5' -GCGTTGCTCATACCCGTTTT-3'	5' -GCAGCACTTAGCACTGGATGAA-3'	90	FY841562
Cytochrome P450	5' -AACATGTCTGCAGCACGAA-3'	5' -GTGCACCGCAAGTAAACCAA-3'	95	FY841345
Extensin	5' -CGAATGTAATCCGAAGTTGCA-3'	5' -CCATCCCAAACCACAGTCT-3'	110	FY844277
Heat shock protein 70	5' -AACACCACCATTCCCACCAA-3'	5' -CGAATTTGCCGAGCAGGTTA-3'	130	FY841300
Hydroxyproline-rich glycoprotein precursor	5' -GAGAAACTGGCACCGTCTTAGGA-3'	5' -ACCTCCCCTCCATCTCACA-3'	140	FY843962
Metallothionein-like protein	5' -TCAGGCTGCTGCGTTATTG-3'	5' -TGTCAGCGCAGTCACAATTTG-3'	120	FY842178
xyloglucan endotransglycosylase	5' -TCTGCGCCCTACTTTTCC-3'	5' -AGCTGGGCGATTGATCATGT-3'	121	FY842425
Elongation factor-1 alpha	5' -GGGAAGCCACCCAAAGTTTT-3'	5' -TACATGGGAAGACGCCGAAT-3'	160	FY842441

The putative functional genes from (a) to (h) were clearly discernible ESTs in susceptible libraries. The putative functional genes from (i) to (p) were clearly discernible ESTs in resistant trees. Elongation factor 1-alpha (EF1a) was used as the reference gene, and the data were calibrated relative to the transcript levels in resistant trees prior to nematode infection (at 0 dpi). The data are presented as the mean \pm S.D. of three replicates. Means designed by the same letter did not significantly differ at $P < 0.05$ according to Tukey's HSD test

samples were taken at 7 dpi because the susceptible trees had died by day 14. SSH libraries were constructed using a SuperSMART cDNA Synthesis kit (Clontech) and a Clontech PCR-Select cDNA Subtraction kit (Clontech). The SSH products were purified using a QIAquick PCR purification kit (QIAGEN) and ligated into the pT7Blue vector (Merck4Biosciences). Blue/white selection was conducted on plates containing ampicillin, isopropyl-D-thiogalactopyranoside and X-gal. Clones were randomly selected and single-pass sequenced using a primer that recognizes vector sequences from the 5' -end of the inserts. On average, 500 clones were sequenced per library using an ABI 3130xl DNA Analyzer (Applied Biosystems). The resulting sequences were trimmed and edited manually to identify the cloning vector sequences, adaptor sequences used in the SSH

procedure, and regions of low-quality sequence using Sequencher 4.10.1 software (Gene Codes Corp.). Quality sequences greater than 100 bp were selected for further analysis. A total of 3,299 sequences were at least 100 bp and these sequences were submitted to the GenBank EST database with the GenBank accession numbers FY841122 to FY844420. To determine the number of contigs and EST singletons in each library, the ESTs were assembled using Sequencher 4.10.1 by the requirement for at least 98% identity over each 20 bp continuous sequence.

EST similarity search and functional assignments

A similarity search and functional annotation were performed for the EST singletons in each library using online version of the BLAST2GO program (BLAST2GO

2006; [15]). The thresholds used with the BLAST algorithms were as follows: (1) blastx comparison with the non-redundant protein database of NCBI with a threshold value of e-6; (2) tblastx comparison with the nucleotide database of NCBI with a threshold value of e-6; (3) tblastx comparison with the EST database of NCBI with a threshold value of e-6; (4) blastn comparison with the EST database of NCBI with a threshold value of e-10. Additionally, the ESTs in each library were compared with 69,968 EST sequences in the DFCI Pinus Gene index release 8.0 using an E value cut-off of e-10. The ESTs were assigned to functional categories using the Blast2GO program, and manual annotations were based on the result of BLAST analysis using the PGI database. The Blast2GO program extracts the GO terms associated with homologies identified with NCBI's QBLAST and returns a list of GO annotations represented as hierarchical categories of increasing specificity. Therefore, the "level" presented in this study corresponds to the depth of hierarchical categories in each principal GO category, with the topmost Level 1 representing the most general classification (principal GO categories) of biological process, molecular function, and cellular component.

Real-time qRT-PCR

Primer pairs were designed for each sequence using Primer Express software v3.0 (Applied Biosystems) and following the manufacturer's guidelines for primer design (Table 3). For SYBR Green real-time RT-PCR assays, the amplification efficiency of all primer pairs was optimized with genomic DNA from resistant and susceptible trees using the StepOnePlus Real-Time PCR System (Applied Biosystems).

RT-PCR was performed using the total RNA used to make the SSH libraries. For the mock samples at each time point and for the reference sample (without nematodes or water), total RNA was extracted from stem tissues and used for RT-PCR. Total RNA (500 ng in a final volume of 20 μ L) was reverse-transcribed using the PrimeScript II 1st strand cDNA synthesis kit (TaKaRa) according to the manufacturer's protocol. Real-time quantitative PCR was performed using the *Power* SYBR Green PCR Master Mix (Applied Biosystems) on the StepOnePlus Real-Time PCR System (Applied Biosystems). PCR mixtures were prepared according to the manufacturer's instructions and contained 300 nM of both the forward and reverse gene-specific primers and 4 μ L of the 20-fold diluted reverse transcription reaction (total 5 ng) in a final volume of 20 μ L. All reactions were heated to 95°C for 10 min; this denaturation step was followed by 40 cycles of 95°C for 15 s and 60°C for 1 min. The PCR products were subjected to melting

curve analysis; the conditions were incubation at 60-95°C with a temperature increment of 0.3°C s⁻¹. Elongation factor 1-alpha was used as the reference gene for normalizing the transcript profiles. The real-time PCR data were calibrated against the transcript levels in resistant tree stem samples prior to nematode exposure; the 2^{- $\Delta\Delta$ Ct} method was used to quantify relative transcript abundance [67]. All assays were carried out in triplicate, and the data are presented as means \pm S.D. of three replicates. The specificity of each amplification was checked by melting analysis and agarose gel electrophoresis of the amplified products.

Additional material

Additional file 1: Summary of BLAST search results for ESTs among 7 SSH libraries.

Additional file 2: Summary of BLAST search results for specific ESTs in the expression analysis.

Additional file 3: Quantitative real-time PCR of transcripts differentially expressed in uninfected resistant and susceptible clones. The putative functional genes from (a) to (h) were clearly discernible ESTs in susceptible libraries. The putative functional genes from (i) to (p) were clearly discernible ESTs in resistant trees. Elongation factor 1-alpha (EF1a) was used as the reference gene, and the data were calibrated relative to the transcript levels in resistant trees prior to nematode infection (at 0 dpi). The data are presented as the mean \pm S. D. of three replicates. Means designed by the same letter did not significantly differ at $P < 0.05$ according to Tukey's HSD test.

Acknowledgements

We thank Dr. Dai Kusumoto and Dr. Toshihiro Yamada at Tokyo University for helpful discussions, comments, and advice on our research; we also thank our colleagues in the field management section of FTBC for the production and cultivation of plant materials. This work was funded by a grant from the Japan Society for the Promotion of Science (No. 20380085).

Author details

¹Forest Bio-research Center, Forestry and Forest Products Research Institute, 3809-1 Ishi, Juo, Hitachi, Ibaraki 319-1301, Japan. ²Kyushu Regional Breeding Office, Forest Tree Breeding Center, Forestry and Forest Products Research Institute, 2320-5 Suya, Goshi, Kumamoto 860-0081, Japan. ³Forest Tree Breeding Center, Forestry and Forest Products Research Institute, 3809-1 Ishi, Juo, Hitachi, Ibaraki 319-1301, Japan.

Authors' contributions

TH prepared the RNA preparation, sequenced the ESTs, analyzed the EST data, and drafted the manuscript. EF helped with data analysis of the ESTs and prepared the DDBJ GenBank submissions. AW conceived of the project, carried out the inoculation of PWN and sampling, and helped to draft the manuscript. All authors assisted with manuscript preparation and read and approved the final draft.

Received: 11 September 2011 Accepted: 24 January 2012

Published: 24 January 2012

References

1. Tokushige Y, Kiyohara T: *Bursaphelenchus* sp. in the wood of dead pine trees. *J Jpn Forest Soc* 1969, **51**:193-195.
2. Wang Z, Wang CY, Fang ZM, Zhang DL, Liu L, Lee MR, Li Z, Li JJ, Sung CK: Advances in research of pathogenic mechanism of pine wilt disease. *Afr J Microbiol Res* 2010, **4**:437-442.

3. Forestry Agency, Ministry of Agriculture, Forestry and Fisheries of Japan: **Annual Report on Trends of Forest and Forestry-Fiscal Year 2009**. Tokyo, 2010, (in Japanese).
4. Toda T: **Studies on the breeding for resistance to the pine wilt disease in *Pinus densiflora* and *P. thunbergii***. *Bull Forest Tree Breed Center* 2004, **20**:83-217, (In Japanese with English summary).
5. Kishi Y: *The Pine Wood Nematode and the Japanese Pine Sawyer* Tokyo: Thomas Co. Ltd; 1995.
6. Fukuda K: **Physiological process of the symptom development and resistance mechanism in pine wilt disease**. *J Forest Res* 1997, **2**:171-181.
7. Nose M, Shiraishi S: **Comparison of the gene expression profiles of resistant and non-resistant Japanese black pine inoculated with pine wood nematode using a modified LongSAGE technique**. *Forest Pathol* 2010, **41**:143-155.
8. Thompson GA, Goggin FL: **Transcriptomics and functional genomics of plant defense induction by phloem-feeding insects**. *J Exp Bot* 2006, **57**:755-766.
9. Diatchenko L, Lau YF, Campbell AP, Chenchik A, Moqadam F, Huang B, Lukyanov S, Lukyanov K, Gurskaya N, Sverdlov ED, Siebert PD: **Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries**. *Proc Natl Acad Sci USA* 1996, **93**:6025-6030.
10. Gibly A, Bonshtien A, Balaji V, Debbie P, Martin GB, Sessa G: **Identification and expression profiling of tomato genes differentially regulated during a resistance response to *Xanthomonas campestris* pv. *vesicatoria***. *Mol Plant Microbe Interact* 2004, **17**:1212-1222.
11. Degenhardt J, Al-Masri AN, Kurcuoglu S, Szankowski I, Gau AE: **Characterization by suppression subtractive hybridization of transcripts that are differentially expressed in leaves of apple scab-resistant and susceptible cultivars of *Malus domestica***. *Mol Gen Genomics* 2005, **273**:326-335.
12. Lin H, Doddapaneni H, Takahasi Y, Walker MA: **Comparative analysis of ESTs involved in grape responses to *Xylella fastidiosa* infection**. *BMC Plant Biol* 2007, **7**:8.
13. Tirumalaraju SV, Jain M, Gallo M: **Differential gene expression in roots of nematode-resistant and -susceptible peanut (*Arachis hypogaea*) cultivars in response to early stage of peanut root-knot nematode (*Meloidogyne arenaria*) parasitization**. *J Plant Physiol* 2011, **168**:481-492.
14. Shin H, Lee H, Woo KS, Noh EW, Koo YB, Lee KI: **Identification of genes upregulated by pinewood nematode inoculation in Japanese red pine**. *Tree Physiol* 2009, **29**:411-421.
15. Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M: **Blast2GO: a universal tool for annotation, visualization, and analysis in functional genomics research**. *Bioinformatics* 2005, **21**:3674-3676.
16. van Loon LC, Rep M, Pieterse CMJ: **Significance of inducible defense-related proteins in infected plants**. *Annu Rev Phytopathol* 2006, **44**:135-162.
17. Ohtsubo N, Mitsuhashi H, Koga M, Seo S, Ohashi Y: **Ethylene promotes the necrotic lesion formation and basic PR gene expression in TMV-infected tobacco**. *Plant Cell Physiol* 1999, **40**:808-817.
18. Glazebrook J: **Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens**. *Annu Rev Phytopathol* 2005, **43**:205-227.
19. Sobczak M, Avrova A, Jupowicz J, Phillips MS, Ernst K, Kumar A: **Characterization of susceptibility and resistance responses to potato cyst nematode (*Globodera* spp.) infection of tomato lines in the absence and presence of the broad-spectrum nematode resistance *Hero* gene**. *Mol Plant Microbe Interact* 2005, **18**:158-168.
20. Uehara T, Sugiyama S, Matsuura H, Arie T, Masuta C: **Resistant and susceptible responses in tomato to cyst nematode are differentially regulated by salicylic acid**. *Plant Cell Physiol* 2010, **51**:1524-1536.
21. Wubben MEJ, Jin J, Baum TJ: **Cyst nematode parasitism of *Arabidopsis thaliana* is inhibited by salicylic acid (SA) and elicits uncoupled SA-independent pathogenesis-related gene expression in roots**. *Mol Plant Microbe Interact* 2008, **21**:424-432.
22. Hückelhoven R: **Cell wall-associated mechanisms of disease resistance and susceptibility**. *Annu Rev Phytopathol* 2007, **45**:101-127.
23. Deepak S, Shailasree S, Kini RK, Muck A, Mithöfer A, Shetty SH: **Hydroxyproline-rich glycoproteins and plant defense**. *J Phytopathol* 2010, **158**:585-593.
24. Pereira CS, Ribeiro JML, Vatulescu AD, Findley K, MacDougall AJ, Jackson PAP: **Extensin network formation in *Vitis vinifera* callus cells in an essential and causal event in rapid H₂O₂-induced reduction in primary cell wall hydration**. *BMC Plant Biol* 2011, **11**:106.
25. Barceló AR, Gómez Ros LV: **Reactive oxygen species in plant cell walls**. In *Reactive Oxygen Species in Plant Signaling*. Edited by: Del Rio LA, Puppo A. Verlag Berlin Heidelberg: Springer; 2009:73-93.
26. Dixon RA, Harrison MJ: **Activation, structure, and organization of genes involved in microbial defense in plants**. *Adv Genet* 1990, **28**:165-234.
27. Esquerré-Tugayé Marie-Thérèse, Campargue C, Mazau D: **The response of plant cell wall hydroxyproline-rich glycoproteins to microbial pathogens and their elicitors**. In *Pathogenesis-Related Proteins in Plants*. Edited by: Datta SK, Muthukrishnan S. Boca Raton, London, New York: CRC Press LLC; 1999:157-170.
28. Niebel A, De Almeida Engler J, Türe C, Engler G, Van Montagu M, Gheysen G: **Induction patterns of an extensin gene in tobacco upon nematode infection**. *Plant Cell* 1993, **5**:1697-1710.
29. Matsye PD, Kumar R, Hosseini P, Jones CM, Tremblay A, Alkharouf NW, Matthews BF, Klink VP: **Mapping cell fate decisions that occur during soybean defense responses**. *Plant Mol Biol* 2011, **77**:513-528.
30. Lambert KN, Ferrie BJ, Nombela G, Brenner ED, Williamson VM: **Identification of genes whose transcripts accumulate rapidly in tomato after root-knot nematode infection**. *Physiol Mol Plant Pathol* 1999, **55**:341-348.
31. Schaff JE, Nielsen DM, Smith CP, Scholl EH, Bird DM: **Comprehensive transcriptome profiling in tomato reveals a role for glycosyltransferase in *Mi*-mediated nematode resistance**. *Plant Physiol* 2007, **144**:1079-1092.
32. Ishida K, Hogetsu T, Fukuda K, Suzuki K: **Cortical responses in Japanese black pine to attack by the pine wood nematode**. *Can J Bot* 1993, **71**:1399-1405.
33. Kusumoto D, Yonemichi T, Murata M, Watanabe A, Isoda K, Hirao T, Yamada T: **Histological observations on host responses and nematode distribution in resistant pine trees infected with pine wood nematode, *Bursaphelenchus xylophilus***. *Tree Forest Health* 2010, **14**:98-100, (in Japanese).
34. Xie YR, Chen ZY, Brown RL, Bhatnagar D: **Expression and functional characterization of two pathogenesis-related protein 10 genes from *Zea mays***. *J Plant Physiol* 2010, **167**:121-130.
35. Xu P, Blancaflor EB, Roossinck MJ: **In spite of induced multiple defense responses, tomato plants infected with cucumber mosaic virus and D satellite RNA succumb to systemic necrosis**. *Mol Plant Microbe Interact* 2003, **16**:467-476.
36. Robert N, Ferran J, Breda C, Coutos-Thevenot P, Boulay M, Buffard D, Esnault R: **Molecular characterization of the incompatible interaction of *Vitis vinifera* leaves with *Pseudomonas syringae* pv. *pis*: Expression of genes coding for stilbene synthase and class 10 PR protein**. *Eur J Plant Pathol* 2001, **107**:249-261.
37. Jwa NS, Agrawal GK, Rakwal R, Park CH, Agrawal VP: **Molecular cloning and characterization of a novel jasmonate-inducible pathogenesis-related class10 protein gene, *JlOsPR10*, from rice (*Oryza sativa* L.) seedling leaves**. *Biochem Biophys Res Commun* 2001, **286**:973-983.
38. Steiner-Lange S, Fischer A, Boettcher A, Rouhara I, Liedgens H, Schmelzer E, Knogge W: **Differential defense reactions in leaf tissues of barley in response to infection by *Rhynchosporium secalis* and to treatment with a fungal avirulence gene product**. *Mol Plant Microbe Interact* 2003, **16**:893-902.
39. Mould MJ, Xu T, Barbara M, Iscove NN, Heath MC: **cDNAs generated from individual epidermal cells reveal that differential gene expression predicting subsequent resistance or susceptibility to rust fungal infection occurs prior to the fungus entering the cell lumen**. *Mol Plant Microbe Interact* 2003, **16**:835-845.
40. Chen ZY, Brown RL, Damann KE, Cleveland TE: **PR10 expression in maize and its effect on host resistance against *Aspergillus flavus* infection and aflatoxin production**. *Mol Plant Pathol* 2010, **11**:69-81.
41. Divol F, Vilaine F, Thibivilliers S, Amselem J, Palauqui JC, Kusiak C, Dinant S: **Systemic response to aphid infestation by *Myzus persicae* in the phloem of *Apium graveolens***. *Plant Mol Biol* 2005, **57**:517-540.
42. Zhu-Salzman K, Salzman RA, Ahn JE, Koiwa H: **Transcriptional regulation of sorghum defense determinants against a phloem-feeding aphid**. *Plant Physiol* 2004, **134**:420-431.
43. Andrade LB, Oliveria AS, Ribeiro JK, Kiyota S, Vasconcelos IM, de Oliveria JT, de Sales MP: **Effects of a novel pathogenesis-related class 10 (PR-10) protein from *Crotalaria pallida* roots with papain inhibitory activity**

- against root-knot nematode *Meloidogyne incognita*. *J Agric Food Chem* 2010, **58**:4145-4152.
44. Kikuchi T, Jones JT, Aikawa T, Kosaka H, Ogura N: A family of glycosyl hydrolase family 45 cellulases from the pine wood nematode *Bursaphelenchus xylophilus*. *FEBS Lett* 2004, **572**:201-205.
45. Kikuchi T, Shibuya H, Jones JT: Molecular and biochemical characterization of an endo-beta-1,3-glucanase from the pinewood nematode *Bursaphelenchus xylophilus* acquired by horizontal gene transfer from bacteria. *Biochem J* 2005, **389**:117-125.
46. Kikuchi T, Shibuya H, Aikawa T, Jones JT: Cloning and characterization of pectate lyases expressed in the esophageal gland of the pine wood nematode *Bursaphelenchus xylophilus*. *Mol Plant Microbe Interact* 2006, **19**:280-287.
47. Kanzaki H, Saitoh H, Ito A, Fujisawa S, Kamoun S, Katou S, Yoshioka H, Terauchi R: Cytosolic HSP90 and HSP70 are essential components of INF1-mediated hypersensitive response and non-host resistance to *Pseudomonas cichorii* in *Nicotiana benthamiana*. *Mol Plant Pathol* 2003, **4**:383-391.
48. Klink VP, Overall CC, Alkharouf NW, MacDonald MH, Matthews BF: Laser capture microdissection (LCM) and comparative microarray expression analysis of syncytial cells isolated from incompatible and compatible soybean (*Glycine max*) roots infected by the soybean cyst nematode (*Heterodera glycines*). *Planta* 2007, **226**:1389-1409.
49. Schuler MA: Plant cytochrome P450 monooxygenases. *Crit Rev Plant Sci* 1996, **15**:235-284.
50. Keeling CJ, Bohlmann J: Genes, enzymes and chemicals of terpenoid diversity in the constitutive and induced defense of conifers against insects and pathogens. *New Phytol* 2006, **170**:657-675.
51. Keeling CJ, Bohlmann J: Diterpene resin acids in conifers. *Phytochemistry* 2006, **67**:2415-2423.
52. Ro DK, Bohlmann J: Diterpene resin acid biosynthesis in loblolly pine (*Pinus taeda*): Functional characterization of abietadiene/levopimaradiene synthase (*PtTPS-LAS*) cDNA and subcellular targeting of *PtTPS-LAS* and abietadienol/abietadienol oxidase (*PtAO*, *CYP720B1*). *Phytochemistry* 2006, **67**:1572-1578.
53. Funk C, Croetau R: Diterpenoid resin acid biosynthesis in conifers: characterization of two cytochrome P450-dependent monooxygenases and an aldehyde dehydrogenase involved in abietic acid biosynthesis. *Arch Biochem Biophys* 1994, **308**:258-266.
54. Ro DK, Arimura GI, Lau SYW, Piers E, Bohlmann J: Loblolly pine abietadienol/abietadienol oxidase *PtAO* (*CYP702B1*) is a multifunctional, multisubstrate cytochrome P450 monooxygenase. *Proc Natl Acad Sci USA* 2005, **102**:8060-8065.
55. Wong HL, Sakamoto T, Kawasaki T, Umemura K, Shimamoto K: Down-regulation of metallothionein, a reactive oxygen scavenger, by the small GTPase *OsRac1* in rice. *Plant Physiol* 2004, **135**:1447-1456.
56. Akimoto-Tomiya C, Sakata K, Yazaki J, Nakamura K, Fujii F, Shimbo K, Yamamoto K, Sasaki T, Kishimoto N, Kikuchi S, Shibuya N, Minami E: Rice gene expression in response to N-acetylchitoooligosaccharide elicitor: comprehensive analysis by DNA microarray with randomly selected ESTs. *Plant Mol Biol* 2003, **52**:537-551.
57. Fry SC, Smith RC, Renwick KF, Martin DJ, Hodge SK, Matthews KJ: Xyloglucan endotransglycosylase, a new wall-loosening enzyme activity in plants. *Biochem J* 1992, **282**:821-828.
58. Pritchard J, Hetherington PR, Fry SC, Tomos AD: Xyloglucan endotransglycosylase activity, microfibril orientation and profiles of cell wall properties along growing regions of maize roots. *J Exp Bot* 1993, **44**:1281-1289.
59. Palmer SJ, Davies WJ: An analysis of relative elemental growth rate, epidermal cell size and xyloglucan endotransglycosylase activity through the growing zone of ageing maize leaves. *J Exp Bot* 1996, **47**:339-347.
60. Burstin J: Differential expression of two barley XET-related genes during coleoptile growth. *J Exp Bot* 2000, **51**:847-852.
61. Uozu S, Tanaka-Ueguchi M, Kitano H, Hattori K, Matsuoka M: Characterization of XET-related genes of rice. *Plant Physiol* 2000, **122**:853-859.
62. Vissenberg K, Martinez-Vilchez IM, Verbelen J-P, Miller JG, Fry SC: In vivo colocalization of xyloglucan endotransglycosylase activity and its donor substrate in the elongation zone of *Arabidopsis* roots. *Plant Cell* 2000, **12**:1229-1237.
63. Vissenberg K, Fry SC, Verbelen JP: Root hair initiation is coupled to a highly localized increase of xyloglucan endotransglycosylase action in *Arabidopsis* roots. *Plant Physiol* 2001, **127**:1125-1135.
64. Lee J, Burns TH, Light G, Sun Y, Fokar M, Kasukabe Y, Fujisawa K, Maekawa Y, Allen RD: Xyloglucan endotransglycosylase/hydrolase genes in cotton and their role in fiber elongation. *Planta* 2010, **232**:1191-1205.
65. Myers RF: Pathogenesis in pine wilt caused by pinewood nematode, *Bursaphelenchus xylophilus*. *J Nematol* 1988, **20**:236-244.
66. Futai K: Pine wilt is an epidemic disease in forests: notes on the interrelationship of forest microbes (tentative translation by authors). Tokyo: Bun-ichi sogo shuyuppan; 2003, (in Japanese).
67. Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCt} method. *Methods* 2001, **25**:402-408.

doi:10.1186/1471-2229-12-13

Cite this article as: Hirao et al.: Characterization of resistance to pine wood nematode infection in *Pinus thunbergii* using suppression subtractive hybridization. *BMC Plant Biology* 2012 **12**:13.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

