

[Chem. Pharm. Bull.]
25(11)3122—3124(1977)

UDC 547.751.02.04 : 543.422.25.06

New Indole Trimer¹⁾

When treated with *p*-toluenesulfonic acid in benzene, indole itself formed a trimer which was different from the known trimer.

The structure of the new trimer was established.

Keywords—trimerization; structure establishment; acid catalyzed rearrangement; indole; Vilsmeier reaction

It is well known that indole itself was easily polymerized under various acid conditions to give a dimer and/or a trimer.²⁾ A sole compound (**1**) has been known as a indole trimer until now, and its structure (**1**) was proposed by Smith³⁾ and Noland.⁴⁾ Recently, in the course of studies on the advanced Fischer indolization,⁵⁾ we occasionally obtained another new indole trimer (**2**). In this paper, we wish to report the structural establishment of **2**.

Refluxing a benzene solution of indole in the presence of TsOH for 3 hr gave colourless dimorphic crystals, mp 105—107° or mp 186.5—187.2°, [IR(CHCl₃): 3475 (NH)] in 39.7% yield. Dimorphism of **2** was confirmed by the cross-seeding experiment. The molecular formula C₂₄H₂₁N₃ of **2** was shown by the facts that the elemental analysis agreed with the empirical formula (C₈H₇N)_n and that the mass spectrum of **2** shows a parent peak at *m/e* 351 corresponding to the molecular ion of the trimeric indole.

2 provided the monoacetate (**3**), colourless needles, mp 168.5—171°, [C₂₆H₂₃N₃O⁶⁾; IR (KBr): 3400, 3355 (NH), 1680 (C=O)] in 70.1% yield by treatment with Ac₂O at 100° for 3 min.

In order to compare with the properties of **2** and **3**, the known trimer⁷⁾ (**1**), mp 173—177° (lit.⁸⁾ mp 169°), and its monoacetate (**4**), mp 209—212° (lit.⁹⁾ mp 202°) were prepared according to the reported method.^{8,9)} Non-identity of **2** with **1** was confirmed by comparison of their infrared (IR) spectra in a solution and *R_f* values on TLC [SiO₂; ether-*n*-hexane (3:1)]. A clue for structural elucidation of **2** came from the comparative inspection of the spectral data on these four compounds mentioned above. As shown in Table I, the nuclear magnetic resonance (NMR) spectra of **1** and **2** show the presence of an ArCH₂CH< and two NH functions. The mass spectra of both trimers have a common base peak at *m/e* 245 corresponding to a bisindolylmethine ion. These data indicate that these two trimers should be allocated to two of three possible isomers, (**1**), (**2**) or (**5**).

In the trial of the Vilsmeier reaction of **3** and **4** using DMF and POCl₃, the former (**3**) gave the monoformyl derivative (**6**), mp 252—255°, [C₂₇H₂₃N₃O₂⁶⁾; IR (Nujol): 3370, 3170

1) This paper forms Part XI of "Fischer Indolization and Its Related Compounds," by H. Ishii.

2) For a review of polymerization of indole with acid, see R.J. Sundberg, "The Chemistry of Indoles," Academic Press, New York, 1970, pp. 6—8; W.A. Remers and R.K. Brown, "Indoles" (ed. by W.J. Houlihan), Part 1, Wiley-Interscience, New York, 1972, pp. 66—70; G.F. Smith, "Advances in Heterocyclic Chemistry," (ed. by A.R. Katritzky), Vol. 2, Academic Press, New York, 1963, pp. 300—309.

3) G.F. Smith, *Chem. and Ind.* (London), **1954**, 1451.

4) W.E. Noland and W.C. Kuryla, *J. Org. Chem.*, **25**, 486 (1960).

5) H. Ishii, Y. Murakami, T. Furuse, K. Hosoya, H. Takeda, and N. Ikeda, *Tetrahedron*, **29**, 1991 (1973); H. Ishii, Y. Murakami, K. Hosoya, T. Furuse, H. Takeda, and N. Ikeda, *Chem. Pharm. Bull.* (Tokyo), **20**, 1088 (1972).

6) The compound gave satisfactory elemental analysis for the formula given.


7) This compound (**1**) also shows dimorphism. Another crystal form melted at 110—113°. This fact was confirmed by the cross-seeding experiment.

8) O. Schmitz-Dumont, B. Nicolajannis, E. Schnorrenberg, and H.H. Saenger, *J. Prakt. Chem.*, **131**, 146 (1931).

9) K. Keller, *Chem. Ber.*, **46**, 726 (1913).

TABLE I. The NMR Spectra of Indole Trimers (in CDCl_3 , δ)

	The known trimer (1)	The new trimer (2)
$-\text{CH}_2-\text{CH}<$	3.33(2H, d, $J=7.0$ Hz)	3.35(2H, diff. d, $J=7.0$ Hz)
$-\text{CH}_2-\text{CH}<$	4.78(1H, t, $J=7.0$ Hz)	4.62(1H, t, $J=7.0$ Hz)
Arom. protons	6.38—6.68(2H, m)	6.35—6.72(3H, m)
	6.76—7.30(10H, m)	6.77—7.40(10H, m)
	7.41(2H, d, $J=7.8$ Hz)	7.43—7.62(1H, m)
NH	3.20(2H, br. s)	2.98(2H, br. s)
	7.66(2H, br. s)	7.77(2H, br. s)



(NH), 1680, 1620 ($\text{C}=\text{O}$)] in 81.9% yield, while the latter (4) resulted in the recovery of the starting material. These evidences indicate that there is no unsubstituted C_3 position of indole moieties in 1, but one in 2, because it is well known that the Vilsmeier reaction easily takes place at the C_3 position of indole nucleus, when remained unsubstituted. The chemical deduction was also supported by the spectral evidences. In the NMR spectrum, there is no signal attributable to the C_3 -proton of an indole part in 4 and 6, while it can be observed at 6.19δ as 1H singlet in the starting 3. Furthermore, in 6, the 1H multiplet appeared at 8.00δ attributable to the C_4 -proton of the formylated skeleton. The down-field shift of this signal should be caused by an anisotropic effect of the peri C_3 -formyl group. Finally, in the mass spectrum, 6 has a base peak at m/e 273 corresponding to the formylated bisindolylmethine ion.

TABLE II. The NMR Spectra of Derivatives of the Indole Trimers (in $\text{DMSO}-d_6$, δ)

	4	3	6
$-\text{CH}_2-\text{CH}<$	3.45(2H, d, $J=7.8$ Hz)	3.46(2H, d, $J=7.5$ Hz)	3.36—3.84(2H, m)
$-\text{CH}_2-\text{CH}<$	4.71(1H, t, $J=7.8$ Hz)	4.66(1H, t, $J=7.5$ Hz)	5.34(1H, diff. t, $J=7.5$ Hz)
Arom. H	$\text{C}_3\text{-H}$	6.19(1H, s)	—
	Others	6.73—7.52(13H, m)	6.76—7.50(12H, m)
	Ha	—	8.00(1H, m)
NH	8.93(1H, s)	9.08(1H, s)	9.35(1H, s)
	10.60(2H, s)	10.72(2H, s)	10.91(1H, s)
COCH_3	1.88(3H, s)	1.96(3H, s)	11.71(1H, s)
CHO	—	—	2.04(3H, s)
			10.04(1H, s)

In 1960, Noland⁴⁾ condensed *o*-nitrophenylacetaldehyde¹⁰⁾ (7) with indole itself in AcOH to give the *o*-nitro-bisindolyl product (8), which was converted to 1 by the reduction of the nitro group. We, however, found that the similar treatment of the same reactants in the presence of anhydr. ZnCl_2 gave a mixture of 8 and an isomeric nitro derivative (9) as a labile crystalline mass, mp 85° [IR(Nujol): 3400 (NH), 1522, 1348 (NO_2); NMR ($\text{DMSO}-d_6$) δ : 3.70 (2H, d, $J=8.0$ Hz, $\text{ArCH}_2\text{CH}<$), 4.64 (1H, t, $J=8.0$ Hz, $\text{ArCH}_2\text{CH}<$), 6.16 (1H, s, indolic $\text{C}_3\text{-H}$), 6.62—7.00 (4H, m, arom. H), 7.00—7.45 (8H, m, arom. H), 7.65—7.80 (1H, m, arom. Hx), 10.66 (2H, br. s, NH, exchangeable); MS m/e : 381 (M^+ , 8.8%), 245 (bisindolylmethine ion, base peak)] in 13.0 and 32.2% yields, respectively. Formation of 9 should be explained by the acid catalyzed rearrangement of the initially formed 8 to 9 during the reaction, because treatment of the pure 8 with anhydr. ZnCl_2 in AcOH under the same condition mentioned above afforded the same product mixture. Catalytic reduction of the pure 9 provided 2.




Chart 2

The evidence described so far indicates that the structure of the new indole trimer should be depicted by the formula (2). Although the mechanism for the formation of 2 could be supposed by the acid catalyzed rearrangement of 1 on the basis of the rearrangement of 8 to 9, the precise discussion on this matter will be described in our full paper in the near future.

Acknowledgement We wish to thank Prof. W.E. Noland, University of Minnesota, for his gift of the known trimer and information about it, and Prof. T. Hino and Dr. M. Nakagawa for their heartfelt discussion.

Faculty of Pharmaceutical Sciences
Chiba University, 1-33, Yayoi-cho,
Chiba, 280, Japan

HISASHI ISHII¹¹⁾
KEIKO MURAKAMI
YASUOKI MURAKAMI
KATSUHIRO HOSOYA

Received June 30, 1977

10) a) M. Mousseron-Canet and J.-P. Boca, *Bull. Soc. Chim. France*, **1967**, 1296; e) R.A. Weerman, *Ann.*, **401**, 1 (1913); c) W.E. Noland and J.H. Sellstedt, *J. Org. Chem.*, **31**, 345 (1966). We prepared 7 according to the reported method in the reference.^{10a)}

11) The author to whom correspondence should be addressed.