

An Efficient *p*-Thiocyanation of Phenols and Naphthols Using a Reagent Combination of Phenyliodine Dichloride and Lead(II) Thiocyanate

Yasuyuki KITA,* Yoshifumi TAKEDA, Takayuki OKUNO, Masahiro EGI, Kiyosei IIO, Ken-ichi KAWAGUCHI, and Shuji AKAI

Faculty of Pharmaceutical Sciences, Osaka University, 1-6, Yamada-oka, Suita, Osaka 565, Japan.

Received May 12, 1997; accepted August 4, 1997

A combination of PhICl_2 and $\text{Pb}(\text{SCN})_2$ is effective for the *p*-selective thiocyanation of various types of *p*-unsubstituted phenols and naphthols **1** to give *p*-thiocyanatophenols and naphthols **3**. The reaction proceeded at 0 °C to room temperature in good to quantitative yields. Twenty-five examples are given, in which various functional groups, such as chloro, allyl, carbonyl, ester, amide, and primary hydroxyl groups, are shown to be compatible with this reaction.

Key words thiocyanation; phenol; hypervalent iodine reagent; lead(II) thiocyanate

Because the thiocyanato group can be transformed into various sulfur functional groups and sulfur-containing heterocycles, the thiocyanation of aromatic compounds is an important subject.^{1,2)} Many thiocyanation methods have been developed using electrophilic thiocyanato species, generated by the chemical¹⁾ or electrochemical³⁾ oxidation of thiocyanate anions. The thiocyanation of aromatic amines was reported to occur rapidly to give high yields of the products. On the other hand, the thiocyanation of phenols has not been as extensively studied, and the yields did not appear to be as high as those with aromatic amines. In particular, the presence of electron-withdrawing substituents on the aromatic rings reduced the yields.^{1,4–6)} In connection with our project on the development of novel transformation reactions of the *p*-sulfinyl group of phenol derivatives through the Pummerer-type reaction,⁷⁾ we briefly reported an efficient *p*-thiocyanation of various phenol compounds by using the combination of PhICl_2 and $\text{Pb}(\text{SCN})_2$.⁸⁾ Here we describe this reaction in detail.

Results and Discussion

We reported that hypervalent iodine(III) reagents, phenyliodine diacetate and phenyliodine bis(trifluoroacetate) (PIFA), are useful for the oxidative addition of alcohols, water, and carboxylic acids at the *para*-position

of phenols (Chart 1).⁹⁾ Similar types of *p*-substitution reactions have been widely reported in the presence of inter- and intramolecular nucleophiles, including amides, oximes, fluoride ion, electron-rich olefins, and electron-rich aromatic compounds.¹⁰⁾ Therefore, we expected that a similar reaction using a reagent combination of a hypervalent iodine(III) compound and a thiocyanate anion ($\text{Nu} = ^-\text{SCN}$) would be useful for the *p*-thiocyanation of phenols **1**.

We anticipated that the appropriate reagent combination of known iodine(III) reagents would undergo ligand-exchange reaction to generate a novel iodine(III) compound, suitable for thiocyanation. We preliminarily tried three types of reagent combinations (runs 1–3 in Table 1); however, **2** or related iodine(III) compounds having one or two thiocyanato ligands could not be isolated or even characterized by spectroscopic means. The compounds identified by IR and ¹H-NMR spectroscopies were iodobenzene and thiocyanogen, even by rapid measurement of ¹H-NMR data between –78 and 0 °C. However, we found that the addition of phenol **1a** to each reaction mixture immediately gave the *p*-thiocyanatophenol **3a** in high to quantitative yield (Table 1).

For example, under a nitrogen atmosphere, PhICl_2 (1.2 mmol) and $\text{Pb}(\text{SCN})_2$ (1.5 mmol) were stirred in dry




Chart 1

Table 1. Combination of the Hypervalent Iodine(III) Reagent and the Thiocyanate and Its Reaction with Phenol (**1a**)

$\text{PhIX}_2 + \text{M-SCN} \xrightarrow[\text{0 °C, 20 min}]{\text{CH}_2\text{Cl}_2} [\text{PhI}(\text{SCN})_2 + 2 \text{MX}]$			
Run	PhIX ₂	M-SCN	Yield (%) of 3a ^{a)}
1	$\text{PhI}(\text{OCOCF}_3)_2$	TMS-NCS (2 eq)	95
2	$\text{PhI}=\text{O}$	TMS-NCS (2 eq)	56
3	PhICl_2	$\text{Pb}(\text{SCN})_2$ (1.2 eq)	93

a) Isolated yield after SiO_2 chromatography.

CH_2Cl_2 at 0°C for 20 min. Phenol **1a** (1.0 mmol) was added to the resultant suspension, and the whole was stirred for 30 min. The reaction mixture was filtered, concentrated, and purified by column chromatography on SiO_2 to give *p*-thiocyanatophenol **3a** in 93% yield (run 3). Similarly, the use of PIFA and TMS-NCS gave 95% yield of **3a** (run 1), while that of iodosobenzene and TMS-NCS was not satisfactory (run 2). Among the former combina-

tions, the use of PhICl_2 and $\text{Pb}(\text{SCN})_2$ seemed to be favorable to that of PIFA and TMS-NCS because the former two reagents are readily available and easily handled solids.

Using this method, various phenols, **1b–m**, and naphthols, **1n–y**, were thiocyanated to give the corresponding products **3b–y**. The results are summarized in Table 2. Several features are noteworthy. i) Usually the reaction was completed at 0°C within 1 h. ii) In most cases, the reaction exclusively occurred at the *para*-position of the phenol ring when the *para*-position was free.¹¹⁾ Even in the cases of 3,5-dimethylphenol **1f** and the congested tricyclic naphthol **1x**, the *p*-thiocyanates **3f**, **x** were obtained as single products in excellent yields (runs 6, 24). However, the similar reaction of the oxo-tricyclic naphthol **1y** was not regiospecific, giving a mixture of the *p*-thiocyanate **3y** (61%) and its regio isomer **4** (18%) (run 25). iii) All products were thiocyanates (**3a–y**, **4**), and the corresponding isothiocyanates were not obtained at all. The structures were determined from the IR absorption spectra, because the products showed IR absorptions at $2172\text{--}2151\text{ cm}^{-1}$, while those of the isothiocyanates are reported to be in the range of $2140\text{--}1990\text{ cm}^{-1}$. iv) Reactive functional groups such as allyl and primary hydroxyl groups were compatible (runs 9, 10, 12). v) The phenols having electron-withdrawing cyano, chloro, carbonyl, ester, and amido groups were converted to the corresponding thiocyanates **3b**, **e**, **k**, **o**, **q–w**, and **y** in good yields (runs 2, 5, 11, 15, 17–23, 25). However, *o*-nitrophenol did not react even when the reaction mixture was stirred at room temperature for 1 d.

As we previously mentioned, spectroscopic measurement of the ligand-exchange reaction of PhICl_2 and $\text{Pb}(\text{SCN})_2$ revealed complete formation of iodobenzene and thiocyanogen. However, a comparative examination of our reagent system (run 1 in Table 3) with the ordinary method using thiocyanogen (run 3) revealed that the former was more effective than the latter. The addition of PhI to the reaction mixture of run 3 slightly increased the yield of **3u** (runs 4, 6). These results may suggest that there is an equilibrium between the novel iodine **2** and the mixture of iodobenzene and thiocyanogen lying far to the right (Chart 2) and that **2** takes part in the thiocyanation

Table 2. Thiocyanation of Various Phenols (**1**) Using $\text{PhICl}_2\text{--Pb}(\text{SCN})_2$

Run	Phenol 1	Product 3 Yield (%) ^{a)}
	$\text{R}^1 = \quad \text{R}^2 = \quad \text{R}^3 = \quad \text{R}^4 =$	
1	1a H H H H	3a 93
2	b CN H H H	b 61
3	c Me H H Me	c 78
4	d ^t Bu H H ^t Bu	d 97
5	e Cl H H Cl	e 64
6	f H Me Me H	f 95
7	g H Me Me Me	g 81
8	h Me Me Me Me	h 94
9	i allyl Me Me Me	i 79
10	j allyl Me Me allyl	j 89
11	k COMe Me Me Me	k 78
12	l $(\text{CH}_2)_3\text{OH}$ Me Me Me	l 91
13		65
	$\text{R}^1 = \quad \text{R}^2 =$	
14	1n H H	3n 88
15	o COMe H	o 97
16	p Me Me	p 67
17	q COMe Me	q 86
18	r COMe Ph	r 85
19	s CO_2Et Me	s 97
20	t CO_2Et CH_2OCOMe	t 88
21	u CO_2Et CO_2Et	u 58
22	v CONEt_2 Me	v 98
23	w CONEt_2 CONEt_2	w 91
	$\text{X} =$	
24	1x H_2	3x 94
25	y O	y 61 ^{b)}

a) Isolated yield after SiO_2 chromatography. b) The by-product **4** was isolated.




Table 3. Thiocyanation of **1u** under Various Conditions

Run	Thiocyanation reagent	Yield (%) of 3u ^{a)}
1	PhICl_2 (1.2 eq), $\text{Pb}(\text{SCN})_2$ (1.5 eq)	58
2	PhICl_2 (2 eq), $\text{Pb}(\text{SCN})_2$ (3 eq) ^{b)}	44
3	$(\text{SCN})_2$ ^{c)} (2 eq)	30
4	$(\text{SCN})_2$ ^{c)} (2 eq), PhI (2 eq)	40
5	$(\text{SCN})_2$ ^{c)} (2 eq), PbCl_2 (2 eq)	39
6	$(\text{SCN})_2$ ^{c)} (2 eq), PhI (2 eq), PbCl_2 (2 eq)	45

a) Isolated yield after SiO_2 chromatography. b) Run after filtration of the by-product, PbCl_2 . c) Generated from bromine and $\text{Pb}(\text{SCN})_2$ in CH_2Cl_2 at 0°C for 20 min and used after filtration of the by-product, PbBr_2 .




Chart 2

as a reactive molecule. The presence of a weak Lewis acid, PbCl_2 , seems to accelerate the reaction, because removal of PbCl_2 from the reaction mixture of run 1 decreased the yield of **3u** (run 2), while addition of PbCl_2 increased the yield (runs 5, 6).

Although elucidation of the reaction mechanism would require a more detailed study, it is clear that the use of the mixture of PhICl_2 and Pb(SCN)_2 without filtration of the by-product PbCl_2 provides a convenient thiocyanation of phenols.^{12–15} This method features i) good to high yields of the products from various types of phenols having electron-withdrawing groups and other reactive functional groups, ii) exclusive *para*-selectivity, and iii) an easy procedure using solid and stable PhICl_2 and Pb(SCN)_2 .

Experimental

All melting points are uncorrected. IR absorption spectra were recorded on a Shimadzu FT-IR-8100 spectrometer by diffuse reflectance measurement of samples dispersed in KBr powder. $^1\text{H-NMR}$ spectra were recorded on Varian VXR-200 (200 MHz), Hitachi R-250HT (250 MHz), and JEOL JNM-EX270 (270 MHz) spectrometers with SiMe_4 as an internal standard. High-resolution mass spectra (HR-MS) were recorded at 70 eV with a direct inlet system on JEOL JMS-D300 and JEOL JMS-HX100 spectrometers. E. Merck Silica gel 60 (70–230 mesh ASTM) was used for column chromatography. Phenylodine(III) dichloride was prepared according to the reported procedure.¹⁶ Pb(SCN)_2 was a commercial product, used as supplied.

General Procedure for the Thiocyanation of Phenols (1) Under a nitrogen atmosphere, PhICl_2 (330 mg, 1.2 mmol) was added to an ice-cooled suspension of Pb(SCN)_2 (485 mg, 1.5 mmol) in dry CH_2Cl_2 (10 mL). The reaction mixture was stirred at the same temperature for 20 min, then a solution of the phenol **1** (1.0 mmol) in dry CH_2Cl_2 (2 mL) was added. The whole was stirred for 30 min, filtered through a Celite pad, and concentrated to half volume. Silica gel (2 g) for column chromatography was added, and the solvent was evaporated *in vacuo*. The product absorbed on SiO_2 was charged on an SiO_2 column and eluted with AcOEt -hexane to give the *p*-thiocyanatophenol **3**. The yields are summarized in Table 2.

4-Thiocyanatophenol (3a): Colorless crystals. mp 59–60 °C (AcOEt -hexane) (lit.¹³ mp 59–60 °C). IR 3350, 2161, 1599, 1584, 1497 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 5.69 (1H, brs), 6.88 (2H, d, $J=9.5$ Hz), 7.45 (2H, d, $J=9.5$ Hz).

2-Cyano-4-thiocyanatophenol (3b): Colorless crystals. mp 140–145 °C (CH_2Cl_2). IR 3250, 2230, 2172, 1597, 1493 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 7.02 (1H, d, $J=9.0$ Hz), 7.63 (1H, dd, $J=2.5, 9.0$ Hz), 7.72 (1H, d, $J=2.5$ Hz). HR-MS Calcd for $\text{C}_8\text{H}_4\text{N}_2\text{OS}$: 176.0044. Found 176.0044.

2,6-Dimethyl-4-thiocyanatophenol (3c): Colorless crystals. mp 103.5–104 °C (CH_2Cl_2 -hexane) (lit.¹⁷ mp 100–101 °C). IR 3450, 2155, 1584, 1480 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.26 (6H, s), 4.90 (1H, s), 7.21 (2H, s).

2,6-Di-*tert*-butyl-4-thiocyanatophenol (3d): Colorless crystals. mp 54–55 °C (pentane) [lit.¹⁸ mp 64–66.5 °C (pentane)]. IR 3629, 2157, 1428 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.44 (18H, s), 5.50 (1H, brs), 7.37 (2H, s).

2,6-Dichloro-4-thiocyanatophenol (3e): Colorless crystals. mp 101–102 °C (CH_2Cl_2 -hexane). IR 3299, 2172, 1563, 1478, 1460 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 6.17 (1H, brs), 7.53 (2H, s). Anal. Calcd for $\text{C}_7\text{H}_3\text{Cl}_2\text{NOS}$: C, 38.20; H, 1.37; N, 6.36. Found: C, 38.24; H, 1.56; N, 6.32.

3,5-Dimethyl-4-thiocyanatophenol (3f): Colorless crystals. mp 132–133 °C (CH_2Cl_2 -hexane) (lit.¹⁷ mp 127–129 °C). IR 3347, 2159, 1588,

1458 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.54 (6H, s), 6.66 (2H, s).

2,3,5-Trimethyl-4-thiocyanatophenol (3g): Colorless crystals. mp 161–162 °C (CH_2Cl_2 -hexane). IR 3400–3100 br, 2166, 1582 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.17 (3H, s), 2.50 (3H, s), 2.55 (3H, s), 5.33 (1H, brs), 6.62 (1H, s). Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{NOS}$: C, 62.15; H, 5.74; N, 7.25; S, 16.59. Found: C, 62.02; H, 5.73; N, 7.22; S, 16.59.

2,3,5,6-Tetramethyl-4-thiocyanatophenol (3h): Colorless crystals. mp 146.5–147 °C (CH_2Cl_2 -hexane) [lit.¹⁹ mp 144.5–145 °C (Et_2O)]. IR 3415, 2163, 1561 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.21 (6H, s), 2.57 (6H, s), 5.03 (1H, s).

2-Allyl-3,5,6-trimethyl-4-thiocyanatophenol (3i): Colorless crystals. mp 93–93.5 °C (CH_2Cl_2 -hexane). IR 3450, 2153, 1638, 1550, 1449, 1402 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.21 (3H, s), 2.58 (6H, s), 3.46 (2H, d, $J=4.0$ Hz), 5.02 (1H, d, $J=18.0$ Hz), 5.13 (1H, d, $J=10.0$ Hz), 5.19 (1H, s), 5.84–6.06 (1H, m). Anal. Calcd for $\text{C}_{13}\text{H}_{15}\text{NOS}$: C, 66.92; H, 6.48; N, 6.00; S, 13.74. Found: C, 66.81; H, 6.30; N, 5.98; S, 13.58.

2,6-Diallyl-3,5-dimethyl-4-thiocyanatophenol (3j): Colorless crystals. mp 95–97 °C (AcOEt -hexane). IR 3420, 2164, 1553 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.58 (6H, s), 3.47 (4H, brd, $J=5.5$ Hz), 5.00 (2H, dd, $J=2.0, 18.0$ Hz), 5.11 (2H, dd, $J=1.5, 10.0$ Hz), 5.29 (1H, s), 5.86–6.01 (2H, m). Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{NOS}$: C, 69.49; H, 6.61; N, 5.40; S, 12.36. Found: C, 69.22; H, 6.60; N, 5.33; S, 12.19.

2-Acetyl-3,5,6-trimethyl-4-thiocyanatophenol (3k): Colorless crystals. mp 59.5–60 °C (CH_2Cl_2 -hexane). IR 3400, 2153, 1697, 1626, 1582, 1559 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.22 (3H, s), 2.61 (3H, s), 2.64 (3H, s), 2.82 (3H, s), 11.80 (1H, s). Anal. Calcd for $\text{C}_{12}\text{H}_{13}\text{NO}_2\text{S}$: C, 61.25; H, 5.57; N, 5.95; S, 13.63. Found: C, 61.21; H, 5.50; N, 5.93; S, 13.39.

2-(3-Hydroxypropyl)-3,5,6-trimethyl-4-thiocyanatophenol (3l): Colorless crystals. mp 91–92 °C (CH_2Cl_2 -hexane). IR 3700–3100 br, 2153, 1557 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.84 (2H, tt, $J=5.5, 6.5$ Hz), 2.20 (3H, s), 2.54 (3H, s), 2.56 (3H, s), 2.82 (2H, t, $J=6.5$ Hz), 2.92 (1H, brs), 3.59 (2H, t, $J=5.5$ Hz), 7.95 (1H, brs). HR-MS Calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_2\text{S}$: 251.0980. Found 251.0986.

5-Hydroxy-1-methyl-8-thiocyanato-3,4-dihydroquinolin-2(1H)-one (3m): Colorless crystals. mp 175–177 °C (CH_2Cl_2 -hexane). IR 3250, 2205, 1650, 1609 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.65 (1H, d, $J=7.0$ Hz), 2.70 (1H, d, $J=6.0$ Hz), 2.98 (1H, d, $J=6.0$ Hz), 3.03 (1H, d, $J=7.0$ Hz), 3.36 (3H, s), 6.78 (1H, d, $J=9.0$ Hz), 7.13 (1H, d, $J=9.0$ Hz). HR-MS Calcd for $\text{C}_{11}\text{H}_{10}\text{N}_2\text{O}_2\text{S}$: 234.0460. Found 234.0460.

4-Thiocyanato-1-naphthol (3n): Colorless crystals. mp 112.5–113 °C (CH_2Cl_2 -hexane) (lit.^{4a} mp 112–113 °C). IR 3300, 2161, 1592, 1570, 1514 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 6.03 (1H, s), 6.81 (1H, d, $J=8.0$ Hz), 7.60 (1H, t, $J=8.0$ Hz), 7.73 (1H, t, $J=8.0$ Hz), 7.75 (1H, d, $J=8.0$ Hz), 8.29 (2H, d, $J=8.0$ Hz).

2-Acetyl-4-thiocyanato-1-naphthol (3o): Colorless crystals. mp 160–161 °C (CH_2Cl_2 -hexane). IR 3069, 2155, 1622, 1590, 1568, 1501 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.72 (3H, s), 7.67 (1H, t, $J=7.5$ Hz), 7.86 (1H, t, $J=7.5$ Hz), 8.14 (1H, s), 8.26 (1H, d, $J=8.5$ Hz), 8.54 (1H, d, $J=8.5$ Hz). Anal. Calcd for $\text{C}_{13}\text{H}_9\text{NO}_2\text{S}$: C, 64.18; H, 3.73; N, 5.76; S, 13.18. Found: C, 64.12; H, 3.85; N, 5.75; S, 13.14.

2,3-Dimethyl-4-thiocyanato-1-naphthol (3p): Colorless crystals. mp 150–153 °C (AcOEt -hexane) [lit.¹⁹ mp 151–153 °C (ligroin)]. IR 3750, 2145, 1592, 1557, 1499 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.38 (3H, s), 2.80 (3H, s), 5.56 (1H, s), 7.53 (1H, t, $J=8.0$ Hz), 7.65 (1H, t, $J=8.0$ Hz), 8.14 (1H, d, $J=8.0$ Hz), 8.44 (1H, d, $J=8.0$ Hz).

2-Acetyl-3-methyl-4-thiocyanato-1-naphthol (3q): Colorless crystals. mp 122–123 °C (CH_2Cl_2 -hexane). IR 3200–2200 br, 2153, 1615, 1572 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.73 (3H, s), 3.02 (3H, s), 7.60 (1H, t, $J=8.0$ Hz), 7.81 (1H, t, $J=8.0$ Hz), 8.47 (1H, d, $J=8.0$ Hz), 8.51 (1H, d, $J=8.0$ Hz), 13.90 (1H, s). Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{NO}_2\text{S}$: C, 65.35; H, 4.31; N, 5.44; S, 12.44. Found: C, 65.16; H, 4.41; N, 5.38; S, 12.44.

2-Acetyl-3-phenyl-4-thiocyanato-1-naphthol (3r): Colorless crystals. mp 195–197 °C (CH_2Cl_2 -hexane). IR 3200–2100 br, 2153, 1615, 1566 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.78 (3H, s), 7.37–7.39 (2H, m), 7.54–7.55 (3H, m), 7.69 (1H, ddd, $J=1.0, 7.5, 8.0$ Hz), 7.91 (1H, ddd, $J=1.0, 7.5, 8.0$ Hz), 8.47 (1H, d, $J=8.0$ Hz), 8.62 (1H, d, $J=8.0$ Hz), 14.61 (1H, s). Anal. Calcd for $\text{C}_{19}\text{H}_{13}\text{NO}_2\text{S}$: C, 71.45; H, 4.10; N, 4.39; S, 10.04. Found: C, 71.25; H, 4.13; N, 4.35; S, 9.95.

2-Ethoxycarbonyl-3-methyl-4-thiocyanato-1-naphthol (3s): Colorless crystals. mp 134–135 °C (CH_2Cl_2 -hexane). IR 3200–2500 br, 2157, 1653, 1617, 1570 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.48 (3H, t, $J=7.0$ Hz), 3.02 (3H, s), 4.52 (2H, q, $J=7.0$ Hz), 7.57 (1H, t, $J=8.0$ Hz), 7.78 (1H, t, $J=8.0$ Hz), 8.40 (1H, d, $J=8.0$ Hz), 8.45 (1H, d, $J=8.0$ Hz), 12.93 (1H, s). HR-MS Calcd for $\text{C}_{15}\text{H}_{13}\text{NO}_3\text{S}$: 287.0613. Found 287.0608.

3-Acetoxyethyl-2-ethoxycarbonyl-4-thiocyanato-1-naphthol (**3t**): Colorless crystals. mp 118–119 °C (CH₂Cl₂–hexane). IR 2157, 1741, 1655, 1619, 1570, 1493 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.44 (3H, t, *J* = 7.5 Hz), 2.10 (3H, s), 4.50 (2H, q, *J* = 7.5 Hz), 5.91 (2H, s), 7.68 (1H, t, *J* = 8.0 Hz), 7.87 (1H, t, *J* = 8.0 Hz), 8.49 (1H, d, *J* = 8.0 Hz), 8.52 (1H, d, *J* = 8.0 Hz), 12.75 (1H, s). *Anal.* Calcd for C₁₇H₁₅NO₅S: C, 59.12; H, 4.38; N, 4.06; S, 9.28. Found: C, 59.04; H, 4.39; N, 4.09; S, 9.18.

2,3-Bis(ethoxycarbonyl)-4-thiocyanato-1-naphthol (**3u**): Colorless crystals. mp 125.5–126 °C (CH₂Cl₂–hexane). IR 2159, 1740, 1665, 1619, 1493 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.43 (3H, t, *J* = 7.5 Hz), 1.50 (3H, t, *J* = 7.0 Hz), 4.42–4.58 (4H, m), 7.71 (1H, t, *J* = 8.0 Hz), 7.90 (1H, t, *J* = 8.0 Hz), 8.43 (1H, d, *J* = 8.0 Hz), 8.54 (1H, d, *J* = 8.0 Hz), 13.04 (1H, s). *Anal.* Calcd for C₁₇H₁₅NO₅S: C, 59.12; H, 4.38; N, 4.06; S, 9.28. Found: C, 59.38; H, 4.50; N, 3.97; S, 9.16.

2-(*N,N*-Diethylcarbamoyl)-3-methyl-4-thiocyanato-1-naphthol (**3v**): Colorless crystals. mp 171–173 °C (CH₂Cl₂–hexane). IR 3000–2900 br, 2153, 1593 cm⁻¹. ¹H-NMR (CDCl₃) δ: 0.81–1.50 (6H, m), 2.44 (3H, s), 2.92–4.02 (4H, m), 7.37 (1H, dd, *J* = 7.0, 7.5 Hz), 7.50 (1H, dd, *J* = 7.0, 7.5 Hz), 7.90–7.94 (2H, m), 10.37 (1H, s). *Anal.* Calcd for C₁₇H₁₈N₂O₂S: C, 64.94; H, 5.77; N, 8.91; S, 10.20. Found: C, 64.99; H, 5.84; N, 8.90; S, 10.19.

2,3-Bis(*N,N*-diethylcarbamoyl)-4-thiocyanato-1-naphthol (**3w**): Colorless crystals. mp 170–173.5 °C (CH₂Cl₂–hexane). IR 2157, 1636, 1617, 1599, 1576, 1561, 1487 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.07 (3H, t, *J* = 7.5 Hz), 1.23–1.36 (9H, m), 3.05–3.93 (8H, m), 7.36 (1H, t, *J* = 8.0 Hz), 7.55 (1H, t, *J* = 8.0 Hz), 7.71 (1H, d, *J* = 8.0 Hz), 8.18 (1H, d, *J* = 8.0 Hz), 9.92 (1H, br s). *Anal.* Calcd for C₂₁H₂₅N₃O₃S: C, 63.13; H, 6.31; N, 10.52; S, 8.03. Found: C, 63.20; H, 6.33; N, 10.35; S, 8.07.

9-Hydroxy-8-methoxy-3,3-dimethyl-10-thiocyanato-1,2,3,4-tetrahydroanthracene (**3x**): Colorless crystals. mp 155–156 °C (AcOEt–hexane). IR 3400–3300 br, 2151, 1620, 1603, 1561, 1374, 1354 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.06 (6H, s), 1.65 (2H, t, *J* = 7.0 Hz), 2.85 (2H, t, *J* = 7.0 Hz), 3.01 (2H, s), 4.09 (3H, s), 6.84 (1H, d, *J* = 8.0 Hz), 7.45 (1H, t, *J* = 8.0 Hz), 8.10 (1H, d, *J* = 8.0 Hz), 10.13 (1H, s). HR-MS Calcd for C₁₈H₁₉NO₂S: 313.1136. Found 313.1148.


9-Hydroxy-8-methoxy-3,3-dimethyl-10-thiocyanato-1,2,3,4-tetrahydroanthracen-1-one (**3y**): Yellow crystals. mp 142–145 °C (CH₂Cl₂–hexane). IR 2957, 2153, 1615, 1576 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.16 (6H, s), 2.64 (2H, s), 3.29 (2H, s), 4.05 (3H, s), 6.99 (1H, d, *J* = 8.0 Hz), 7.73 (1H, t, *J* = 8.0 Hz), 8.06 (1H, d, *J* = 8.0 Hz), 15.89 (1H, s). HR-MS Calcd for C₁₈H₁₇NO₃S: 327.0929. Found 327.0930.

9-Hydroxy-8-methoxy-3,3-dimethyl-5-thiocyanato-1,2,3,4-tetrahydroanthracen-1-one (**4**): Yellow crystals. mp 184–187 °C (CH₂Cl₂–hexane). IR 2950, 2161, 1620, 1565, 1372, 1281, 1107 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.12 (6H, s), 2.62 (2H, s), 2.97 (2H, s), 4.06 (3H, s), 6.82 (1H, d, *J* = 8.5 Hz), 7.49 (1H, s), 7.95 (1H, d, *J* = 8.5 Hz), 15.12 (1H, s). HR-MS Calcd for C₁₈H₁₇NO₃S: 327.0929. Found 327.0932.

Acknowledgments This study was supported by Special Coordination Funds of the Science and Technology Agency of the Japanese Government.

References and Notes

- Reviews: a) Wood J. L., "Organic Reactions," Vol. 3, ed. by Adams R., John Wiley & Sons, New York, 1946, Chapter 6, pp. 240–266; b) Guy R. G., "The Chemistry of Cyanates and Their Thio Derivatives," Part 2, ed. by Patai S., John Wiley & Sons, New York, 1977, Chapter 18, pp. 819–886; c) Harusawa S., Shioiri T., *Yuki Gosei Kagaku Kyokaiishi*, **39**, 741–753 (1981) [*Chem. Abstr.*, **95**, 167873k (1981)].
- Recent reports on thiocyanation: Wilkerson W. W., Galbraith W., Gans-Brangs K., Grubb M., Hewes W. E., Jaffee B., Kenney J. P., Kerr J., Wong N., *J. Med. Chem.*, **37**, 988–998 (1994); Toste F. D., Stefano V. D., Still I. W. J., *Synth. Commun.*, **25**, 1277–1286 (1995). Recent reports on the conversion of the aryl thiocyanates: Toste F. D., LaRonde F., Still I. W. J., *Tetrahedron Lett.*, **36**, 2949–2952 (1995); Still I. W. J., Toste F. D., *J. Org. Chem.*, **61**, 7677–7680 (1996).
- Krishnan P., Gurjar V. G., *Synth. Commun.*, **22**, 2741–2744 (1992).
- a) Kaji A., *Nippon Kagaku Zasshi*, **82**, 382–385 (1961) [*Chem. Abstr.*, **56**, 10054h (1962)]; b) Pohloudek-Fabini R., Lüß K.-D., *Arch. Pharm. (Weinheim, Ger.)*, **299**, 866–877 (1966).
- Although the use of thiocyanogen in the presence of AlCl₃ may improve the thiocyanation yields for less reactive phenols, its general applicability is unclear.⁶⁾
- Söderbäck E., *Acta Chem. Scand.*, **8**, 1851–1858 (1954).
- Akai S., Takeda Y., Iio K., Yoshida Y., Kita Y., *J. Chem. Soc., Chem. Commun.*, **1995**, 1013–1014; Akai S., Iio K., Takeda Y., Ueno H., Yokogawa K., Kita Y., *ibid.*, **1995**, 2319–2320; Kita Y., Takeda Y., Iio K., Yokogawa K., Takahashi K., Akai S., *Tetrahedron Lett.*, **37**, 7545–7548 (1996); Akai S., Iio K., Takeda Y., Ueno H., Kita Y., *Synlett*, **1997**, 310–312; Kita Y., Takeda Y., Matsugi M., Iio K., Gotanda K., Murata K., Akai S., *Angew. Chem., Int. Ed. Engl.*, **36**, 1529–1531 (1997).
- Kita Y., Okuno T., Egi M., Iio K., Takeda Y., Akai S., *Synlett*, **1994**, 1039–1040.
- Tamura Y., Yakura T., Haruta J., Kita Y., *J. Org. Chem.*, **52**, 3927–3930 (1987); Tamura Y., Yakura T., Tohma H., Kikuchi K., Kita Y., *Synthesis*, **1989**, 126–127; Kita Y., Tohma H., Kikuchi K., Inagaki M., Yakura T., *J. Org. Chem.*, **56**, 435–438 (1991); Kita Y., Tohma H., Yakura T., *Trends in Organic Synthesis*, **3**, 113–128 (1992).
- Recent reviews: Stang P. J., Zhdkankin V. V., *Chem. Rev.*, **96**, 1123–1178 (1996); Varvoglis A., *Tetrahedron*, **53**, 1179–1255 (1997).
- The similar reaction of 2,4,6-trimethylphenol **1z** gave the 3-thiocyanatophenol **3z** in 49% yield, probably through migration of the thiocyanato group from the initial product **5**.



- Use of an equimolar mixture of PhICl₂ and Pb(SCN)₂ for the thiocyanation of phenols and anilines was briefly reported by Neu in 1939.¹³⁾ He used the filtrate of a mixture of PhICl₂ and Pb(SCN)₂, and the reported yields of the products were not satisfactory. This was probably due to the sensitivity of PhI(SCN)₂ to moisture and/or oxygen. Direct addition of phenols to the mixture without the filtration process in our method afforded *p*-thiocyanatophenols in good to quantitative yields.
- Neu R., *Chem. Ber.*, **72**, 1505–1512 (1939).
- The use of a 2 : 1 combination of PhICl₂ and Pb(SCN)₂ was reported to generate thiocyanogen chloride,^{15a)} which reacted with the silyl enol ether to give the α-thiocyanatone.^{15b)} Because thiocyanogen chloride reacts with excess Pb(SCN)₂ to give thiocyanogen,^{15a)} we think the formation of thiocyanogen chloride is negligible in our case, in which a 1 : 1.2 mixture of PhICl₂ and Pb(SCN)₂ was employed.
- a) Bacon R. G. R., Guy R. G., *J. Chem. Soc.*, **1960**, 318–324; b) Back T. G., Dyck B. P., Parvez M., *J. Org. Chem.*, **60**, 703–710 (1995).
- Karele B., Neilans O., *Latv. PSR Zinat. Akad. Vestis, Kim. Ser.*, **1970**, 587–590 [*Chem. Abstr.*, **74**, 42033w (1971)].
- Konishi K., *Takeda Kenkyusho Nempo*, **24**, 233–239 (1965) [*Chem. Abstr.*, **64**, 8077a (1966)].
- Müller E., Stegmann H. B., Scheffler K., *Justus Liebigs Ann. Chem.*, **645**, 79–91 (1961).
- Wieland T., Bäuerlein E., *Chem. Ber.*, **97**, 2103–2108 (1964).