

Biotransformation of Hinesol Isolated from the Crude Drug *Atractylodes lancea* by *Aspergillus niger* and *Aspergillus cellulosa*

Toshihiro HASHIMOTO,^{a)} Yoshiaki NOMA,^{b)} Sumie KATO,^{a)} Masami TANAKA,^{a)} Shigeru TAKAOKA,^{a)} and Yoshinori ASAKAWA^{*,a)}

Faculty of Pharmaceutical Sciences, Tokushima Bunri University,^{a)} Japan; Faculty of Domestic Sciences, Tokushima Bunri University,^{b)} Yamashiro-cho, Tokushima 770-8514, Japan.

Received March 5, 1999; accepted April 13, 1999

Biotransformation of the sesquiterpene alcohol hinesol (1) with spasmolytic activity, which was prepared from the rhizome of *Atractylodes lancea*, was carried out by *Aspergillus niger* and *Aspergillus cellulosa* IFO 4040. Compound 1 was easily converted to compounds 2—9 by *A. niger*, and compounds 10 and 11 by *A. cellulosa*, respectively. Their stereostructures were established by a combination of high-resolution NMR spectral analysis, X-ray crystallographic analysis, and chemical reactions such as epoxidation.

Key words *Atractylodes lancea*; hinesol; biotransformation; *Aspergillus niger*; *Aspergillus cellulosa* IFO 4040

We are continuing to study the biotransformation of secondary plant metabolites by microorganisms¹⁾ and mammals²⁾ from the pharmacological point of view. Previously, we reported the biotransformation of three germacrane-type sesquiterpenoids³⁾ from the crude drug *Curcuma aromatica*, and 6-gingerol and 6-shogaol⁴⁾ from the crude drug *Zingiber officinale* belonging to the Zingiberaceae by the fungus *Aspergillus niger*. In continuation of the biotransformation of the chemical constituents isolated from crude drugs into biologically active compounds, the biotransformation of hinesol (1), which has spasmolytic activity,⁵⁾ from *Atractylodes lancea* was examined by *A. niger* and *A. cellulosa*. This paper deals with the structure elucidation of 10 metabolites (2—11) of 1 biotransformed by two fungi.

A. niger was inoculated and cultivated under rotation (100 rpm) in Czapek-peptone medium⁶⁾ at 30 °C, pH 7.0, for 2 days. (–)-Hinesol (1) (150 mg/200 ml) was added to the medium and further cultivated for 10 days. The crude metabolites obtained from the culture broth by ether extraction were chromatographed on silica gel (*n*-hexane–EtOAc gradient) and a Sephadex LH-20 column (CHCl₃–MeOH = 1 : 1) to give eight metabolites, 2 (15%), 3 (11%), 4 (12%), 5 (8%), 6 (7%), 7 (9%), 8 (6%), and 9 (12%). (–)-Hinesol (1) (150 mg/200 ml) was cultivated for 8 days by *A. cellulosa* in the same medium to afford 10 (26%) and 11 (14%).

The IR, UV, and ¹H-NMR spectra of compound 2, C₁₅H₂₄O₂ (HRMS; [M]⁺ *m/z* 236.1857), showed the presence of an α, β-conjugated ketone (IR; 1664 cm^{−1}; UVλ_{max} 240 nm [log ε = 3.99]; δ_H 5.76 [1H, br s, 1-H]). From careful analysis of its 2D NMR spectrum, the structure of the metabolite 2 was formulated as 2-oxo-hinesol.⁵⁾

Compounds 3 and 4 showed the same molecular formula, C₁₅H₂₆O₂, and similar spectral data. Acetylation (Ac₂O, Py) of 3 and 4 afforded the monoacetates 12 and 13, respectively.

The NaBH₄ reduction of 2 afforded 3 (2%) as a minor product and 4 (97%) as a major product, whereas the biotransformation of 1 by *A. niger* afforded 3 and 4 in almost the same yield. Compounds 3 and 12 showed the NOE between H-2 and H-15 in the NOESY spectra (Fig. 1). On the other hand, compounds 4 and 13 showed the NOE between H-2 and H-4 (Fig. 1). Thus the stereostructures of 3 and 4 were formulated as 2α-hydroxyhinesol and 2β-hydroxyhinesol.

Compounds 5 and 6 showed the same molecular formula, C₁₅H₂₆O₃, and similar spectral data. The ¹H-NMR spectra of 5 and 6 showed the presence of an exomethylene (5: δ_H 4.93, 5.16 [br s, H-14]; 6: δ_H 4.58, 4.98 [br s, H-14]). The relative structure of 5 was established by X-ray crystallographic analysis⁷⁾ of 5, as shown in Fig. 2. The structure of 6 was determined to be the 1β, 2α-dihydroxyl isomer of 5 by the NOESY spectrum (Fig. 1) of 6.

Compounds 7 and 8 showed the same molecular formula, C₁₅H₂₄O₃, and similar spectral data. Acetylation of 7 and 8 afforded the diacetates 14 and 15, respectively. The relative structure of 7 was deduced by the NOESY spectrum (Fig. 3) of 14 and finally established by X-ray crystallographic analysis⁸⁾ of 14, as shown in Fig. 4, indicating that 7 contained an intramolecular ether linkage between C-10 and C-11. The structure of 8 was determined to be the 12-hydroxyl isomer of 7 by the NOESY spectrum (Fig. 3) of 15.

The ¹H-NMR spectrum of 9, C₁₅H₂₄O₄ showed the pres-




Fig. 1. The Metabolites of (–)-Hinesol (1) by *A. niger* and *A. cellulosa*

* To whom correspondence should be addressed.




Fig. 2. ORTEP Drawing of Compound 5




Fig. 3. NOESY Spectra of Compounds 14 and 15





Fig. 4. ORTEP Drawing of Compound 14

ence of an epoxy ring (δ_{H} 3.37 [d, $J=3.6$ Hz, H-1]; 3.46 [dd, $J=2.7, 3.6$ Hz, H-2]). Acetylation of **9** afforded the diacetate **16**. Epoxidation with MCPBA of **14** gave **16** in 68% yield, indicating that the structure of **9** was the epoxide of **7**.

The spectral data of **10**, $\text{C}_{15}\text{H}_{26}\text{O}_2$, resembled those of **1**, except for the presence of a primary hydroxyl group (δ_{H} 4.04, 4.16 [d, $J=12.6$ Hz, H-14]) in place of the vinyl methyl group observed in **1**, establishing that the structure of **10** was the C-14-hydroxylated derivative of **1**.

The ^{13}C -NMR spectrum of **11**, $\text{C}_{15}\text{H}_{26}\text{O}_3$, showed the presence of two tertiary alcohols (δ_{C} 74.5, 87.6 [each s]). Acetylation of **11** gave the monoacetate **17**, followed by reaction with 2, 2-dimethoxypropane and *p*-TsOH to give the acetonide **18**. The structure of **11** was determined to be the 7 α , 14-dihydroxyl derivative of **1** by the NOESY spectrum (Fig. 1) of **18**.

In the time course of biotransformation of (–)-hinesol (**1**), it was converted into 2-hydroxyhinesols (**3** and **4**) after 24 h, and both compounds were successively converted into 2-oxohinesol (**2**) and compounds **5**–**9** by rearrangement and intramolecular etherification. 1-Aminobenzotriazole, an inhibitor of cytochrome P-450, inhibited the oxidation process of **1** into **3** and **4**. Possible metabolic pathways of **1** by *A. niger* might be as shown in Fig. 5. It is noteworthy that the

Fig. 5. Possible Metabolic Pathways of Hinesol (**1**) by *A. niger*

metabolic pathways of **1** are strikingly different between *A. niger* and *A. cellulosa*. Intramolecular etherification and rearrangement occurred in *A. niger*, and hydroxydations of the five-membered ring at C-7 and the vinyl methyl group at C-14 of **1** occurred in *A. cellulosa*. The biotransformation of **1** by *A. niger* is very similar to that of oral administration to mammals since **1** was mainly converted into **2**–**4** by rabbits.⁹⁾

References and Notes

- 1) a) Noma Y., Asakawa Y., *Biotechnology in Agriculture and Forestry*, Vol. 28, ed. Bajaj Y.P.S., Springer, Berlin, 1994, pp. 185–202; b) *Idem, ibid.*, Vol. 33, 1995, pp. 62–96; c) *Idem, ibid.*, Vol. 41, 1998, pp. 194–237.
- 2) a) Matsumoto T., Hayashi N., Ishida T., Asakawa Y., *J. Pharm. Sci.*, **79**, 540–547 (1990). b) Matsumoto T., Ishida T., Yoshida T., Terao H., Takeda Y., Asakawa Y., *Chem. Pharm. Bull.*, **40**, 1721–1726 (1992).
- 3) Asakawa Y., Takahashi H., Toyota M., *Phytochemistry*, **30**, 3993–3997 (1991).
- 4) Takahashi H., Hashimoto T., Noma Y., Asakawa Y., *Phytochemistry*, **34**, 1497–1550 (1993).
- 5) Morita M., Nakanishi H., Morita H., Mihashi S., Itokawa H., *Chem. Pharm. Bull.*, **44**, 1603–1604 (1996).
- 6) Asakawa Y., Takahashi H., Toyota M., Noma Y., *Phytochemistry*, **30**, 3981–3987 (1991).
- 7) The crystal data for **5** are as follows: monoclinic; space group $P2_1$ with $a=7.642$ (1), $b=18.353$ (8), $c=10.688$ (5) Å, $\beta=92.50$ (5)°, $V=1497.65$ (1) Å³, $Z=4$, and μ (Cu K- α)=5.765 mm^{–1} by Mac Science MXC 18 instrument. Final R value was 0.067 for 1741 reflections.
- 8) The crystal data for **14** are as follows: monoclinic; space group $P2_1$ with $a=11.349$ (2), $b=7.857$ (2), $c=10.601$ (2) Å, $\beta=99.53$ (1)°, $V=932.2$ (2) Å³, $Z=2$, and μ (Cu K- α)=7.244 mm^{–1} by Mac Science MXC 18 instrument. Final R value was 0.065 for 1604 reflections.
- 9) Lie K., Toyota M., Asakawa Y., unpublished data.