

RESEARCH ARTICLE

Salt stress responsiveness of a wild cotton species (*Gossypium klotzschianum*) based on transcriptomic analysis

Yangyang Wei^{1,2}, Yanchao Xu¹*, Pu Lu¹*, Xingxing Wang¹, Zhenqing Li¹, Xiaoyan Cai¹, Zhongli Zhou¹, Yuhong Wang¹, Zhenmei Zhang¹, Zhongxu Lin^{2*}, Fang Liu^{1*}, Kunbo Wang^{1*}

1 State Key Laboratory of Cotton Biology, / Institute of Cotton Research of Chinese Academy of Agricultural Science (ICR-CAAS), Anyang, Henan, China, **2** National Key Laboratory of Crop Genetic Improvement (Wuhan), Huazhong Agricultural University, Wuhan, Hubei, China

* These authors contributed equally to this work.

* linzhongxu@mail.hzau.edu.cn (ZL); liufcrl@163.com (FL); wkbcri@163.com (KW)



OPEN ACCESS

Citation: Wei Y, Xu Y, Lu P, Wang X, Li Z, Cai X, et al. (2017) Salt stress responsiveness of a wild cotton species (*Gossypium klotzschianum*) based on transcriptomic analysis. PLoS ONE 12(5): e0178313. <https://doi.org/10.1371/journal.pone.0178313>

Editor: Baohong Zhang, East Carolina University, UNITED STATES

Received: March 29, 2017

Accepted: May 11, 2017

Published: May 26, 2017

Copyright: © 2017 Wei et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: RNA-Seq data in this study have been deposited at the NCBI under the accessions: SRR5500539

Funding: This program was financially sponsored by the National Natural Science Foundation of China (31530053), the National Natural Science Foundation of China for Young Scholars (31401424), and the Innovation Scientists and Technicians Troop Construction Projects of Henan Province. The funders had no role in study design,

Abstract

Cotton is a pioneer of saline land crop, while salt stress still causes its growth inhibition and fiber production decrease. Phenotype identification showed better salt tolerance of a wild diploid cotton species *Gossypium klotzschianum*. To elucidate the salt-tolerant mechanisms in *G. klotzschianum*, we firstly detected the changes in hormones, H₂O₂ and glutathione (GSSH and GSH), then investigated the gene expression pattern of roots and leaves treated with 300 mM NaCl for 0, 3, 12, 48 h, and each time control by RNA-seq on the Illumina-Solexa platform. Physiological determination proved that the significant increase in hormone ABA at 48 h, while that in H₂O₂ was at 12 h, likewise, the GSH content decrease at 48 h and the GSSH content increase at 48 h, under salt stress. In total, 37,278 unigenes were identified from the transcriptome data, 8,312 and 6,732 differentially expressed genes (DEGs) were discovered to be involved in salt stress tolerance in roots and leaves, respectively. Gene function annotation and expression analysis elucidated hormone biosynthesis and signal transduction, reactive oxygen species (ROS), and salt overly sensitive (SOS) signal transduction related genes revealed the important roles of them in signal transmission, oxidation balance and ion homeostasis in response to salinity stress. This is a report which focuses on primary response to highly salty stress (upto 300 mM NaCl) in cotton using a wild diploid *Gossypium* species, broadening our understanding of the salt tolerance mechanism in cotton and laying a solid foundation of salt resistant for the genetic improvement of upland cotton with the resistance to salt stress.

Introduction

Abiotic stresses such as salt excess (NaCl), drought, and temperature are among factors most limiting to plant productivity [1]. Plants have to evolve strategies to survive from abiotic and biotic stresses because they cannot escape from stress by moving to places that are more

data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

favorable. Therefore, the complete systems have been formed over the long course of evolution by plants to perceive, transduce, and respond to stresses at the molecular, cellular, and physiological levels. Salt stress is the most serious abiotic stress which threat to agriculture and environment in many parts of the world. It could destroy ion homeostasis, osmotic homeostasis, and lead to redox imbalances, further cause the ion toxicity, oxidative damage, and water-deficit, as a consequence of inhibition of photosynthesis, metabolic dysfunction, and damage to cellular structures within plant cells. In order to survive under such conditions, plants have developed the ability of perceiving and responding to these stresses rapidly via signal transduction pathways [2]. Although the molecular mechanisms are relatively complex comparing to the physiological and biochemical process, an increasing number of studies have focused on this field and have achieved a consensus in model plants such as *Arabidopsis* and rice [3]. The stress information was perceived by an unknown sensor in the plasma, while the signal with transmission result in the activation of the stress-activated transcription factors initiates transcriptional reprogramming. Eventually, some active proteins directly or indirectly participate in the maintenance of osmotic, ion, and redox balance [4, 5].

High concentrations of Na^+ causes osmotic imbalance, membrane disorganization, reduction in growth, inhibition of cell division and expansion, which also leads to reduction in photosynthesis and production of reactive oxygen species [6, 7]. It is clear that various hormones, Ca^{2+} -related and reactive oxygen species (ROS) signaling pathways play the key role in signaling transmission [8]. It is well known that abscisic acid (ABA), ethylene (ET), salicylic acid (SA) and jasmonates (JA) these four kinds of phytohormones play major roles in mediating plant defense response against pathogens and abiotic stresses [9–11]. Especially ABA, which is responsible for plant defense against abiotic stresses because environmental conditions such as drought, salinity, cold, heat stress and wounding are known to trigger increase in ABA levels [12, 13]. Briefly, salt overly sensitive pathway (SOS), which was known as a novel pathway linking the Ca^{2+} signaling in response to salt stress [14, 15], results in the exclusion of excess Na^+ ions out of the cell via the plasma membrane Na^+/H^+ antiporter and helps in reinstating cellular ion homeostasis. A rapid increase in the rate of ROS production, known as ‘the oxidative burst’, occurs as a response to stress conditions, which always depends on the respiratory burst oxidase homolog protein (RBOH) [16]. High ROS levels not only result in oxidative damage to proteins, DNA, and lipids, but also act as signaling molecules to control various processes including pathogen defense, programmed cell death, and stomatal behavior [17–19]. To cope with the oxidative damage resulting from ROS, plants have developed a complex scavenging system, components of this system include enzymatic antioxidants (SOD, APX, PRX, GPX, CAT, GRX and TRX) and non-enzymatic scavengers (AsA, GSH, tocopherols, carotenoids and phenolic compounds) [20, 21]. Many signal transduction pathways involved ROS stress have been reported, such as abscisic acid (ABA) signaling pathway [22, 23], Ca^{2+} signaling pathway [24]. H_2O_2 , one kind of mild ROS, also plays as the signal molecule in the antioxidation [25]. These signaling pathways work together to adaptive the salt stress through activating complex regulatory networks, which control global gene expression, protein modification and metabolite composition [26].

Cotton is a relatively salt tolerant species, but growth inhibition still occurs when the plant is exposed to saline stress. Germination and seedling growth are the two growing procedures particularly with salt sensitive [27, 28]. As such, in order to breed salt-tolerant cotton cultivars, researchers have focused on investigating the key molecular factors involved in the response to salt stress. Previous studies have shown that ectopic expression of cotton CBL-interacting protein kinase gene (*GhCIPK6*) and *SnRK2* could enhance abiotic stress tolerance [29, 30]. Furthermore, over-expressing the vacuolar location *AtNHX1* gene in cotton can also improve salt tolerance [31]. Evidence from transgenic plants has demonstrated the important role of

transcription factors under salt stress in cotton. Such as *GhWRKY39-1* and *GhDREB1* enhance abiotic stress tolerance in transgenic plant [32, 33] and over-expression of the rice NAC gene, *SNAC1*, could also improve salt tolerance in transgenic cotton [34]. Overexpression ROS scavengers, such as *GhSOD1*, *GhCAT1*, and *GhMT3a*, showed the high salt stress tolerance in cotton [35, 36].

In addition, the completion of the *G. raimondii* [37, 38], *G. arboreum* [39], *G. hirsutum* [40, 41], and *G. barbadense* [42] whole genome sequencing offers a new model system for better understanding the adaptive mechanisms in extreme environments. Transcriptomic analysis provides detailed information about gene expression at the mRNA level and is widely used to screen candidate genes involved in stress responses. Recently, several transcriptome studies have contributed to our knowledge of the molecular regulatory pathways to salt stress tolerance and adaptation in cotton, such as miRNA networks by comprehensive analysis of two contrasting cotton genotypes under salt stress [43], such as the protein network of early salt stress responsiveness has also been revealed by iTRAQ in upland cotton [44]. However, major RNA-Seq studies on cultivated cotton have contributed to the knowledge of the mRNA regulatory pathways to salt stress tolerance [45–49], there still is insufficient knowledge of cotton salt tolerance mechanism. So it is necessary that the studies on non-cultivated cotton species, in order to supplement cotton salt tolerance mechanism. Two wild cotton species, *G. aridum* [50] and *G. davidsonii* [51] showed their tolerance to salt stress, whereas the responses were tested on the low NaCl concentration (200 mM) and the response knowledge of high NaCl concentration, e.g. 300 mM NaCl, needs to be understood. Furthermore, few studies took concentrations on the plant hormones response to salt stress in cotton. Hence, we designed the experiment using the wild diploid cotton species *G. klotzschianum* to decipher the mechanism of cotton's rapid response to 300 mM NaCl salinity stress.

Here, a cotton D-genome diploid species, *G. klotzschianum*, was selected for transcriptome analysis to identify the regulation networks involved in responses to salinity stress. On account of its sister species *G. davidsonii* has already been proved salt tolerance by physiological and RNA-Seq analysis under mild stress [51], the same phenotype of *G. klotzschianum* and *G. davidsonii* in various concentration salt stresses demonstrated the salt tolerance of *G. klotzschianum*. Likewise, the contents of four kinds of phytohormones (ABA, ETH, GA, IAA), H₂O₂, AsA, and glutathione in leaves were measured to exam the response mechanism of high salinity tolerance in cotton. The transcriptome analysis of leaves and roots over 48 h (0, 3, 12, and 48) in response to 300 mM NaCl stress were designed to investigate regulatory network at early stage under severe salt stress. The study not only provides supplement data for cotton salt tolerance research but also lays a solid foundation on engineering breeding for salt resistance in cotton.

Materials and methods

Plant materials and salt stress treatments

The plant materials in the study were two diploid D genome wild cotton species *G. klotzschianum* (accession D_{3k}-01) and *G. davidsonii* (accession D_{3d}-01), which were grown in the greenhouses at the Institute of Cotton Research, Chinese Academy of Agricultural Sciences (ICR-CAAS) (Anyang, Henan, China) and also perennially maintained in the National Wild Cotton Nursery which locates at Sanya City, Hainan Island, China. The accession seeds were delinted and sterilized by using 1% of sodium hypochlorite for 15 min, followed by washing with sterile water for three times. Sterilized seeds were germinated at 28°C under long day conditions in a 16 h light/8 h dark cycle with a light intensity of 150 μmol m⁻²s⁻¹ on 17% water content sands. Three days after germination, the properly plants were potted in soil and placed in a growth room at 28°C, 60–70% relative humidity, a photoperiod of 16 h/8 h (day/night) and

light intensity of $150\mu\text{mol m}^{-2}\text{s}^{-1}$. Seedlings containing two simple leaves and one heart-shaped leaf were randomly selected to four groups, one group with three pots, then the plants were watered with 0, 200mM, 250mM, and 300 mM NaCl for 3 days, three biological replicates were prepared for each time point. The leaf and root samples of 300 mM NaCl treatment were harvested for varying durations (0 h, 3 h, 12 h, and 48 h) after exposing to salt stress, the control plants were also harvested at the same time point, three plants were collected and mixed, to minimize the effect of transcriptome unevenness among plants. For the treated leaves and roots, parts of them were used to RNA-seq, and parts of them were used to quantitative real-time PCR (qRT-PCR), parts of leaves were used to detect hormones (ABA, ETH, GA and IAA) and other biochemistry indexes (H_2O_2 , GSH, GSSH and AsA). All the samples were frozen and stored in liquid nitrogen and stored at -70°C for further use (named GKLC0, GKLC3, GKLC12, GKLC48, GKLS3, GKLS12, GKLS48, GKRC0, GKRC3, GKRC12, GKRC48, GKRS3, GKRS12, GKRS48, where GK, L, R, C, and S correspond to *G. klotzschianum*, leaf, root, control and salt treatment, respectively).

Measurement of the hormones, glutathione, H_2O_2 , and AsA contents

The concentration of abscisic acid (ABA), ethylene (ETH), gibberellic acid (GA), and indole acetic acid (IAA) in leaves were determined using the ELISA method. ABA (CK-E00005P, DG, Beijing), ETH (CK-E00021P, DG, Beijing), GA (CK-E00001P, DG, Beijing), IAA (CK-E00015P, DG, Beijing), these four kind of kits were utilized for the determination of each hormone contents. 0.5 gram for each sample applied for the measurement ELISA (Infinite M1000, TECAN, Switzerland). Three replicated experiments (two-time technical repeats per biological replicate) were performed.

The contents of GSH (reduced glutathione) and GSSG (oxidized glutathione) were measured using a GSH and GSSG assay kit (Nanjing Jiancheng Bioengineering Institute). The reaction was initiated by the addition of H_2O_2 . A series of enzymatic reactions were activated by GSH in the homogenate which subsequently led to the conversion of GSH (reduced glutathione) to oxidize glutathione (GSSG). The change in absorbance during the conversion of GSH to GSSG was recorded spectrophotometrically at 405 nm. Three replicated experiments (two-time technical repeats per biological replicate) were performed.

The content of H_2O_2 in leaves was assessed using a commercially available kit (Nanjing Jiancheng Bioengineering Institute). H_2O_2 bound with molybdenic acid to form a complex, which was measured at 405 nm and the content of H_2O_2 was then calculated. The content of AsA in leaves was assessed using a commercially available kit (Nanjing Jiancheng Bioengineering Institute), which was measured at 265 nm and the content of AsA was then calculated. Three replicated experiments (two-time technical repeats per biological replicate) were performed.

RNA extraction, cDNA library construction, and RNA-Seq

Total RNA of each sample was extracted from the cotton samples according to the instruction manual of the TRIzol Reagent (Life technologies, California, USA). RNA integrity and concentration were checked using an Agilent 2100 Bioanalyzer (Agilent Technologies, Inc., Santa Clara, CA, USA). The mRNA was isolated by NEB Next Poly (A) mRNA Magnetic Isolation Module (NEB, E7490). The cDNA library was constructed following the manufacturer's instructions of NEB Next Ultra RNA Library Prep Kit for Illumina (NEB, E7530) and NEB Next Multiplex Oligos for Illumina (NEB, E7500). In briefly, the enriched mRNA was fragmented into approximately 200nt RNA inserts, which were used to synthesize the first-strand cDNA and the second cDNA. The double-stranded cDNA were performed end-repair/dA-tail

and adaptor ligation. The suitable fragments were isolated by Agencourt AMPure XP beads (Beckman Coulter, Inc.), and enriched by PCR amplification. Finally, the constructed cDNA libraries of the cotton were sequenced on a flow cell using an Illumina HiSeq™ 2500 sequencing platform.

Mapping and sequence annotation

Low quality reads, such as only adaptor, unknown nucleotides >5%, or Q20 <20% (percentage of sequences with sequencing error rates <1%), were removed by per script. The clean reads that were filtered from the raw reads were mapped to cotton genome (*G. raimondii*; JGI v2.1) using Tophat2 software [52]. The aligned records from the aligners in BAM/SAM format were further examined to remove potential duplicate molecules. Gene expression levels were estimated using FPKM values (fragments per kilobase of exon per million fragments mapped) by the Cufflinks software [52]. Genes were compared against various protein databases by BLASTX, including the National Center for Biotechnology Information (NCBI) non-redundant protein (Nr) database, Swiss-Prot database with a cut-off E-value of 10^{-5} . Furthermore, genes were searched against the NCBI non-redundant nucleotide sequence (Nt) database using BLASTn by a cut-off E-value of 10^{-5} . Genes were retrieved based on the best BLAST hit (highest score) along with their protein functional annotation. To annotate the gene with gene ontology (GO) terms, the Nr BLAST results were imported into the Blast2GO program [53]. GO annotations for the genes were obtained by Blast2GO. This analysis mapped all of the annotated genes to GO terms in the database and counted the number of genes associated with each term. Perl script was then used to plot GO functional classification for the unigenes with a GO term hit to view the distribution of gene functions. The obtained annotation was enriched and refined using TopGo (R package). The gene sequences were also aligned to the Clusters of Orthologous Group (COG) database to predict and classify functions [54]. KEGG pathways were assigned to the assembled sequences by perl script.

Identification of differential gene expression

DESeq and Q-value were employed and used to evaluate differential gene expression between control and treatment [55]. After that, gene abundance differences between those samples were calculated based on the ratio of the FPKM values. The false discovery rate (FDR) control method was used to identify the threshold of the P-value in multiple tests in order to compute the significance of the differences. Here, only gene with an absolute value of \log_2 ratio ≥ 2 and FDR significance score < 0.01 were used for subsequent analysis.

Validation of RNA-Seq by qRT-PCR

Real-time RT-PCR (qRT-PCR) was performed on a new set of 3 replicates for each sample. A set of 38 genes was chosen randomly (S1 Table) from roots and leaves by dividing their expression levels at different time points with their observed FPKM. In addition, to further verify the correctness of the analysis results, nineteen SOS-, ABA- and ROS-related genes were selected and detected by qRT-PCR (S1 Table-Further Examination). Reverse transcription was conducted with the GoScript™ Reverse Transcription System (Promega, Madison, USA). NCBI primer-BLAST was employed to design specific primers for the chosen genes. Using Fast Start Universal SYBR Green Master (Rox) mix (Roche), real-time PCR was conducted in a 7500 fast Real-Time PCR system (Applied Biosystems), and the results were analysed via the $\Delta\Delta C_t$ method *Gractin7* (F: ATCCTCCGTCTTGACCTTG, R: TGTCCGTCAGGCAACTCAT) gene was used as a control. Each reaction was carried out in a final volume of 20 μ L, containing 10 μ L of SYBR Green PCR master mix, 0.5 μ L of each gene-specific primer and 2 μ L of diluted cDNA.

The PCR thermal cycling conditions were as follows: 95°C for 10 min; 40 cycles of 95°C for 5 s, 60°C for 30 s and 72°C for 30 s. Data were collected during the extension step: 95°C for 15 s, 60°C for 1 min, 95°C for 30 s and 60°C for 15 s. Three biological replicates were performed, and three technical replicates were designed per cDNA sample.

Results

Phenotypic and biochemistry responses to salt stress in *G. klotzschianum*

G. klotzschianum was considered as sister species of *G. davidsonii* in *Gossypum* taxonomy and *G. davidsonii* has been proved with better salt-tolerances by the previous study [51]. Our results showed that *G. klotzschianum* owned the same phenotypic feature to *G. davidsonii* after post in various NaCl concentrations of salt stress for 3 days (Fig 1A). The old leaves of 200 mM and 250 mM NaCl treatments became yellow comparing to control, and dropped down after post at the 300 mM NaCl after 3 days posted salt stress. Furthermore, the contents of ABA were significantly increased at three time points in leaves with 300 mM NaCl treatment as compared to control (Fig 1B-a), and got the maximum value at 48h. In addition the contents




Fig 1. Phenotypic, biochemistry analysis of *G. klotzschianum* in response to salt stress. **A:** Phenotype of *G. klotzschianum* and *G. davidsonii* post salt stress (200 mM, 250 mM, 300 mM NaCl) for three days. **B a-d:** the changes of ABA, ETH, GA, and IAA concentration at different times post 300 mM NaCl stress. **C a-d:** the changes of GSSH, GSH, H₂O₂, and AsA concentration at different times post 300 mM NaCl stress. *G. klotzschianum* seedlings grown under normal conditions were used as controls. Three biological replicates were used. Multiple comparisons were performed with significant difference in different letter at P < 0.05 level; Error bars represent SD.

<https://doi.org/10.1371/journal.pone.0178313.g001>

of ETH and GA also kept increases after post at 300 mM NaCl salt stress at three time points, with the same tendency of ABA (Fig 1B b-c), but the contents of IAA kept decreases lower than control after post at salt stress at three time points, which revealed the completely opposite tendency with ABA (Fig 1B d), all these results suggesting that the different roles of hormones in responding to salt stress. A significant increase in H₂O₂ was observed at 3, 12, and 48 h compared to each of their corresponding controls (Fig 1C a), which showed a statistically significant increase at 12 h. Likewise the GSSH contents were keeping increase under salt stress and maximizing at 48 h, while the GSH contents showed a completely opposite pattern compared with GSSH at the same time point (Fig 1C b-c). Further, the AsA contents remain unchanged within 3 h and 12 h salt treatments, while increases 48 h (Fig 1C d). These findings suggested that these time points might show a substantial change in the expression of salt responsive genes, especially ROS/hormone-related genes. Therefore, RNA-seq was followed to profile the gene expression at each time point to deeply understand the salt stress response in cotton.

Processing and mapping of Illumina reads

Here, we used high-throughput RNA sequencing (RNA-seq) to investigate the changes in gene expression of the diploid D-genome species *G. klotzschianum* at 300 mM NaCl concentration. Three plants were collected and mixed with each sample in this study. In total 14 sequencing libraries were constructed from both the roots and leaves at 0, 3, 12 and 48 h post salt stress and control conditions. All the samples were sequenced by using an Illumina HiSeq™ 2500 sequencing platform. On average of 61.5 million raw reads for the 14 libraries were obtained (Table 1). After the process of adaptor deletion, junk filtering and low copy filtering, > 95% of the sequences were confirmed as clean data. Then the clean reads were mapped to cotton whole genome (*G. raimondii*) using Tophat2 software [52], and 64.39–71.52% of the total reads were mapped to the reference genome, and the unique mapped reads were 62.49%–69.02%. The mapped sequences were assembled with Cufflinks software guided by a reference annotation from JGI Genomes (*G. raimondii* 2.1) [38]. The RNA-Seq assays revealed that a total of 37,278 unigenes and 797 novel genes were found in our data. Among them, the unigenes (18,855, 50.83%) with length exceeds 1000bp showed the more enrichment than the number of unigene (15,591, 42.03%) with the length exceeds 300bp and less-than1000bp.

Table 1. Summary of RNA-Seq results and their matches in the *G. raimondii* genome.

| Samples | Raw Reads(M) | Clean Reads(M) | GC Content | ≥Q30 | Mapped Reads(M) | Uniq Mapped Reads(M) | Multiple Mapped Reads(M) |
|----------------|--------------|----------------|------------|--------|-----------------|----------------------|--------------------------|
| GKLC0 | 71.6 | 71.4 | 44.73% | 96.42% | 71.52% | 69.02% | 2.49% |
| GKLC3 | 60.8 | 60.6 | 44.95% | 96.33% | 71.28% | 68.77% | 2.51% |
| GKLS3 | 53.8 | 53.6 | 44.41% | 96.46% | 69.96% | 68.07% | 1.89% |
| GKLC12 | 51.7 | 51.6 | 44.98% | 94.24% | 66.74% | 64.05% | 2.68% |
| GKLS12 | 58.5 | 58.3 | 44.92% | 94.12% | 65.97% | 62.82% | 3.15% |
| GKLC48 | 48.5 | 48.4 | 44.23% | 94.21% | 66.96% | 64.97% | 1.99% |
| GKLS48 | 63.0 | 60.2 | 44.50% | 85.06% | 64.39% | 62.49% | 1.90% |
| GKRC0 | 66.6 | 66.4 | 44.26% | 96.29% | 70.72% | 67.91% | 2.82% |
| GKRC3 | 49.1 | 49.0 | 44.62% | 96.05% | 69.45% | 66.05% | 3.41% |
| GKRS3 | 77.7 | 77.5 | 44.43% | 95.85% | 69.19% | 66.51% | 2.69% |
| GKRC12 | 72.3 | 72.1 | 44.50% | 94.32% | 66.67% | 63.55% | 3.12% |
| GKRS12 | 70.6 | 70.3 | 44.19% | 95.85% | 67.25% | 65.14% | 2.10% |
| GKRC48 | 60.8 | 60.7 | 44.27% | 94.24% | 65.72% | 63.45% | 2.27% |
| GKRS48 | 55.6 | 55.4 | 44.08% | 94.62% | 66.73% | 65.08% | 1.65% |
| Average | 61.5 | 61.1 | 44.51% | 94.58% | 68.04% | 65.56% | 2.48% |

<https://doi.org/10.1371/journal.pone.0178313.t001>

Exploration of DEGs in roots and leaves in response to salt stress

As the first step in the characterization of *G. klotzschianum* transcriptional responses to salt stress, we carried out the identification of the unigenes with expression level significantly changed upon NaCl-treatments. The transcript abundance of each gene was estimated by FPKM value. DESeq and Q-value were employed and used to evaluate differential gene expression pattern. A cutoff P-value < 0.01 adjusted by false discovery rate (FDR) and fold change ≥ 2 was used for identifying differentially expressed genes. A total of 11417 genes showed differential expression under salt stress conditions (L3 GKLS3 vs. GKLC3, L12 GKLS12 vs. GKLC12, L48 GKLS48 vs. GKLC48, R3 GKRS3 vs. GKRC3, R12 GKRS12 vs. GKRC12, R48 GKRS48 vs. GKRC48). In total, respectively, 8,312 and 6,732 DEGs were found in roots and leaves in response to salt stress. To identify candidate genes that respond to salt stress in *G. klotzschianum*, we adapted the DEGs identified by comparing the gene expression levels under salt stress vs. control conditions at the same time points. Meanwhile, 3,603 DEGs were commonly identified in the two tissue types, which highlights that 1,406 and 782 common genes with different expressions at three time points in roots (RSC RS vs. RC) and leaves (LSC LS vs. LC), respectively. The distribution of these genes is shown in a Venn diagram (Fig 2A). The results revealed that the induction of severe NaCl concentration would be more sensitive than mild NaCl concentration. Interestingly, there were several more DEGs in the roots than the leaves, and the up-regulated DEGs and down-regulated were with roughly the same number in roots, while, the down-regulated genes were more than the up-regulated genes in leaves in response to the salt stress (Fig 2B), which showed the significant inhibition of genes expression in leaves under salt stress. Likewise, 193 common genes were different expressed in the two organs, within them there were 95 genes up-regulated and 88 ones




Fig 2. Summary of DEGs in leaves and roots of *G. klotzschianum* upon salt stress. **A:** Number of DEGs by salt stress under different time points (L3 GKLS3 vs. GKLC3, L12 GKLS12 vs. GKLC12, L48 GKLS48 vs. GKLC48, R3 GKRS3 vs. GKRC3, R12 GKRS12 vs. GKRC12, R48 GKRS48 vs. GKRC48), in leaves and roots. **B:** Comparison of up-/down-regulated genes in roots and leaves. **C:** Number of regulated genes between different conditions (LSC LS vs. LC, RSC RS vs. RC).

<https://doi.org/10.1371/journal.pone.0178313.g002>

down-regulated (Fig 2C). These results are also in line with a transcriptionally higher responsivity of the roots system when compared to leaves', as previously reported, e.g., in the case of two kinds of cotton [50, 51], which indicates tissue-specific responses to salt stress in *G. klotzschianum*.

Differentially regulated genes were also identified by comparing both of adjacent stages and every stage to 0h. "LC3/LC0" indicates a comparison between the gene expression in the GKLC3 library with that in the GKLC0 library, and the same describes the other labels (S2 Table). Interestingly, the up-regulated genes outnumbered the down-regulated genes between two time points under the same treatment.

A cluster analysis of the common different expression genes at three time points in roots (1406) and leaves (782), respectively, resulted a dynamic change of gene expression by a heat map (Fig 3A and 3B), which implied the inhabitation of severe salt concentration in cotton. With comparing the 1406 and 782 DEGs in common at three time points in leaves (LSC LS vs. LC) and roots (RSC RS vs. RC), 193 common DEGs expressed in the two organs at three time points in total (S3 Table). We further observed a general conservation of expression patterns by the heat-map of these genes (S1 Fig), while 183 of the genes were regulated in the same way at all three time points. Among them, nine genes involved in ABA signal pathway were significant up-regulated, include two biosynthesis genes *NCED3* and seven signal transmission genes (six *PP2Cs* and one *ABF*). The results revealed that ABA plays the important role in cotton under salt stress.

In order to validate the differential expression analysis by RNA-seq, 38 genes were chosen randomly from roots and leaves to confirm the veracity of RNA-seq data (S1 Table), including 19 genes for roots and 19 genes for leaves, respectively. Real-time PCR was performed on the same RNA pools that had been previously used for the next-generation sequencing. To corroborate the expression levels measured by RNA-Seq, the ratio of expression levels between salt stressed tissues and controls using RNA-Seq was compared to the ratio of expression as measured by qRT-PCR. A good correlation between RNA-Seq and real-time PCR results (coefficient of determination $R^2 = 0.801$ and 0.905) indicates the reliability of RNA-Seq quantification of gene expression (Fig 4). The validation experiments support the accurateness of the relative values provided by the RNA-Seq analysis.




Fig 3. A: A heat map of 1406 differentially expressed genes from seven root samples. **B:** A heat map of 782 differentially expressed genes from seven leaf samples.

<https://doi.org/10.1371/journal.pone.0178313.g003>




Fig 4. qRT-PCR validation of transcript levels evaluated by RNA-Seq in roots and leaves of *G. klotzschianum* under 300 mM NaCl stress conditions. (A, B) represent the correlation of the fold change analyzed by RNA-Seq platform (x-axis) with data obtained using real time PCR (y-axis) in roots and leaves of *G. klotzschianum*. 19 independent genes for each organ were randomly selected, three biological replicates were used from each sample at three time points post salt stress.

<https://doi.org/10.1371/journal.pone.0178313.g004>

GO and KEGG analysis of the DEGs

To understand DEGs' function, GO category and KEGG pathway enrichment analysis were performed using P-value of 0.01 adjusted by false discovery rate (FDR) as the cutoff. 6,698 (80.58%) and 5,535 (82.22%) of the DEGs in roots and leaves, respectively, were annotated to GO terms. Further GO category enrichment analysis was performed using these DEGs. Respectively, 6,698 and 5,535 DEGs in roots and leaves were classified into 51 and 51 GO terms based on biological process, cellular component and molecular function (Fig 5A and 5B). In the molecular function, as comparing with the background genes, DEGs involved with antioxidant activity (90, 38.30%) nucleic acid binding transcription factor activity (763, 33.54%) were significantly enriched in roots, followed by two GO terms, electron carrier activity (161, 31.76%) and transporter activity (623, 28.53%). Likewise, the genes sorted to the two GO terms electron carrier activity (130, 25.64%) and guanyl-nucleotide exchange factor activity (10, 25.64%) were revealed most different expression as comparing with the background genes in leaves, followed with antioxidant activity (60, 25.53%) and nucleic acid binding transcription factor activity (575, 25,27%). The results demonstrated the important role of the antioxidation under salt stress, which was consistent with the high level of glutathione (Fig 1C). Furthermore, for the number of DEGs, binding (4,072; 3,377) and catalytic activity (3,648; 2,961) showed the significantly enrichment in response to salt stress. For the biological process, genes involved with cell killing (5, 50%) and signaling (1825, 31.39%) were most different expression comparing with




Fig 5. A: GO functional classification of DEGs in roots. B: GO functional classification of DEGs in leaves.

<https://doi.org/10.1371/journal.pone.0178313.g005>

Table 2. GO terms related to stimulus or stress of DEGs.

| GO Term | R3 | R12 | R48 | RSC | L3 | L12 | L48 | LSC |
|---------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|
| response to osmotic stress | 92 | 91 | 69 | 29 | 63 | 35 | 53 | 9 |
| response to oxidative stress | 126 | 140 | 120 | 46 | 101 | 56 | 85 | 18 |
| response to starvation | 14 | 16 | 9 | 4 | 14 | 4 | 5 | 0 |
| response to salt stress | 473 | 505 | 392 | 253 | 424 | 233 | 320 | 129 |
| response to water deprivation | 511 | 536 | 424 | 185 | 461 | 208 | 286 | 80 |
| cation transport | 33 | 34 | 32 | 7 | 27 | 10 | 24 | 2 |
| inorganic anion transport | 9 | 9 | 6 | 3 | 11 | 1 | 5 | 0 |
| monovalent inorganic cation transport | 5 | 5 | 4 | 2 | 4 | 1 | 1 | 0 |
| metal ion transport | 134 | 127 | 117 | 52 | 132 | 68 | 87 | 26 |
| nitrate transport | 178 | 182 | 186 | 75 | 151 | 77 | 127 | 29 |
| response to abscisic acid stimulus | 31 | 36 | 20 | 12 | 31 | 18 | 17 | 7 |
| response to ethylene stimulus | 22 | 20 | 19 | 9 | 16 | 5 | 9 | 2 |
| response to hormone stimulus | 6 | 4 | 3 | 0 | 6 | 2 | 4 | 0 |
| response to jasmonic acid stimulus | 9 | 9 | 8 | 4 | 4 | 2 | 2 | 2 |
| response to salicylic acid stimulus | 9 | 10 | 7 | 5 | 8 | 5 | 6 | 4 |
| response to chitin | 398 | 462 | 291 | 121 | 338 | 134 | 227 | 32 |
| response to sucrose stimulus | 5 | 5 | 3 | 2 | 4 | 3 | 2 | 1 |

<https://doi.org/10.1371/journal.pone.0178313.t002>

background genes in roots under salt stress. While in leaves, cell killing (6, 60%) and growth (1080, 25.63%) showed the most enriched, signaling term was at the third level (1474, 25.35%).

Further GO category enrichment analysis was performed using common DEGs. In our data, 1,191 (84.71%) and 680 (86.96) of the common DEGs in roots and leaves were classified into 44 and 46 GO terms based on biological process, cellular component and molecular function. For the molecular function, most DEGs were sorted into binding (724 genes, 51.49% of the 1406 DEGs; 411 genes, 52.55% of the 782 DEGs) and catalytic activity (684, 48.65%; 355, 45.40%). The GO term of response to stimulus in the biological process section, was significantly enriched with the number of 941 genes, 66.93% of the 1406 DEGs, and 510 genes, 65.22% of the 782 DEGs, respectively. Furthermore, in order to understand the dynamic expression patterns of DEGs under salt stress, several key terms were investigated (Table 2). The results revealed that mass of DEGs enriched in the “response to stress”, “ion transport”, “response to hormone” along with “response to oxidative stress” these four terms at three time points in roots and leaves. Moreover, according to the number of DEGs in the categories of “cation transport”, “inorganic anion transport” and “monovalent inorganic cation transport”, ion stress was not in significantly different expression in leaves, only 2 common DEGs in the category of “cation transport” in leaves, whereas 7, 3, and 2 common DEGs in roots. The results are consistent with previous studies which showed that the salt stress would induce osmotic stress, ion stress, and ROS stress in plants [43, 49].

For the KEGG pathway enrichment analysis, 1452 (17.47%) and 1270 (18.87%) of the DEGs in roots and leaves, respectively, were annotated to KEGG pathway. A set of these DEGs were mapped onto KEGG pathways in *Arabidopsis thaliana* and *Oryza sativa*, highlighting the involvement of several hormone and ROS-related pathways in cotton (Fig 6). The plant hormone signal transduction pathway showed the most significant enrichment in roots (187, 12.88%) and leaves (151, 11.89%). Likewise, the biosynthetic pathways of seven hormones were also enriched in the DEGs (“carotenoid biosynthesis”, “zeatin biosynthesis”, “diterpenoid biosynthesis”, “brassinosteroid biosynthesis”, “cysteine and methionine metabolism”, “alpha-linolenic acid metabolism”, and “tryptophan metabolism”). Our data were in line with the




Fig 6. Hormones and ROS-related pathways under salt stress.

<https://doi.org/10.1371/journal.pone.0178313.g006>

previous studies that plant hormones play crucial roles in a diverse set of developmental processes, as well as in the response to biotic and abiotic stress [56–58]. Further, glutathione metabolism, peroxisome, ascorbate and aldarate metabolism, and oxidative phosphorylation of the four ROS-related pathways were also remarkably different expressed in roots and leaves.

In order to comprehend the dynamic change of gene expression among the two organs, 117, 120, 118, 119, 102, and 115 pathways were individual categorized in R3, R12, R48, L3, L12, and L48, respectively (S4 Table). Based on the ranks of the top 30 pathways from R3, the biosynthetic pathways of seven hormones were enriched in the DEGs (“carotenoid biosynthesis”, “zeatin biosynthesis”, “diterpenoid biosynthesis”, “brassinosteroid biosynthesis”, “cysteine and methionine metabolism”, “alpha-linolenic acid metabolism”, and “tryptophan metabolism”), as well as one pathway for “plant hormone signal transduction”. Interestingly, these eight pathways showed the most enrichment at 3h after post at salt stress, especially in roots. Likewise, we observed that the DEG numbers of hormone synthetic pathways showed the decrease with the duration exposure in at salt stress. The results were consistent with the previous transcriptome researches [59, 60]. Meanwhile, photosynthesis-antenna proteins and photosynthesis from these two photosynthesis-related pathways were observed in decrease at three time points in leaves, suggesting the severe damage of photosynthesis caused by salt stress. The results were implying the rapid response of plant hormone under salt stress.

Analyses on both the GO and KEGG enrichments demonstrated that the genes related to ion transport, ROS, hormones and signal transmission were significantly affected during the early stage of salt stress in cotton. Hence, we further explored that DEGs were involved in the metabolism of ROS and hormones, along with the signal transduction pathways systematically.




Fig 7. Expression profiling of hormone synthesis (A), hormone transduction (B), ROS (C), ion homeostasis (D) related genes.

<https://doi.org/10.1371/journal.pone.0178313.g007>

DEGs involved in hormone biosynthesis and hormone signal transduction

Plant could perceive and respond to stresses rapidly via signal transduction pathways mediated by stress hormones [10, 11]. Hormones, in particular ABA along with cytokinins and ethylene, have been implicated in the root–shoot signaling, the long distance signaling may be mediated particularly via ABA as well as ROS [61]. To examine systematically the effect of salt stress on the pathways of hormone biosynthesis and plant hormone signal transduction, the seven hormones related KEGG pathways (biosynthesis of ABA, ethylene, BRs, JA, auxin, cytokinin, and GA) and plant hormone signal transduction pathway, were further examined manually (Fig 7A, S5 Table).

Under the salt stress, three phytoene synthase genes (*PSY*), two beta-carotene hydroxylase genes (*BCH*), nine 9-cis-epoxycarotenoid dioxygenase genes (*NCED*), and seven abscisic acid 8 hydroxylase genes (*CYP707A*) were significant different expression in the ABA biosynthetic pathway. Furthermore, most of *NCED* genes, an important biosynthetic gene which has been proved the important function on ABA biosynthesis [62], continuous increased at three time points in roots, as well as two *NCED* genes (*Gorai.002G038100*; *Gorai.013G177100*) were

completely activated in two tissues. By contrast, the *CYP707A* genes, which were the key ABA catabolic genes, maintained higher expression at early stage under salt stress, while keep unchanged or down-regulated after exposure in salt stress 48h. Twenty-four genes involved in the ethylene biosynthetic pathway were detected as DEGs. Among them, two S-adenosyl-methionine synthase (*SAM*) genes (*Gorai.008G135900*; *Gorai.005G071300*), precursor synthesis, were continuously upregulated. Meanwhile all seven 1-aminocyclopropane-1-carboxylate (*ACC*) synthase genes (*ACS*) were significantly induced, while half of ten 1-aminocyclopropane-1-carboxylate oxidase genes (*ACO*) were induced, and other five were inhibited gradually. These results showed that the key regulatory components of the biosynthetic pathways of two stress hormones, ABA and ethylene, were changed significantly during cotton's response to salt stress (Fig 1B).

Many studies reported that BRs and JA play essential roles in plant responses to abiotic stress. The expression of thirteen genes involved in BRs biosynthetic pathways changed under salt stress (Fig 7A). Ten cytochrome P450 (*CYP*) genes participating in BRs biosynthesis in cotton, two *CYP749A22* (*Gorai.012G185300*; *Gorai.001G217300*) continuous expression in roots, while *CYP90A1* was only up-regulated before 12 h treatment in roots, and one of two *CYP85A* was activated. Three *CYP734A1* genes and two *CYP724B1* genes significant decreased. Only one genes encoding steroid 5-alpha-reductase (*DET2*), a major rate-limiting enzyme in BRs biosynthesis, showed decrease to more than 2-fold expression levels at 12h treatment in roots. Twenty-seven DEGs were identified involved JA biosynthesis pathway. And the expression of five allene oxide synthase (*AOS*) genes, which are the first enzyme in the branch pathway leading to JA biosynthesis, were significant increase after 3h treatment. These findings indicated that BRs and JA also actively participate in the stress response of cotton.

In addition, 14, 16, and 8 EDGs were also found in the biosynthetic pathway of GA, cytokinin, and auxin, which were widely considered to participate in regulating plant development. Interestingly, many of their genes showed increased transcript abundances in response to salt (Fig 7A).

In *Arabidopsis*, PP2C family genes were demonstrated to regulate ABA signaling pathways negatively by opposing the action of particular protein kinases and eight *PP2Cs* were characterized as key factors in ABA signaling transduction, otherwise *PP2Cs* could inactivate *SnRK2* via dephosphorylation, and this inactivation is inhibited by *PYR/PYL/RCRA*, ABA receptors, in an ABA-dependent manner [63–66]. In our data, thirty-four DEGs were participated in ABA signaling pathway and nine of eleven *PYLs* including *PYL1*, 4 and 8 were significantly down-regulated, especially with 3 members of the *PYL4* continuously down-regulated at both two organs under salt stress (Fig 7B, S6 Table). Fourteen *PP2Cs* were remarkably up-regulated, even six *PP2Cs* were de novo induced at the two tissues. Similarly, three of four *SnRK2s* were induced, the other one *SnRK2* was repressed at 3h time point in roots. Likewise, five of six *ABFs* expression increased, another one just decreased in roots at 12 h and 48 h under salt stress. These results were concord with early report [67], which showed the important role of ABA in the mechanism of salt tolerance in cotton. For ethylene signal pathway, one of two *ETRs* and five *EIN3s* were significant induced, while another *ETR* just decreased at 48 h in leaves under salt stress. Five *ERFs* were found to be changed by NaCl treatments, and most of them were up regulated. As for BRs and JA signal pathways, 16 and 19 genes were detected under salt stress (Fig 7B). Among DEGs in BRs signal pathways, one of each two *BAK1s*, *BKII1s* and *BZR1/2s* was inhibited and the other one in the three DEGs was induced, while two *BRI1s* and eight *CYCD3s* were all decreased. In the JA signal pathway, only one *COI1*, which is major node gene, was found and showed stable decreases. The expressions of nine *JAZs* (TIFY family) and three *MYC2s* increased significantly. In particular, most of the *JAZs*' expressions increased at 3h in roots (Fig 7B).

In addition, 9 DEGs in GA signal transduction were detected. Among them, three *GID1s* were induced, especially the one *Gorai.008G007200* was in continuous expression at three time points in the two organs, whereas four *DELLA* proteins and two *PIFs* were totally inhibited under salt stress. Besides, the DEGs in cytokinin and IAA signal pathways showed similar trends with almost all genes repressed (Fig 7B).

DEGs related with the oxidation-reduction process respond to salt stress

Salt stress induces reactive oxygen species (ROS), which leads to secondary oxidation stress, disturbs cellular redox homeostasis and damages cell components and structures. The alleviation of oxidative damage and increased resistance to salt stresses was often correlated with the balance between ROS producing and ROS scavenging [18]. In the present study, nine *RBOHs* (respiratory burst oxidase homologs), which take the responsibility for the accumulation of H_2O_2 , were significantly up-regulated in both root and leaf tissues when suffering salt stress (Fig 7C; S7 Table). Simultaneously, ROS scavenging systems were markedly influenced, showing different expression patterns under salt treatment and the same models with the *G. davidsonii* transcriptome results [51]. In total, twenty-three members of peroxidase (*POD*) family were up-regulated in the two organs and most of them were up-regulated at 12h treatment in roots. In addition, two catalase (*CAT*) genes were up-regulated one in roots and one in leaves, respectively, and the superoxide dismutase (*SOD*) maintained no change. Most of the twenty-one glutathione S-transferase (*GST*) genes were significantly upregulated. Seven glutaredoxins (*GRXs*) genes and one Glutathione reductase (*GR*) were also upregulated, whereas a completely opposite expression pattern of two *PRX* genes with one upregulated in roots and another down-regulated in leaves under salt treatments was examined. These results were consistent with the high level of GSH (Fig 1C) and suggested that the enzymatic pathways of *POD*, *GST*, and *CAT* gene families play particularly important roles in protecting cotton against oxidative damage under salt stress.

DEGs involved in ion homeostasis and Ca^{2+} -related genes

The damage of ion homeostasis by excess Na^+ will immediately activated Ca^{2+} signal pathway, Ca^{2+} acts as a secondary messenger in salt stress responses [68]. The cytosolic free Ca^{2+} concentration changes within seconds in plants subjected to salt stress. The salt overly sensitive (SOS) signaling pathway, a kind of Ca^{2+} -related pathway, has been well documented at the cellular level, play the vital role in maintain ion homeostasis. In our study, six *SOS2* were upregulated in roots and leaves after post in salt stress (Fig 7D; S8 Table). Whereas, the level of *SOS3* remain unchanged, two *CBL10* genes were significant activated at three time points in roots. *CBL10* (also known as SOS3-like calcium binding protein 8 SCaBP8), is more remarkable in roots in *Arabidopsis* than *SOS3* [69]. Interesting, our result is consistent with the *G. davidsonii* transcriptome consequence that the significant activation of *CBL10* under salt stress in cotton [51]. Further, the Na^+/H^+ antiporter is an important protein in excluding Na^+ at cellular level. In our data, *NHX7/SOS1* remain unchanged, while the vacuolar location gene *NHX2* (*Gorai.007G264500*) was found up-regulated in both tissues, especially in roots. Moreover, other genes involving the ion homeostasis were also found in our data. Such as the Na^+ transporter gene *HKT6* (*Gorai.004G096400*; *Gorai.012G180500*) were down-regulated in roots. Such as the potassium transporters *KUP* (*Gorai.011G051700*; *Gorai.008G049100*; *Gorai.006G029100*; *Gorai.012G171200*; *Gorai.005G267900*), the plasma membrane ATPase (*Gorai.006G023600*; *Gorai.009G125200*) were also found to be up-regulated under salty stress.

In addition, four *CIPK6* genes were upregulated in roots (S9 Table). As the previous studies reported that the tomato *CIPK6* has been reported the interaction with *CBL10* in transmission




Fig 8. Heatmap analysis of the expression levels of ABA, ROS and SOS-related genes. A: A Heatmap by qRT-PCR Log₂ Fold Change in roots and leaves of *G. klotzschianum*. B: A Heatmap by RNA-Seq Log₂ Fold Change in roots and leaves of *G. klotzschianum*. The results were performed by the Fold Change ≥ 2 .

<https://doi.org/10.1371/journal.pone.0178313.g008>

plant immunity signal in transgenic *N. benthamiana* [70], Furthermore, fourteen *CIPKs*, eighteen *CMLs* and eight Ca^{2+} -ATPase genes showed significant activating by salt stress, and most of them up-regulated in roots, which revealed the important role of roots in perception and transmission salt signal.

Examination of SOS-, ABA- and ROS-related genes by qRT-PCR

To further verify the correctness of the analysis results, nineteen typical genes were selected and performed by RT-PCR, which contained ten ABA-related genes included five biosynthesis genes and five signal transduction genes, six ROS-related genes including one ROS-producing gene and five ROS-scavenging genes, three SOS-related genes (S1 Table-Further Examination). The qRT-PCR results revealed a good correlation with RNA-Seq data (Fig 8), which implied the reliability of our analysis results. All the chosen genes were five ABA biosynthesis related genes included one *PSY*, one *BCH*, two *NCEDs*, and one *CYP707A*, five ABA signal transduction genes included one *PYL*, two *PP2Cs*, one *SnRK2*, and one *ABF*, six ROS-related genes included one *RBOH*, one *CAT*, two *PODs*, one *GR*, and one *GST*, and three SOS-related genes consisted of one *CBL10*, one *SOS2*, and one *NHX2*. These genes have been proved the important roles in response to salt stress by the previous studies [51, 62, 63, 67, 69], and the results in our data were in accordance with the previous, which showed the further support of the data veracity and the important roles of these genes in responding to salt stress in cotton.

Identification of transcription factors responsive to salt stress in *G. klotzschianum*

Transcriptional modulation is vital aspects of the complex genetic and biochemical networks that plants use to respond to stress. TFs are master regulators that control gene clusters involved ABA-independent pathway and ABA-dependent pathway [71]. The increased level of ABA promotes the activity of downstream transcription factors to modulate the expression of

various ABA-responsive genes [22], and stress-responsive transcription factors which have been extensively characterized in studies on plant stress tolerance [72]. Members of the *AP2/ERF*, *MYB*, *WRKY*, and *NAC* families were shown to regulate salt tolerance and many studies have shown that TFs could enhance salt tolerance through ABA signaling and the modulation of ROS production [73, 74]. In our data, 2348 TFs were discovered in total, 601 and 796 different expression TFs were identified in leaves and roots, respectively. These TFs were classified into 49 families (S10 Table). The most represented *G. klotzschianum* differentially expressed TFs families were *MYB*, *bHLH*, *ERF*, *NAC*, *bZIP*, and *WRKY*, which are known to mediate stress responses in plants, and the results were consistent with previous studies [75, 76]. The *NAC* family TFs, which is closely related with ABA signal pathway, showed significant up-regulated at the organs under salt stress. Moreover, six tissue-specific regulated TFs *YABBY* family proteins were also found to be altered in our data, as like the previous studies.

Discussion

When salt stress occurs, two main factors are responsible for salt stress-induced inhibition of plant growth: osmotic stress and ionic stress [26]. As it is well known, the stress signal is first perceived by receptors, which results in generation of many secondary signal molecules, such as Ca^{2+} , ROS and hormones [23, 77]. The signal molecules are then transduced into the nucleus, which leads to the activation of related genes to induce responses to salt stress [78]. In present study, RNA-seq was utilized to explore the time course of the response mechanism in cotton under salt stress. Various kinds of salt stress-related DEGs were found and the changes in expression under salt stress were consistent with previous reports. However, only three early time points were investigated in our study, we did concentrate on the ABA-, ROS-, and SOS-related signaling pathways induced by salt stress in the two organs.

In response to salt stress, the generation of ROS (Fig 1C c) may be achieved by activation of ROS-producing genes, such as *RBOHs* (Fig 7C). It has been demonstrated that ROS are produced by *RBOH* in a Ca^{2+} -dependent manner [79]. However, plants have evolved a series of mechanisms to maintain the homeostasis of ROS [18]. In this study, we found that most of the *GPX*, *GST*, and *GRX* genes were expressed at higher levels under salt stress (Fig 7C). The enzymes encoded by these genes play important roles in the reaction of peroxide detoxification by catalyzing GSH to GSSG [80]. The involvement of this process was confirmed by the determination of GSH and GSSG contents (Fig 1C), which showed that the dramatically decrease of the GSH contents, and increase of the GSSG contents. The decrease in the AsA contents (Fig 1C) would also due to ROS scavenging. In addition, *POD* and *CAT* were found to be up-expressed, which was also consistent with previous report in which GhCAT1 acted as a ROS scavenger and participated in salt stress in cotton [36]. No *SOD* with different expression was found, suggesting that *SODs* may not be the main scavenger of ROS under short-term salt stress in cotton. These results suggest that *GST*, *CAT*, and *POD* may be the major scavengers of ROS in salt stress. Prior to their detoxication, genetic evidence suggests that ROS can also act as a signaling molecule in regulating diverse functions in plants [79]. However ROS function, not independently, but synergistically with other signaling pathways, can be regulated by transcription factors.

Hormones, in particular ABA, have been implicated in the stress signal transmission. ABA leads to activation of ion channels in guard cells and stomatal closure [72]. Ma et al. (2012) revealed that ABA would also induce H_2O_2 by activation of NADPH oxidases such as *AtRBOHD* and *AtRBOHF* in *Arabidopsis* [81]. ABA-induced H_2O_2 and then H_2O_2 -activated Ca^{2+} channels are important mechanisms for ABA-induced stomatal closure [82]. Our transcriptional data demonstrated that seven *RBOHs* were upregulated (Fig 7C) and the accumulation

of H_2O_2 was maintained at a higher level (Fig 1C), which in turn triggers increased ABA biosynthesis, leading to further increases in H_2O_2 accumulation. Our results showed that many ABA biosynthetic genes (*PSY*, *BCH*, and *NCED*) were activated under salt stress (Fig 7A), but the ABA contents did not show a significant increase from initial stage (Fig 1C). This might be related to the high expression of four *ABAH4/CYP707A* genes (Fig 7A), which are key genes of ABA degradation to control ABA level [83]. In addition, ROS can affect the ABA signaling pathway by regulating *ABI1* and *ABI2*. In the presence of ABA, PYR/RCARs, as ABA receptors, interact with PP2Cs and inhibit phosphatase activity, allowing SnRK2 activation and phosphorylation of target proteins to control *ABI5* and *RBOH* gene expression [72, 84, 85]. Accordingly, we observed that five members of *RBOHA*, *B*, and *C* were significantly induced under salt stress (Fig 7C), which might be related to ABA signaling. Moreover, various other TFs belonging to several classes including *AP2/ERF*, *MYB*, *NAC*, and *HD-ZF* have been reported to be engaged in ABA-mediated gene expression. These results revealed the crosstalk of ABA and ROS in response to salt stress in cotton.

Once Na^+ enters the cytosol, it can potentially be excluded (back to the soil) by Na^+/H^+ exchangers located in the plasma membrane, or sequestered into the vacuole by Na^+/H^+ exchangers (e.g. NHX proteins) located in the tonoplast [86–88]. The SOS signaling pathway was a well document CBL–CIPK pathway, which plays vital roles in the maintenance of ion homeostasis. Briefly, external stresses lead to the cytosolic free Ca^{2+} concentration changes within seconds, then SOS3 appeared to function as a primary calcium sensor by binding Ca^{2+} and activating SOS2, and the latest the SOS3–SOS2 complex stimulated the Na^+/H^+ exchanger activity of SOS [14], resulting in exclusion of excess Na^+ . In present study, seven antiporting ATPases were up-regulated. Whereas, the expression of SOS3 and SOS1 was unchanged, and several SOS2, two *CBL10*, and one tonoplast location Na^+/H^+ antiporter gene *NHX2* were up-regulated. The results were consistent with those in *G. davidsonii* at the early stage under salt stress [51]. As previous studies, increased calcium was perceived by two calcium-binding proteins, SOS3 and calcineurin B-like10 (CBL10, known as SOS3-like calcium-binding protein8, SCaBP8), which interact with and activate the SOS2 serine/threonine protein kinase [69, 89, 90]. Further, CBL10 has been shown to be an alternative regulator of SOS2 [91, 92]. It was suggested that tonoplast Na^+/H^+ NHX antiporters are activated by AtCIPK24/SOS2 through the mechanism related to the AtCBL10 to sequester intracellular extra Na^+ in the vacuole and AtCBL10–AtCIPK24/SOS2 was reported to regulate Na^+ homeostasis in the vacuolar membrane in shoots and leaves [90, 93]. Therefore, we speculated that CBL10–SOS2 is paramount in the regulation of Na^+ exclusion and cellular ion homeostasis at early stage in cotton. In addition, *HKT6*, which encodes a K^+ transporter, was down-regulated at late stages under salt stress treatment in roots, likely leading to a decrease in Na^+ and K^+ uptake. Interestingly, all *KUPs* in roots and *KUP11* in leaves were up-regulated, likely leading to further K^+ influx and maintaining the K^+ balance of the cell. Together, these signaling pathways are involved in maintaining ion homeostasis to keep the plant away from ion toxicity.

All of the signaling and metabolic pathways described could regulate gene expression involved in protecting plants from salt stress at different levels. Combined with previous studies, our results implied that there might be a connection among H_2O_2 , ABA signal molecules transductions and SOS-related genes to mediate salinity stress in cotton (Fig 9). However, the response to salt stress is an all-dimensional and multitier event. We are still far from determining the exact nature of the full mechanisms of the response to salt stress in cotton.

In conclusion, we presented the first comprehensive transcriptome data from the wild diploid cotton species *G. klotzschianum*. The transcriptome revealed that 8,312 and 6,732 DEGs were discovered to be involved in salt stress tolerance in roots and leaves. Furthermore, the DEGs related to ROS, hormones and ion balance were significantly involved in the response to




Fig 9. Hypothetical interaction network at early stage in response to salt stress in cotton. Black arrow: salt stress induced hormone biosynthesis and signal transduction, likewise, the SOS pathway in the early part of the treatment, which leads to H₂O₂ accumulation; **Blunt arrow:** negative regulation.

<https://doi.org/10.1371/journal.pone.0178313.g009>

salt stress, especially at the initial stage of the salt treated two tissues, which may benefit cotton rapid response to cope with high salinity damage. Based on these findings, the crosstalk between ROS, ABA and Ca²⁺-related genes will be further elucidated in future cotton studies and also be very useful for salt resistance genetic improvement of the cultivated upland cotton.

Supporting information

S1 Fig. Heat map of 193 DEGs from fourteen samples.
(TIF)

S1 Table. Primer list of transcripts for real-time RT-PCR.
(XLSX)

S2 Table. Summary of differentially regulated genes during the time course.
(XLSX)

S3 Table. The common DEGs in roots and leaves.
(XLSX)

S4 Table. KEGG pathway enrichment of differentially expressed genes.
(XLSX)

S5 Table. Differentially expressed genes for hormone synthesis.
(XLSX)

S6 Table. Differentially expressed genes for hormone signal pathways.
(XLS)

S7 Table. Differentially expressed genes for ROS production and scavenging.
(XLSX)

S8 Table. Differentially expressed genes for ion balance.

(XLSX)

S9 Table. Differentially expressed genes for calcium-related.

(XLSX)

S10 Table. Differentially expressed TFs in roots and leaves. Data Section. RNA-Seq data in this study have been deposited at the NCBI under the accessions: SRR5500539.

(XLSX)

Acknowledgments

This program was financially sponsored by the National Natural Science Foundation of China (31530053), the National Natural Science Foundation of China for Young Scholars (31401424), and the Innovation Scientists and Technicians Troop Construction Projects of Henan Province. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Conceptualization: KW FL XW Z. Lin.

Data curation: Y. Wei XC Z. Li PL.

Formal analysis: Y. Wei XC Z. Lin PL.

Funding acquisition: KW FL XW.

Investigation: Y. Wei YX Z. Li PL.

Methodology: KW FL XW.

Project administration: KW FL XW.

Resources: KW FL XW Z. Lin.

Software: Y. Wei.

Supervision: XW XC Z. Zhou Y. Wang Z. Zhang.

Validation: Y. Wei PL YX Z. Li.

Visualization: Y. Wei.

Writing – original draft: Y. Wei.

Writing – review & editing: KW XW Y. Wei.

References

1. Bohnert HJ, Nelson DE, Jensen RG. Adaptations to Environmental Stresses. *Plant Cell*. 1995; 7: 1099–1111. <https://doi.org/10.1105/tpc.7.7.1099> PMID: 12242400
2. Chinnusamy V, Zhu J, Zhu JK. Salt Stress Signaling and Mechanisms of Plant Salt Tolerance. *Springer US*. 2006; 27: 141–177 p.
3. Zhang JL, Shi H. Physiological and molecular mechanisms of plant salt tolerance. *Photosynthesis Research*. 2013; 115: 1–22. <https://doi.org/10.1007/s11120-013-9813-6> PMID: 23539361
4. Atkinson NJ, Urwin PE. The interaction of plant biotic and abiotic stresses: from genes to the field. *Journal of Experimental Botany*. 2012; 63: 3523–3543. <https://doi.org/10.1093/jxb/ers100> PMID: 22467407

5. Huang GT, Ma SL, Bai LP, Zhang L, Ma H, Jia P, et al. Signal transduction during cold, salt, and drought stresses in plants. *Molecular Biology Reports*. 2012; 39: 969–987. <https://doi.org/10.1007/s11033-011-0823-1> PMID: 21573796
6. Niu X, Pardo JM. Ion Homeostasis in NaCl Stress Environments. *Plant Physiology*. 1995; 109: 735–742. PMID: 12228628
7. Parida AK, Das AB. Salt tolerance and salinity effects on plants: a review. *Ecotoxicology & Environmental Safety*. 2005; 60(3): 324–349.
8. Gollack D, Li C, Mohan H, Probst N. Tolerance to drought and salt stress in plants: Unraveling the signaling networks. *Frontiers in Plant Science*. 2014; 5: 151. <https://doi.org/10.3389/fpls.2014.00151> PMID: 24795738
9. Ryu H, Cho YG. Plant hormones in salt stress tolerance. *Journal of Plant Biology*. 2015; 58(3):147–155.
10. Bari R, Jones JD. Role of plant hormones in plant defence responses. *Plant Mol Biol*. 2009; 69(4): 473–488. <https://doi.org/10.1007/s11103-008-9435-0> PMID: 19083153
11. Nakashima K, Yamaguchi-Shinozaki K. ABA signaling in stress-response and seed development. *Plant Cell Rep*. 2013; 32(7): 959–970. <https://doi.org/10.1007/s00299-013-1418-1> PMID: 23535869
12. Lata C, Prasad M. Role of DREBs in regulation of abiotic stress responses in plants. *J Exp Bot*. 2011; 62(14): 4731–4748. <https://doi.org/10.1093/jxb/err210> PMID: 21737415
13. Zhang J, Jia W, Yang J, Ismail AM. Role of ABA in integrating plant responses to drought and salt stresses. *Field Crops Res*. 2006; 97(1): 111–119.
14. Zhu JK, Liu J, Xiong L. Genetic analysis of salt tolerance in *Arabidopsis*. Evidence for a critical role of potassium nutrition. *Plant Cell*. 1998; 10: 1181–1191. PMID: 9668136
15. Zhu JK. Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology*. 2002; 53(1): 247.
16. Baxter A, Mittler R, Suzuki N. ROS as key players in plant stress signalling. *Journal of Experimental Botany*. 2014; 65(5):1229. <https://doi.org/10.1093/jxb/ert375> PMID: 24253197
17. Brosché M, Overmyer K, Wrzaczek M, Kangasjärvi J, Kangasjärvi S. Stress Signaling III: Reactive Oxygen Species (ROS). *Abiotic Stress Adaptation in Plants*. Springer Netherlands. 2009; pp. 91–102.
18. Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R. Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant Cell & Environment*. 2010; 33: 453–467.
19. Petrov V, Hille J, Mueller-Roeber B, Gechev TS. ROS-mediated abiotic stress-induced programmed cell death in plants. *Frontiers in Plant Science*. 2015; 6: 69. <https://doi.org/10.3389/fpls.2015.00069> PMID: 25741354
20. Foyer CH, Noctor G. Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell*. 2005; 17: 1866–1875. <https://doi.org/10.1105/tpc.105.033589> PMID: 15987996
21. Pang CH, Wang BS. *Oxidative Stress and Salt Tolerance in Plants*: Springer Berlin Heidelberg. 2008; 69: 231–245
22. Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park SY, et al. In vitro reconstitution of an abscisic acid signalling pathway. *Nature*. 2009; 462: 660–664. <https://doi.org/10.1038/nature08599> PMID: 19924127
23. Chater CC, Oliver J, Casson S, Gray JE. Putting the brakes on: abscisic acid as a central environmental regulator of stomatal development. *New Phytologist*. 2014; 202: 376–391. <https://doi.org/10.1111/nph.12713> PMID: 24611444
24. Dubiella U, Seybold H, Durian G, Komander E, Lassig R, Witte CP, et al. Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for rapid defense signal propagation. *Proceedings of the National Academy of Sciences of the United States of America*. 2013; 110: 8744–8749. <https://doi.org/10.1073/pnas.1221294110> PMID: 23650383
25. Bianka Steffens SHMS. Reactive oxygen species mediate growth and death in submerged plants. *Frontiers in Plant Science*. 2013; 4: 179. <https://doi.org/10.3389/fpls.2013.00179> PMID: 23761805
26. Julkowska MM, Testerink C. Tuning plant signaling and growth to survive salt. *Trends in Plant Science*. 2015; 20: 586–594. <https://doi.org/10.1016/j.tplants.2015.06.008> PMID: 26205171
27. Elzahab AAA. Salt tolerance of eight Egyptian cotton varieties. I. At germination stage. *Z Acker—und Pflanz*. 1971.
28. Zhong H, Läuchli A. Spatial and Temporal Aspects of Growth in the Primary Root of Cotton Seedlings: Effects of NaCl and CaCl₂. *Journal of Experimental Botany*. 1993; 44: 763–771.
29. He L, Yang X, Wang L, Zhu L, Zhou T, Deng J. et al. Molecular cloning and functional characterization of a novel cotton CBL-interacting protein kinase gene (*GhCIPK6*) reveals its involvement in multiple

- abiotic stress tolerance in transgenic plants. *Biochemical & Biophysical Research Communications*. 2013; 435: 209–215.
30. Bello B, Zhang X, Liu C, Yang Z, Wang Q, et al. Cloning of *Gossypium hirsutum* sucrose non-fermenting 1-related protein kinase 2 gene (*GhSnRK2*) and its overexpression in transgenic *Arabidopsis* escalates drought and low temperature tolerance. *Plos One*. 2014; 9: e112269. <https://doi.org/10.1371/journal.pone.0112269> PMID: 25393623
 31. He C, Yan J, Shen G, Fu L, Holaday AS, Blumwald E, et al. Expression of an *Arabidopsis* vacuolar sodium/proton antiporter gene in cotton improves photosynthetic performance under salt conditions and increases fiber yield in the field. *Plant Cell Physiol*. 2005; 46: 1848–1854. <https://doi.org/10.1093/pcp/pci201> PMID: 16179357
 32. Shi W, Hao L, Li J, Liu D, Guo X, Li H. The *Gossypium hirsutum* WRKY gene *GhWRKY39-1* promotes pathogen infection defense responses and mediates salt stress tolerance in transgenic *Nicotiana benthamiana*. *Plant Cell Reports*. 2014; 33: 483–498. <https://doi.org/10.1007/s00299-013-1548-5> PMID: 24337818
 33. Huang JG, Yang M, Liu P, Yang GD, Wu CA, Zheng CC. *GhDREB1* enhances abiotic stress tolerance, delays GA-mediated development and represses cytokinin signalling in transgenic *Arabidopsis*. *Plant Cell & Environment*. 2009; 32: 1132–1145.
 34. Liu G, Li X, Jin S, Liu X, Zhu L, Nie Y, et al. Overexpression of rice NAC gene *SNAC1* improves drought and salt tolerance by enhancing root development and reducing transpiration rate in transgenic cotton. *Plos One*. 2014; 9: e86895. <https://doi.org/10.1371/journal.pone.0086895> PMID: 24489802
 35. Xue T, Li X, Zhu W, Wu C, Yang G, Zheng C. Cotton metallothionein *GhMT3a*, a reactive oxygen species scavenger, increased tolerance against abiotic stress in transgenic tobacco and yeast. *Journal of Experimental Botany*. 2009; 60: 339–349. <https://doi.org/10.1093/jxb/em291> PMID: 19033550
 36. Luo X, Wu J, Li Y, Nan Z, Xing G, Wang Y, et al. Synergistic Effects of *GhSOD1* and *GhCAT1* Overexpression in Cotton Chloroplasts on Enhancing Tolerance to Methyl Viologen and Salt Stresses. *Plos One*. 2013; 8: e54002. <https://doi.org/10.1371/journal.pone.0054002> PMID: 23335985
 37. Wang K, Wang Z, Li F, Ye W, Wang J, Song G, et al. The draft genome of a diploid cotton *Gossypium raimondii*. *Nature Genetics*. 2012; 44: 1098–1103. <https://doi.org/10.1038/ng.2371> PMID: 22922876
 38. Paterson AH, Wendel JF, Gundlach H, Guo H, Jenkins J, Jin D, et al. Repeated polyploidization of *Gossypium* genomes and the evolution of spinnable cotton fibres. *Nature*. 2012; 492: 423–427. <https://doi.org/10.1038/nature11798> PMID: 23257886
 39. Li F, Fan G, Wang K, Sun F, Yuan Y, Song G, et al. Genome sequence of the cultivated cotton *Gossypium arboreum*. *Nature Genetics*. 2014; 46: 567–572. <https://doi.org/10.1038/ng.2987> PMID: 24836287
 40. Li F, Fan G, Lu C, Xiao G, Zou C, Kohel RJ, et al. Genome sequence of cultivated Upland cotton (*Gossypium hirsutum* TM-1) provides insights into genome evolution. *Nature Biotechnology*. 2015; 33: 524–530. <https://doi.org/10.1038/nbt.3208> PMID: 25893780
 41. Zhang T, Hu Y, Jiang W, Fang L, Guan X, Chen J, et al. Sequencing of allotetraploid cotton (*Gossypium hirsutum* L. acc. TM-1) provides a resource for fiber improvement. *Nature Biotechnology*. 2015; 33: 531–537. <https://doi.org/10.1038/nbt.3207> PMID: 25893781
 42. Yuan D, Tang Z, Wang M, Gao W, Tu L, Jin X, et al. The genome sequence of Sea-Island cotton (*Gossypium barbadense*) provides insights into the allopolyploidization and development of superior spinnable fibres. *Scientific Reports*. 2015; 5: 17662. <https://doi.org/10.1038/srep17662> PMID: 26634818
 43. Peng Z, He S, Gong W, Sun J, Pan Z, Xu F, et al. Comprehensive analysis of differentially expressed genes and transcriptional regulation induced by salt stress in two contrasting cotton genotypes. *BMC Genomics*. 2014; 15: 760. <https://doi.org/10.1186/1471-2164-15-760> PMID: 25189468
 44. Wu L, Zhao F, Fang W, Xie D, Hou J, Yang X, et al. Identification of early salt stress responsive proteins in seedling roots of upland cotton (*Gossypium hirsutum* L.) employing iTRAQ-based proteomic technique. *Frontiers in Plant Science*. 2015; 6: 732. <https://doi.org/10.3389/fpls.2015.00732> PMID: 26442045
 45. Park W, Scheffler BE, Bauer PJ, Campbell BT. Genome-wide identification of differentially expressed genes under water deficit stress in upland cotton (*Gossypium hirsutum* L.). *BMC Plant Biology*. 2012; 12: 90. <https://doi.org/10.1186/1471-2229-12-90> PMID: 22703539
 46. Yin Z, Li Y, Yu J, Liu Y, Li C, Han X, et al. Difference in miRNA expression profiles between two cotton cultivars with distinct salt sensitivity. *Molecular Biology Reports*. 2012; 39: 4961–4970. <https://doi.org/10.1007/s11033-011-1292-2> PMID: 22160515
 47. Rodriguez-Urbe L, Higbie SM, Stewart JM, Wilkins T, Lindemann W, Sengupta-Gopalan C, et al. Identification of salt responsive genes using comparative microarray analysis in Upland cotton (*Gossypium hirsutum* L.). *Plant Science*. 2011; 180: 461–469. <https://doi.org/10.1016/j.plantsci.2010.10.009> PMID: 21421393

48. Zhu YN, Shi DQ, Ruan MB, Zhang LL, Meng ZH, Liu J, et al. Transcriptome Analysis Reveals Crosstalk of Responsive Genes to Multiple Abiotic Stresses in Cotton (*Gossypium hirsutum* L.). Plos One. 2013; 8: e80218. <https://doi.org/10.1371/journal.pone.0080218> PMID: 24224045
49. Zhang X, Yao D, Wang Q, Xu W, Wei Q, Wang CC, et al. mRNA-seq analysis of the *Gossypium arbor-eum* transcriptome reveals tissue selective signaling in response to water stress during seedling stage. Plos One. 2013; 8: e54762. <https://doi.org/10.1371/journal.pone.0054762> PMID: 23382961
50. Xu P, Liu Z, Fan X, Gao J, Zhang X, Yang Y, et al. De novo transcriptome sequencing and comparative analysis of differentially expressed genes in *Gossypium aridum* under salt stress. Gene. 2013; 525: 26–34. <https://doi.org/10.1016/j.gene.2013.04.066> PMID: 23651590
51. Zhang F, Zhu G, Du L, Shang X, Cheng C, Yang B, et al. Genetic regulation of salt stress tolerance revealed by RNA-Seq in cotton diploid wild species, *Gossypium davidsonii*. Scientific Reports. 2016; 6: 20582. <https://doi.org/10.1038/srep20582> PMID: 26838812
52. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biology. 2013; 14(4): R36. <https://doi.org/10.1186/gb-2013-14-4-r36> PMID: 23618408
53. Götz S, Garcíagómez JM, Terol J, Williams TD, Nagaraj SH, José M, et al. High-throughput functional annotation and data mining with the Blast2GO suite. Nucleic Acids Research. 2008; 36: 3420–3435. <https://doi.org/10.1093/nar/gkn176> PMID: 18445632
54. Tatusov RL, Galperin MY, Natale DA, Koonin EV. The COG database: a tool for genome-scale analysis of protein functions and evolution. Nucleic Acids Research. 2000; 28: 33–36. PMID: 10592175
55. Anders S, Huber W. Differential expression analysis for sequence count data. Genome Biology. 2010; 11: R106. <https://doi.org/10.1186/gb-2010-11-10-r106> PMID: 20979621
56. Vaadia Y. Plant Hormones and Water Stress. Philosophical Transactions of the Royal Society of London. 1976; 273: 513–522.
57. Kaya C, Tuna AL, Yokas I. The Role of Plant Hormones in Plants Under Salinity Stress: Springer Netherlands. 2009; 45–50 p.
58. Fahad S, Nie L, Chen Y, Wu C, Xiong D, Saud S, et al. Crop Plant Hormones and Environmental Stress: Springer International Publishing. 2015; 371–400 p.
59. Zheng L, Meng Y, Ma J, Zhao X, Cheng T, Ji J, et al. Transcriptomic analysis reveals importance of ROS and phytohormones in response to short-term salinity stress in *Populus tomentosa*. Frontiers in Plant Science. 2015; 6: 678. <https://doi.org/10.3389/fpls.2015.00678> PMID: 26442002
60. Dang ZH, Zheng LL, Wang J, Gao Z, Wu SB, Qi Z, et al. Transcriptomic profiling of the salt-stress response in the wild recretahalophyte *Reaumuria trigyna*. BMC Genomics. 2013; 14: 29. <https://doi.org/10.1186/1471-2164-14-29> PMID: 23324106
61. Lake JA, Woodward FI, Quick WP. Long-distance CO₂ signalling in plants. Journal of Experimental Botany. 2002; 53: 183–193. PMID: 11807121
62. North HM, De AA, Boutin JP, Frey A, To A, Botran L, et al. The *Arabidopsis* ABA-deficient mutant *aba4* demonstrates that the major route for stress-induced ABA accumulation is via neoxanthin isomers. Plant Journal. 2007; 50: 810–824. <https://doi.org/10.1111/j.1365-313X.2007.03094.x> PMID: 17470058
63. Fujita Y, Fujita M, Shinozaki K, Yamaguchi-Shinozaki K. ABA-mediated transcriptional regulation in response to osmotic stress in plants. Journal of Plant Research. 2011; 124: 509–525. <https://doi.org/10.1007/s10265-011-0412-3> PMID: 21416314
64. Lee SC, Luan S. ABA signal transduction at the crossroad of biotic and abiotic stress responses. Plant Cell & Environment. 2011; 35: 53–60.
65. Ye N, Jia L, Zhang J. ABA signal in rice under stress conditions. Rice. 2012; 5: 1–9. <https://doi.org/10.1186/1939-8433-5-1> PMID: 24764501
66. Yoshida T, Mogami J, Yamaguchishinozaki KABA-dependent and ABA-independent signaling in response to osmotic stress in plants. Current Opinion in Plant Biology. 2014; 21: 133–139. <https://doi.org/10.1016/j.pbi.2014.07.009> PMID: 25104049
67. Finkelstein R. Abscisic Acid synthesis and response. *Arabidopsis* Book. 2013; 11: e166.
68. Luan S. The CBL-CIPK network in plant calcium signaling. Trends in Plant Science. 2009; 14: 37. <https://doi.org/10.1016/j.tplants.2008.10.005> PMID: 19054707
69. Quan R, Lin H, Mendoza I, Zhang Y, Cao W, Yang YQ, et al. SCABP8/CBL10, a putative calcium sensor, interacts with the protein kinase SOS2 to protect *Arabidopsis* shoots from salt stress. Plant Cell. 2007; 19: 1415. <https://doi.org/10.1105/tpc.106.042291> PMID: 17449811
70. De la Torre F, Gutiérrez-Beltrán E, Pareja-Jaime Y, Chakravarthy S, Martin GB, del Pozo O. The Tomato Calcium Sensor Cbl10 and Its Interacting Protein Kinase Cipk6 Define a Signaling Pathway in

- Plant Immunity. *The Plant Cell*. 2013; 25(7): 2748–2764. <https://doi.org/10.1105/tpc.113.113530> PMID: 23903322
71. Yamaguchishinozaki K, Shinozaki K. Organization of cis-acting regulatory elements in osmotic- and cold-stress-responsive promoters. *Trends in Plant Science*. 2005; 10: 88–94. <https://doi.org/10.1016/j.tplants.2004.12.012> PMID: 15708346
 72. Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. Abscisic Acid: Emergence of a Core Signaling Network. *Annual Review of Plant Biology*. 2010; 61: 651–679. <https://doi.org/10.1146/annurev-arplant-042809-112122> PMID: 20192755
 73. Jung J, Kim M. The barley ERF-type transcription factor HvRAF confers enhanced pathogen resistance and salt tolerance in *Arabidopsis*. *Planta*. 2007; 225: 575–588. <https://doi.org/10.1007/s00425-006-0373-2> PMID: 16937017
 74. Yun KY, Park MR, Mohanty B, Herath V, Xu F, Mauleon R, et al. Transcriptional regulatory network triggered by oxidative signals configures the early response mechanisms of japonica rice to chilling stress. 2010; 10: 16. <https://doi.org/10.1186/1471-2229-10-16> PMID: 20100339
 75. Ma J, Lu J, Xu J, Duan B, He X, Liu JQ. Genome-wide Identification of *WRKY* Genes in the Desert Poplar *Populus euphratica* and Adaptive Evolution of the Genes in Response to Salt Stress. *Evolutionary Bioinformatics Online*. 2015; 11: 47–55. <https://doi.org/10.4137/EBO.S22067> PMID: 26309388
 76. Liu Y, Ji X, Zheng L, Nie X, Wang Y. Microarray Analysis of Transcriptional Responses to Abscisic Acid and Salt Stress in *Arabidopsis thaliana*. *International Journal of Molecular Sciences*. 2013; 14(5): 9979–9998. <https://doi.org/10.3390/ijms14059979> PMID: 23665901
 77. Cui JX, Zhou YH, Ding JG, Xia XJ, Shi K, Shen SC, et al. Role of nitric oxide in hydrogen peroxide-dependent induction of abiotic stress tolerance by brassinosteroids in cucumber. *Plant Cell & Environment*. 2011; 34: 347–358.
 78. Chen JH, Jiang HW, Hsieh EJ, Chen HY, Chien CT, Hsieh HL. Drought and Salt Stress Tolerance of an *Arabidopsis* Glutathione S-Transferase U17 Knockout Mutant Are Attributed to the Combined Effect of Glutathione and Abscisic Acid. *Plant Physiology*. 2012; 158: 340–351. <https://doi.org/10.1104/pp.111.181875> PMID: 22095046
 79. Kurusu T, Kuchitsu K, Tada Y. Plant signaling networks involving Ca²⁺ and Rboh/Nox-mediated ROS production under salinity stress. *Frontiers in Plant Science*. 2015; 6: 427. <https://doi.org/10.3389/fpls.2015.00427> PMID: 26113854
 80. Marí M, Morales A, Colell A, García Ruiz C, Fernándezchecha JC. Mitochondrial glutathione, a key survival antioxidant. *Antioxidants & Redox Signaling*. 2009; 11 (11): 2685–2700.
 81. Ma L, Zhang H, Sun L, Jiao Y, Zhang G, Chen M, et al. NADPH oxidase AtrbohD and AtrbohF function in ROS-dependent regulation of Na⁺/K⁺ homeostasis in *Arabidopsis* under salt stress. *Journal of Experimental Botany*. 2012; 63: 305–317. <https://doi.org/10.1093/jxb/err280> PMID: 21984648
 82. Pei ZM, Murata Y, Benning G, Thomine S, Klüsener B, Allen GJ, et al. Calcium channels activated by hydrogen peroxide mediate abscisic acid signaling in guard cells. *Nature*. 2000; 406, 731–734. <https://doi.org/10.1038/35021067> PMID: 10963598
 83. Matakadiadis T, Alboresi A, Jikumaru Y, Tatematsu K, Pichon O, Renou JP, et al. The *Arabidopsis* abscisic acid catabolic gene *CYP707A2* plays a key role in nitrate control of seed dormancy. *Plant Physiology*. 2009; 149: 949–960. <https://doi.org/10.1104/pp.108.126938> PMID: 19074630
 84. Hubbard KE, Nishimura N, Hitomi K, Getzoff ED, Schroeder JI. Early abscisic acid signal transduction mechanisms: newly discovered components and newly emerging questions. *Genes & Development*. 2010; 24: 1695.
 85. Peleg Z, Blumwald E. Hormone balance and abiotic stress tolerance in crop plants. *Current Opinion in Plant Biology*, 2011, 14(3):290–295. <https://doi.org/10.1016/j.pbi.2011.02.001> PMID: 21377404
 86. Apse MP, Aharon GS, Snedden WA, Blumwald E. Salt Tolerance Conferred by Overexpression of a Vacuolar Na⁺/H⁺ Antiport in *Arabidopsis*. *Science*. 1999; 285: 1256–1258. PMID: 10455050
 87. Qiu QS, Guo Y, Dietrich MA, Schumaker KS, Zhu JK. Regulation of SOS1, a plasma membrane Na⁺/H⁺ exchanger in *Arabidopsis thaliana*, by SOS2 and SOS3. *Proceedings of the National Academy of Sciences of the United States of America*. 2002; 99: 8436–8441. <https://doi.org/10.1073/pnas.122224699> PMID: 12034882
 88. Yokoi S, Quintero FJ, Cubero B, Ruiz MT, Bressan RA, Hasegawa PM, et al. Differential expression and function of *Arabidopsis thaliana* NHX Na⁺/H⁺ antiporters in the salt stress response. *Plant Journal*. 2002; 30: 529–539. PMID: 12047628
 89. Halfter U, Ishitani M, Zhu JK. The *Arabidopsis* SOS2 protein kinase physically interacts with and is activated by the calcium-binding protein SOS3. *Proc Natl Acad Sci USA*. 2000; 97: 3735. <https://doi.org/10.1073/pnas.040577697> PMID: 10725350

90. Kim BG, Waadt R, Cheong YH, Pandey GK, Dominguez-Solis JR, Stefanie S, et al. The calcium sensor CBL10 mediates salt tolerance by regulating ion homeostasis in *Arabidopsis*. *Plant Journal*. 2007; 52: 473–484. <https://doi.org/10.1111/j.1365-313X.2007.03249.x> PMID: 17825054
91. Du W, Lin H, Chen S, Wu Y, Zhang J, Fuglsang AT, et al. Phosphorylation of SOS3-like calcium-binding proteins by their interacting SOS2-like protein kinases is a common regulatory mechanism in *Arabidopsis*. *Plant Physiology*. 2011; 156: 2235. <https://doi.org/10.1104/pp.111.173377> PMID: 21685179
92. Lin H, Guo Y. Phosphorylation of SOS3-LIKE CALCIUM BINDING PROTEIN8 by SOS2 Protein Kinase Stabilizes Their Protein Complex and Regulates Salt Tolerance in *Arabidopsis*. *Plant Cell*. 2009; 21: 1607–1619. <https://doi.org/10.1105/tpc.109.066217> PMID: 19448033
93. Cheng NH, Pittman JK, Zhu JK, Hirschi KD. The protein kinase SOS2 activates the *Arabidopsis* H⁺/Ca²⁺ antiporter CAX1 to integrate calcium transport and salt tolerance. *The Journal of biological chemistry*. 2004; 279: 2922–2926. <https://doi.org/10.1074/jbc.M309084200> PMID: 14583601