

Malaysian Journal of Microbiology

Published by Malaysian Society for Microbiology (In SCOPUS since 2011)



Characteristics of a red-pigmented thermotolerant bacteria (GSB-001) isolated from Gedong Songo hot spring, Bandungan, Semarang – Indonesia

Endang Kusdiyantini^{1*}, Iffan Alif¹, Salma Fuadiyah^{1,2}, Muhammad Fadhil Randa Putra¹, Dyah Wulandari² and Anto Budiharjo^{2,3}

¹ Department of Biology, Faculty of Science and Mathematics, Diponegoro University, Indonesia.
²Molecular and Applied Microbiology Laboratory-Central Laboratory of Research and Service-Diponegoro University, Indonesia.

³Biotechnology Study Program, Faculty of Sciences and Mathematics, Diponegoro University, Indonesia. Email: <u>ekusdiyantini@gmail.com</u>

Received 26 February 2019; Received in revised form 13 June 2019; Accepted 8 July 2019

ABSTRACT

Aims: Pigments used as food additives for a long time and people prefer natural pigment than synthetic ones because of their safety. Microorganisms are interesting for pigment production and many of them have been used in food industry. Many research have been conducted to find out for natural pigment sources. The objective of this research was to investigate the characteristics of a red-pigmented bacteria isolated from Gedong Songo hot spring, Bandungan-Indonesia.

Methodology and results: Bacterial isolates were grown on Nutrient Agar for 24 h, and the morphology of the colonies and of the cells were identified. Biochemical tests included indole, methyl red, catalase, urease tests and carbohydrate fermentation. Molecular identification was based on 16S rRNA sequence. The isolate was a rod shape and Gramnegative. Biochemical tests showed that the isolate was indole negative, catalase positive, methyl red negative and urease negative. This isolate was glucose, maltose and sucrose positive and negative for lactose. 16S rRNA sequence was BLAST and it matched with *Serratia marcescens* strain S823. The red pigment antioxidant activity showed the highest DPPH radical scavenging of 49.11% obtained from 48 h of incubation. Functional group of *S. marcescens* pigment on Fourier Transform InfraRed Spectroscopy (FTIR) displayed specific peak at 1740 cm⁻¹ represented of C=O (carbonyl) stretching group.

Conclusion, significant and impact of study: Based on morphological, biochemical and moleculer identifications, it showed that the bacteria isolated from Gedong Songo hot spring Bandungan-Indonesia had 94% homology with *Serratia marcescens* strain S823. Based on DPPH radical scavenging test, it demonstrated that *S. marcescens* had potency as an antioxidant.

Keywords: Red-pigmented thermotolerant bacteria, antioxidant, Serratia marcescens

INTRODUCTION

Carotenoids are pigment consisting of C_{40} in living organisms, they are synthesized by all photosynthetic organisms and few non photosynthetic organisms such as bacteria and fungi. Carotenoids are various molecules containing carbon and hydrogen such as β -carotene, γ -carotene, lycopene and xanthophylls such as astaxanthin, cantaxanthin, astacene. Among natural resources of pigments, microbial resources have an immense potential to produce good alternative ones to synthetic pigments. Microbial pigments are more promising producers than other natural resources of pigments because they are considered as natural, having no seasonal production problems, growing quikly in cheap culture media and

showing high productivity. They grow faster compared to plants and animals and are relatively easy to be manipulated and their growth conditions are easy to produce high pigment content. Thus microbial pigments production are nowadays one of the emerging fields of research to demonstrate their potential for various industrial applications (Kumar al., et 2015). Microorganisms produce many natural pigments such as β-carotene, astaxanthin, canthanxanthin, prodigiosin (Elkenawy *et al.*, 2017; Bathgare *et al.*, 2018). Considering the important use of pigments in various industries, especially food, cosmetics and health, the search for natural pigments from extreme environments such as hot springs is encouraged (Manimala and Murugesan, 2017; Mukherjee et al., 2017). Pigment

*Corresponding author

compounds contains bioactive compounds functioned as antioxidant. This is due to the pigment structure can prevent oxidation or neutralizing oxidized compound by adding hydrogen or electron.

Microorganisms from different environment have been isolated to substitute synthetic pigments for industries, like fungal Monascus for its monascorubramine redpigment (Srianta et al., 2016) and the yeast Phaffia rhodozyma for its astaxantin used as natural feed supplement in aquaculture (Cheng et al., 2018). The others microorganisms pigment producer are the yeasts Rhodotorula sp., Rhodosporidium sp. for synthesis of torulen, torulahordin and β-carotene (Tkačova, 2015; Bonadio et al., 2018;). Many bacteria producing pigment are: Rhodococcus sp., Serratia sp., Rhodotorula sp. (Hernández-Almanza et al., 2014; Zang et al., 2014; Chen et al., 2018). Microalgae of Dunaliella sp., Haematococcus pluvialis, Chlorella sp. are also commercial pigment producing (Kleinegris et al., 2011; Shah et al., 2016).

Our laboratory has isolated red-pigmented thermotolerant bacteria from sediment of hot springs at Gedong Songo in Bandungan – Semarang. Bandungan is one of subdistricts in Semarang Regency, where it is a highland tourist destination with 1200 m above sea level. It has hot springs and geothermal steam vent aerosol, which makes Bandungan a unique environment. Hence, it has microorganisms which are different from other places. Therefore, identification of the isolate found in this region is needed to determine the species.

The purposes of this research were to isolate and identify pigmented thermotolerant bacteria based on 16S rRNA, biochemical and morphological characteristics.

MATERIALS AND METHODS

The red pigmented bacteria was cultivated on Nutrient Agar (NA) at pH 6 and temperature at 28 °C. This culture was conserved at temperature 4 °C.

Morphological and biochemical identifications

The isolate was cultivated on Nutrient Agar media, incubated at ambiant temperature, for 24 h. The colony was identified for its colony characters, cell shape, and Gram staining. Biochemical tests included indole, methyl red, catalase, urease test and carbohydrate fermentation using carbon sources of glucose, lactose, sucrose and maltose (Talaeikhozani *et al.*, 2015).

Molecular Identification

DNA extraction

DNA extraction was done by Chelex method (Walsh *et al.*, 2013). Briefly, 3 loops overnight culture of endosporeforming bacteria were put into a tube with contains 100 μ L ddH₂O and it was added 1 mL saponin 0.5% for overnight at 4 °C. The solution was centrifuged for 10 min with 12,000 rpm and then the supernatant was removed. The pellet was added with 100 μ L ddH₂O and 50 μ L chelex 20%. The mixture was boiled for 10 min and vortex every 5 min and centrifuged. The supernatant was the DNA and kept at 4 °C.

DNA amplification

DNA Amplification was done using the following program: pre denaturation at 95 °C for 3 min, denaturation at 94 °C for 1 min, annealing at 55 °C for 1 min, extention at 72 °C for 1 min and the last final extention at 72 °C for 7 min. used were 27F Primers (5'-AGAGTTTGATCMTGGCTCAG-3') and primer 1492R (5'-GGYTACCTTGTTACGACTT-3') (Osborne et al., 2005). The PCR mixture were 3 µL of DNA template, 1.5 µL of each appropriate primers, 25 µL of KAPA 2GFAST Kits and 19 μL of ddH₂O. Visualization of PCR product was using gel electrophoresis with agarose done concentration at 1% and run with 100V for 30 min. The band of PCR product was seen on gel documentation.

Sequences analysis

Sequenced of DNA product was done at Genetics Science of Indonesia, Inc. Sequence was analyzed used base alignment by Basic Local Alignment Search Tool (BLAST) to establish percentage of base pairs similarity with reference of isolate which be found in gene bank.

Phylogenetic tree

Phylogenetic tree was made using MEGA 5 software. Sequences of bacteria isolate were analyzed and compared with sequence of bacteria which had highest of homology percentage compared to sequence of other bacteria species. Phylogenetic tree was constructed by test of Neighbour-joining tree with Bootstrap method.

Pigment extraction

Serratia marcescens culture at 48, 72 and 96 h were centrifuged, and the pellet was washed with distillated water. Methanol was added and vortexed until pigment extracted. Pigment extract was assayed its antioxidant activity using DPPH methods and the functional groups were also identified by FTIR (Fourier Transform InfraRed Spectroscopy).

Antioxidant activity

Diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity assay was carried out by (Molyneux, 2004) methods and vitamin C commercial used as control. To 0.1 mL pigment extract was added 3.9 mL of DPPH 0.05 mM solution was prepared in methanol, vortexed and followed by incubation at 20 min for 30 °C. DPPH radical scavenging was measured at $\lambda = 517$ nm using spectrophotometer. Antioxidant activity represented the percentage of DPPH radical scavenging ability was calculated the following formula as below:

DPPH radical scavenging (%) = $[(A_0 - A_s/A_0)] \times 100$, where A_0 was the control absorbance, A_s was the sample absorbance.

RESULTS AND DISCUSSION

Thermotolerant red-pigmented bacteria isolate GSB-001

Red-pigmented bacteria isolate GSB-001was isolated from sediment of hot spring Gedong-Songo, Bandungan, Semarang-Indonesia. This isolate has been adapted to grow at 30 °C on Nutrient Agar medium and produced red colony (Figure 1).





The morphological characteristics of the redpigmented bacteria isolate had moderate size, irregulare shape, raised elevation and entire margin. This isolate was Gram-negative and rod. Biochemical tests showed that the isolate had negative indole test, positive catalase, negative methyl red and negative urease. This result was the same with the identification carried out by Abdullah *et al.* (2017). Positive catalase on biochemical test showed that this bacteria had catalase enzyme having role in breaking down H₂O₂ to H₂O and O₂.

This isolate was able to ferment glucose, sucrose and maltose, but there was no gas production, so the bacteria isolate could produce acid only. Lactose could not be fermented by this isolate (Table 1). The biochemical test of the isolate was confirmed with the molecular identification to determine the species.

Molecular identification was started with DNA extraction which produced DNA with concentration of 39.8 ng/ul, and purity of 1.42. DNA amplification for molecular identification was done using 27F and 1492R primers which amplified 16S rRNA gene. Low mass ladder marker was used to determine the size of PCR product which had 1500 bp length (Figure 2). Galkiwiecz and Kellog (2008) stated that 27F primer targeted bacteria whereas 1492R was universal primer. Osborne *et al.* (2005), added that primers 27F and 1492R amplified 16S rRNA gene in almost all bacteria known and has been used to study bacteria in many habitats.

The PCR product was then sequenced to find out its sequence and resulted in 1469 bp length. The sequence was analyzed to find its homology sequence from data bank in National Center for Biotechnology Information (NCBI) using Basic Local Alignment Search Tool (BLAST). The BLAST showed that the isolate had 94% homology with Serratia marcescens strain S823.

 Table 1: Morphological and biochemical characteristics of red-pigmented thermotolerant bacteria isolate GSB-001

Tests	Morphological		Biochemical
	Colony	Cells	Diochemical
Color	Red	-	-
Size	Moderate	Rod	-
Shape	Irregular	-	-
Elevation	Raised	-	-
Margin	Entire	-	-
Gram	-	Negative	-
Indole test	-	-	Negative
Catalase test	-	-	Positive
Methyl red test	-	-	Negative
Urease test	-	-	Negative
Carbohydrate Fermentation			
Glucose	-	-	Positive
Lactose	-	-	Negative
Sucrose	-	-	Positive
Maltose	-	-	Positive



Figure 2: PCR product of 16S rRNA amplification of redpigmented thermotolerant bacteria isolate GSB-001 (M=marker; I =isolate).

Phylogenetic tree

Phylogenetic tree was constructed using *neighbor-joining* (NJ) and showed that the high relationship between GSB-001 isolate with pigment-producing bacteria (Figure 3). Phylogenetic tree showed that bacteria isolate GSB-001 had closely relation with *S. marcescens* strain S823.

Serratia marcescens is a Gram-negative bacteria that can be found in many habitats such as soil, water and plant. This bacteria belongs to the opportunistic pathogenic bacteria in the family Enterobacteriaceae. It produces a red pigment that is a secondary metabolite known as prodigiosin. It grows from 5 - 40 °C and pH 5-9.



Figure 3: Phylogenetic tree between red-pigmented bacteria isolate GSB-001 and *S. marcescens* strain S82, based on 16S rRNA has 94% similarity and gaps 1%.

Pigment of S. marcescens

Red pigment extration was done using methanol solvent 70% (Figure 4). Pigment production increased in accordance with the incubation time. The red pigment produced by *S. marcescens* is prodigiosin (Srimanthi *et al.*, 2017) and the bacterial cell has to be broken to get out its intracellulare pigment. Result showed that 48 h of culture produced the lowest pigment concentration compared to 72 and 96 h of culture. On the contrary, Rakh *et al.* (2017) showed that *Serratia rubidaea* JCM 124OT produced the highest prodigiosin at 48 h of incubation. Production of prodigiosin pigment is related with the increasing of biomass cells (Haddix and Shanks, 2018).



Figure 4: Red pigment of *S. marcescens*: (A) Extract pigment in methanol 70% (B) Pigment after evaporation.

The FTIR spectrum of compounds showed wave lenght between 1500-3500 cm⁻¹. Specific peak at 1740 cm⁻¹ represented of C=O (carbonyl) stretching group. This result matched with the research of Charttejee *et al.* 2017. Based on theory, the carbonyl group present in 1820-1600 cm⁻¹. The other peak at 3364 cm⁻¹ (stretching in N-H) represented amine functional group. The wave number at 3015 cm⁻¹ showed C–H (from C=C) (Figure 5).

Antioxidant activity

DPPH is stable radical and frequently used to test the antioxidant activity of pigment. It changes its color from purple at free radical state to yellow at stable compound by antioxidant reaction, where antioxidant gives one electron to DPPH hence the free radical is neutralized. An electron not coupled with DPPH gives strong absorbance, maximal at $\lambda = 517$ nm and purple coloured.

The potency of antioxidant from red pigmented bacteria was measured with DPPH and the pigment extracts in methanol were incubated at different time. Result showed that the antioxidant activity of cultures incubated at 48, 72 and 96 h were 49,11%; 46,59% and 37,53%, respectively (Figure 7). The C vitamin used as a control showed at 24,43%. The data proved that the longer incubation time, the lower antioxidant activity was. Pawar *et al.* (2015) stated that pigment of *Serratia* sp. had the potential as antioxidant. Ibrahim *et al.* (2014) showed that *S. marcescens* IBRL USM 84 pigment has the ability as antibacterial against several bacteria.



Figure 5: FTIR spectrum for crude pigment of S. marcescens.



Figure 6: Antioxidant activity of *S. marcescens* with DPPH method.



Figure 7: DPPH radical scavenging of *S. marcescens* cultured at 48, 72 and 96 h.

CONCLUSION

Based on molecular, biochemical and morphological identification, the red-pigmented bacteria isolated from Gedong Songo hot spring, Semarang was *S. marcescens*. Based on DPPH radical scavenging test, it showed that *S. marcescens* had potency as antioxidant.

ACKNOWLEDGEMENTS

This research was supported by PNBP grant – Diponegoro University for professorship program, No: 278-27/UN7.5.1/PG/2017.

REFERENCES

- Abdullah, A. H., Nadhom, B. N. and Al-Ammiri, H. H. (2017). Isolation and identification of *Serratia marcescens* from Bovine Mastitis infections in Iraq and their susceptibility to antibiotics. *Journal of Entomology and Zoology Studies* 5(2), 489-492.
- Bathgare, A. H., Singh, N. and Moholkar, V. S. (2018). Investigations in ultrasound-induced enhancement of astaxanthin production by wild strain *Phaffia rhodozyma* MTCC 7536. *Bioresource Technology* 254, 166-173.
- Bonadio, M. P., Aline de Freita, L. and Mutton, M. J. R. (2018). Carotenoid production in sugarcane juice and synthetic media supplemented with nutrients by *Rhodotorula rubra* 102. *Brazilian Journal of Microbiology* 49, 872-878.
- Chen, Y., Xie, B., Yang, J., Chen, J. and Sun, Z. (2018). Identification of microbial carotenoids and isoprenoid quinones from *Rhodococcus* sp. B7740 and its stability in the presence of iron in model gastric conditions. *Food Chemistry* 240, 204-211.

- Cheng, C-H., Guo, Z-X., Ye, C-X. and Wang, A-L. (2018). Effect of dietary astaxanthin on the growth performance, non-specific immunity, and antioxidant capacity of pufferfish (*Takifugu obscurus*) under high temperature stress. *Fish Physiology and Biochemistry* 44(1), 209-218.
- Elkenawy, N. M., Yassin, A. S., Elhifnawy, H. N. and Amin, M. A. (2017). Optimization of prodigiosin production by *Serratia marcescens* using crude glycerol and enhancing production using gamma radiation. *Biotechnology Reports* 14, 47-53.
- Galkiewicz, J. P. and Kellogg, C. A. (2008). Crosskingdom amplification using bacterial-specific primers: Complications for coral microbial ecology. *Applied and Environmental Microbiology* 74 (24), 7828-7831.
- Haddix, P. L. and Skanks, R. M. (2018). Prodigiosin pigment of Serratia marcescens is associated with increased biomass production. Archives of Microbiology 200, 989-999.
- Hernández-Almanza A., Montanez J. C., Aguilar-González M. A., Martinez-Ávila, Rodríguez-Herrera R. and Aguilar C. N. (2014). *Rhodotorula glutinis* as source of pigments and metabolites for food industry. *Food Bioscience* 5, 64-72.
- Ibrahim, D., Nazari T. F., Kassim, J. and Lim, S-H. (2014). Prodigiosin - an antibacterial red pigment produced by Serratia marcescens IBRL USM 84 associated with a marine sponge Xestospongia testudinaria. Journal of Applied Pharmaceutical Science 4(10), 001-006.
- Kleinegris, D. M. M., Jassen M., Brandenburg, W. A. and Wijffels R. H. (2011). Continuous production of carotenoids from *Dunaliella salina*. *Enzyme and Microbial Technology* **48**, **254-259**.
- Kumar, A., Vishwakarma, H. S., Singh, J., Dwivedi, S. and Kumar M. (2015). Microbial pigments: Production and their applications in various industries. International Journal of Pharmaceutical, Chemical and Biological sciences (IJPCBS) 5(1), 203-212.
- Manimala, M. R. A. and Murugesan, R. (2017). Carotenoid pigment production from yeast: Health benefits and their industrial applications. *International Journal of Chemical Studies* 5(6), 392-395.
- **Molyneux, P. (2004).** The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology* **26(2) : 211-219.**
- Mukherjee, T., Bose, S. and Mukhopadhyay, S. K. (2017). Antioxidant properties of the carotenoid extracts of three Deinococcus–Thermusphylum bacteria, *Meiothermus* sp. strains RP and TP and *Thermus* sp. strain YY from Paniphala hot spring, India. *Nutrire* 42(7), 2-11.
- Osborne, C. A., Galic M., Sagwan P. and Jansen P. H. (2005). PCR generated artifact from 16S rRNA genespecific primers. *FEMS Microbiology Letters* 248, 183-187.
- Pawar, R., Mohandass, C., Sivaperumal, E., Sabu, E., Rajasabapathy, R. and Jagtap, T. (2015). Epiphytic marine pigmented bacteria: A prospective source of

natural antioxidants. *Brazilian Journal of Microbiology* **46(1), 29-39.**

- Rakh, R. R., Dalvi, S. M., Muscle, B. B. and Raut, L. S. (2017). Production, extraction and characterization of red pigment produced by Serratia rubidaea JCM 1240T isolated from soil. International Journal of Current Microbiology and Applied Sciences 6(1), 143-154.
- Shah, Md. M. R., Liang, Y., Cheng, J. J. and Daroch, M. (2016). Astaxanthin-producing green microalga Haematococcus pluvialis: From single cell to high value commercial products. Frontiers in Plant Science 7.
- Srianta I., Zuabaedah E., Estiasih T., Yamada M. and Harijono. (2016). Comparition of Monascus purpureus growth, pigment production and composition on different cereal substrates with solid state fermentation. Biocatalysis and Agricultural Biotechnology 7, 181-186.
- Srimathi, R., Priya R., Nirmala, M. and Malarvizhi, A. (2017). Isolation, identification, optimization of prodigiosin pigment produced by Serratia Marcescens and its applications. International Journal of Latest Engineering and Management Research (IJLEMR) 2(9), 11-21.
- Talaeikhozani, A., Alaee, S. and Pinraj, M. (2015). Guidelines for quick application of biochemical tests to identify unknown bacteria. *Accounts of Biotechnology Research* 2(2), 065-082.
- Tkačova, J., Furdikova, K., Klempova, T., Ďurčanska, K. and Čertik M. (2015). Screening of carotenoidproducing *Rhodotorula* strains isolated from natural sources. *Acta Chimica Slovaca* 8(1), 34-38.
- Walsh, P. S., Metzger, D. A. and Higuchi, R. (2013). Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques* 54(3), 134-139.
- Zang, C-Z., Yeh C-W., Chang, W-F., Lin, C-C., Kan, S-C., Shieh, C-J. and Liu, Y-C. (2014). Identification and enhanced production of prodigiosin isoform pigment from Serratia marcescens N10612. Journal of the Taiwan Institute of Chemical Engineers 45, 1133-1139.