

Original Paper

Effects of an eHealth Literacy Intervention for Older Adults

Bo Xie, BS, MS, PhD

University of Maryland, College of Information Studies, College Park, MD, United States

Corresponding Author:

Bo Xie, BS, MS, PhD

University of Maryland

College of Information Studies

2117G Hornbake, South Wing

College Park, MD, 20740

United States

Phone: 1 301 405 8617

Fax: 1 301 314 9145

Email: boxie@umd.edu

Abstract

Background: Older adults generally have low health and computer literacies, making it challenging for them to function well in the eHealth era where technology is increasingly being used in health care. Little is known about effective interventions and strategies for improving the eHealth literacy of the older population.

Objective: The objective of this study was to examine the effects of a theory-driven eHealth literacy intervention for older adults.

Methods: The experimental design was a 2×2 mixed factorial design with learning method (collaborative; individualistic) as the between-participants variable and time of measurement (pre; post) as the within-participants variable. A total of 146 older adults aged 56–91 (mean 69.99, SD 8.12) participated in this study during February to May 2011. The intervention involved 2 weeks of learning about using the National Institutes of Health's SeniorHealth.gov website to access reliable health information. The intervention took place at public libraries. Participants were randomly assigned to either experimental condition (collaborative: $n = 72$; individualistic: $n = 74$).

Results: Overall, participants' knowledge, skills, and eHealth literacy efficacy all improved significantly from pre to post intervention ($P < .001$ in all cases; effect sizes were >0.8 with statistical power of 1.00 even at the .01 level in all cases). When controlling for baseline differences, no significant main effect of the learning method was found on computer/Web knowledge, skills, or eHealth literacy efficacy. Thus, collaborative learning did not differ from individualistic learning in affecting the learning outcomes. No significant interaction effect of learning method and time of measurement was found. Group composition based on gender, familiarity with peers, or prior computer experience had no significant main or interaction effect on the learning outcomes. Regardless of the specific learning method used, participants had overwhelmingly positive attitudes toward the intervention and reported positive changes in participation in their own health care as a result of the intervention.

Conclusions: The findings provide strong evidence that the eHealth literacy intervention tested in this study, regardless of the specific learning method used, significantly improved knowledge, skills, and eHealth literacy efficacy from pre to post intervention, was positively perceived by participants, and led to positive changes in their own health care. Collaborative learning did not differ from individualistic learning in affecting the learning outcomes, suggesting the previously widely reported advantages of collaborative over individualistic learning may not be easily applied to the older population in informal settings, though several confounding factors might have contributed to this finding (ie, the largely inexperienced computer user composition of the study sample, potential instructor effect, and ceiling effect). Further research is necessary before a more firm conclusion can be drawn. These findings contribute to the literatures on adult learning, social interdependence theory, and health literacy.

(*J Med Internet Res* 2011;13(4):e90) doi: [10.2196/jmir.1880](https://doi.org/10.2196/jmir.1880)

KEYWORDS

Health literacy; lifelong learning; aged; technology

Introduction

Health literacy is “the degree to which individuals have the capacity to obtain, process, and understand basic health information and services needed to make appropriate health decisions” [1]. This concept has drawn much attention recently [2-6], with increasing well-documented evidence of both the negative impact of poor health literacy on health outcomes and health care costs [7] and the alarmingly low levels of health literacy among American adults: a national survey showed only 12% of the adults in the United States have proficient health literacy, and this proportion drops to 3% among older adults [8].

Recently, information and communication technologies (ICTs) are increasingly being widely used in health care [9,10], presenting both opportunities and challenges for developing and implementing effective health literacy interventions. As government agencies such as the US National Institutes of Health (NIH), nonprofit organizations such as medical associations, and for-profit organizations alike are increasingly putting health information online, the Internet has already become an invaluable resource for high-quality health information [11-13]. This resource, however, can only be useful if the user has adequate eHealth literacy, or “the ability to seek, find, understand, and appraise health information from electronic sources and apply the knowledge gained to addressing or solving a health problem” [14]. Individuals who have low health literacy—for instance, older adults—are likely to also have low computer and Internet literacy [15-17], thus facing a double jeopardy in the eHealth era.

Existing literature provides little scientific evidence about effective health literacy interventions [18], and even less about effective interventions for improving the eHealth literacy of the older population. Existing interventions focus predominantly on simplifying medical materials and instructions [19-21]. This “lowering-the-bar” approach is useful but limited given the complexity of medical terminology and knowledge. Education and training is another key approach to addressing the health illiteracy crisis [22]. This approach requires an understanding of health literacy as an active, lifelong learning process that features continuous learning of new, valid health information and unlearning of outdated, harmful information [22]. Such an understanding is especially important in the context of ICTs being increasingly used by health consumers, professionals, and policy makers alike in health care [9,10]. As ICTs change at a rapid rate, so do the requirements for health literacy skills [23].

Electronic Health Information for Lifelong Learners

The present study is a part of the Electronic Health Information for Lifelong Learners (eHiLL) research project that aims to address these gaps in the literature [4-6,24]. The goal of the larger eHiLL research project is to generate scientific knowledge about optimal learning conditions and strategies that can effectively and efficiently improve older adults’ learning and use of eHealth applications. To achieve this goal, the eHiLL research project consists of a series of experimental studies designed to examine the effects of various learning conditions

and strategies through theory-driven, hypothesis-testing, rigorous experiments. The eHiLL experimental studies build on an understanding of health literacy as an active, lifelong learning process that goes beyond the formal educational settings in early life stages [22]. Importantly, guided by the literature on older adults’ learning of computer technology, the eHiLL interventions all feature key elements designed specifically to accommodate older computer learners’ needs and preferences [4]. These include (1) providing step-by-step, detailed instructions and avoiding technical jargon [24,25], (2) providing hands-on practice and encouraging questions [26], (3) making sure each lesson builds on previous lessons and increases complexity gradually [25,27], (4) ensuring the learners experience at least some level of success at the initial stage of the training [25,28,29], (5) conducting the training in a familiar, relaxed, and supportive environment [27,29], and (6) offering the training in the early morning hours, which is generally the optimal time of day for older learners [30].

These key elements were fully incorporated in prior eHiLL interventions and proven to be effective in improving older adults’ eHealth literacy [4-6,24]. Building on and expanding the success of these prior studies, the present study fully incorporates these key elements while adding a new aspect—that is, collaborative versus individualistic learning. Existing literatures on adult learning and cognitive development in later life provide a theoretical foundation for the collaborative versus individualistic versions of the intervention tested in the present study. The present study differs from prior eHiLL studies in important ways. First, the only prior eHiLL study [6] that involved both collaborative and individualistic learning had a brief intervention time (the experimental session lasted 2 hours) and it used a 16-minute-long video tutorial as the curriculum. In comparison, the present study involved 2 weeks of intervention for a total of 8 hours, using a paper-based curriculum. Second, while the present study and prior eHiLL studies [4,5,24] all used instructional materials drawn from the same tutorial, the Xie and Bugg [4] and Xie [24] studies focused on individualistic learning and the Xie [5] study was only on collaborative learning. The present study is the first eHiLL study that compares the relative effects of collaborative versus individualistic learning over an extended period of intervention time.

Collaborative Versus Individualistic Learning

Collaborative learning is one of the most common forms of active learning. It can be defined as “any instructional method in which students work together in small groups toward a common goal,” ([31] p 223). Collaborative learning requires learners to actively engage in the learning process by engaging in meaningful activities and reflecting on what they are learning from doing those activities [32]. Collaborative learning is often contrasted with individualistic learning that features students working on their own with little or no interaction with peer students [33].

The superiority of collaborative learning over individualistic learning is predicted by social interdependence theory, which emphasizes the interdependence among group members by arguing that the group is a “dynamic whole,” such that any

change in the state of a group member changes that of other group members [33]. The social interdependence among group members can be positive, negative, or nonexistent. Positive interdependence (collaboration) facilitates learning by promoting collaboration among group members. It can be found when individuals recognize that the only way they can achieve their goals is when other group member also achieve their goals. Negative interdependence (competition) exists when individuals recognize that the only way they can achieve their goals is when others fail to do so. It often results in obstructive interactions impeding learning. Nonexistent interdependence (individualistic efforts) exists when individuals perceive that their achievement is not affected by others' performance. It features no interaction among group members as each member learns independently [33,34].

Extensive empirical evidence supports the effectiveness of collaborative learning. A meta-analysis of over 300 studies shows that collaborative learning outperforms individualistic and competitive learning in postsecondary and professional settings [34]. However, there are still major gaps in the literature that require further examination. Collaborative learning research within the social interdependence tradition is predominantly based on formal education of younger adults [33,34]. Whether these findings can be generalized to older age groups in informal educational settings is yet to be answered.

Recently, the cognitive-developmental literature has begun to examine collaborative learning as a mechanism for improving cognitive abilities in later life [35-37]. Some (non-computer-related) studies find a positive impact of collaborative learning on older adults' performance [38,39]. However, there is also evidence that, compared with individualistic learning, collaborative learning has no, or even negative, impact [40-42]. A possible reason might be that existing research within the cognitive-developmental tradition generally does not provide detailed instructions to ensure collaboration [42]. Instead, participants are simply instructed to "work together" [43] or "collaborate as much as possible" [40]. To ensure collaborative learning for older adults in informal settings, it is critical to develop effective strategies that really work.

Group Composition

Group composition may affect the "dynamic whole" of a group and, subsequently, the learning process and outcomes [33,34]. Evidence suggests there is more collaboration in groups with either female- or male-gender majority than in groups with equal gender composition [44,45], and more collaboration in same-gender groups than in mixed ones [44,46].

The time and effort spent on getting familiar with each other and coordinating may negatively affect the learning process and outcomes [43]. This argument finds support in research reporting collaborative learning with familiar partners (typically defined as related family members such as a spouse) being more effective than that with unfamiliar partners in enhancing cognitive performance [47] or in reducing the negative effects of collaboration [40,41]. Yet evidence exists that, even with familiar partners, collaborative learning does not generate more

benefits than individualistic learning in improving older adults' cognitive performance [42].

Ample evidence suggests that prior computer experience is a strong predictor for older adults' computer task performance and learning outcomes [48-50]. Cody et al [28] in their study of older computer learners found that "the same [computer training] program was too challenging to some and insufficiently stimulating to others" (p 282). Some researchers [25] suggest forming homogeneous groups based on prior computer experience to ensure the success of computer training for older adults. This suggestion should be taken with caution, given that to date research on computer training for older adults has focused predominantly on individualistic learning while paying little attention to collaborative learning (an important exception is the Zandri and Charness study [51], which found promising signs for the superiority of collaborative learning over individualistic learning). While homogeneous groups based on prior computer experience might work better than heterogeneous groups in individualistic learning, this might not be the case for the collaborative learning condition.

Research Questions and Hypotheses

The present study asked the following primary research question: what impact might the intervention have on older adults' computer and Web knowledge, procedural skills, eHealth literacy efficacy, attitudes, and participation in their own health care?

Under this primary research question, 2 subresearch questions were asked: (1) what impact might the learning method (collaborative; individualistic) have on the learning outcomes?, and (2) what impact might group composition (based on gender, prior familiarity with peers, and prior computer experience) have on the learning outcomes?

The following hypotheses were tested: (1) hypothesis 1: computer/Web knowledge, skills, and eHealth literacy efficacy increase significantly from pre to post intervention (at the .05 level; same for all hypotheses), (2) hypothesis 2: collaborative learning is significantly more effective than individualistic learning in improving the learning outcomes, and (3) hypothesis 3: collaborative learning is significantly more effective than individualistic learning in heterogeneous group compositions, while individualistic learning is more effective than collaborative learning in homogeneous group compositions.

Methods

Design

We used a 2 × 2 mixed factorial design with learning method (collaborative; individualistic) as the between-participants variable and time of measurement (pre; post) as the within-participants variable.

Research Sites

The Hyattsville and New Carrollton branch libraries of the Prince George's County Memorial Library System served as the primary research sites for this study. The Library System is a publicly funded large, urban library system serving over 830,000 residents in Prince George's County, Maryland, USA.

It has been serving as the primary research site for the larger *eHiLL* research project since 2007 [4-6,24]. This Library System was selected as the key site for the *eHiLL* research project because it serves a large population of ethnic minorities, particularly African American/black people. According to the US Census Bureau, 66% of Prince George's County residents are African American/black, much higher than the 30% overall rate of African American/black residents in Maryland or the 12% rate nationwide (<http://www.census.gov>). Partnering with this Library System ensures the reach of *eHiLL* interventions to individuals from this underserved minority group. The Hyattsville and New Carrollton branch libraries of the Library System provided free networked computers, space, and staff support to facilitate the implementation of this study. The geographic location of these branch libraries is convenient for potential research participants and the researchers. Both are

within 10 miles of the University of Maryland and easily accessible by car or public transportation.

Participants

Standard recruitment techniques were used to recruit participants. These included posting recruitment flyers in the branch libraries and other local organizations (eg, senior centers, community centers, and churches) and advertising in the Library System's newsletter. The inclusion criterion was age 60 years and above, though on request and in cases where seats were available, we accommodated individuals a few years younger. A total of 146 older adults aged 56–91 years (mean 69.99, SD 8.12) participated in the present study during a 4-month period (February to May 2011). We randomly assigned 72 participants to the collaborative learning experimental condition and 74 to the individualistic learning condition. Table 1 summarizes participants' basic characteristics, including demographics, health status, and prior computer experience.

Table 1. Participants' basic characteristics

Variable	n	%
Gender		
Female	96	69
Male	44	31
Highest level of education		
Less than high school graduate	15	11
High school graduate/GED ^a	39	28
Vocational training	11	8
Some college/associate's degree	33	23
Bachelor's degree	25	18
Master's degree or other postgraduate training	16	11
Doctoral degree	2	1
Ethnic group		
Asian	8	6
African American/black	90	64
White	30	21
Other	13	4
Household annual income range (US \$)		
<20,000	38	28
20,000–29,999	28	20
30,000–39,999	14	10
40,000–49,999	8	6
50,000–59,999	9	6
60,000–69,999	6	4
70,000–99,999	4	3
≥100,000	3	2
Do not know for certain	12	9
Do not wish to answer	18	13
Health status		
Poor	7	5
Fair	24	17
Good	75	53
Very good	25	18
Excellent	11	8
English as primary language		
Yes	125	88
No	16	12
Frequency of computer use		
Every day	12	9
Every 2–3 days	27	20
Once a week	18	13
More than once a month	13	9

Variable	n	%
Less than once a month	20	15
Never	48	35

^a General equivalency diploma.

Measures

Adapted from existing outcome measures of collaborative learning (that focus on younger learners in formal educational settings) and with necessary modifications based on the results of a prior eHiLL study [5], outcome measures (dependent variables) for the present study covered the following categories: knowledge gains, skill gains, efficacy, attitudes, and changes in participation in own health care. Learning preference, familiarity with peers, prior computer experience, and basic demographics were measured to serve as control variables. (Copies of the instrument are available on request to the author.)

Computer/Web Knowledge

This was measured by objective tests of knowledge about components of the computer and the Web. Participants were shown an image of a computer and a screenshot of the NIHSeniorHealth.gov website, and were instructed to write down names of the main components of each image (eg, keyboard, mouse, link, scroll bar). Computer knowledge and Web knowledge were each measured by 5 items; each item scored 1 point if answered correctly and 0 points if answered incorrectly with a scoring range of 0–5.

Computer/Web Skills

These were measured by procedural tests of the abilities to carry out specific computer and Web operations. Each participant had one computer to use during the testing. Participants were asked to perform a number of operations on their computers independently. Participants had up to 1 minute to complete each operation. Sample operations were to open a Web browser; go to the NIHSeniorHealth.gov Web site; increase text size; find information on the Falls and Older Adults health topic; and open a video. There were a total of 20 operations. Each operation scored 1 point if successfully completed and 0 points if unsuccessful, with a scoring range of 0–20.

eHealth Literacy Efficacy

This construct was measured by the eHealth literacy scale [52], which was built on the self-efficacy concept [53] to measure perceived skills at and comfort with using the Internet for health information and decision making. The main scale has 8 items. Each item is on a 1- to 5-point Likert scale with the following anchors: 1: strongly disagree; 2: disagree; 3: undecided; 4: agree; 5: strongly agree. Higher score indicates higher eHealth literacy efficacy. A sample item is “I know **how** to find helpful health resources on the Internet” (bold original). This scale has been used in multicultural samples and has shown excellent internal

consistency reliability (scale alpha = .89–.97) with good test–retest reliability [23].

Attitude Toward the Class

This was measured by the following 5 items (items 3 and 4 were modified from Pace and Kuh [54]):

1. Overall, what would you say about the instructor’s teaching? (Anchors: 1: very poor; 2: poor; 3: fair; 4: good; 5: excellent)
2. Overall, was this computer class useful to you? (1: completely useless; 2: useless; 3: somewhat useful; 4: useful; 5: very useful)
3. How would you evaluate your entire experience in this computer class? (1: extremely dissatisfied; 2: dissatisfied; 3: neither satisfied nor dissatisfied; 4: satisfied; 5: extremely satisfied)
4. If you could start over again, would you go to the same computer class you are now attending? (1: definitely not; 2: probably not; 3: not sure; 4: yes; 5: definitely yes)
5. Would you want to recommend this training class to other people similar to your age? (1: definitely not; 2: probably not; 3: not sure; 4: yes; 5: definitely yes)

Changes in Participation in Own Health Care

This was measured by 12 items, including 6 items modified from a Kaiser survey study [16], 5 items modified from a Pew survey study [55], and an additional item added to supplement the Kaiser and Pew items. (These items are detailed in Table 6 in the Results section where the results are reported.)

Attitude Toward the Individualistic Versus Collaborative Learning Method

This was measured by the following item: “When I have to learn a new skill, I prefer to learn alone, rather than with others” [51]. It was scored on a 1- to 5-point Likert scale with the following anchors: 1: disagree strongly; 2: disagree; 3: undecided; 4: agree; and 5: agree strongly.

Prior Experience

Prior experience (familiarity) with peers was measured by the following item: “Are you related to or familiar with at least one person taking this same computer class? (eg, spouse, sibling, friend, acquaintance)”. Prior computer experience was measured by the frequency of computer use.

Basic Demographics

Age, gender, education, health status, race/ethnicity, income, and primary language were recorded.

Table 2 summarizes these measures and time(s) of measurement.

Table 2. Measures used in the present study and the time(s) of measurement

Variable	Pre	Post
Computer/Web knowledge	X	X
Computer/Web skills	X	X
eHealth literacy efficacy	X	X
Changes in health behavior/decision making		X
Attitude toward the class		X
Attitude toward the individualistic versus collaborative learning method	X	
Prior experience with peers	X	
Prior computer experience	X	
Basic demographics	X	

Instructional Materials

This study used a set of instructional materials developed by the National Institute on Aging (NIA) of the NIH, “Helping Older Adults Search for Health Information Online: A Toolkit for Trainers” [56]. This freely available toolkit is designed to improve older adults’ ability to access and use NIH online health information resources (eg, the NIHSeniorHealth.gov website). By focusing on only NIH resources, this eHiLL intervention avoids potential problems associated with the quality of online health information [57,58]. The toolkit features detailed lesson plans, in-class interactive exercises, take-home practice exercises, and other supportive handouts (eg, a glossary of computer terms). This toolkit was chosen because, first, it contains key elements that, as addressed in the Introduction

section above, facilitate older adults’ learning of computer technology [24-30]; and second, it was tested in prior eHiLL studies and proven to be effective [4,5,24].

The toolkit contains lesson plans (modules) designed to be used independently or in any combination. We used 4 modules in the present study to help older adults learn about (1) basic computer terms (1 module), (2) NIHSeniorHealth.gov (2 modules), a website designed to accommodate age-related changes in cognitive, physical, and sensory abilities [59], and (3) evaluating the quality of online health information (1 module). Together, these 4 selected modules provide a good coverage of the eHealth literacy skills as defined by Norman and Skinner [14]. Table 3 [56] outlines the lesson plans and goals of these 4 modules.

Table 3. Lesson plans and goals included in the National Institute on Aging (NIA) toolkit (extracted from the toolkit [56]).

Class session	Lesson goals
Session 1: Internet Basics (NIA Module #1)	<ol style="list-style-type: none"> 1. Learn basic computer terms 2. Practice using the mouse 3. Learn basic Internet terms 4. Learn how to get to a website 5. Learn how to explore a website 6. Learn how to use a search box 7. Learn how to use a site map
Session 2: Introduction to NIHSeniorHealth (NIA Module #2)	<ol style="list-style-type: none"> 1. Use the Home Page to find health topics on NIHSeniorHealth 2. Use the Table of Contents of a health topic to find specific information 3. Navigate through a health topic 4. Enlarge, view, and close images 5. Find answers to health questions of personal interest
Session 3: NIHSeniorHealth Quizzes and Videos (NIA Module #3)	<ol style="list-style-type: none"> 1. Recall how to use the Home Page of the NIHSeniorHealth website 2. Recall how to use the All Topics A–Z page to find health topics on the NIHSeniorHealth website 3. Recall how to use the special features (optional) 4. Learn how to take online quizzes 5. Learn how to open, watch, and close a video 6. Learn how to open, read, and close a video transcript 7. Learn how to find answers to health questions of personal interest
Session 4: Evaluating Health Websites (NIA Module #9)	<ol style="list-style-type: none"> 1. Reliable health information websites 2. The sponsor of a health website 3. The purpose of a health website 4. The authors of the health information 5. The reviewers of the health information 6. The most recent update of the health information 7. The privacy policy of a health website 8. Clues about the accuracy of a website's health information 9. The contact information for a health website

Procedure

The general procedures of the present study are similar to those of prior *eHiLL* studies [4-6,24]. In the first session, participants first signed the consent form (approved by the Institutional Review Board of the University of Maryland). The pre intervention survey questionnaire was then administered, followed by the pre skill testing. The intervention began with the completion of the pre testing. At the end of the last session, the post intervention survey questionnaire was administered, followed by the post skill testing. Each class met twice a week, 2 hours each time between 9:00 and 11:00 am, for a total of 2 weeks, at a library site. Class size was small (no more than 8 participants per class). The instructors, trained Master of Library Science students, frequently provided immediate, positive, and useful feedback when needed. Each participant had one computer to work on during each session. The instructors

emphasized hands-on practice and provided relevant handouts during each session. These procedural components were carefully designed based on the literature on older adults' computer learning [24-30] and proven to be effective in prior *eHiLL* studies [4-6,24].

Compared with prior *eHiLL* studies [4-6,24], a unique procedural aspect of the present study was the use of both collaborative and individualistic learning methods. Building on prior work on computer class structure for older learners that focused primarily on individualistic learning [26] and the common strategies used to promote collaborative learning among younger adults in formal learning settings [33,34], and adapting those strategies to accommodate the special needs and preferences of older adult learners in the public library setting, we used several strategies during the sessions to promote individualistic or collaborative learning (Table 4).

Table 4. Class structure and strategies for the learning conditions

Activity/time	Individualistic learning	Collaborative learning
Housekeeping: 5 minutes	<ul style="list-style-type: none"> Welcome Instructor self-introduction Participants self-introduction Practical information How long the class session will last Where the restrooms are Environment check Everyone has a computer Everyone can see instructor Everyone can hear 	<ul style="list-style-type: none"> Same as the individualistic condition
Overview: 5 minutes	<ul style="list-style-type: none"> Goal statement What the participants will know or be able to do after this class session Agenda What will happen during this session Steps and procedure What instructor will do and what participants will do 	<ul style="list-style-type: none"> Same as the individualistic condition
Explanation of learning method: 1 minute	<ul style="list-style-type: none"> Explain explicitly that everyone in the class is expected to learn independently Encourage participants to ask the instructor any questions that they might have 	<ul style="list-style-type: none"> Explain explicitly that the class is expected to learn together as a group Encourage participants to share with and help peers with any questions that they might have (and explain that instructor will answer any remaining question)
Introduction to the specific topic of this class session: 5 minutes	<ul style="list-style-type: none"> Definitions, scope, background information 	<ul style="list-style-type: none"> Same as the individualistic condition
Lecture and demonstration, step-by-step instruction (part 1): 20 minutes	<ul style="list-style-type: none"> Present material and demonstrate processes, following the instructions and examples used in the National Institute on Aging toolkit Encourage questions Get confirmation after each step is explained and demonstrated Check frequently to make sure everyone is on the same page 	<ul style="list-style-type: none"> Same as the individualistic condition
Brief reflection: 2 minutes	<ul style="list-style-type: none"> Pause briefly and instruct participants specifically to check their own notes and reflect independently 	<ul style="list-style-type: none"> Pause briefly and instruct each participant specifically to compare notes with a peer sitting next to him or her and reflect together with peer

Activity/time	Individualistic learning	Collaborative learning
Continuation of lecture and demonstration: 20 minutes	<ul style="list-style-type: none"> • Same as part 1 above 	<ul style="list-style-type: none"> • Same as the individualistic condition
Brief reflection: 2 minutes	<ul style="list-style-type: none"> • Same as the first independent reflection session above 	<ul style="list-style-type: none"> • Same as the first collaborative reflection session above
Break: 5 minutes	<ul style="list-style-type: none"> • Distribute handouts, which have in-class practice exercises and detailed, step-by-step instructions for completing the exercise 	<ul style="list-style-type: none"> • Same as the individualistic condition
Hands-on practice: 40 minutes	<ul style="list-style-type: none"> • Participants perform the hands-on practice activity independently • Encourage participants to ask instructor questions about the specific steps of the exercise • Help to answer each participant's questions 	<ul style="list-style-type: none"> • Each participant pairs up with a peer to do the hands-on practice activity together • Encourage participant to ask peers questions about the specific steps of the exercise • Participants discuss with peers and try to answer peers' questions (if, after peer discussion and exploration, questions remain, then instructor will answer the remaining questions)
Practice vs group reflection: 10 minutes	<ul style="list-style-type: none"> • Same as above (participants continue to engage in hands-on practice independently) 	<ul style="list-style-type: none"> • All participants (and instructor) sit in a circle to discuss, share, and reflect together
Closing: 5 minutes	<ul style="list-style-type: none"> • Summarize content covered in this class session • Distribute handouts for take-home exercises, which have detailed, step-by-step instructions for completing the exercises • Point out opportunities for coming back to use the library's computers to practice • Preview the topic of next class session • Thank participants for coming to this class session and remind them to come to the next class session 	<ul style="list-style-type: none"> • Same as the individualistic condition

Data Analysis

The main statistical analyses conducted to test the hypotheses were various techniques of analysis of variance (ANOVA). These included multivariate repeated measures analyses, one-way ANOVA, and univariate analysis of covariance. Two-independent-samples tests (Mann-Whitney U) and chi-square tests were used to compare the collaborative versus individualistic samples.

Results

Comparing the Collaborative Versus Individualistic Experimental Groups

Two-independent-samples tests (Mann-Whitney U) found no significant difference in age ($P = .13$), education ($P = .11$), health ($P = .85$), income ($P = .32$), and computer use frequency ($P = .06$), and chi-square tests found no significant difference in gender ($P = .66$), ethnicity ($P = .07$), primary language ($P = .81$), and familiarity with peers ($P = 1.00$) among participants

in the collaborative and individualistic experimental conditions. Chi-square test also found no significant difference in the retention rate of the 2 experimental conditions ($P = .56$); overall, a total of 108 participants completed both the pre and post testing, resulting in a 74% retention rate for this 2-week intervention. These results suggest the 2 experimental groups were comparable in these aspects.

Comparing Participants Who Completed the Intervention Versus Those Who Did Not

Mann-Whitney U tests found no significant difference in age ($P = .51$), education ($P = .41$), health ($P = .42$), income ($P = .78$), and baseline computer knowledge ($P = .80$), Web knowledge ($P = .81$), skills ($P = .70$), and eHealth literacy efficacy ($P = .12$) between participants who completed both the pre and post testing and those who completed only the pre testing. Significant difference was found in computer use frequency, with participants who completed both the pre and post testing reported having *less* prior use of computers than those who completed only the pre testing ($P = .007$).

Changes From Pre to Post Intervention

Multivariate repeated measures analyses found that, overall, computer knowledge, Web knowledge, procedural skills, and eHealth literacy efficacy improved significantly from pre to post intervention ($F_{4,90} = 119.60$, $P < .001$). Univariate repeated measures analyses revealed significant improvement from pre to post intervention on each of these 4 measures ($P < .001$ in

all 4 cases; computer knowledge: $F_{1,93} = 60.60$; Web knowledge: $F_{1,93} = 54.92$; procedural skills: $F_{1,93} = 264.40$; and eHealth literacy efficacy: $F_{1,93} = 229.31$). Hypothesis 1 was strongly supported. Further, effect sizes (measured by Cohen's d) with regard to gains from pre to post intervention in computer and Web knowledge, skills, and eHealth literacy efficacy ranged from 0.88 to 2.25. The statistical power of these measures reached 1.00 even at the $\alpha = .01$ level (Table 5).

Table 5. Effect sizes and statistical power

Variable	Cohen's d	Percentile standing	Statistical power ($\alpha = .01$)
Computer knowledge	1.05	84	1.00
Web knowledge	0.88	80	1.00
Procedural skill	1.70	95	1.00
eHealth literacy efficacy	2.25	99	1.00

Attitudes

Participants in both the collaborative and individualistic learning conditions had overwhelmingly positive attitudes toward the intervention. Across the 2 conditions, 99% (94/95) of participants felt the instructor's teaching was good or excellent; 95% (90/95) of participants felt the intervention was useful or very useful; 98% (92/94) were satisfied or extremely satisfied with the entire experience participating in the intervention; 84% (79/94) said they would attend the same class if it started over; and 98% (93/95) would recommend the intervention to their age peers. Multivariate ANOVA found no significant difference between participants in the collaborative and individualistic conditions in all measures of attitude.

It is worth noting, though, that 1 attitudinal measure, the instructor's teaching, *approached* a statistically significant difference between the 2 experimental conditions ($F_{1,104} = 3.34$; $P = .07$). While participants in both conditions had very positive assessment of the instructor's teaching, participants in the individualistic learning condition had a slightly more positive attitude toward the instructor's teaching than those in the collaborative learning condition (individualistic: mean 4.87, SD 0.35; collaborative: mean 4.72, SD 0.45).

Changes in Participation in Own Health Care

Across the collaborative and individualistic learning conditions, a notable number of participants reported changes in various aspects of participation in their own health care (Table 6).

Table 6. Changes in participation in own health care as a result of the intervention

	Yes		No	
	n	%	n	%
Since you started taking this computer class, have you had a conversation with a friend or family member about the health information you found on NIHSeniorHealth?	55	59	39	42
Since you started taking this computer class, have you talked with a doctor or other health care provider about the information you found on NIHSeniorHealth?	9	10	84	90
Have you changed your behavior because of the health information you found on NIHSeniorHealth?	51	55	41	45
Have you made a decision about how to treat an illness or condition because of the information you found on NIHSeniorHealth?	56	61	36	39
Have you changed your health insurance plan because of the information you found on NIHSeniorHealth?	2	2	86	98
Have you changed your overall approach to maintaining your health or the health of someone you help take care of because of the information you found on NIHSeniorHealth?	58	64	33	36
Has the information you learned from NIHSeniorHealth led you to ask a doctor new questions or to get a second opinion from another doctor?	54	62	33	38
Has the information you found on NIHSeniorHealth changed the way you think about diet, exercise, or stress management?	69	76	22	24
Has the information you found on NIHSeniorHealth changed the way you cope with a chronic condition or manage pain?	51	57	38	43
Has the information you found on NIHSeniorHealth affected a decision about whether to see a doctor?	37	42	52	58
Have you changed the way you take medicine because of the information you found on NIHSeniorHealth?	32	36	58	64

One-way ANOVA found no significant difference in the total number of reported changes in participation in own health care between participants in the collaborative and individualistic learning conditions ($P = .45$).

Collaborative Versus Individualistic Learning Method

Multivariate repeated measures analyses found a significant main effect of the learning method on computer knowledge, Web knowledge, procedural skills, and eHealth literacy ($F_{4,90} = 4.56$, $P = .002$). To examine on which specific outcome measure(s) the learning method showed a main effect, univariate repeated measures analyses were performed and revealed a significant main effect of the learning method on procedural skills ($F_{1,93} = 7.81$; $P = .006$) and eHealth literacy ($F_{1,93} = 8.64$; $P = .004$). Univariate analyses revealed no significant main effect of the learning method on either computer knowledge ($P = .51$) or Web knowledge ($P = .47$).

Interestingly, one-way ANOVA found a significant difference in pre intervention procedural skills ($F_{1,142} = 7.17$; $P = .008$) and eHealth literacy ($F_{1,140} = 6.18$; $P = .01$) between participants in the individualistic and collaborative learning groups. Pre intervention computer knowledge ($P = .90$) and Web knowledge ($P = .94$) did not differ significantly between the 2 experimental groups. To examine whether the significant main effects of the learning method on procedural skills and eHealth literacy as revealed by the univariate analyses were due to significant differences in these variables at the baseline, univariate analysis of covariance was performed and, after controlling for pre intervention procedural skills and eHealth literacy, respectively, the significant effects of the learning method on these variables

both disappeared (procedural skills: $P = .36$; eHealth literacy: $P = .06$). These findings suggest that, when controlling for these variables at the baseline, the learning method had no main effect on knowledge, skills, and eHealth literacy efficacy. Hypothesis 2 was not supported.

Multivariate repeated measures analyses found no significant interaction effect of the learning method (collaborative; individualistic) and time of measurement (pre; post): $P = .73$.

Group Compositions

Groups in this study had 4 gender compositions (there was no male-only group): female-only (14 participants in this group composition), female majority ($n = 96$), equal number of female and male ($n = 10$), and male majority ($n = 13$). Multivariate repeated measures analyses found no significant 3- or 2-way interaction effect of group gender composition, learning method, and time of measurement (learning condition \times gender composition: $P = .40$; time \times gender composition: $P = .74$; time \times learning condition \times gender composition: $P = .79$). Further, group gender composition had no significant main effect on the outcome measures ($P = .68$).

Groups had 2 compositions based on familiarity with peers: familiar with at least one other person in the same session ($n = 37$) and not familiar with anyone in the same session ($n = 106$). Multivariate repeated measures analyses found no significant 3- or 2-way interaction effect of group familiarity composition, learning method, and time of measurement (learning condition \times familiarity composition: $P = .37$; time \times familiarity composition: $P = .80$; time \times learning condition \times familiarity

composition: $P = .88$). Further, no significant main effect of group familiarity composition was found ($P = .53$).

Similar to 2 prior *eHiLL* studies [5,6], computer use frequency was used to categorize group composition based on prior computer experience. “Experienced computer users” were defined as individuals who use the computer every day or every 2–3 days a week, and “inexperienced computer users” were defined as individuals who use the computer less than every 2–3 days a week. Using this criterion, we coded each session into 1 of the following 5 groups: inexperienced user only ($n = 36$), inexperienced user majority ($n = 88$), equal number of experienced and inexperienced users ($n = 4$), experienced user majority ($n = 10$), and experienced user only ($n = 1$). These groups were further recoded into 2 groups: groups with mixed users ($n = 102$) and groups with either all inexperienced users or all experienced users ($n = 37$). Multivariate repeated measures analyses found no significant 3- or 2-way interaction effect of group composition based on prior computer experience, learning method, and time of measurement (learning condition \times prior computer experience composition: $P = .40$; time \times prior computer experience composition: $P = .20$; time \times learning condition \times prior computer experience composition: $P = .53$). Further, no significant main effect of group composition based on prior computer experience was found ($P = .54$). Hypothesis 3 was not supported.

Learning Preferences

To examine the potential impact of attitude toward the individualistic versus collaborative learning method (learning preference), participants were recoded into a “matched” group, in which participants’ learning preferences matched their collaborative versus individualistic group assignments, and a “no-match” group, where the preference and the group assignment did not match. Multivariate repeated measures analyses found neither an interaction effect among learning preference matching, learning method, and time of measurement ($P = .18$) nor a significant main effect of the learning preference matching factor ($P = .41$).

Discussion

Older adults are in great need of health literacy interventions, given that their needs for health information and services are typically high [17,60-62] and yet their health literacy levels are low [8]. Due to age-related changes in social environments and individual abilities [63], interventions that target younger age groups are unlikely to reach or have similar impact on older adults. Further complicating the situation is that the requirement for health literacy skills is a moving target, particularly in the context of eHealth becoming increasingly prominent in contemporary society [9,10]. As Norman [23] correctly points out, as technology changes, so do the requirements for health literacy skills.

Impact of the Overall Intervention

This study aimed to generate new scientific knowledge about effective eHealth literacy interventions for the older population. The primary research question was “What impact might the intervention have on the learning outcomes?” The analyses

revealed that computer knowledge, Web knowledge, procedural skills, and eHealth literacy efficacy all improved significantly from pre to post intervention ($P < .001$ in all cases). The effect sizes of these improvements ranged from 0.88 to 2.25, suggesting that the magnitude of these improvements was large [64]. What these effect sizes mean is that, for instance, with respect to improvements in eHealth literacy efficacy, an effect size of 2.25 meant a learner increased from the 50th percentile on the pre test to the 99th percentile on the post test on this measure. These results strongly support the magnitude of the effects of the intervention. Further adding to the strength of these positive results is that the statistical power of these measures was strong: it reached 1.00 even at the $\alpha = .01$ level in all cases.

These findings are even more impressive when interpreted in the context of the literature showing that “effect sizes of 0.8 are rare for any [learning] intervention and require truly impressive gains” ([31] p 224). Also, as summarized in several meta-analyses, the effect sizes of prior collaborative learning interventions (that focused on younger learners in formal educational settings) ranged from 0.29 to 0.70 [33,65,66]. In the present study, the effect size of the intervention on all knowledge, skill, and efficacy measures was greater than 0.8, suggesting the intervention has indeed resulted in “truly impressive gains.” These results provide strong support that the intervention, regardless of the specific learning method used, was effective in improving older adults’ eHealth literacy.

Across the 2 experimental conditions, participants had overwhelmingly positive views of the intervention. Notable percentages of participants also reported changes in various aspects of participation in their own health care. These findings suggest the health information these participants obtained from the intervention had affected their health behavior and decision making, which is a key component of eHealth literacy [14]. These findings further suggest that the intervention, regardless of the specific learning method used, was effective in improving older adults’ eHealth literacy. These findings are particularly meaningful in the context of the contemporary health care system increasingly promoting shared medical decision making, where patients are expected to participate more in their own health care [60,67-71].

An important reason for the effectiveness of the intervention tested in this study was that it fully incorporated the key elements of successful computer training for older adults [24-30], as outlined in the Introduction section above. These key elements were proved effective in prior *eHiLL* studies [4-6,24]. A unique aspect of this study was to compare the effectiveness of the collaborative and individualistic learning methods built into the intervention. The analyses yielded interesting findings, as discussed in the next subsection.

Collaborative Versus Individualistic Learning Method

The analyses found neither an interaction effect of the learning method and time of measurement nor a main effect of the learning method on any of the outcome measures, suggesting the collaborative and individualistic learning methods did not differ in their relative effects. This finding deserves careful consideration. As reviewed above, the superiority of

collaborative learning over individualistic learning, as is well documented in the literature, is based on studies of younger learners in formal educational settings [33,34]. It is possible that collaborative learning may simply not work as well for the older population in informal settings. If this is the case, then this study contributes to the literature by identifying some key limits of the social interdependency theory.

In particular, the sample of this study consisted of primarily inexperienced computer users: less than 9% of participants used computers every day, while 35% of participants had never used a computer before (participants who used computers less often than every 2–3 days made up 72% of the study sample). Engaging in collaborative learning might have been too challenging for most participants who had to focus on their own activities with little attention to spare to interact with and help others, which is in line with the findings of prior research [25,28]. Thus, the previously widely reported advantages of collaborative learning over individualistic learning may not be easily realized among individuals who have limited prior computer experience.

Another way to look at this matter, however, is that perhaps different collaborative learning strategies could be used to better promote collaboration among inexperienced computer users. The collaborative learning strategies used in this study were carefully developed based on prior research on younger adult learning in formal educational settings [33,34] and modified to accommodate the older population in informal settings (see Table 4). In reflection, however, there might not have been sufficient consideration for the largely inexperienced user composition of the study sample.

For logistical reasons, multiple instructors were hired on an hourly basis to provide the instructions. It is likely that individual differences in these instructors (eg, teaching style, commitment, personality) may have affected the learning outcomes. One indication of this possibility is that the measure of the instructor's teaching *approached* a statistically significant difference between the 2 experimental conditions ($P = .07$). Participants in the individualistic learning condition expressed more positive views of the instructor's teaching than those in the collaborative learning condition. It is possible that this factor might have helped at least partially offset any hypothesized advantage of collaborative learning over individualistic learning.

The ceiling effect might have also affected the results: due to the positive impact of either version of the intervention, it may have been difficult to differentiate between the relative effects of the collaborative versus individualistic versions of the intervention. A possible solution for future research is to make the knowledge and skill tests more challenging so that the measures can be more sensitive to potential differences in learning outcomes.

Group Composition

As reviewed above, the literature suggests that group composition factors (based on gender, prior familiarity with peers, and prior computer experience) may affect the learning outcome. This study, however, found neither an interaction effect of any of these group composition factors with the

collaborative versus individualistic learning method nor a main effect of any of these group composition factors. These findings are in line with those of 2 earlier eHiLL studies [5,6]. Replicating the same findings in these 3 independent studies, which differed in multiple ways (eg, intervention duration, instructional materials, procedures, and participants), lends some support to the generalizability of these findings. Note, though, that the study samples of these 3 independent eHiLL studies were similar: in each study the majority of the participants were women, unfamiliar with their study peers, and inexperienced computer users. While these findings might be generalized to populations with similar characteristics, they may not be so to populations with different characteristics. Also, as discussed above, if there were insufficient strategies to fully promote collaborative learning, then the potential effect of group composition might have also been affected.

Participants who completed both the pre and post testing reported *less* prior use of computers than those who completed only the pre testing ($P < .01$), suggesting that participants who used computers more often were more likely to drop out. One possible reason is that the intervention, by design, started from basic computer terms and increased in complexity gradually (see Table 3). Yet this might not have been made clear to the participants and some of them, after the first session, might have gotten the impression that the class was too “basic” for them and thus left. In future research, it will be necessary to fully communicate to participants, during the very first session, all the topics that will be covered in the remaining sessions.

Practical Implications

As in the earlier eHiLL interventions [4-6,24], the intervention tested in the present study also involved productive partnerships among local public libraries, a library and information science academic program at a state university, and the NIH. These local, state, and federal organizations bring complementary resources to the project and, in doing so, help each organization to achieve its mission [4]. The local public libraries provide the facility and staff support for the project, helping the libraries better serve socially, economically, and technologically underserved library patrons. The library and information science academic program provides the human resources through well-trained and dedicated faculty and graduate students and, in doing so, better achieves its research and educational missions. The NIH provides reliable online health information resources, and its involvement in this study helps promote the use of these resources. Tapping into these well-established public infrastructures ensures the intervention's capacity for scaling up (eg, it can be easily rolled out to other communities across the country to improve older adults' eHealth literacy).

Limitations and Future Directions

This study has some limitations. First, the sample was a convenience one, consisting of mostly African American/black people (64%), women (69%), and inexperienced computer users (72%). The findings may not be representative of the older population as a whole and should not be generalized without caution. Future research will benefit from examining the issues in a representative sample.

Second, potential confounding factors might have affected the relative effects of collaborative versus individualistic learning. These include insufficient consideration for the inexperienced user composition of the sample, instructor differences, and the ceiling effect. It will be necessary in future research to test strategies that can more fully promote collaboration among participants who have limited computer experience, control for instructor differences, and use more sensitive measurements to eliminate a potential ceiling effect.

Third, eHealth literacy skills may involve different levels of skills, with some skills being relatively easier to obtain and others requiring more effort. In future research, it will be necessary to develop more refined measures to assess changes in skills on different levels (eg, skills in not only finding a particular health topic on the NIHSeniorHealth.gov site but also determining the quality of information on any health website).

Fourth, this experimental study did not have a qualitative component. Future research can include qualitative data collection and analysis, which may generate additional insights into the learning process and the relative effects of collaborative versus individualistic learning.

Fifth, this study did not include measures of participants' potential practice of the skills outside of the intervention (eg, at home or other locations). Future research may include these measures to determine whether and how outside practice might affect the effects of the intervention.

Finally, changes in participation in own health care were measured in this study by self-report and at only one time point (post intervention) with no follow-up beyond the intervention period. In future research, it will be necessary to add more objective measures (eg, physicians' reports) and measure this variable over time at multiple time points.

Conclusions

The findings of this experimental study contribute to the literatures on adult learning, social interdependence theory, and health literacy. This study used both objective and self-report measures. The findings from both types of measures are consistent and, together, provide strong evidence that the eHealth literacy intervention tested in this study, regardless of the specific learning method used, significantly improved knowledge, skills, and efficacy. Participants also had overwhelmingly positive attitudes toward the intervention. Participants reported changes in participation in their own health care as a result of the intervention, further supporting the effectiveness of the intervention on improving older adults' eHealth literacy. Collaborative learning did not differ from individualistic learning in affecting the learning outcomes, suggesting that the previously widely reported advantages of

collaborative learning over individualistic learning may not be easily applied to the older population in informal settings, though several confounding factors might have contributed to this finding. Further research is necessary before a more firm conclusion can be drawn. Finally, group composition based on gender, familiarity with peers, and prior computer experience demonstrated no significant interaction or main effect on the learning outcomes.

The study addressed an important social problem: the health "illiteracy" problem among older adults, particularly those who have low incomes, limited education, limited prior computer/Internet experience, and/or belong to ethnic minority groups. The findings of this study contribute to scientific knowledge by advancing theory in older adult learning, particularly the generalizability and application of the collaborative versus individualistic learning method to the older population in an informal setting, and the use of these learning methods as an effective eHealth literacy intervention. By focusing the content of learning on eHealth literacy knowledge and skills, this study broadens current understanding of the health literacy concept and interventions to address the increasing importance of technology in health care. By developing and testing the effectiveness of an eHealth literacy intervention, this study shapes this newly emerging component of health literacy (ie, eHealth literacy) that has increasing significance in contemporary health care.

This study broadens current paradigms in health literacy by using concepts and approaches novel to the field of health literacy. First, while health literacy has been promoted as a lifelong learning process [22], little attention has been paid to examining the relative benefits of different instructional methods (eg, individualistic versus collaborative learning) on older adults' learning of health literacy knowledge and skills. Second, while some prior interventions have involved the use of computers, their primary approach is presenting medical materials on a specific topic through a specially designed localized computer-based system [21]. While such an approach has its advantages (eg, targeting a very specific problem), it also has limitations: it requires extensive resources (to develop and update), and the knowledge and skills learned through this approach are often difficult to generalize to other areas or computer systems. In contrast, in the present study, we used the high-quality Internet health information resources maintained and updated by the NIH that focus on a broad range of medical knowledge, and provide training to improve general knowledge about and skills in finding information on any health topic that might be of interest to an individual—and from a common computer system. Thus, this eHiLL intervention is cost effective and easily transferrable.

Acknowledgments

The author thanks the administrators and staff of the Prince George's County Memorial Library System for facilitating the on-site recruitment and logistics, the instructors for carrying out the instructions, graduate assistant Ivan Watkins for scheduling and administering the data collection instruments, and graduate assistant Man Huang for administering the data collection instruments and data entry. Special thanks to the reviewers and the editor for their constructive and helpful feedback on earlier versions of this manuscript. The Electronic Health Information for Lifelong Learners (eHiLL) research project is funded by a Faculty Early

Career Development Award from the Institute of Museum and Library Services (RE-04-09-0067-09; PI: Bo Xie) and an R01 grant from the National Institute on Aging and the Office of Behavioral and Social Sciences Research (OBSSR) in the Office of the Director, National Institutes of Health (1R01AG041284-01; PI: Bo Xie).

Conflicts of Interest

None declared

References

1. US Department of Health and Human Services. Healthy People 2010: Understanding and Improving Health. 2nd edition. Washington, DC: US Government Printing Office; 2000.
2. Stvilia B, Mon L, Yong JY. A model for online consumer health information quality. *J Am Soc Inf Sci Technol* 2009;60(9):1781-1791. [doi: [10.1002/asi.21115](https://doi.org/10.1002/asi.21115)]
3. Lustria MLA. Can interactivity make a difference? Effects of interactivity on the comprehension of and attitudes toward online health content. *J Am Soc Inf Sci Technol* 2007;58(6):766-776. [doi: [10.1002/asi.20557](https://doi.org/10.1002/asi.20557)]
4. Xie B, Bugg JM. Public library computer training for older adults to access high-quality Internet health information. *Libr Inf Sci Res* 2009;31(3):155-162. [doi: [10.1016/j.lisr.2009.03.004](https://doi.org/10.1016/j.lisr.2009.03.004)] [Medline: [20161649](https://pubmed.ncbi.nlm.nih.gov/20161649/)]
5. Xie B. Older adults, e-health literacy, and collaborative learning: an experimental study. *J Am Soc Inf Sci Technol* 2011;62(5):933-946. [doi: [10.1002/asi.21507](https://doi.org/10.1002/asi.21507)]
6. Xie B. Experimenting on the impact of learning methods and information presentation channels on older adults' e-health literacy. *J Am Soc Inf Sci Technol* 2011;62(9):1797-1807. [doi: [10.1002/asi.21575](https://doi.org/10.1002/asi.21575)]
7. Nielsen-Bohman L, Panzer AM, Kindig DA, Institute of Medicine. Health Literacy: A Prescription to End Confusion. Washington, DC: National Academies Press; 2004.
8. Kutner M, Greenberg E, Jin Y, Paulsen C. National Center for Education Statistics. 2006. The Health Literacy of America's Adults: Results From the 2003 National Assessment of Adult Literacy (NCES 2006-483) URL: <http://nces.ed.gov/pubs2006/2006483.pdf> [accessed 2011-08-17] [WebCite Cache ID 610sP11KU]
9. Oh H, Rizo C, Enkin M, Jadad A. What is eHealth?: a systematic review of published definitions. *J Med Internet Res* 2005;7(1):e1 [FREE Full text] [Medline: [15829471](https://pubmed.ncbi.nlm.nih.gov/15829471/)] [doi: [10.2196/jmir.7.1.e1](https://doi.org/10.2196/jmir.7.1.e1)]
10. Hernandez LM. Health Literacy, eHealth, and Communication: Putting the Consumer First: Workshop Summary. Washington, DC: National Academies Press; 2009.
11. Bylund CL, Sabee CM, Imes RS, Sanford AA. Exploration of the construct of reliance among patients who talk with their providers about internet information. *J Health Commun* 2007;12(1):17-28. [doi: [10.1080/10810730601091318](https://doi.org/10.1080/10810730601091318)] [Medline: [17365346](https://pubmed.ncbi.nlm.nih.gov/17365346/)]
12. Fox S. E-patients with a disability or chronic disease. Washington, DC: Pew Internet & American Life Project; 2007 Oct 08. URL: http://www.pewinternet.org/~media/Files/Reports/2007/EPatients_Chronic_Conditions_2007.pdf.pdf [accessed 2011-10-29] [WebCite Cache ID 62o2c4anF]
13. Huntington P, Nicholas D, Jamali HR, Russell C. Health information for the consumer: NHS vs the BBC. *Aslib Proc* 2007;59(1):46-67. [doi: [10.1108/00012530710725205](https://doi.org/10.1108/00012530710725205)]
14. Norman CD, Skinner HA. eHealth literacy: essential skills for consumer health in a networked world. *J Med Internet Res* 2006;8(2):e9 [FREE Full text] [doi: [10.2196/jmir.8.2.e9](https://doi.org/10.2196/jmir.8.2.e9)] [Medline: [16867972](https://pubmed.ncbi.nlm.nih.gov/16867972/)]
15. Jones S, Fox S. Generations online in 2009. Washington, DC: Pew Internet & American Life Project; 2009 Jan 28. URL: http://www.pewinternet.org/~media/Files/Reports/2009/PIP_Generations_2009.pdf [accessed 2011-10-29] [WebCite Cache ID 62o2oy9hv]
16. Rideout V, Neuman T, Kitchman M, Brodie M. Kaiser Family Foundation. 2005 Jan. e-Health and the Elderly: How Seniors Use the Internet for Health Information URL: <http://www.kff.org/entmedia/upload/e-Health-and-the-Elderly-How-Seniors-Use-the-Internet-for-Health-Information-Key-Findings-From-a-National-Survey-of-Older-Americans-Survey-Report.pdf> [accessed 2011-10-29] [WebCite Cache ID 62o3DMlkl]
17. Xie B. Older adults, health information, and the Internet. *Interactions* 2008;15(4):44-46.
18. Mika VS, Kelly PJ, Price MA, Franquiz M, Villarreal R. The ABCs of health literacy. *Fam Community Health* 2005;28(4):351-357. [Medline: [16166862](https://pubmed.ncbi.nlm.nih.gov/16166862/)]
19. Parker R, Kreps GL. Library outreach: overcoming health literacy challenges. *J Med Libr Assoc* 2005 Oct;93(4 Suppl):S81-S85. [Medline: [16239962](https://pubmed.ncbi.nlm.nih.gov/16239962/)]
20. Andrus MR, Roth MT. Health literacy: a review. *Pharmacotherapy* 2002 Mar;22(3):282-302. [Medline: [11898888](https://pubmed.ncbi.nlm.nih.gov/11898888/)]
21. Schaefer CT. Integrated review of health literacy interventions. *Orthop Nurs* 2008;27(5):302-317. [doi: [10.1097/01.NOR.0000337283.55670.75](https://doi.org/10.1097/01.NOR.0000337283.55670.75)] [Medline: [18832992](https://pubmed.ncbi.nlm.nih.gov/18832992/)]
22. Kickbusch I. Kickbusch Health Consult. 2004. Background Paper: Improving Health Literacy in the European Union: Towards a Europe of Informed and Active Health Citizens URL: <http://www.ilonakickbusch.com/health-literacy/health-literacy-gastein.pdf> [accessed 2011-08-17] [WebCite Cache ID 610sfpbYK]

51. Zandri E, Charness N. Training older and younger adults to use software. *Educ Gerontol* 1989;15(6):615-631. [doi: [10.1080/0380127890150606](https://doi.org/10.1080/0380127890150606)]
52. Norman CD, Skinner HA. eHEALS: the eHealth Literacy Scale. *J Med Internet Res* 2006;8(4):e27 [FREE Full text] [doi: [10.2196/jmir.8.4.e27](https://doi.org/10.2196/jmir.8.4.e27)] [Medline: [17213046](https://pubmed.ncbi.nlm.nih.gov/17213046/)]
53. Bandura A. *Social Foundations of Thought and Action: A Social Cognitive Theory*. Englewood Cliffs, NJ: Prentice-Hall; 1986.
54. Pace RC, Kuh GD. Indiana University Center for Postsecondary Research and Planning, School of Education. 1988. College Student Experiences Questionnaires URL: <http://dpb.cornell.edu/documents/1000093.pdf> [accessed 2011-10-29] [WebCite Cache ID 62o4A8Umw]
55. Fox S. Online health search 2006. Washington, DC: Pew Internet & American Life Project; 2006 Oct 29. URL: http://www.pewinternet.org/~media/Files/Reports/2006/PIP_Online_Health_2006.pdf.pdf [accessed 2011-08-17] [WebCite Cache ID 610snOh4j]
56. NIHSeniorHealth. National Institute on Aging. 2010 Oct 12. Helping Older Adults Search for Health Information Online: A Toolkit for Trainers URL: <http://nihseniorhealth.gov/toolkit/toolkit.html> [accessed 2011-08-19] [WebCite Cache ID 613iGk3mp]
57. Childs S. Judging the quality of Internet-based health information. *Perform Meas Metr* 2005;6(2):80-96. [doi: [10.1108/14678040510607803](https://doi.org/10.1108/14678040510607803)]
58. Kunst H, Groot D, Latthe PM, Latthe M, Khan KS. Accuracy of information on apparently credible websites: survey of five common health topics. *BMJ* 2002 Mar 9;324(7337):581-582 [FREE Full text] [Medline: [11884323](https://pubmed.ncbi.nlm.nih.gov/11884323/)]
59. Morrell RW, Dailey SR, Rousseau GK. Applying research: the NIHSeniorHealth.gov project. In: Charness N, Warner Schaie K, editors. *Impact of Technology on Successful Aging*. New York, NY: Springer; 2003:134-161.
60. Xie B. Older adults' health information wants in the internet age: implications for patient-provider relationships. *J Health Commun* 2009 Sep;14(6):510-524. [doi: [10.1080/10810730903089614](https://doi.org/10.1080/10810730903089614)] [Medline: [19731124](https://pubmed.ncbi.nlm.nih.gov/19731124/)]
61. Williamson K. Discovered by chance: the role of incidental information acquisition in an ecological model of information use. *Libr Inf Sci Res* 1998;20(1):23-40. [doi: [10.1016/j.lisr.2009.03.004](https://doi.org/10.1016/j.lisr.2009.03.004)]
62. Asla T, Williamson K, Mills J. The role of information in successful aging: the case for a research focus on the oldest old. *Libr Inf Sci Res* 2006;28(1):49-63. [doi: [10.1016/j.lisr.2005.11.005](https://doi.org/10.1016/j.lisr.2005.11.005)]
63. Birren J, Warner Schaie K. *Handbook of the Psychology of Aging*. 3rd edition. San Diego, CA: Academic Press; 1990.
64. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd edition. Hillsdale, NJ: L Erlbaum Associates; 1988.
65. Johnson DW, Johnson RT, Smith KA. *Active Learning: Cooperation in the College Classroom*. Edina, MN: Interaction Book Co; 1991.
66. Springer L, Stanne M, Donovan S. Effects of small-group learning on undergraduates in science, mathematics, engineering and technology: a meta-analysis. *Rev Educ Res* 1999;69(1):21-52. [doi: [10.3102/00346543069001021](https://doi.org/10.3102/00346543069001021)]
67. Brody DS. The patient's role in clinical decision-making. *Ann Intern Med* 1980 Nov;93(5):718-722. [Medline: [7212484](https://pubmed.ncbi.nlm.nih.gov/7212484/)]
68. Jones JA, Phillips GM. *Communicating With Your Doctor: Rx for Good Medical Care*. Carbondale, IL: Southern Illinois University Press; 1988.
69. Ballard-Reisch DS. A model of participative decision making for physician-patient interaction. *Health Commun* 1990;2(2):91-104. [doi: [10.1207/s15327027hc0202_3](https://doi.org/10.1207/s15327027hc0202_3)]
70. McNutt RA. Shared medical decision making: problems, process, progress. *JAMA* 2004 Nov 24;292(20):2516-2518. [doi: [10.1001/jama.292.20.2516](https://doi.org/10.1001/jama.292.20.2516)] [Medline: [15562133](https://pubmed.ncbi.nlm.nih.gov/15562133/)]
71. Benbassat J, Pilpel D, Tidhar M. Patients' preferences for participation in clinical decision making: a review of published surveys. *Behav Med* 1998;24(2):81-88. [doi: [10.1080/08964289809596384](https://doi.org/10.1080/08964289809596384)] [Medline: [9695899](https://pubmed.ncbi.nlm.nih.gov/9695899/)]

Abbreviations

ANOVA: analysis of variance

eHiLL: Electronic Health Information for Lifelong Learners

Edited by G Eysenbach; submitted 29.06.11; peer-reviewed by ML Lustria, D Kaufman; comments to author 04.08.11; revised version received 14.09.11; accepted 27.09.11; published 03.11.11

Please cite as:

Xie B

Effects of an eHealth Literacy Intervention for Older Adults

J Med Internet Res 2011;13(4):e90

URL: <http://www.jmir.org/2011/4/e90/>

doi: [10.2196/jmir.1880](https://doi.org/10.2196/jmir.1880)

PMID: [22052161](https://pubmed.ncbi.nlm.nih.gov/22052161/)

©Bo Xie. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 03.11.2011. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.