

NIH Public Access

Author Manuscript

Crit Rev Biochem Mol Biol. Author manuscript; available in PMC 2011 December 1

Published in final edited form as:

Crit Rev Biochem Mol Biol. 2010 December ; 45(6): 463-487. doi:10.3109/10409238.2010.502516.

The ESCRT Complexes

James H. Hurley

Laboratory of Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, U.S. Department of Health and Human Services, Bethesda, MD 20892.

Abstract

The ESCRT machinery consists of the peripheral membrane protein complexes, ESCRT-0, -I, -II, -III, and Vps4-Vta1, and the ALIX homodimer. The ESCRT system is required for degradation of unneeded or dangerous plasma membrane proteins; biogenesis of the lysosome and the yeast vacuole; the budding of most membrane enveloped viruses; the membrane abscission step in cytokinesis; macroautophagy; and several other processes. From their initial discovery in 2001-2002, the literature on ESCRTs has grown exponentially. This review will describe the structure and function of the six complexes noted above and summarizes current knowledge of their mechanistic roles in cellular pathways and in disease.

INTRODUCTION

Historical background

The discovery and history of the ESCRT complexes is intimately connected to the characterization of multivesicular bodies (MVBs) (Piper and Katzmann, 2007). The term MVB was first applied to features in electron micrographs of cells in the 1950s (Sotelo and Porter, 1959). The role of MVBs in the lysosomal degradation of activated signaling receptors became clear two decades later from electron microscopic analysis of the internalization of epidermal growth factor (Gorden et al., 1978; Haigler et al., 1979) (Fig. 1). The power of yeast genetics to dissect intracellular transport pathways became clear at the start of the 1980s (Novick et al., 1980). The vacuole is the yeast counterpart of the lysosome of animal cells. By the late 1990s, more than 60 yeast vacuolar protein sorting (VPS) genes involved in vacuole biogenesis had been identified (Bryant and Stevens, 1998). Thirteen of these genes fell into the class E morphological subgroup, and were shown to be involved in the transport of proteins into the MVB en route to the vacuole. These genes code for most of the core subunits of the ESCRT complexes-0 (Bilodeau et al., 2002; Katzmann et al., 2003), ESCRT-I (Katzmann et al., 2001), ESCRT-II (Babst et al., 2002b), ESCRT-III (Babst et al., 2002a). Soon thereafter, similar functions in receptor sorting to lysosomes via the MVB pathway were demonstrated for the mammalian counterparts of the yeast ESCRT complexes (Bache et al., 2003a; Bache et al., 2003b; Lu et al., 2003; Raiborg et al., 2002). In the past decade, the ESCRTs have been found to act in viral budding, cytokinesis, autophagy, and other pathways in addition to MVB biogenesis (Fig. 2).

Scope of review

As we approach the ten-year anniversary of the discovery and naming of the ESCRT system, it seems appropriate to take a comprehensive look at what has been learned about the system. With a literature of over 500 publications that is growing at a rate of about three per week, this may also be one of the last opportunities to even attempt a comprehensive review. Even in a

Correspondence: hurley@helix.nih.gov.

broad overview, it is impractical to cite and discuss every paper in the field. For more detailed information on the structure, function, and biochemistry of the ESCRTs, see (Alam and Sundquist, 2007; Babst, 2005; Hurley, 2008; Hurley and Emr, 2006; Saksena et al., 2007; Williams and Urbe, 2007). The membrane mechanics of ESCRT-mediated budding and scission are reviewed by Hurley & Hanson (Hurley and Hanson, 2010). For more background on the biogenesis of multivesicular bodies, see (Gruenberg and Stenmark, 2004; Hanson et al., 2009; Piper and Katzmann, 2007). Further details on the sorting of ubiquitinated receptors and other cargo by the ESCRTs can be found in (Hurley and Ren, 2009; Katzmann et al., 2002; Luzio et al., 2009; Piper and Luzio, 2007; Raiborg and Stenmark, 2009). For reviews on the role of ESCRTs in viral budding, see (Bieniasz, 2006; Bieniasz, 2009; Carter and Ehrlich, 2008; Chen and Lamb, 2008; Freed and Mouland, 2006; Fujii et al., 2007; Morita and Sundquist, 2004; Usami et al., 2009). The role of ESCRTs in cytokinesis is reviewed by (Carlton and Martin-Serrano, 2009; McDonald and Martin-Serrano, 2009; Samson and Bell, 2009; Steigemann and Gerlich, 2009). The possible roles of ESCRTs in autophagy are reviewed by (Rusten and Stenmark, 2009). The role of ESCRTs in disease are covered by (Saksena and Emr, 2009; Stuffers et al., 2009).

Most of the newcomers to the ESCRT field are drawn in by discovering interactions between their favorite (usually metazoan) proteins and ESCRTs, or that the ESCRTs function in the metazoan cellular processes or human diseases that they study. One of the most confounding aspects for such newcomers is that much of the fundamental cell biology, biochemistry, and structural biology have focused on the yeast ESCRTs. Yeast and human ESCRT nomenclature differs. Unfortunately, ESCRT researchers have thus far been unable to reach agreement on a systematic nomenclature across species. Generally yeast-centric terminology will be used in this review, although an exception will be made for ALIX and for the subunits of ESCRT-0, for which the metazoan literature is substantially more extensive. See Table 1 for the conversion between yeast and metazoan nomenclature. While the main functions of the ESCRTs are conserved between yeast and humans, there are also important differences. One of the main goals of this review will be to highlight the similarities and differences between the most important aspects of the yeast and human ESCRTs.

STRUCTURE AND FUNCTION OF THE ESCRT COMPLEXES

ESCRT-0

The ESCRT-0 complex is required for sorting plasma membrane proteins into the MVB pathway in animal cells and for MVB biogenesis in yeast. ESCRT-0 binds to and clusters ubiquitinated cargo for delivery into MVBs, and recruits clathrin, ubiquitin ligases, and deubiquitinating enzymes, and almost certainly has other functions as well. ESCRT-0 functions as a 1:1 heterodimer (Ren et al., 2009) of the subunits Vps27 and Hse1 (yeast; Fig. 3a)) (Bilodeau et al., 2002) and Hrs and STAM (metazoa; Fig. 3b) (Asao et al., 1997; Bache et al., 2003b). In humans there are two isoforms of the latter, STAM1 and STAM2. Both the Vps27 and Hse1 subunits contain N-terminal ubiquitin-binding VHS domains (Ren and Hurley). The VHS domain of Vps27 is followed by a FYVE domain (Burd and Emr, 1998; Gaullier et al., 1998; Kutateladze et al., 1999; Lohi and Lehto, 1998; Misra and Hurley, 1999) (Misra et al., 2001). The FYVE domains of Vps27 and Hrs bind phosphatidylinositol 3-phosphate with tens of nM affinity (Stahelin et al., 2002) and are responsible for targeting ESCRT-0 to early endosomes (Raiborg et al., 2001b). Two ubiquitin-binding UIM motifs follow the FYVE domain of Vps27 (Bilodeau et al., 2002; Fisher et al., 2003; Shih et al., 2002; Swanson et al., 2003). The domain structure of human Hrs mirrors that of Vps27 in most respects, but the two UIMs of Vps27 are replaced by a single double-sided UIM (DUIM) in Hrs (Hirano et al., 2006a) such that both the human and yeast subunits bind equal numbers of ubiquitin moieties, a total of five each. The multiplicity of ubiquitin binding domains allows ESCRT-0 to bind polyubiquitin chains with high avidity (Ren and Hurley) and cluster ubiquitinated cargo

(Wollert and Hurley, 2010) in vitro. Hrs can undergo UIM-dependent autoubiquitination that is thought to result in autoinhibition of the protein's ability to bind ubiquitin (Hoeller *et al.*, 2006). Subsequent to the VHS domain, STAM (and Hse1) contain an SH3 domain that is involved in recruiting deubiquitinating enzymes such as UBPY (Kato *et al.*, 2000).

The two subunits of the ESCRT-0 heterodimerize through an elongated, rigid core consisting of an antiparallel coiled coil and two domain-swapped GAT domains (Prag et al., 2007; Ren et al., 2009). In HeLa cells all of the endogenous Hrs and STAM proteins are incorporated into the ESCRT-0 complex, leaving no pool of free subunits (Ren et al., 2009). When taken out of the context of the assembled ESCRT-0 complex, free Hrs and STAM expose a large region of unpartnered coiled coil that binds promiscuously to many potentially non-physiological partners and leads to the formation of large non-physiological aggregates. As with subunits of ESCRT-I and -II, which are also constitutively assembled, caution must be exercised in drawing conclusions from studies of isolated subunits. The GAT domains involved in heterodimerization do not bind ubiquitin and lack the consensus residues for ubiquitin binding (Prag et al., 2007). The regions of ESCRT-0 subunits C-terminal to the core region are unstructured and contain a number of interaction motifs for other partners, although the function of most of these sequences (apart from a role a spacer) is unknown. The best-studied of these short motifs are P(S/T)XP motifs of Vps27 that bind to the Vps23 subunit of ESCRT-I (Bilodeau et al., 2003; Katzmann et al., 2003; Lu et al., 2003; Pornillos et al., 2003) and a clathrin-binding sequence at the C-terminus of Hrs (Raiborg et al., 2001a). The C-terminal portion of Hse1 is involved in recruiting the ubiquitin ligase Rsp5 (Ren et al., 2007) via an interaction whose counterpart in STAM is uncertain.

ESCRT-I

The ESCRT-I complex co-assembles with ESCRT-II on membranes (Kostelansky *et al.*, 2007) (Fig. 4), and these two complexes appear to function as a 1:1 supercomplex (Gill *et al.*, 2007) to bud the limiting membrane of the MVB into its lumen (Wollert and Hurley, 2010). In HIV-1 budding and cytokinesis, where the ESCRTs are not needed for bud formation, ESCRT-I appears to be able act independently of ESCRT-II. In these latter pathways, ESCRT-I is probably important mainly for recruiting ESCRT-III. ESCRT-I is a heterotetramer of one copy each of the subunits Vps23, Vps28, Vps37 (Katzmann et al., 2001) and Mvb12 (Audhya *et al.*, 2007; Chu *et al.*, 2006; Curtiss *et al.*, 2006; Kostelansky et al., 2007; Morita *et al.*, 2007a; Oestreich *et al.*, 2006b). The subunits heterotetramerize through two contiguous but distinct core regions. Vps23, Vps37, and Mvb12 assemble into a 13 nm-long stalk that consists in part of an unusual antiparallel coiled-coil (Kostelansky et al., 2007). Vps23, Vps28, and Vps37 form a headpiece region consisting of three pairs of antiparallel helices, one pair from each subunit, spread out in the shape of a fan (Kostelansky *et al.*, 2006; Teo *et al.*, 2006). Together the stalk and headpiece form a single rigid 18 nm-long structure.

Vps23 (known as TSG101 in humans) has an N-terminal Ubiquitin E2 Variant (UEV) domain responsible for binding ubiquitinated cargo (Katzmann et al., 2001; Sundquist *et al.*, 2004; Teo *et al.*, 2004b) and P(S/T)XP motifs of ESCRT-0, viral proteins, and other proteins (Bilodeau et al., 2003; Katzmann et al., 2003; Lu et al., 2003; Pornillos et al., 2003). There is a Pro-rich linker region after the UEV domain with a GPPX₃Y motif that targets ESCRT-I to the midbody during cytokinesis (Carlton and Martin-Serrano, 2007; Lee *et al.*, 2008; Morita *et al.*, 2007b). The core region of Vps28 is followed by a short linker and a C-terminal four helix bundle domain (CTD) (Gill et al., 2007; Pineda-Molina *et al.*, 2006). The yeast Vps28-CTD is primarily responsible for the 1:1 interaction with ESCRT-II (Gill et al., 2007; Kostelansky et al., 2006); the binding site on human ESCRT-I for ESCRT-II is unknown. The Vps28-CTD contains a conserved hydrophobic patch of unknown function; a possible role in interactions with the ESCRT-III subunit Vps20 has been suggested (Pineda-Molina *et al.*, 2006). The N-

terminus of yeast Vps37 consists of a basic helix that contributes to membrane binding by ESCRT-I (Kostelansky et al., 2007). Four VPS37 isoforms are present in the human proteome. In addition to two characterized MVB12 isoforms in humans and one in nematodes (Audhya et al., 2007; Morita et al., 2007a), two more putative human MVB12 isoforms have been proposed based on bioinformatics (de Souza and Aravind, 2010). Their sequence similarity to yeast Mvb12 is so low that their identification depended on proteomics and functional characterization, rather than homology. Despite a few clues, much remains to be learned about the functional significance of the multiple VPS37 and MVB12 isoforms in human cells.

ESCRT-II

The ESCRT-II complex is the essential partner of ESCRT-I in MVB biogenesis and bud formation (Babst et al., 2002b; Wollert and Hurley, 2010). Indeed, ESCRT-II is probably the more important of the two complexes in MVB biogenesis, since overexpression of ESCRT-II can rescue deletions of ESCRT-I genes in yeast, but not vice versa (Babst *et al.*, 2002b). Despite its central role in MVB biogenesis and membrane budding, ESCRT-II appears to be non-essential for HIV-1 budding (Langelier *et al.*, 2006) and cytokinesis (Morita et al., 2007b). In addition to playing a central role in membrane bud formation, Like ESCRT-0 and –I, ESCRT-II binds ubiquitinated cargo, although only at one site (Alam *et al.*, 2004). ESCRT-II has a pivotal role in MVB biogenesis in bridging the upstream ubiquitin-binding ESCRT-II is probably responsible for connecting MVBs to microtubules via RILP, Rab7, and dynein (Progida *et al.*, 2007; Wang and Hong, 2006).

ESCRT-II is a Y-shaped 1:2:1 heterotetramer of the subunits Vps22, Vps25, and Vps36 (Hierro et al., 2004; Im and Hurley, 2008; Teo et al., 2004a) (Fig. 4). Despite lacking sequence homology to each other, the bulk of each of these subunits consists of two tandem wingedhelix (WH) motifs. In other contexts, WH motifs are usually involved in DNA binding, however there is no evidence that ESCRT-II binds DNA. ESCRT-II does bind RNA (Irion and St Johnston, 2007), but the RNA-binding activity, described below, does not involve the WH domains. The WH domain core regions of Vps22 and Vps36 form an extensive interface with one another, and these two subunits probably require one another for folding and stability. The two Vps25 subunits are more loosely associated with the Vps22-Vps36 subcomplex, and they do not contact each other. The tips of the second WH domain (WH2) of the Vps25 subunits are the locus for binding to the ESCRT-III subunit Vps20 (Im et al., 2009). Both copies of Vps25 are essential for function (Hierro et al., 2004; Teis et al.). Similar to Vps37 of ESCRT-I, the N-terminus of Vps22 consists of a basic helix that is important for membrane targeting (Im and Hurley, 2008), albeit without specificity for any particular endosomal lipid. The Nterminus of yeast Vps36 is complex. This region contains a variant pleckstrin homology (PH) domain that is referred to as a "GLUE" domain (Slagsvold et al., 2005; Teo et al., 2006). The GLUE domain binds preferentially to phosphatidylinositol 3-phosphate (Teo et al., 2006), although it also binds to other phosphoinositides, as detailed below. Two Npl4 type zinc fingers, NZF1 and NZF2, are inserted within a loop in the yeast GLUE domain. NZF1 is the locus for binding to the ESCRT-I Vps28-CTD (Gill et al., 2007), while NZF2 binds to a single ubiquitin moiety (Alam et al., 2004). Mammalian VPS36 contains a GLUE domain as well, but without any zinc finger insertion. In mammalian VPS36, the GLUE domain itself binds directly to ubiquitin (Alam et al., 2006; Hirano et al., 2006b; Slagsvold et al., 2005). In yeast Vps36, the GLUE-WH1 linker comprises a secondary binding site for ESCRT-I (Im and Hurley, 2008). This region is conserved in human VPS36 and could be the locus for binding to human ESCRT-I, although this has not been verified.

ESCRT-III

The ESCRT-III complex is the central membrane scission machine at the heart of the ESCRT system (Wollert *et al.*, 2009a). Of all the ESCRT complexes, only homologs of ESCRT-III and Vps4 (described below) are present in Archaeal proteomes, where they have a role in cell division (Lindas *et al.*, 2008; Samson *et al.*, 2008). In contrast to ESCRTs 0-II, ESCRT-III subunits contain no known ubiquitin binding domains, and exist in cytosol as inactive monomers (or possibly heterodimers) (Babst et al., 2002a). The subunits polymerize into the active ESCRT-III complex only on the membrane (Babst et al., 2002a). This assembly is detergent insoluble and relatively intractable to most biochemical methods.

In yeast, the core subunits required for function are Vps20, Snf7, Vps24, and Vps2, which assemble in that order (Teis et al., 2008) (Fig. 5). Vps20, Snf7, and Vps24 alone are sufficient for membrane scission (Wollert et al., 2009a), while Vps2 is required for coupling to the Vps4 recycling machinery. The other yeast ESCRT-III subunits, Did2 (Nickerson et al., 2006), Ist1 (Dimaano et al., 2008; Rue et al., 2008), and Vps60 are not strictly essential for function and appear to assemble with the rest of ESCRT-III at a late stage. Did2 (Nickerson et al., 2006) and Vps60 help recruit and activate the Vps4-Vta1 complex for recycling, while Ist1 inhibits Vps4 activity (Dimaano et al., 2008). Vps20-Snf7, Vps24-Vps2 (Babst et al., 2002a), and Did2-Ist1 (Rue et al., 2008; Xiao et al., 2009) former preferential binary pairings with one another. Vps60 is the odd man out in that it binds more tightly to Vta1 (see below) than to any of its fellow ESCRT-III proteins (Azmi et al., 2008; Bowers et al., 2004; Shiflett et al., 2004; Shim et al., 2008). These pairings do not appear to dictate the stoichiometry of assembly. The precise stoichiometry of ESCRT-III is not known, and it is not clear if it is strictly defined or not. At least in yeast, Snf7 is the dominant subunit, present at several-fold higher copy number than any other subunit, including Vps20 (Teis et al., 2008). Certain ESCRT-III subunits either alone (Snf7) or in binary combinations (Vps24+Vps2; Ist1+Did2) form helical tubes of dimensions comparable to those of the necks of HIV-1 buds or the intralumenal vesicles (ILVs) of MVBs (Bajorek et al., 2009b; Hanson et al., 2008; Lata et al., 2008b).

All ESCRT-III subunits contain an electrically polarized five-helix core (Muziol *et al.*, 2006). The first two helices are basic and bind strongly to acidic membranes, while helices 3-5 are acidic. More variable regions are found C-terminal to the five-helix core. In all cases studied, the C-terminal region autoinhibits assembly of the ESCRT-III complex and helps maintain the soluble monomeric state (Lata *et al.*, 2008a; Shim et al., 2007; Zamborlini et al., 2006). Truncations of these regions strongly promote complex assembly (Shim et al., 2007; Zamborlini et al., 2006), as does membrane binding and ESCRT-II binding to Vps20 (Saksena *et al.*, 2009). The extreme C-terminal regions of various subunits contain the MIT-domain-interacting motifs-1 and -2 (MIM1) (Vps24, Vps2, Did2, Ist1) and (MIM2) (Vps20, Snf7, Ist1) motifs that bind to two distinct faces of the Vps4 mIT domain. Human DID2B contains a specialized MIM1 variant that binds to both Vps4 and the MIT domain of the microtubule severing enzyme spastin. These regions also contain sequences that are less well defined at this point and bind a spectrum of MIT domains, described below. The C-terminus of Snf7 contains a motif that binds to the Bro1 domain of Bro1 (McCullough *et al.*, 2008).

Vps4-Vta1

Disassembly of the membrane-bound ESCRT-III complex is required to finish the ESCRT cycle and replenish the cytosolic pool of ESCRT-III subunits. The AAA ATPase Vps4 solubilizes ESCRT-III subunits at the cost of ATP hydrolysis (Fig. 6). ATP consumption by Vps4 is the main thermodynamic driving force for the ESCRT cycle. The domain structure of Vps4 consists of an N-terminal ESCRT-III-binding MIT domain (Scott *et al.*, 2005b), a flexible linker, a large ATPase mixed- α/β domain, a small ATPase helical domain, a β -domain, and a C-terminal helix (Gonciarz *et al.*, 2008;Scott *et al.*, 2005a;Xiao *et al.*, 2007). Vps4 functions

as an oligomer that under most conditions and by most reports is a dodecamer (Inoue *et al.*, 2008;Landsberg *et al.*, 2009;Yu *et al.*, 2008), although a tetradecameric form has been reported as well (Hartmann et al., 2008). The dodecamer consists of two conformationally distinct hexameric rings (Yu et al., 2008). The lower ring has a constricted pore and has been modeled based on the structure of the p97 D1 domain hexamer. The upper ring has a wider pore but its structure is not known in detail due to the limited resolution of the EM study and the lack of a closely related crystal structure. The central pore is required for function (Gonciarz et al., 2008;Scott et al., 2005a). ESCRT-III subunits physically contact pore residues during disassembly, but whether they pass all the way through the pore is uncertain.

Vta1 is an ESCRT protein that binds to Vps4 and promotes its oligomerization, activity, and ESCRT-III binding (Lottridge et al., 2006; Shestakova et al., 2010; Shiflett et al., 2004; Ward et al., 2005; Yeo et al., 2003). Although Vta1 is not constitutively associated with Vps4 in cytosol as are the subunits of ESCRT-0, -I, and -II, for practical purposes Vps4 and Vta1 function together is a complex in MVB biogenesis, HIV-1 budding, and probably other Vps4dependent pathways. The deletion of Vta1 does not cause as severe of a defect in cargo sorting as Vps4 (Lottridge et al., 2006; Shiflett et al., 2004; Yeo et al., 2003). Vta1 accelerates Vps4 ATPase activity (Azmi et al., 2006; Lottridge et al., 2006) and oligomerization in vitro, and promotes ESCRT-III disassembly in vitro (Azmi et al., 2008). Vta1 binds to Vps4 via the ß domain of the latter protein (Scott et al., 2005a; Yang and Hurley, 2010). Vta1 contains two MIT domains at its N-terminus (Xiao et al., 2008), both so divergent in sequence from other MIT domains that they were only identified after their crystal structures were solved. Vta1 is a dimer, with the subunits associating via the Vps4-binding VSL (Vps4, SBP1, LIP5) domain at the C-terminus of the protein (Azmi et al., 2006; Xiao et al., 2008). The Vta1 dimer appears to be tight and constitutive, and the dimer contact residues are required for its function. A long flexible linker connects the MIT domains to the VSL region. The Vta1 MIT domains interact with various ESCRT-III proteins, but especially strongly with the late-acting ESCRT-III protein Vps60 (Bowers et al., 2004) (Azmi et al., 2008; Shiflett et al., 2004; Shim et al., 2008). Given that Vps60 appears to bind more tightly to Vta1 than to any of the ESCRT-III subunits, it seems reasonable to think of Vps60 as an adaptor for Vta1 to interact with the ESCRT-III assembly (Nickerson et al., 2010).

The assembly of the Vps4-Vta1 complex has been intensively studied, yet it is still a source of mystery. A 37 Å resolution EM reconstruction shows that the two hexameric rings of Vps4 are non-equivalent, and that the lower ring surrounds a large pore (Yu *et al.*, 2008). Incorporation of Vta1 into the assembly leads to the appearance of additional densities at several regions around the structure, but it is not clear which densities belong to Vta1 itself and which reflect conformational changes in Vps4 induced by Vta1 binding. The β domain of Vps4, to which Vta1 binds, projects outward from the six tips of the lower ring of the hexamer, providing potentially up to 6 binding sites for Vta1 dimers at the outer edges of the two rings. The orientation of the β domain in the upper ring is not known. The reported 6:12 stoichiometry of Vta1 to Vps4 is hard to reconcile with current understanding of the symmetry of the two molecules and the number of binding site. It has been suggested that Vta1 dimers could crosslink multiple Vps4 dodecamers in a lattice like arrangement (Yang and Hurley, 2010), but direct confirmation of this idea is lacking. Thus a higher resolution structure of the complete Vps4-Vta1 assembly is urgently needed.

The mechanism of ESCRT-III disassembly is understood only in the most general outlines. The Vps4 MIT domains bind to the substrates. The high affinity interaction between the Vps2 MIM1 and the Vps4 MIT appears to be particularly critical. Recycling cannot be observed in vitro without either Vps2 (Lata et al., 2008b; Wollert et al., 2009a) or at least a chimeric Vps24 bearing a Vps2 MIM1 at its C-terminus (Ghazi-Tabatabai *et al.*, 2008). Even the very low affinity interactions of the Vps4 MIT domain with the Snf7 MIM2 are important for function

(Shestakova et al., 2010). After all, Snf7 is the most abundant ESCRT-III subunit in the lattice. Residues of the central pore of Vps4 are required for the disassembly function (Scott et al., 2005a), but more direct evidence that ESCRT-III subunit physically occupy the pore at some point during disassembly is lacking. However it is hard to envision how the disassembly reaction could proceed without the bulk of the dodecamer acting as a rigid fulcrum to pry apart the ESCRT-III subunits. This might involve direct ESCRT-III pore interactions, or could be less direct if the MIT domain were physically in contact with both the pore and the ESCRT-III subunits Did2 and Ist1 offer further questions. These two subunits preferentially associate, yet Did2 is a key activator of disassembly, while Ist1 seems to inhibit Vps4 via an unknown mechanism. Mutations in these subunits are among the few defects in the ESCRT pathway that perturb the size of ILVs (Nickerson et al., 2010; Nickerson et al., 2006), and the mechanism behind their unusual phenotypes is obscure. Given these open questions, the ESCRT-III disassembly mechanism is likely to remain one of the most active areas of mechanistic ESCRT research in the next few years.

Bro1/ALIX

In yeast, Bro1 is not strictly required for MVB biogenesis, and seems to be mainly involved in promoting the recruitment and activity of the deubiquitinating enzyme Doa4 (Luhtala and Odorizzi, 2004) to the assembling ESCRT-III complex. By the same token, the mammalian counterpart of Bro1, ALIX, seems to be dispensable for the MVB targeting of the canonical cargo EGFR. However, ALIX has a central and required role in targeting the ESCRT machinery to the midbody for membrane abscission in cytokinesis, and is essential for the budding of certain viruses. ALIX was initially isolated based on its association with an apoptosis-related protein, ALG-2 (Missotten *et al.*, 1999; Vito *et al.*, 1999). It remains a mystery whether ALIX in some way serves to connect the apoptosis and MVB pathways. ALIX also has a range of interactions with endocytic and cytoskeletal components whose connection to the ESCRT machinery is unclear (Odorizzi, 2006). ALIX seems to be connected somehow to the lipid lysobisphosphatidic acid (LBPA), which is enriched in mammalian late endosomes and lysosomes (Matsuo *et al.*, 2004), although direct binding of ALIX and LBPA has not been shown.

ALIX has a tripartite domain structure consisting of an N-terminal Bro1 domain, a central V domain, and a flexible C-terminal proline-rich domain (PRD) (Fig. 7). The Bro1 domain is shaped like a banana (Kim *et al.*, 2005), and binds to the C-terminal region of Snf7 through a conserved patch near the center of its concave face (Kim et al., 2005;McCullough et al., 2008). The V domain consists of two helical arms joined so as to form the shape of the letter "V". The V domain binds to YPXL motif sequences at a site on one of the arms facing into the crevice of the V-shape (Fisher *et al.*, 2007;Lee *et al.*, 2007b;Zhai *et al.*, 2008). The isolated V domain undergoes a monomer-dimer equilibrium (Lee et al., 2007b) and is responsible for dimerizing the intact ALIX molecule (Pires *et al.*, 2009).

The Alix PRD comprises a remarkably concentrated zone of interaction motifs. The overlapping motifs that bind the midbody protein CEP55 (Lee et al., 2008) and the apoptotic protein ALG-2 (Suzuki *et al.*, 2008) have been co-crystallized and visualized with the interaction partners. The Alix PRD contains a P(S/T)XP motif that interacts with ESCRT-I, but this motif is dispensable for function, at least in the setting of HIV-1 budding (Fisher et al., 2007). The polyproline tract responsible for endophilin binding is also dispensable for the HIV-1 budding activity (Fisher et al., 2007) (Usami *et al.*, 2007). Finally, the PRD autoinhibits the ability of the intact protein to bind SNF7 (Zhou *et al.*, 2008).

ESCRT INTERACTIONS AND FUNCTIONAL DOMAINS

ESCRT-ubiquitin interactions

By recent counts, roughly twenty classes of ubiquitin binding domains (UBDs) have been identified (Dikic *et al.*, 2009; Hurley *et al.*, 2006) in eukaryotic proteomes. The ESCRT system is a microcosm of this richness, making use of no fewer than six types of UBD: the VHS domain; the UIM and its variant the DUIM; the UEV domain; the Mvb12 C-terminal domain; the GLUE domain; and the NZF domain (Fig. 8). All of these domains interact with the low affinity (Kd > 100 μ M) with monoubiquitin via Ile44 hydrophobic patch on the ubiquitin surface. This implies that the binding of any one ESCRT UBD to an ubiquitin moiety excludes the binding of that moiety to another ESCRT UBD. It also implies that the UBDs must cooperate either with one another or with other domains to increase their affinity for cargo, since ubiquitinated substrate proteins are present at bulk concentrations far lower than the 100s of μ M in cells.

VHS domains are eight-helix bundles that contain an ubiquitin binding site formed by hydrophobic residues on helices 2 and 4 (Hong *et al.*, 2009). These residues, in particular a Leu and a Trp from helix 2, are conserved in the VHS domains of ESCRT-0 subunits from yeast to mammals. ESCRT-0 VHS domain binds ubiquitin at a single site with affinities of $K_d = ~100 \mu$ M to ~2 mM. Yeast ESCRT-0 contains a total of three ubiquitin interacting motifs (UIMs). UIMs are among the simplest class of UBD, consisting of a single Leu and Ala-rich α -helix. The yeast ESCRT-0 UIMs bind ubiquitin with ~200 to 300 μ M affinities (Fisher et al., 2003). Human ESCRT-0 contains one conventional UIM and one double-sided UIM (DUIM). DUIMs contain two interlaced ubiquitin binding motifs that allow a single α -helix to bind two different ubiquitin moieties, one on each side of the helix (Hirano et al., 2006a). Thus both yeast and human ESCRT-0 contain a total of five ubiquitin binding sites. None of the ESCRT-0 UBDs are individually essential for function, but inactivation of more than two out of the five cripples the sorting function of ESCRT-0 in yeast.

ESCRT-I from all species contains a UEV domain at the N-terminus of the Vps23 subunit, which binds one ubiquitin moiety (Sundquist et al., 2004; Teo et al., 2004b). Yeast Mvb12 has a short C-terminal sequence that binds ubiquitin (Shields *et al.*, 2009), again in close proximity in three dimensions to the Vps23 UEV domain such that it could potentially cooperate in polyubiquitin binding. Finally, yeast and human ESCRT-II both contain a single ubiquitin-binding site, but they bind ubiquitin in different ways. The yeast Vps36 GLUE domain contains two Npl4 zinc fingers inserted within the $\beta6-\beta7$ loop. The second zinc finger (NZF2) of Vps36 binds a single ubiquitin moiety (Alam et al., 2004). Human VPS36 has a simpler structure than its yeast ortholog, and is missing the two NZF domains. However, the GLUE domain of human VPS36 has acquired the ability to bind ubiquitin directly, with an affinity of 105 μ M (Alam et al., 2006), slightly tighter than that of the yeast NZF2. None of the UBDs of yeast ESCRT-I and –II is individually essential, but they contribute to cargo sorting in a cooperative manner, and multiple deletions shut down ubiquitinated cargo sorting (Shields et al., 2009). The remaining ESCRT complexes ESCRT-III, Vps4-Vta1, and Bro1, are not known to bind ubiquitin.

The very low affinities of ESCRT UBDs for ubiquitin monomers beg the question as to their function in sorting cargoes that are typically present in cells at nM concentrations. There is growing evidence that in many or most cases, Lys63-linked polyubiquitination is involved in the trafficking of many or most cargoes into the ESCRT pathway. In yeast the vacuolar hydrolase Cps1 requires K63 polyubiquitination to transit from the Golgi into the MVB pathway, and the plasma membrane amino acid transporter Gap1 requires polyubiquitination for its sorting from early endosomes into MVBs (Lauwers *et al.*, 2009). Many other yeast transporters require polyubiquitination for their vacuolar sorting (Lauwers *et al.*, 2010), and

the ESCRT-dependence of this process can be reasonably assumed, even if not demonstrated explicitly. In human cells, the majority of ubiquitin attached to the ESCRT substrate EGFR occurs in the form of K63-linked chains (Huang *et al.*, 2006). While the K63-linked polyubiquitin dependence of ESCRT trafficking has only been rigorously demonstrated in a few cases, the observation of polyubiquitination of most of the relevant cargoes has led a general impression that K63 polyubiquitination is probably a major signal for cargo entry into the pathway. Typically chains lengths of 2-3 ubiquitin moieties have been observed (Lauwers et al., 2010), but the chain length requirements have not been tested rigorously. ESCRT-0 binds to K63-linked di- and tetraubiquitin with affinities in the tens of μ M (Ren and Hurley). The reduction of dimensionality from three to two on the membrane is probably a more compelling solution to the apparent paradox posed by the low affinity of ESCRT UBDs for ubiquitin in solution. When ubiquitin at the biologically plausible bulk concentrations of 65 nM for ubiquitin and 15 nM for ESCRTs (Wollert and Hurley, 2010).

ESCRT-lipid interactions

The endosomal lipid phosphatidylinositol 3-phosphate (PI(3)P) plays a pivotal role in the ESCRT pathway. PI(3)P binds to the FYVE domain of the Hrs/Vps27 subunit of ESCRT-0 with high affinity (Sankaran *et al.*, 2001; Stahelin et al., 2002)(Fig. 9), and is essential for recruitment of ESCRT-0 to endosomes (Katzmann et al., 2003; Raiborg et al., 2001b). ESCRT-I has a low affinity for membranes on its own (Kostelansky et al., 2007), and depends on protein-protein interactions to recruit it to and stabilize it at membranes. ESCRT-II binds preferentially to PI(3)P but- in contrast to the highly specific Hrs-PI(3)P interaction- also binds to other phosphoinositides via its GLUE domain (Slagsvold et al., 2005; Teo et al., 2006). The affinity of ESCRT-II for membranes is further enhanced by a non-specific but functionally important interaction between the basic N-terminal helix of the Vps22 subunit and acidic membranes lipids (Im and Hurley, 2008). ESCRT-III subunits bind to acidic membrane lipids with little or no specificity through a broad basic patch (Lata et al., 2008b; Muziol et al., 2006). There has been one report that Vps24 binds preferentially to PI(3,5)P₂ (Whitley *et al.*, 2003), but in vitro, this lipid has a minimal effect on the activity of Vps24 (Wollert et al., 2009a).

MIT domains and MIT-interacting motifs

Microtubule-interacting and transport (MIT) domains were so named prior to the discovery of their main function in the ESCRT pathway: as modules that interact with C-terminal MIT domain-interacting motifs (MIMs) of ESCRT-III subunits (Hurley and Yang, 2008; Kieffer et al., 2008; Obita et al., 2007; Stuchell-Brereton et al., 2007; Yang et al., 2008). MIT domains allow the Vps4-Vta1 recycling machinery to engage with its substrate, the ESCRT-III complex. They also provide the basis for a host of ESCRT-III effectors to bind to activated, openconformation ESCRT-III subunits. The effectors include microtubule severing enzymes (Connell et al., 2009; Yang et al., 2008), and deubiquitinating enzymes (Agromayor and Martin-Serrano, 2006; Ma et al., 2007; Row et al., 2007; Tsang et al., 2006), among many others (Hurley and Yang, 2008; Tsang et al., 2006). MIT domains are three-helix bundles (Scott et al., 2005b) that bind to ESCRT-III MIMs via grooves between the helices (Kieffer et al., 2008; Obita et al., 2007; Stuchell-Brereton et al., 2007; Yang et al., 2008) (Fig. 10). The MIT domain family is remarkably divergent in sequence space. Some MIT domains are so divergent in sequence that they have only been identified as such by three-dimensional structure determination (Xiao et al., 2008) or by exceptionally sensitive bioinformatics analyses (Tsang et al., 2006).

The MIT domain of Vps4 binds the helical MIM1 motif between helices $\alpha 2$ and $\alpha 3$ (Obita et al., 2007; Stuchell-Brereton et al., 2007) and the extended (non-helical) MIM2 motif between

 α 1 and α 3 (Kieffer et al., 2008). The microtubule-severing enzyme spastin contains an Nterminal MIT domain that binds only to DID2B and IST1 (Connell et al., 2009; Yang et al., 2008). DID2B and IST1 contain MIM1 motifs that have an additional two turns of helix Nterminal to the canonical MIM1. These two turns contain several key small amino acid residues that are sterically compatible with a constricted binding site on the spastin MIT domain. The extended MIM1 binding site on spastin-MIT is located between helices α 1 and α 3, like the Vps4-MIT MIM2 binding site. These differences show how hard it will be to predict the recognition specificity of MIT domains from sequence-gazing. Despite their small size in a structural sense, the substrate binding repertoire of the MIT domain family is remarkably rich. The large number of MIT domain-containing effector proteins in mammalian proteome explains, in part, the expansion of the number of ESCRT-III isoforms in mammalian cells. In terms of the details, most of the MIT-MIM recognition code, and its functional significance, remains to be elucidated.

Bro1 domains

Bro1 domains have several parallels to MIT domains. Structurally, they consist mainly of a tetratricopeptide repeat (TPR) -like helical solenoid (Kim et al., 2005), although the Bro1 domain is far larger than the MIT domain, at ~370 residues vs. ~90 residues for the MIT domain (Fig. 11). They both bind to C-terminal helical motifs of ESCRT-III subunits (Kim et al., 2005; McCullough et al., 2008). In contrast to the diversity in structure, sequence, and specificity of MIT domains, the Bro1 domain family is well conserved and has only one known ESCRT-III interaction partner, Snf7. A hydrophobic patch surrounding a conserved lle on the Bro1 domain (Kim et al., 2005) contacts exposed hydrophobic residues from the C-terminal helix of Snf7 (McCullough et al., 2008) (Fig. 11). Bro1 domains have a second conserved patch centered on a Tyr that protrudes from one end of the structure (Kim et al., 2005). The Tyr is phosphorylated in human cells, leading to the binding of this region to the SH2 domain of Src. Src phoshorylation of the PRD modulates interactions with endocytic and signaling proteins (Schmidt et al., 2005), but whether it modulates the ESCRT pathway is unknown. Bro1 domains occur in two proteins in yeast, Bro1 and Rim20. The only identified function of Bro1 in yeast is to cooperate with Snf7 in the recruitment and activation of the deubiquitinating enzyme Doa4 to deubiquitinate cargo prior to ILV scission. The pH-sensing pathway in yeast converges with the ESCRT pathway at Snf7, which interacts with the Bro1 domain of Rim20 (Boysen and Mitchell, 2006). The human proteome includes the Bro1 domain proteins ALIX, discussed above, and rhophilin, Brox, and HD-PTP.

CELLULAR FUNCTIONS OF THE ESCRTs

ESCRTs sort ubiquitinated plasma membrane proteins to the lysosome

The central observation from the earliest analyses of ESCRTs is that all of the complexes, with the exception of ESCRT-III, contain ubiquitin-binding domains (Alam et al., 2004; Bache et al., 2003b; Bilodeau et al., 2002; Katzmann et al., 2001; Katzmann et al., 2003; Mizuno *et al.*, 2003; Raiborg et al., 2002; Shih et al., 2002; Urbe *et al.*, 2003). This binding provided a pivotal link between genes involved in transport through MVBs and the then-emerging role of ubiquitination as the major signal for sorting to the yeast vacuole or mammalian lysosome (Hicke, 2001). While animal cells appear to have multiple mechanisms for the formation of MVBs and MVB-like structures, some independent of the ESCRTs (Theos *et al.*, 2006; Trajkovic *et al.*, 2008), the only known route for lysosomal degradation of ubiquitinated plasma membrane proteins is the ESCRT pathway.

Historically, the initial observation that eventually led to the discovery of the ESCRTs was the ligand-dependent sorting of the EGF receptor (EGFR) in MVBs. Direct confirmation of a role for the ESCRTs in sorting EGFR to MVBs came from analysis of the human ESCRT-I subunit

VP23 (a.k.a. TSG101, Table 1) (Babst *et al.*, 2000;Lu et al., 2003) and VPS28 (Bishop *et al.*, 2002), and the ESCRT-0 subunit Hrs (Bishop et al., 2002;Chin *et al.*, 2001;Lu et al., 2003). Precise criteria for ESCRT cargoes have not been defined. The sorting of the best-characterized cargoes, exemplified by EGFR, has been examined by electron microscopy in the context of multiple cell types and the knockdown and overexpression of various ESCRT components. In many other cases, the turnover of the cargo is affected by depletion or overexpression of a handful of ESCRT proteins for which reagents have been widely disseminated, included Hrs, TSG101, and VPS4 in animal cells, or by deletion of class E *VPS* genes in yeast. For an even larger number of less well-characterized putative cargoes, ubiquitin-dependent turnover of plasma membrane proteins is reasonably assumed to signify ESCRT-dependent lysosomal degradation, since no other mechanism is known for the turnover of such proteins (Lauwers *et al.*, 2010). The range of plasma membrane proteins that are confirmed ESCRT cargoes spans receptor tyrosine kinases (RTKs) such as the EGF receptor, G-protein coupled receptors (GPCRs), ion channels, cadherins, permeases, gap junction proteins, and miscellaneous classes of receptors.

PDGF receptor was shown early on to be downregulated via the ESCRT-0 complex (Takata et al., 2000). In Drosophila, mutation of the ESCRT-0 subunit Hrs impedes downregulation of the EGFR, PVR (PDGFR and VEGFR-related), and Torso RTKs (Jekely and Rorth, 2003; Lloyd et al., 2002), as well as the non-RTK receptors Notch (Herz et al., 2006; Jekely and Rorth, 2003; Thompson et al., 2005; Vaccari and Bilder, 2005), Hedgehog receptor (Jekely and Rorth, 2003), and Dpp (TGF- β -related) receptor (Jekely and Rorth, 2003). The TGF- β receptor is an ESCRT substrate in human cells (Shim et al., 2006). A role for ESCRT-0 in downregulating E-cadherin has been proposed (Toyoshima et al., 2007). The interleukin-2 receptor is downregulated by ESCRTs via the β chain of the receptor (Yamashita *et al.*, 2008). The polycystin-1 and -2 proteins form a mechanosensory receptor-channel complex, and their C. elegans homologs are downregulated by ESCRT-0 (Hu et al., 2007). The GABA (B) receptor is another example of a channel that has been reported to be downregulated via ESCRTs (Kantamneni et al., 2008). The δ-opiod receptor (Hislop et al., 2004), calcitonin-like receptor (Hasdemir et al., 2007), protease-activated receptor-2 (Hasdemir et al., 2007), and chemokine (C-X-C motif) receptor-4 (CXCR4) (Marchese et al., 2003) are examples of human GPCRs that are downregulated through the ESCRT pathway. Transporters, such as ferroportin, are downregulated via the ESCRTs (De Domenico et al., 2007). Virally ubiquitinated class I MHC is degraded via the ESCRTs (Hewitt et al., 2002). The gap junction protein connexin-43 undergoes regulated turnover that is ESCRT-dependent (Auth et al., 2009; Worsdorfer et al., 2008). In yeast, the GPCRs Ste2 (Odorizzi et al., 1998) and Ste3 (Piper et al., 1995) are substrates of the ESCRT pathway. The yeast general amino acid permease Gap1 is an ESCRT substrate (Nikko et al., 2003; Rubio-Texeira and Kaiser, 2006). The preceding are examples where a direct dependence of trafficking on ESCRT proteins has been shown, but the actual repertoire of ESCRT substrates is almost certainly far greater. Many other yeast proteins are sorted to the vacuole in a polyubiquitination-dependent manner, and are therefore presumed to be ESCRT substrates (Lauwers et al., 2010). In plants, degradation of the PIN proteins and AUX1, which are plasma membrane proteins involved in transport of the hormone auxin are regulated via the ESCRTs (Spitzer et al., 2009).

Arrestin-like proteins have emerged as major players in selecting cell surface receptors for ubiquitination (Lin *et al.*, 2008), but their roles might go beyond this. The fungal PalF/Rim8 arrestin-like protein binds directly to the Vps23 subunit of ESCRT-I through an SXP motif that mimics the mode of binding of ESCRT-0 to ESCRT-I (Herrador *et al.*, 2010). Other arrestin-like proteins, Art1 and Art2, target ubiquitin ligases to plasma membrane receptors destined for downregulation. This suggests that PalF/Rim8 might be an adaptor that brings the heptahelical receptor PalH/Rim21 into the MVB pathway (Lin et al., 2008).

ESCRT sort resident vacuolar and lysosomal proteins from the Golgi

The ESCRTs are involved in the sorting of newly synthesized degradative enzymes to the lysosome/vacuole. The mannose 6-phosphate receptors (MPRs) and their yeast counterpart Vps10 are not themselves ESCRT substrates and do not travel to the lysosome. These receptors deliver their cargo to the limiting membrane of the endosome, where their cargoes dissociate in the lumen and the receptors are recycled to the Golgi. MPR and Vps10 cargoes, notably yeast carboxypeptidase Y (Prc1), are missorted in ESCRT mutant cells, and Prc1 secretion is a convenient diagnostic for ESCRT dysfunction in yeast (Marcusson et al., 1994; Raymond et al., 1992). MPR cargo mis-sorting comes about because the MPR and ESCRT pathways converge at the MVB, and defects in the ESCRT pathway inhibit the maturation of MVBs and their fusion with the vacuole/lysosome. Carboxypeptidase S (Cps1) is another yeast vacuolar hydrolase that depends on the ESCRTs for its proper localization (Odorizzi et al., 1998). In contrast, Cps1 is targeted by ubiquitination, which leads to its direct interaction with the ESCRTs. Similar to Cps1, the polyphosphatase Phm5 and the haem oxygenase Hmx1 appear to be ESCRT substrates based on their ubiquitination-dependent sorting to the vacuole (Reggiori and Pelham, 2001). The vacuolar protein Sna3 (McNatt et al., 2006; Oestreich et al., 2006a; Reggiori and Pelham, 2001; Stawiecka-Mirota et al., 2007; Watson and Bonifacino, 2007) is an ESCRT substrate of unknown function, which by several accounts is ubiquitinindependent. However, the most recent data suggests that it traffics to the vacuole in a canonical ubiquitin-dependent manner (Stawiecka-Mirota et al., 2007). The notion that Sna3 is an active participant in MVB biogenesis, as opposed to being a normal cargo, has been suggested (Piper and Katzmann, 2007). In human cells, lysosomal sorting of the resident protease cathepsin D depends on ESCRT function (Babst et al., 2000).

ESCRTs are required for budding of most membrane enveloped viruses

Most, though not all, membrane enveloped viruses make use of the host ESCRT machinery to facilitate their budding from cells (Bieniasz, 2009; Carlton and Martin-Serrano, 2009; Chen and Lamb, 2008; Demirov and Freed, 2004; Morita and Sundquist, 2004; Usami et al., 2009). Nearly all viruses that make use of this pathway encode proteins that contain one or more socalled late domains (Freed, 2002), short peptide sequences that interact with the ESCRTs or ESCRT-associated ubiquitin ligases. The nuances of ESCRT function may vary somewhat for different viruses. Bearing in mind this variation, a broad working model for the main role of ESCRTs in viral budding posits that 1) viral buds are formed at the membrane by the assembly of viral proteins and do not require the ESCRTs, 2) one or more late domains in a viral protein recruit one or more of ESCRT-I, ALIX, and/or a WW-domain containing ubiquitin ligase, 3) ESCRT-I, ALIX, and/or the ubiquitin ligase initiate a pathway that leads to the assembly of ESCRT-III at the narrow neck connecting the viral bud to the host cell membrane, 4) ESCRT-III constricts the bud neck, leading to its severing, and 5) VPS4 recycles the ESCRT-III subunits, allowing for further rounds of budding. The three classes of late domain involved in ESCRT recruitment are of the form PPXY, P(S/T)XP, and YPX_nL. Despite some similarities in that all contain Pro residues and all are short, the cellular partners for each of these late domains are different.

PPXY late domains

PPXY sequences bind to WW domains (Kay *et al.*, 2000). WW domains are widespread in the human proteome, and bind to a range of PPXY sequences in human proteins as part of their normal function. The WW domain proteins most important in viral budding are a subset of HECT domain ubiquitin ligases (Martin-Serrano *et al.*, 2005). Knockdown experiments indicate that the presence of these ligases is required for the ESCRT-mediated release of PPXY-motif viruses. Despite considerable efforts, direct binding between one of the above-mentioned ubiquitin ligases and ESCRT complexes has not been demonstrated. The activity of the HECT

domain is required for budding via these ligases (Martin-Serrano et al., 2005), so it is possible that ubiquitination itself is enough to recruit ESCRT complexes to the sites of PPXY-directed virus budding. The ESCRT-I dependence of Mason-Pfizer monkey virus, which has a PPXY motif but not P(S/T)AP or YPX_nL , suggests that ESCRT-I could be targeted to budding sites by the ubiquitination of viral proteins (Chung *et al.*, 2008). Overexpression of HECT domain ubiquitin ligases can rescue budding of HIV-1 and Moloney murine leukemia virus mutants lacking late domains (Chung et al., 2008; Jadwin *et al.*; Usami *et al.*, 2008), consistent with this concept. Direct fusion of ubiquitin to the C-terminus of a retroviral Gag protein lacking any late domain is sufficient to direct efficient virus release (Joshi *et al.*, 2008). On the other hand, the PPXY-dependent budding of prototypic foamy virus (PFV) can occur without ubiquitination of viral proteins (Zhadina *et al.*, 2007).

P(S/T)AP late domains

The P(S/T)AP motif has been intensively studied because interference with this motif blocks the budding of the HIV-1. The P(S/T)AP motif binds directly to the UEV domain of the VPS23 subunit of ESCRT-I. This is the same domain the binds to ubiquitin, however, the two sites are non-overlapping, and ubiquitin binding slightly enhances P(S/T)AP motif binding to the UEV domain. Part of the normal function of the VPS23 UEV domain is to bind to P(S/T)AP motifs in host proteins, of which the archetype is the Hrs subunit of ESCRT-0 (Lu et al., 2003; Pornillos et al., 2003). In contrast to humans, in yeast the Ala at the third position in the motif is not required. Other human P(S/T)AP proteins that interact with this site include the ubiquitin ligase Tal (Amit et al., 2004), the GGA and Tom-like protein trafficking adaptors (Puertollano, 2005; Puertollano and Bonifacino, 2004), and ALIX, as described above. The structure of the PTAP motif of HIV-1 Gag in complex with the VPS23 UEV domain has been determined in solution by NMR and shows that the PTAP peptide binds in a groove between two loops and a small C-terminal β -sheet. (Pornillos *et al.*, 2002). Hydrophobic contacts are made between the core PTAP residues (Fig. 12) with Tyr63, Tyr69, and Met95 (Pornillos et al., 2002). The HIV-1 PTAP motif peptide binds to the UEV domain with a higher affinity than host protein motifs such as that of Hrs, 20 μ M vs. 150 μ M (Pornillos et al., 2003). The HIV-1 sequence contains an additional Pro after the canonical tetrapeptide motif that contributes to this higher affinity. Library screens of peptide interactors with the human VPS23 UEV domain have shown that conservation of the Ala and the second Pro is critical; that Ser or Thr are equally favored at the second position; and that the first Pro is favored but not absolutely required for binding (Schlundt et al., 2009).

The biggest mystery surrounding P(S/T)AP-mediated budding is how ESCRT-III proteins are recruited downstream of ESCRT-I. ESCRT-I has been reported to bind VPS20 (Pineda-Molina et al., 2006) the complex also binds to IST1 via the VPS37 subunit (Bajorek et al., 2009a). However, neither VPS20 nor IST1 have been shown to be required for HIV-1 budding. ESCRT-I binds to ALIX, again via its UEV domain, which in turn binds to SNF7 subunits of ESCRT-III that are critical for HIV-1 budding (Fisher et al., 2007; Usami et al., 2007). The ESCRT-I-ALIX interaction seems insufficient to explain how ESCRT-III is recruited, however, since knockdown of ALIX produces a much more modest effect on the efficiency of wild-type HIV-1 budding than knockdown of ESCRT-I or mutation of the HIV-1 PTAP motif. Knockdown of ESCRT-II also fails to impact HIV-1 budding efficiency (Langelier et al., 2006). The effects of double knockdowns of ALIX and ESCRT-II subunits on HIV-1 budding have not been reported, however, and redundancy between these two factors (or additional factors such as Brox (Popov et al., 2009)) cannot be ruled out. Other possibilities include that there may be direct interactions between ESCRT-I and ESCRT-III that have as yet evaded detection, or that as yet unidentified bridging partners are involved. There is considerable interest in the possibility of therapeutic interference with the HIV-1 PTAP/ESCRT-I interaction, and more insight into ESCRT-I-III connectivity is urgently needed.

YPX_nL late domains

The YPX_nL class of late domains is less widespread than the other two, but its mechanism of action is arguably the best understood. YPX_nL late domains bind to one arm of the V domain of ALIX (Fisher et al., 2007; Lee et al., 2007b; Munshi *et al.*, 2007) with ~5 μ M affinity. YPX_nL peptides derived from different viruses contain n =1 or 3 residues. The motif from HIV-1 has n=3, while that of equine infectious anemia virus (EIAV) has n=1. For n=3, the additional two residues fold into a helical conformation (Fig. 12), which is not present for n= 1 (Zhai et al., 2008). This allows the flanking Tyr and Leu residues to bind to the same sites on the V domain. There has been relatively little characterization of host YPX_nL motif proteins. YPX_nL-based late domains provide an elegant mechanism to support the release of viral buds because the Bro1 domain of ALIX directly binds to the SNF7 subunit of ESCRT-III. The SNF7 subunit is thought to be the most critical for membrane scission by analogy to the yeast proteins (Wollert *et al.*, 2009b). The SNF7-binding residues on the Bro1 domain are required for ALIX to support viral budding (Fisher et al., 2007; Usami et al., 2007).

Non-late domain recruitment of ESCRTs by viruses

The HIV-1 nucleocapsid (NC) is part of the Gag open reading frame, and primarily responsible for RNA encapsidation. NC has a second function in binding to the Bro1 domain of ALIX and contributes to recruiting the ESCRT machinery for budding (Dussupt *et al.*, 2009; Popov *et al.*, 2008). This property is not limited to ALIX but is found among three other Bro1 domain-containing proteins, rhophilin, HD-PTP, and Brox (Popov et al., 2009). The multiplicity of Bro1 domain proteins contributes redundancy to this mode of ESCRT recruitment, and may thus have masked the importance of ALIX in knockdown analyses. Bro1 domain-mediated recruitment of SNF7 now appears to be more central to HIV-1 budding than previously appreciated.

ESCRTs are required for the membrane abscission step in cytokinesis

ESCRTs localize to midbodies, the structure connecting two daughter cells just prior to the completion of cell division, as first visualized in studies of human VPS23 (Xie *et al.*, 1998). ESCRT mutants in *Arabidopsis* and *S. pombe* have impairments in cell division (Jin *et al.*, 2005; Spitzer *et al.*, 2006). The term "membrane abscission" refers to the cleavage of the narrow membrane neck connecting the two daughter cells. This is the final step in cytokinesis. These observations were developed into a mechanism for the role of the ESCRT machinery in the membrane abscission step in cytokinesis in 2007 by (Carlton and Martin-Serrano, 2007). This may be the most ancient role for the ESCRTs, as it is preserved even in a subset of the Archaea (Lindas et al., 2008; Samson et al., 2008). Curiously, however, ESCRT genes have not been reported as cell division mutants in yeast.

Localization of ESCRTs to the midbody requires the presence of the centrosomal and midbody protein CEP55 (Carlton and Martin-Serrano, 2007; Morita et al., 2007b). CEP55 is a dimeric coiled coil protein that is recruited to midbodies and centrosomes downstream of the microtubule-associated protein MKLP1. The central portion of CEP55 comprises an unconventional coiled coil in which charged and bulky groups replaced the normal small hydrophobic residues at the core a and d position. This leads to local asymmetry in the dimer and the pushing apart of the two coils to create a single binding site for GPPX₃Y motifs of ESCRT proteins (Fig. 13) (Lee et al., 2008). The UEV-stalk linker region of VPS23 and the PRD of ALIX both contain GPPX₃Y motifs that bind to CEP55 with 1 µM affinity (Lee et al., 2008). In a working model for cytokinetic membrane abscission by the ESCRTs, CEP55 recruits ESCRT-I and ALIX. ESCRT-I (directly or indirectly) and ALIX (directly via its Bro1 domain) recruit ESCRT-III, in particular, SNF7 (Carlton *et al.*, 2008). Assembly of ESCRT-III leads to cleavage of the membrane neck by the same mechanism as in MVB biogenesis and detachment of viral buds.

Membrane trafficking and ubiquitination pathways have multiple roles in cytokinesis, and an important challenge is to understand the division of labor between them. A number of factors from the secretory pathway, including the exocyst complex, are required for cytokinesis (Gromley *et al.*, 2005). Components of the recycling endosome pathway are also required (Fielding *et al.*, 2005; Prekeris and Gould, 2008). In models of membrane abscission predating the discovery that ESCRTs were involved, it was postulated that fusion of secretory and/or recycling vesicles could lead to membrane abscission. It now appears that these pathways are individually necessary but not sufficient for membrane abscission. The ubiquitin ligase BRUCE is required for cytokinesis (Pohl and Jentsch, 2008). It is tempting to speculate that BRUCE-dependent ubiquitination could be important for the recruitment and activation of the ESCRTs in cytokinesis, but direct evidence is lacking. As in the endosome and MVB pathways, the lipid PI(3)P is important for ESCRT function in cytokinesis. A FYVE domain containing, kinesin-binding centrosomal protein, FYVE-CENT (Sagona *et al.*, 2010) appears to link these molecules and plays a key role in delivering centrosomes to the site of midbody formation.

ESCRTs are required for autophagy

Macroautophagy, or autophagy for short, is the process whereby cells adapt to starvation by engulfing portions of their own cytosol for degradation, so as to replenish the pool of biosynthetic precursor molecules (Nakatogawa *et al.*, 2009). Beyond the starvation response, autophagy has a central role in signal downregulation, lipid catabolism, and degradation of damaged organelles, including peroxisomes and mitochondria (Levine and Kroemer, 2008). The autophagy pathway, in brief, consists of the formation of a preautophagosomal structure (PAS), the growth of an isolation membrane, the engulfment of cytosol within the isolation membrane (or phagophore), the closure of the phagophore into a double membraned autophagosome that is topologically equivalent to an MVB with a single ILV, and the fusion of the autophagosome with the lysosome/vacuole (Nakatogawa et al., 2009). Like the MVB pathway, the autophagy pathway is initiated in conjunction with the synthesis of PI(3)P by class III PI 3-kinase (Nakatogawa et al., 2009). In another parallel to the MVB pathway, ubiquitination can serve to target proteins into autophagosomes (Kirkin *et al.*, 2009; Pankiv *et al.*, 2007). Finally, the closure of the phagophore neck to form the autophagosome is topologically equivalent to the scission of endosomal bud necks to form ILVs.

Over the past three years it has become clear that the ESCRT machinery is required for autophagy in human cells (Rusten and Simonsen, 2008; Rusten and Stenmark, 2009; Rusten et al., 2007). Stenmark and colleagues have made an excellent summary of the evidence that ESCRTs are involved in autophagy and the possible mechanisms that could be at work (Rusten and Stenmark, 2009). Most of the evidence is derived from EM studies showing autophagosome accumulation when ESCRT genes are silenced or knocked out in HeLa cells (Filimonenko et al., 2007), C. elegans (Roudier et al., 2005), Drosophila (Rusten et al., 2007), and mouse neurons (Lee et al., 2007a). Several possible mechanisms have been put forward (Rusten and Stenmark, 2009). First, in a parallel to the other roles for ESCRTs, ESCRT-III might be required to cleave the membrane neck of the phagophore to yield the closed double-membraned autophagosome. This is a difficult model to test directly as this neck is less amenable to visualization in vivo than the cytokinetic or viral bud necks. Reconstitution such as carried out for MVB biogenesis seems a still remote prospect. Second, ESCRT dysfunction could act indirectly by triggering signals that inhibit autophagy. Third, it might be important to have a pool of ESCRT-generated MVBs available for fusion with autophagosomes. In a variation on this model, the role of ESCRTs may be indirect in that they are required for the biogenesis of the lysosomes with which autophagosomes ultimately fuse. These possibilities are not mutually exclusive.

Other functions of the ESCRTs

Several others functions have been reported for ESCRTs, which may or may not ultimately prove to be connected to their well-characterized membrane remodeling and cargo sorting activities. In Drosophila, ESCRT-II, via its GLUE domain, binds to bicoid mRNA and establishes the mRNA gradient responsible for establishing polarity in the embryo (Irion and St Johnston, 2007). It is hard to rationalize why membrane trafficking machinery would be involved in setting up an mRNA gradient. One hypothesis is that the mRNA uses the microtubule-based transport of MVBs to "hitchhike" towards the minus end of microtubules (Piper and Luzio, 2007). The ESCRT-I subunit VPS23 was shown very recently to be required for the establishment of the cSMAC in the immunological synapse (Vardhana et al., 2010). It is not yet clear if cSMAC organization involves the membrane budding activity of ESCRT-I or not. ESCRT-II subunits in human cells were originally isolated and characterized as binding partners for the ELL proteins, elongation factors of RNA polymerase II (Kamura et al., 2001), but the mechanistic implications of this have not been elucidated. The human ESCRT-III subunit DID2A was first cloned and named "CHMP1" for Chromatin modifying protein 1 for a putative role in gene silencing, which has to date been neither refuted nor further elucidated.

ESCRT DYSFUNCTION AND NON-VIRAL DISEASE

ESCRTs and cancer

The human ESCRT-I subunit VPS23 was first isolated as a results of an antisense RNA screen for factors whose disruption promoted tumorigenesis, and named "tumor susceptibility gene 101" (Li and Cohen, 1996). Given what we now know about the role of ESCRT-I in downregulating proliferative receptor signaling, it is tempting to speculate that this could be connected to its tumor susceptibility phenotype. ESCRT functions in cytokinesis and autophagy could conceivably mediate such a phenotype, as well. Following some early reports that proved erroneous (Li *et al.*, 1998), there have been no confirmed reports of mutations in *TSG101* human tumors. A separate and compelling line of evidence mechanistically linking ESCRT dysfunction to cancer comes from observations in *Drosophila*. Mutation of *Drosophila* VPS23 leads to tumors (Moberg *et al.*, 2005). Mutations in the ESCRT-II subunit VPS25 also leads to cell overproliferation in *Drosophila* and induces tumor formation when apoptosis is blocked (Herz et al., 2006; Thompson et al., 2005; Vaccari and Bilder, 2005). The overproliferation is due at least in part to excess signaling by Notch, consistent with the model that ESCRTs antagonize proliferative signaling by downregulating proliferative receptors.

ESCRTs and neurological diseases

Two main lines of evidence connect the ESCRTs to neurological diseases. First, autosomal dominant mutations of the gene for the ESCRT-III subunit VPS2B lead to the neurodegenerative disorder frontotemporal dementia-3 (FTD3) (Lee et al., 2007a). The mechanism of pathogenesis is thought to be linked to the role of VPS2B as a factor required for autophagy. Second, the hereditary spastic paraplegias (HSPs) are a group of inherited disorders (SPG1-46) characterized by a length dependent axonopathy of corticospinal motor neurons. Two of the proteins encoded by these genes, the SPG4 protein spastin (Connell et al., 2009; Yang et al., 2008) and the SPG15 protein FYVE-CENT (Sagona et al., 2010), have functional interactions with ESCRT-III proteins. Impairment of these interactions leads to cytokinesis defects in cell culture. It is not clear how cytokinesis defects would lead selectively to impaired neuronal development. The midbody structure that coordinates cytokinetic abscission is closely linked to the centrosome, and these two structures have an overlapping protein composition. Indeed, ESCRTs have been visualized at both midbodies and centrosomes (Sagona et al., 2010). It is unclear to what extent neurological pathogenesis in the HSPs is due

to the ESCRT-SPG protein connection. The observation of ESCRT-associated protein localization to centrosomes (Sagona et al., 2010) suggests a potential link to axon development.

CONCLUSION

The structure, function, and interactions of the central ESCRT machinery- that is, the portions conserved from yeast to humans and preserved between different human subunit isoforms- are now relatively well understood. The reason for the diversity of complexes in human cells-with in principle twelve ESCRT-III and sixteen ESCRT-I combinations possible- is only starting to be explored. Recent progress in visualizing ESCRT assemblies on membranes, reconstituting the ESCRT reaction in vitro, and analyzing the budding and scission mechanism computationally, have moved the field from hand-waving to a rough pictorial outline of the reaction with some biophysical understanding. The stage is now set for a deeper quantitative biophysical analysis of the mechanism. Connections between the ESCRTs and other major cellular machineries are only beginning to be understood. We have an initial view of how the ESCRTs could be linked the early endocytic apparatus and the cytoskeleton, but connections to other areas, such as the endosome-vacuole fusion machinery, need much more study. ESCRT-virus interaction may be appealing drug targets, but concerns about interference with normal physiological functions must be overcome. ESCRT dysfunction is linked to neurological diseases, but the mechanistic basis for the connection is largely a mystery. It is hoped that this review will serve as a one-stop sourcebook of information for newcomers who may go on to answer some of these questions.

Acknowledgments

I thank E. Freed and C. Blackstone for comments on the manuscript and members of my group and colleagues in the ESCRT field for many stimulating discussions.

Declaration of interests

Research in my laboratory is supported by the NIDDK and IATAP programs of the NIH intramural research program.

REFERENCES

- Agromayor M, Martin-Serrano J. Interaction of AMSH with ESCRT-III and deubiquitination of endosomal cargo. J Biol Chem 2006;281:23083–23091. [PubMed: 16760479]
- Alam SL, Langelier C, Whitby FG, Koirala S, Robinson H, Hill CP, Sundquist WI. Structural basis for ubiquitin recognition by the human ESCRT-II EAP45 GLUE domain. Nat Struct Mol Biol 2006;13:1029–1030. [PubMed: 17057716]
- Alam SL, Sun J, Payne M, Welch BD, Black BK, Davis DR, Meyer HH, Emr SD, Sundquist WI. Ubiquitin interactions of NZF zinc fingers. Embo J 2004;23:1411–1421. [PubMed: 15029239]
- Alam SL, Sundquist WI. Structural biology ESCRT service. Nature 2007;447:921–922. [PubMed: 17581573]
- Amit I, Yakir L, Katz M, Zwang Y, Marmor MD, Citri A, Shtiegman K, Alroy I, Tuvia S, Reiss Y, et al. Tal, a Tsg101-specific E3 ubiquitin ligase, regulates receptor endocytosis and retrovirus budding. Genes Devel 2004;18:1737–1752. [PubMed: 15256501]
- Asao H, Sasaki Y, Arita T, Tanaka N, Endo K, Kasai H, Takeshita T, Endo Y, Fujita T, Sugamura K. Hrs is associated with STAM, a signal-transducing adaptor molecule - Its suppressive effect on cytokine-induced cell growth. J Biol Chem 1997;272:32785–32791. [PubMed: 9407053]
- Audhya A, McLeod IX, Yates JR, Oegama K. MVB-12, a fourth subunit of metazoan ESCRT-I, functions in receptor downregulation. PLOS ONE 2007;2:e956. [PubMed: 17895996]
- Auth T, Schluter S, Urschel S, Kussmann P, Sonntag S, Hoher T, Kreuzberg MM, Dobrowolski R, Willecke K. The TSG101 protein binds to connexins and is involved in connexin degradation. Exp Cell Res 2009;315:1053–1062. [PubMed: 19210987]

- Azmi I, Davies B, Dimaano C, Payne J, Eckert D, Babst M, Katzmann DJ. Recycling of ESCRTs by the AAA-ATPase Vps4 is regulated by a conserved VSL region in Vta 1. J Cell Biol 2006;172:705–717. [PubMed: 16505166]
- Azmi IF, Davies BA, Xiao J, Babst M, Xu Z, Katzmann DJ. ESCRT-III family members stimulate Vps4 ATPase activity directly or via Vta1. Dev Cell 2008;14:50–61. [PubMed: 18194652]
- Babst M. A Protein's Final ESCRT. Traffic 2005;6:2-9. [PubMed: 15569240]
- Babst M, Katzmann DJ, Estepa-Sabal EJ, Meerloo T, Emr SD. ESCRT-III: An endosome-associated heterooligomeric protein complex required for MVB sorting. Dev Cell 2002a;3:271–282. [PubMed: 12194857]
- Babst M, Katzmann DJ, Snyder WB, Wendland B, Emr SD. Endosome-associated complex, ESCRT-II, recruits transport machinery for protein sorting at the multivesicular body. Dev Cell 2002b;3:283– 289. [PubMed: 12194858]
- Babst M, Odorizzi G, Estepa EJ, Emr SD. Mammalian tumor susceptibility gene 101 (TSG101) and the yeast homologue, Vps23p, both function in late endosomal trafficking. Traffic 2000;1:248–258. [PubMed: 11208108]
- Bache KG, Brech A, Mehlum A, Stenmark H. Hrs regulates multivesicular body formation via ESCRT recruitment to endosomes. J Cell Biol 2003a;162:435–442. [PubMed: 12900395]
- Bache KG, Raiborg C, Mehlum A, Stenmark H. STAM and Hrs are subunits of a multivalent ubiquitinbinding complex on early endosomes. J Biol Chem 2003b;278:12513–12521. [PubMed: 12551915]
- Bajorek M, Morita E, Skalicky JJ, Morham SG, Babst M, Sundquist WI. Biochemical analyses of human IST1 and its function in cytokinesis. Mol Biol Cell 2009a;20:1360–1373. [PubMed: 19129479]
- Bajorek M, Schubert HL, McCullough J, Langelier C, Eckert DM, Stubblefield WMB, Uter NT, Myszka DG, Hill CP, Sundquist WI. Structural basis for ESCRT-III protein autoinhibition. Nat Struct Mol Biol 2009b;16:754–U795. [PubMed: 19525971]
- Bieniasz PD. Late budding domains and host proteins in enveloped virus release. Virology 2006;344:55–63. [PubMed: 16364736]
- Bieniasz PD. The Cell Biology of HIV-1 Virion Genesis. Cell Host Micr 2009;5:550–558.
- Bilodeau PS, Urbanowski JL, Winistorfer SC, Piper RC. The Vps27p-Hse1p complex binds ubiquitin and mediates endosomal protein sorting. Nat Cell Biol 2002;4:534–539. [PubMed: 12055639]
- Bilodeau PS, Winistorfer SC, Kearney WR, Robertson AD, Piper RC. Vps27-Hse1 and ESCRT-I complexes cooperate to increase efficiency of sorting ubiquitinated proteins at the endosome. J Cell Biol 2003;163:237–243. [PubMed: 14581452]
- Bishop N, Horman A, Woodman P. Mammalian class E vps proteins recognize ubiquitin and act in the removal of endosomal protein-ubiquitin conjugates. J Cell Biol 2002;157:91–101. [PubMed: 11916981]
- Bowers K, Lottridge J, Helliwell SB, Goldthwaite LM, Luzio JP, Stevens TH. Protein-protein interactions of ESCRT complexes in the yeast Saccharomyces cerevisiae. Traffic 2004;5:194–210. [PubMed: 15086794]
- Boysen JH, Mitchell AP. Control of Bro1-domain protein Rim20 localization by external pH, ESCRT machinery, and the Saccharomyces cerevisiae Rim101 pathway. Mol Biol Cell 2006;17:1344–1353. [PubMed: 16407402]
- Bryant NJ, Stevens TH. Vacuole biogenesis in Saccharomyces cerevisiae: Protein transport pathways to the yeast vacuole. Microbiol Mol Biol Rev 1998;62:230. [PubMed: 9529893]
- Burd CG, Emr SD. Phosphatidylinositol(3)-phosphate signaling mediated by specific binding to RING FYVE domains. Mol Cell 1998;2:157–162. [PubMed: 9702203]
- Carlton JG, Agromayor M, Martin-Serrano J. Differential requirements for Alix and ESCRT-III in cytokinesis and HIV-1 release. Proc Natl Acad Sci U S A 2008;105:10541–10546. [PubMed: 18641129]
- Carlton JG, Martin-Serrano J. Parallels between cytokinesis and retroviral budding: a role for the ESCRT machinery. Science 2007;316:1908–1912. [PubMed: 17556548]
- Carlton JG, Martin-Serrano J. The ESCRT machinery: new functions in viral and cellular biology. Biochem Soc Trans 2009;37:195–199. [PubMed: 19143630]

- Carter CA, Ehrlich LS. Cell Biology of HIV-1 Infection of Macrophages. Annu Rev Microbiol 2008;62:425–443. [PubMed: 18785842]
- Chen BJ, Lamb RA. Mechanisms for enveloped virus budding: Can some viruses do without an ESCRT? Virology 2008;372:221–232. [PubMed: 18063004]
- Chin LS, Raynor MC, Wei XL, Chen HQ, Li L. Hrs interacts with sorting nexin 1 and regulates degradation of epidermal growth factor receptor. J Biol Chem 2001;276:7069–7078. [PubMed: 11110793]
- Chu T, Sun J, Saksena S, Emr SD. New component of ESCRT-I regulates endosomal sorting complex assembly. J Cell Biol 2006;175:815–823. [PubMed: 17145965]
- Chung HY, Morita E, von Schwedler U, Muller B, Krausslich HG, Sundquist WI. NEDD4L overexpression rescues the release and infectivity of human immunodeficiency virus type 1 constructs lacking PTAP and YPXL late domains. J Virol 2008;82:4884–4897. [PubMed: 18321968]
- Connell JW, Lindon C, Luzio JP, Reid E. Spastin couples microtubule severing to membrane traffic in completion of cytokinesis and secretion. Traffic 2009;10:42–56. [PubMed: 19000169]
- Curtiss M, Jones C, Babst M. Efficient cargo sorting by ESCRT-I and the subsequent release of ESCRT-I from multivesicular bodies requires the subunit Mvb12. Mol Biol Cell 2006;18:636–645. [PubMed: 17135292]
- De Domenico I, Ward DM, Langelier C, Vaughn MB, Nemeth E, Sundquist WI, Ganz T, Musci G, Kaplan J. The molecular mechanism of hepcidin-mediated ferroportin down-regulation. Mol Biol Cell 2007;18:2569–2578. [PubMed: 17475779]
- de Souza RF, Aravind L. UMA and MABP domains throw light on receptor endocytosis and selection of endosomal cargoes. Bioinformatics. 2010 epub ahead of printe.
- Demirov DG, Freed EO. Retrovirus budding. Virus Research 2004;106:87–102. [PubMed: 15567490]
- Dikic I, Wakatsuki S, Walters KJ. Ubiquitin-binding domains from structures to functions. Nat Rev Mol Cell Biol 2009;10:659–671. [PubMed: 19773779]
- Dimaano C, Jones CB, Hanono A, Curtiss M, Babst M. Ist1 regulates Vps4 localization and assembly. Mol Biol Cell 2008;19:465–474. [PubMed: 18032582]
- Dussupt V, Javid MP, Abou-Jaoude G, Jadwin JA, de La Cruz J, Nagashima K, Bouamr F. The Nucleocapsid Region of HIV-1 Gag Cooperates with the PTAP and LYPXnL Late Domains to Recruit the Cellular Machinery Necessary for Viral Budding. Plos Pathogens 2009:5.
- Fabrikant G, Lata S, Riches JD, Briggs JAG, Weissenhorn W, Kozlov MM. Computational model of membrane fission catalyzed by ESCRT-III. PLOS Computational Biology 2009;5:e1000575. [PubMed: 19936052]
- Fielding AB, Schonteich E, Matheson J, Wilson G, Yu XZ, Hickson GRX, Srivastava S, Baldwin SA, Prekeris R, Gould GW. Rab11-FIP3 and FIP4 interact with Arf6 and the Exocyst to control membrane traffic in cytokinesis. Embo J 2005;24:3389–3399. [PubMed: 16148947]
- Filimonenko M, Stuffers S, Raiborg C, Yamamoto A, Malerod L, Fisher EMC, Isaacs A, Brech A, Stenmark H, Simonsen A. Functional multivesicular bodies are required for autophagic clearance of protein aggregates associated with neurodegenerative disease. J Cell Biol 2007;179:485–500. [PubMed: 17984323]
- Fisher RD, Chung HY, Zhai QT, Robinson H, Sundquist WI, Hill CP. Structural and biochemical studies of ALIX/AIP1 and its role in retrovirus budding. Cell 2007;128:841–852. [PubMed: 17350572]
- Fisher RD, Wang B, Alam SL, Higginson DS, Robinson H, Sundquist WI, Hill CP. Structure and ubiquitin binding of the ubiquitin-interacting motif. J Biol Chem 2003;278:28976–28984. [PubMed: 12750381]
- Freed EO. Viral late domains. J Virol 2002;76:4679-4687. [PubMed: 11967285]
- Freed EO, Mouland AJ. The cell biology of HIV-I and other retroviruses. Retrovirology 2006:3. [PubMed: 16409629]
- Fujii K, Hurley JH, Freed EO. Beyond Tsg101: the role of Alix in 'ESCRTing' HIV-1. Nat Rev Microbiol 2007;5:912–916. [PubMed: 17982468]
- Gaullier JM, Simonsen A, D'Arrigo A, Bremnes B, Stenmark H, Aasland R. FYVE fingers bind Ptdins (3)P. Nature 1998;394:432–433. [PubMed: 9697764]

- Ghazi-Tabatabai S, Saksena S, Short JM, Pobbati AV, Veprintsev DB, Crowther RA, Emr SD, Egelman EH, Williams RL. Structure and disassembly of filaments formed by the ESCRT-III subunit Vps24. Structure 2008;16:1345–1356. [PubMed: 18786397]
- Gill DJ, Teo H, Sun J, Perisic O, Veprintsev DB, Emr SD, Williams RL. Structural insight into the ESCRT-I/-II link and its role in MVB trafficking. Embo J 2007;26:600–612. [PubMed: 17215868]
- Gonciarz MD, Whitby FG, Eckert DM, Kieffer C, Heroux A, Sundquist WI, Hill CP. Biochemical and Structural Studies of Yeast Vps4 Oligomerization. J Mol Biol 2008;384:878–895. [PubMed: 18929572]
- Gorden P, Carpentier JL, Cohen S, Orci L. Epidermal growth factor: morphological demonstration of binding, internalization, and lysosomal association in human fibroblasts. Proc Natl Acad Sci U S A 1978;75:5025–5029. [PubMed: 311005]
- Gromley A, Yeaman C, Rosa J, Redick S, Chen CT, Mirabelle S, Guha M, Sillibourne J, Doxsey SJ. Centriolin anchoring of exocyst and SNARE complexes at the midbody is required for secretoryvesicle-mediated abscission. Cell 2005;123:75–87. [PubMed: 16213214]
- Gruenberg J, Stenmark H. The biogenesis of multivesicular endosomes. Nat Rev Mol Cell Biol 2004;5:317–323. [PubMed: 15071556]
- Haigler HT, McKanna JA, Cohen S. Direct visualization of the binding and internalization of a ferritin conjugate of epidermal growth factor in human carcinoma cells A-431. J Cell Biol 1979;81:382–395. [PubMed: 313931]
- Hanson PI, Roth R, Lin Y, Heuser JE. Plasma membrane deformation by circular arrays of ESCRT-III protein filaments. J Cell Biol 2008;180:389–402. [PubMed: 18209100]
- Hanson PI, Shim S, Merrill SA. Cell biology of the ESCRT machinery. Curr Opin Cell Biol 2009;21:568– 574. [PubMed: 19560911]
- Hartmann C, Chami M, Zachariae U, de Groot BL, Engel A, Grutter MG. Vacuolar protein sorting: Two different functional states of the AAA-ATPase Vps4p. J Mol Biol 2008;377:352–363. [PubMed: 18272179]
- Hasdemir B, Bunnett NW, Cottrell GS. Hepatocyte growth factor-regulated tyrosine kinase substrate (HRS) mediates post-endocytic trafficking of protease-activated receptor 2 and calcitonin receptorlike receptor. J Biol Chem 2007;282:29646–29657. [PubMed: 17675298]
- Herrador A, Herranz S, Lara D, Vincent O. Recruitment of the ESCRT Machinery to a Putative Seven-Transmembrane-Domain Receptor Is Mediated by an Arrestin-Related Protein. Mol Cell Biol 2010;30:897–907. [PubMed: 20028738]
- Herz HM, Chen Z, Scherr H, Lackey M, Bolduc C, Bergmann A. vps25 mosaics display non-autonomous cell survival and overgrowth, and autonomous apoptosis. Development 2006;133:1871–1880. [PubMed: 16611691]
- Hewitt EW, Duncan L, Mufti D, Baker J, Stevenson PG, Lehner PJ. Ubiquitylation of MHC class I by the K3 viral protein signals internalization and TSG101-dependent degradation. Embo J 2002;21:2418–2429. [PubMed: 12006494]
- Hicke L. A new ticket for entry into budding vesicles Ubiquitin. Cell 2001;106:527–530. [PubMed: 11551499]
- Hierro A, Sun J, Rusnak AS, Kim J, Prag G, Emr SD, Hurley JH. Structure of the ESCRT-II endosomal trafficking complex. Nature 2004;431:221–225. [PubMed: 15329733]
- Hirano S, Kawasaki M, Ura H, Kato R, Raiborg C, Stenmark H, Wakatsuki S. Double-sided ubiquitin binding of Hrs-UIM in endosomal protein sorting. Nat Struct Mol Biol 2006a;13:272–277. [PubMed: 16462748]
- Hirano S, Suzuki N, Slagsvold T, Kawasaki M, Trambaiolo D, Kato R, Stenmark H, Wakatsuki S. Structural basis of ubiquitin recognition by mammalian Eap45 GLUE domain. Nat Struct Mol Biol 2006b;13:1031–1032. [PubMed: 17057714]
- Hislop JN, Marley A, von Zastrow M. Role of mammalian vacuolar protein-sorting proteins in endocytic trafficking of a non-ubiquitinated G protein-coupled receptor to lysosomes. J Biol Chem 2004;279:22522–22531. [PubMed: 15024011]
- Hoeller D, Crosetto N, Blagoev B, Raiborg C, Tikkanen R, Wagner S, Kowanetz K, Breitling R, Mann M, Stenmark H, Dikic I. Regulation of ubiquitin-binding proteins by monoubiquitination. Nat Cell Biol 2006;8:163–U145. [PubMed: 16429130]

- Hong YH, Ahn HC, Lim J, Kim HM, Ji HY, Lee S, Kim JH, Park EY, Song HK, Lee BJ. Identification of a novel ubiquitin binding site of STAM1 VHS domain by NMR spectroscopy. FEBS Lett 2009;583:287–292. [PubMed: 19111546]
- Hu JH, Wittekind SG, Barr MM. STAM and hrs down-regulate ciliary TRP receptors. Mol Biol Cell 2007;18:3277–3289. [PubMed: 17581863]
- Huang FT, Kirkpatrick D, Jiang XJ, Gygi S, Sorkin A. Differential regulation of EGF receptor internalization and degradation by multiubiquitination within the kinase domain. Mol Cell 2006;21:737–748. [PubMed: 16543144]
- Hurley JH. ESCRT Complexes and the Biogenesis of Multivesicular Bodies. Curr Opin Cell Biol 2008;20:4–11. [PubMed: 18222686]
- Hurley JH, Emr SD. The ESCRT complexes: structure and mechanism of a membrane-trafficking network. Annu Rev Biophys Biomol Struct 2006;35:277–298. [PubMed: 16689637]
- Hurley JH, Hanson PI. Membrane budding and scission by the ESCRT complexes: It's all in the neck. Nat Rev Mol Cell Biol. 2010 in press.
- Hurley JH, Lee S, Prag G. Ubiquitin binding domains. Biochem J 2006;399:361–372. [PubMed: 17034365]
- Hurley JH, Ren XF. The circuitry of cargo flux in the ESCRT pathway. J Cell Biol 2009;185:185–187. [PubMed: 19380875]
- Hurley JH, Yang D. MIT domainia. Dev Cell 2008;14:6-8. [PubMed: 18194647]
- Im YJ, Hurley JH. Integrated structural model and membrane targeting mechanism of the human ESCRT-II complex. Dev Cell 2008;14:902–913. [PubMed: 18539118]
- Im YJ, Wollert T, Boura E, Hurley JH. Structure and Function of the ESCRT-II-III Interface in Multivesicular Body Biogenesis. Dev Cell 2009;17:234–243. [PubMed: 19686684]
- Inoue M, Kamikubo H, Kataoka M, Kato R, Yoshimori T, Wakatsuki S, Kawasaki M. Nucleotide-Dependent Conformational Changes and Assembly of the AAA ATPase SKD1/VPS4B. Traffic 2008;9:2180–2189. [PubMed: 18796009]
- Irion U, Johnston D. bicoid RNA localization requires specific binding of an endosomal sorting complex. Nature 2007;445:554–558. [PubMed: 17268469]
- Jadwin JA, Rudd V, Sette P, Challa S, Bouamr F. Late Domain-Independent Rescue of a Release-Deficient Moloney Murine Leukemia Virus by the Ubiquitin Ligase Itch. J Virol 84:704–715. [PubMed: 19864377]
- Jekely G, Rorth P. Hrs mediates downregulation of multiple signalling receptors in Drosophila. Embo Rep 2003;4:1163–1168. [PubMed: 14608370]
- Jin Y, Mancuso JJ, Uzawa S, Cronembold D, Cande WZ. The fission yeast homolog of the human transcription factor EAP30 blocks meiotic spindle pole body amplification. Dev Cell 2005;9:63–73. [PubMed: 15992541]
- Joshi A, Munshi U, Ablan SD, Nagashima K, Freed EO. Functional Replacement of a Retroviral Late Domain by Ubiquitin Fusion. Traffic 2008;9:1972–1983. [PubMed: 18817521]
- Kamura T, Burian D, Khalili H, Schmidt SL, Sato S, Liu WJ, Conrad MN, Conaway RC, Conaway JW, Shilatifard A. Cloning and characterization of ELL-associated proteins EAP45 and EAP20. a role for yeast EAP-like proteins in regulation of gene expression by glucose. J Biol Chem 2001;276:16528–16533. [PubMed: 11278625]
- Kantamneni S, Holman D, Wilkinson KA, Correa SAL, Feligioni M, Ogden S, Fraser W, Nishimune A, Henley JM. GISP binding to TSG101 increases GABA(B) receptor stability by down-regulating ESCRT-mediated lysosomal degradation. J Neurochem 2008;107:86–95. [PubMed: 18643869]
- Kato M, Miyazawa K, Kitamura N. A deubiquitinating enzyme UBPY interacts with the Src homology 3 domain of Hrs-binding protein via a novel binding motif PX(V/I)(D/N)RXXKP. J Biol Chem 2000;275:37481–37487. [PubMed: 10982817]
- Katzmann DJ, Babst M, Emr SD. Ubiquitin-dependent sorting into the multivesicular body pathway requires the function of a conserved endosomal protein sorting complex, ESCRT-I. Cell 2001;106:145–155. [PubMed: 11511343]
- Katzmann DJ, Odorizzi G, Emr SD. Receptor downregulation and multivesicular-body sorting. Nat Rev Mol Cell Biol 2002;3:893–905. [PubMed: 12461556]

- Katzmann DJ, Stefan CJ, Babst M, Emr SD. Vps27 recruits ESCRT machinery to endosomes during MVB sorting. J Cell Biol 2003;162:413–423. [PubMed: 12900393]
- Kay BK, Williamson MP, Sudol M. The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. Faseb J 2000;14:231–241. [PubMed: 10657980]
- Kieffer C, Skalicky JJ, Morita E, De Domenico I, Ward DM, Kaplan J, Sundquist WI. Two distinct modes of ESCRT-III recognition are required for VPS4 functions in lysosomal protein targeting and HIV-1 budding. Dev Cell 2008;15:62–73. [PubMed: 18606141]
- Kim J, Sitaraman S, Hierro A, Beach BM, Odorizzi G, Hurley JH. Structural basis for endosomal targeting by the Bro1 domain. Dev Cell 2005;8:937–947. [PubMed: 15935782]
- Kirkin V, McEwan DG, Novak I, Dikic I. A Role for Ubiquitin in Selective Autophagy. Mol Cell 2009;34:259–269. [PubMed: 19450525]
- Kostelansky MS, Schluter C, Tam YYC, Lee S, Ghirlando R, Beach B, Conibear E, Hurley JH. Molecular architecture and functional model of the complete yeast ESCRT-I heterotetramer. Cell 2007;129:485–498. [PubMed: 17442384]
- Kostelansky MS, Sun J, Lee S, Kim J, Ghirlando R, Hierro A, Emr SD, Hurley JH. Structural and functional organization of the ESCRT-I trafficking complex. Cell 2006;125:113–126. [PubMed: 16615894]
- Kutateladze TG, Ogburn KD, Watson WT, de Beer T, Emr SD, Burd CG, Overduin M.
 Phosphatidylinositol 3-phosphate recognition by the FYVE domain. Mol Cell 1999;3:805–811.
 [PubMed: 10394369]
- Landsberg MJ, Vajjhala PR, Rothnagel R, Munn AL, Hankamer B. Three-Dimensional Structure of AAA ATPase Vps4: Advancing Structural Insights into the Mechanisms of Endosomal Sorting and Enveloped Virus Budding. Structure 2009;17:427–437. [PubMed: 19278657]
- Langelier C, von Schwedler UK, Fisher RD, De Domenico I, White PL, Hill CP, Kaplan J, Ward D, Sundquist WI. Human ESCRT-II complex and its role in human immunodeficiency virus type 1 release. J Virol 2006;80:9465–9480. [PubMed: 16973552]
- Lata S, Roessle M, Solomons J, Jamin M, Gottlinger H, Svergun DI, Weissenhorn W. Structural basis for autoinhibition of ESCRT-III CHMP3. J Mol Biol 2008a;378:816–825.
- Lata S, Schoehn G, Jain A, Pires R, Piehler J, Gottlinger H, Weissenhorn W. Helical Structures of ESCRT-III are Disassembled by VPS4. Science 2008b;321:1354–1357. [PubMed: 18687924]
- Lauwers E, Erpapazoglou Z, Haguenauer-Tsapis R, Andre B. The ubiquitin code of yeast permease trafficking. Trends Cell Biol. 2010 epub ahead of print.
- Lauwers E, Jacob C, Andre B. K63-linked ubiquitin chains as a specific signal for protein sorting into the multivesicular body pathway. J Cell Biol 2009;185:493–502. [PubMed: 19398763]
- Lee HH, Elia N, Ghirlando R, Lippincott-Schwartz J, Hurley JH. Midbody targeting of the ESCRT machinery by a noncanonical coiled coil in CEP55. Science 2008;322:576–580. [PubMed: 18948538]
- Lee JA, Beigneux A, Ahmad ST, Young SG, Gao FB. ESCRT-III dysfunction causes autophagosome accumulation and neurodegeneration. Curr Biol 2007a;17:1561–1567. [PubMed: 17683935]
- Lee S, Joshi A, Nagashima K, Freed EO, Hurley JH. Structural basis for viral late-domain binding to Alix. Nat Struct Mol Biol 2007b;14:194–199. [PubMed: 17277784]
- Levine B, Kroemer G. Autophagy in the pathogenesis of disease. Cell 2008;132:27–42. [PubMed: 18191218]
- Li LM, Cohen SN. tsg101: A novel tumor susceptibility gene isolated by controlled homozygous functional knockout of allelic loci in mammalian cells. Cell 1996;85:319–329. [PubMed: 8616888]
- Li LM, Francke U, Cohen SN. The TSG101 tumor susceptibility gene is located in chromosome 11 band p15 and is mutated in human breast cancer. Cell 1998;88:143. 1997. 93:U17-U17. [PubMed: 9019400]
- Lin CH, MacGum JA, Chu T, Stefan CJ, Emr SD. Arrestin-Related Ubiquitin-Ligase Adaptors Regulate Endocytosis and Protein Turnover at the Cell Surface. Cell 2008;135:714–725. [PubMed: 18976803]
- Lindas AC, Karlsson EA, Lindgren MT, Ettema TJG, Bernander R. A unique cell division machinery in the Archaea. Proc Natl Acad Sci USA 2008;105:18942–18946. [PubMed: 18987308]

- Lloyd TE, Atkinson R, Wu MN, Zhou Y, Pennetta G, Bellen HJ. Hrs regulates endosome membrane invagination and tyrosine kinase receptor signaling in Drosophila. Cell 2002;108:261–269. [PubMed: 11832215]
- Lohi O, Lehto VP. VHS domain marks a group of proteins involved in endocytosis and vesicular trafficking. FEBS Lett 1998;440:255–257. [PubMed: 9872381]
- Lottridge JM, Flannery AR, Vincelli JL, Stevens TH. Vta1p and Vps46p regulate the membrane association and ATPase activity of Vps4p at the yeast multivesicular body. Proc Natl Acad Sci USA 2006;103:6202–6207. [PubMed: 16601096]
- Lu Q, Hope LWQ, Brasch M, Reinhard C, Cohen SN. TSG101 interaction with HRS mediates endosomal trafficking and receptor down-regulation. Proc Natl Acad Sci USA 2003;100:7626–7631. [PubMed: 12802020]
- Luhtala N, Odorizzi G. Bro1 coordinates deubiquitination in the multivesicular body pathway by recruiting Doa4 to endosomes. J Cell Biol. 2004
- Luzio JP, Piper SC, Bowers K, Parkinson MDJ, Lehner PJ, Bright NA. ESCRT proteins and the regulation of endocytic delivery to lysosomes. Biochem Soc Trans 2009;37:178–180. [PubMed: 19143626]
- Ma YM, Boucrot E, Villen J, Affar el B, Gygi SP, Gottlinger H, Kirchhausen T. Targeting of AMSH to endosomes is required for epidermal growth factor degradation. J Biol Chem 2007;282:9805–9812. [PubMed: 17261583]
- Marchese A, Raiborg C, Santini F, Keen JH, Stenmark H, Benovic JL. The E3 ubiquitin ligase AIP4 mediates ubiquitination and sorting of the G protein-coupled receptor CXCR4. Dev Cell 2003;5:709–722. [PubMed: 14602072]
- Marcusson EG, Horazdovsky BF, Cereghino JL, Gharakhanian E, Emr SD. The Sorting Receptor For Yeast Vacuolar Carboxypeptidase-Y Is Encoded By The Vps10 Gene. Cell 1994;77:579–586. [PubMed: 8187177]
- Martin-Serrano J, Eastman SW, Chung W, Bieniasz PD. HECT ubiquitin ligases link viral and cellular PPXY motifs to the vacuolar protein-sorting pathway. J Cell Biol 2005;168:89–101. [PubMed: 15623582]
- Matsuo H, Chevallier J, Mayran N, Le Blanc I, Ferguson C, Faure J, Blanc NS, Matile S, Dubochet J, Sadoul M, et al. Role of LBPA and Alix in multivesicular liposome formation and endosome organization. Science 2004;303:531–534. [PubMed: 14739459]
- McCullough J, Fisher RD, Whitby FG, Sundquist WI, Hill CP. ALIX-CHMP4 interactions in the human ESCRT pathway. Proc Natl Acad Sci U S A 2008;105:7687–7691. [PubMed: 18511562]
- McDonald B, Martin-Serrano J. No strings attached: the ESCRT machinery in viral budding and cytokinesis. J Cell Sci 2009;122:2167–2177. [PubMed: 19535732]
- McNatt NW, McKittrick I, West M, Odorizzi G. Direct binding to Rsp5 mediates ubiquitin-independent sorting of Sna3 via the multivesicular body pathway. Mol Biol Cell 2006;18:697–706. [PubMed: 17182850]
- Misra S, Hurley JH. Crystal structure of a phosphatidylinositol 3-phosphate- specific membrane-targeting motif, the FYVE domain of Vps27p. Cell 1999;97:657–666. [PubMed: 10367894]
- Misra S, Miller GJ, Hurley JH. Recognizing phosphatidylinositol 3-phosphate. Cell 2001;107:559–562. [PubMed: 11733055]
- Missotten M, Nichols A, Rieger K, Sadoul R. Alix, a novel mouse protein undergoing calcium-dependent interaction with the apoptosis-linked-gene 2 (ALG-2) protein. Cell Death Differentiation 1999;6:124–129.
- Mizuno E, Kawahata K, Kato M, Kitamura N, Komada M. STAM proteins bind ubiquitinated proteins on the early endosome via the VHS domain and ubiquitin-interacting motif. Mol Biol Cell 2003;14:3675–3689. [PubMed: 12972556]
- Moberg KH, Schelble S, Burdick SK, Hariharan IK. Mutations in erupted, the Drosophila ortholog of mammalian tumor susceptibility gene 101, elicit non-cell-autonomous overgrowth. Dev Cell 2005;9:699–710. [PubMed: 16256744]
- Morita E, Sandrin V, Alam SL, Eckert DM, Gygi SP, Sundquist WI. Identification of human MVB12 proteins as ESCRT-I subunits that function in HIV budding. Cell Host Microbe 2007a;2:41–53. [PubMed: 18005716]

- Morita E, Sandrin V, Chung HY, Morham SG, Gygi S, Rodesch CK, Sundquist WI. Human ESCRT and ALIX proteins interact with proteins of the midbody and function in cytokinesis. Embo J 2007b; 26:4215–4227. [PubMed: 17853893]
- Morita E, Sundquist WI. Retrovirus budding. Annu Rev Cell Devel Biol 2004;20:395–425. [PubMed: 15473846]
- Munshi UM, Kim J, Nagashima K, Hurley JH, Freed EO. An Alix fragment potently inhibits HIV-1 budding - Characterization of binding to retroviral YPXL late domains. J Biol Chem 2007;282:3847–3855. [PubMed: 17158451]
- Muziol T, Pineda-Molina E, Ravelli RB, Zamborlini A, Usami Y, Gottlinger H, Weissenhorn W. Structural basis for budding by the ESCRT-III factor CHMP3. Dev Cell 2006;10:821–830. [PubMed: 16740483]
- Nakatogawa H, Suzuki K, Kamada Y, Ohsumi Y. Dynamics and diversity in autophagy mechanisms: lessons from yeast. Nat Rev Mol Cell Biol 2009;10:458–467. [PubMed: 19491929]
- Nickerson DP, West M, Henry R, Odorizzi G. Regulators of Vps4 activity at endosomes differentially influence the size and rate of formation of intralumenal vesicles. Mol Biol Cell. 2010 epub ahead of print.
- Nickerson DP, West M, Odorizzi G. Did2 coordinates Vps4-mediated dissociation of ESCRT-III from endosomes. J Cell Biol 2006;175:715–720. [PubMed: 17130288]
- Nikko E, Marini AM, Andre B. Permease recycling and ubiquitination status reveal a particular role for Bro1 in the multivesicular body pathway. J Biol Chem 2003;278:50732–50743. [PubMed: 14523026]
- Novick P, Field C, Schekman R. Identification of 23 Complementation Groups Required for Post-Translational Events in the Yeast Secretory Pathway. Cell 1980;21:205–215. [PubMed: 6996832]
- Obita T, Saksena S, Ghazi-Tabatabai S, Gill DJ, Perisic O, Emr SD, Williams RL. Structural basis for selective recognition of ESCRT-III by the AAA ATPase Vps4. Nature 2007;449:735–739. [PubMed: 17928861]
- Odorizzi G. The multiple personalities of Alix. J Cell Sci 2006;119:3025–3032. [PubMed: 16868030]
- Odorizzi G, Babst M, Emr SD. Fab1p PtdIns(3)P 5-kinase function essential for protein sorting in the multivesicular body. Cell 1998;95:847–858. [PubMed: 9865702]
- Oestreich AJ, Aboian M, Lee J, Azmi I, Payne J, Issaka R, Davies BA, Katzmann DJ. Characterization of multiple multivesicular sorting determinants within Sna3: A role for the ubiquitin ligase Rsp5. 2006a;18:707–720.
- Oestreich AJ, Davies BA, Payne JA, Katzmann DJ. Mvb12 is a novel member of ESCRT-I involved in cargo selection by the multivesicular body pathway. Mol Biol Cell 2006b;18:646–657. [PubMed: 17151358]
- Pankiv S, Clausen TH, Lamark T, Brech A, Bruun JA, Outzen H, Overvatn A, Bjorkoy G, Johansen T. p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. J Biol Chem 2007;282:24131–24145. [PubMed: 17580304]
- Pineda-Molina E, Belrhali H, Piefer AJ, Akula I, Bates P, Weissenhorn W. The crystal structure of the C-terminal domain of Vps28 reveals a conserved surface required for Vps20 recruitment. Traffic 2006;7:1007–1016. [PubMed: 16749904]
- Piper RC, Cooper AA, Yang H, Stevens TH. Vps27 Controls Vacuolar And Endocytic Traffic Through A Prevacuolar Compartment In Saccharomyces-Cerevisiae. J Cell Biol 1995;131:603–617. [PubMed: 7593183]
- Piper RC, Katzmann DJ. Biogenesis and function of multivesicular bodies. Annu Rev Cell Devel Biol 2007;23:519–547. [PubMed: 17506697]
- Piper RC, Luzio JP. Ubiquitin-dependent sorting of integral membrane proteins for degradation in lysosomes. Curr Opin Cell Biol 2007;19:459–465. [PubMed: 17689064]
- Pires R, Hartlieb B, Signor L, Schoehn G, Lata S, Roessle M, Moriscot C, Popov S, Hinz A, Jamin M, et al. A Crescent-Shaped ALIX Dimer Targets ESCRT-III CHMP4 Filaments. Structure 2009;17:843–856. [PubMed: 19523902]
- Pohl C, Jentsch S. Final stages of cytokinesis and midbody ring formation are controlled by BRUCE. Cell 2008;132:832–845. [PubMed: 18329369]

- Popov S, Popova E, Inoue M, Gottlinger HG. Human immunodeficiency virus Type 1 Gag engages the Bro1 domain of ALIX/AIP1 through the nucleocapsid. J Virol 2008;82:1389–1398. [PubMed: 18032513]
- Popov S, Popova E, Inoue M, Gottlinger HG. Divergent Bro1 Domains Share the Capacity To Bind Human Immunodeficiency Virus Type 1 Nucleocapsid and To Enhance Virus-Like Particle Production. J Virol 2009;83:7185–7193. [PubMed: 19403673]
- Pornillos O, Alam SL, Davis DR, Sundquist WI. Structure of the Tsg101 UEV domain in complex with the PTAP motif of the HIV-1 p6 protein. Nat Struct Biol 2002;9:812–817. [PubMed: 12379843]
- Pornillos O, Higginson DS, Stray KM, Fisher RD, Garrus JE, Payne M, He GP, Wang HE, Morham SG, Sundquist WI. HIV Gag mimics the Tsg101-recruiting activity of the human Hrs protein. J Cell Biol 2003;162:425–434. [PubMed: 12900394]
- Prag G, Watson H, Kim YC, Beach BM, Ghirlando R, Hummer G, Bonifacino JS, Hurley JH. The Vps27/ Hse1 complex is a GAT domain-based scaffold for ubiquitin-dependent sorting. Dev Cell 2007;12:973–986. [PubMed: 17543868]
- Prekeris R, Gould GW. Breaking up is hard to do membrane traffic in cytokinesis. J Cell Sci 2008;121:1569–1576. [PubMed: 18469013]
- Progida C, Malerod L, Stuffers S, Brech A, Bucci C, Stenmark H. RILP is required for the proper morphology and function of late endosomes. J Cell Sci 2007:120. [PubMed: 18073238]
- Puertollano R. Interactions of Tom1L1 with the multivesicular body sorting machinery. J Biol Chem 2005;280:9258–9264. [PubMed: 15611048]
- Puertollano R, Bonifacino JS. Interactions of GGA3 with the ubiquitin sorting machinery. Nat Cell Biol 2004;6:244–251. [PubMed: 15039775]
- Raiborg C, Bache KG, Gillooly DJ, Madshus IH, Stang E, Stenmark H. Hrs sorts ubiquitinated proteins into clathrin-coated microdomains of early endosomes. Nat Cell Biol 2002;4:394–398. [PubMed: 11988743]
- Raiborg C, Bache KG, Mehlum A, Stang E, Stenmark H. Hrs recruits clathrin to early endosomes. Embo J 2001a;20:5008–5021. [PubMed: 11532964]
- Raiborg C, Bremnes B, Mehlum A, Gillooly DJ, D'Arrigo A, Stang E, Stenmark H. FYVE and coiledcoil domains determine the specific localisation of Hrs to early endosomes. J Cell Sci 2001b; 114:2255–2263. [PubMed: 11493665]
- Raiborg C, Stenmark H. The ESCRT machinery in endosomal sorting of ubiquitylated membrane proteins. Nature 2009;458:445–452. [PubMed: 19325624]
- Raymond CK, Howald-Stevenson I, Vater CA, Stevens TH. Morphological Classification of the Yeast Vacuolar Protein Sorting Mutants - Evidence for a Prevacuolar Compartment in Class-E Vps Mutants. Mol Biol Cell 1992;3:1389–1402. [PubMed: 1493335]
- Reggiori F, Pelham HRB. Sorting of proteins into multivesicular bodies: ubiquitin-dependent and independent targeting. Embo J 2001;20:5176–5186. [PubMed: 11566881]
- Ren J, Kee Y, Huibergtse JM, Piper RC. Hse1, a Component of the Yeast Hrs-STAM Ubiquitin Sorting Complex, Associates with Ubiquitin Peptidases and a Ligase to Control Sorting Efficiency into Multivesicular Bodies. Mol Biol Cell 2007;18:324–335. [PubMed: 17079730]
- Ren X, Kloer DP, Kim YC, Ghirlando R, Saidi LF, Hummer G, Hurley JH. Hybrid structural model of the complete human ESCRT-0 complex. Structure 2009;17:406–416. [PubMed: 19278655]
- Ren XF, Hurley JH. VHS domains of ESCRT-0 cooperate in high-avidity binding to polyubiquitinated cargo. Embo J 2010;29:1045–1054. [PubMed: 20150893]
- Roudier N, Lefebvre C, Legouis R. CeVPS-27 is an endosomal protein required for the molting and the endocytic trafficking of the low-density lipoprotein receptor-related protein 1 in Caenorhabditis elegans. Traffic 2005;6:695–705. [PubMed: 15998324]
- Row PE, Lui H, Hayes S, Welchman R, Charalabous P, Hofmann K, Claugue MJ, Sanderson CM, Urbe S. The MIT domain of UBPY constitutes a CHMP binding and endosomal localization signal required for efficient EGF receptor degradation. J Biol Chem 2007;282:30929–30937. [PubMed: 17711858]
- Rubio-Texeira M, Kaiser CA. Amino acids regulate retrieval of the yeast general amino acid permease from the vacuolar targeting pathway. Mol Biol Cell 2006;17:3031–3050. [PubMed: 16641373]

- Rue SM, Mattei S, Saksena S, Emr SD. Novel Ist1-Did2 complex functions at a late step in multivesicular body sorting. Mol Biol Cell 2008;19:475–484. [PubMed: 18032584]
- Rusten TE, Simonsen A. ESCRT functions in autophagy and associated disease. Cell Cycle 2008;7:1166– 1172. [PubMed: 18418046]
- Rusten TE, Stenmark H. How do ESCRT proteins control autophagy? J Cell Sci 2009;122:2179–2183. [PubMed: 19535733]
- Rusten TE, Vaccari T, Lindmo K, Rodahl LMW, Nezis IP, Sem-Jacobsen C, Wendler F, Vincent JP, Brech A, Bilder D, Stenmark H. ESCRTs and Fab1 regulate distinct steps of autophagy. Curr Biol 2007;17:1817–1825. [PubMed: 17935992]
- Sagona AP, Nezis IP, Pedersen NM, Liestol K, Poulton J, Rusten TE, Skotheim RI, Raiborg C, Stenmark H. PtdIns(3)P controls cytokinesis through KIF13A-mediated recruitment of FYVE-CENT to the midbody. Nat Cell Biol 12:362–U140. [PubMed: 20208530]
- Saksena S, Emr SD. ESCRTs and human disease. Biochem Soc Trans 2009;37:167–172. [PubMed: 19143624]
- Saksena S, Sun J, Chu T, Emr SD. ESCRTing proteins in the endocytic pathway. Trends Biochem Sci 2007;32:561–573. [PubMed: 17988873]
- Saksena S, Wahlman J, Teis D, Johnson AE, Emr SD. Functional Reconstitution of ESCRT-III Assembly and Disassembly. Cell 2009;136:97–109. [PubMed: 19135892]
- Samson RY, Bell SD. Ancient ESCRTs and the evolution of binary fission. Trends Microbiol 2009;17:507–513. [PubMed: 19783442]
- Samson RY, Obita T, Freund SM, Williams RL, Bell SD. A role for the ESCRT system in cell division in Archaea. Science 2008;322:1710–1713. [PubMed: 19008417]
- Sankaran VG, Klein DE, Sachdeva MM, Lemmon MA. High-affinity binding of a FYVE domain to phosphatidylinositol 3-phosphate requires intact phospholipid but not FYVE domain oligomerization. Biochemistry 2001;40:8581–8587. [PubMed: 11456498]
- Schlundt A, Sticht J, Piotukh K, Kosslick D, Jahnke N, Keller S, Schuemann M, Krause E, Freund C. Proline-rich Sequence Recognition - II. PROTEOMICS ANALYSIS OF Tsg101 UBIQUITIN-E2-LIKE VARIANT (UEV) INTERACTIONS. Mol Cell Proteomics 2009;8:2474–2486. [PubMed: 19542561]
- Schmidt MHH, Dikic I, Bogler O. Src phosphorylation of Alix/AIP1 modulates its interaction with binding partners and antagonizes its activities. J Biol Chem 2005;280:3414–3425. [PubMed: 15557335]
- Scott A, Chung HY, Gonciarz-Swiatek M, Hill GC, Whitby FG, Gaspar J, Holton JM, Viswanathan R, Ghaffarian S, Hill CP, Sundquist WI. Structural and mechanistic studies of VPS4 proteins. Embo J 2005a;24:3658–3669. [PubMed: 16193069]
- Scott A, Gaspar J, Stuchell-Brereton MD, Alam SL, Skalicky JJ, Sundquist WI. Structure and ESCRT-III protein interactions of the MIT domain of human VPS4A. Proc Natl Acad Sci USA 2005b; 102:13813–13818. [PubMed: 16174732]
- Shestakova A, Hanono A, Drosner S, Curtiss M, Davies BA, Katzmann DJ, Babst M. Assembly of the AAA ATPase Vps4 on ESCRT-III. Mol Biol Cell. 2010 epub ahead of print.
- Shields SB, Oestreich AJ, Winistorfer S, Nguyen D, Payne JA, Katzmann DJ, Piper R. ESCRT ubiquitin binding domains function cooperatively during MVB cargo sorting. J Cell Biol 2009;185:213–224. [PubMed: 19380877]
- Shiflett SL, Ward DM, Huynh D, Vaughn MB, Simmons JC, Kaplan J. Characterization of Vta1p, a class E Vps protein in Saccharomyces cerevisiae. J Biol Chem 2004;279:10982–10990. [PubMed: 14701806]
- Shih SC, Katzmann DJ, Schnell JD, Sutanto M, Emr SD, Hicke L. Epsins and Vps27p/Hrs contain ubiquitin-binding domains that function in receptor endocytosis. Nat Cell Biol 2002;4:389–393. [PubMed: 11988742]
- Shim JH, Xiao CC, Hayden MS, Lee KY, Trombetta ES, Pypaert M, Nara A, Yoshimori T, Wilm B, Erdjument-Bromage H, et al. CHMP5 is essential for late endosome function and down-regulation of receptor signaling during mouse embryogenesis. J Cell Biol 2006;172:1045–1056. [PubMed: 16567502]

- Shim S, Kimpler LA, Hanson PI. Structure/Function Analysis of Four Core ESCRT-III Proteins Reveals Common Regulatory Role for Extreme C-terminal Domain. Traffic 2007;8:1068–1079. [PubMed: 17547705]
- Shim S, Merrill SA, Hanson PI. Novel interactions of ESCRT-III with LIP5 and VPS4 and their implications for ESCRT-III disassembly. Mol Biol Cell. 2008 epub ahead of print.
- Slagsvold T, Aasland R, Hirano S, Bache KG, Raiborg C, Trambaiano D, Wakatsuki S, Stenmark H. Eap45 in mammalian ESCRT-II binds ubiquitin via a phosphoinositide-interacting GLUE domain. J Biol Chem 2005;280:19600–19606. [PubMed: 15755741]
- Sotelo JR, Porter KR. An Electron Microscope Study of the Rat Ovum. J Biophys Biochem Cytol 1959;5:327. [PubMed: 13654454]
- Spitzer C, Reyes FC, Buono R, Sliwinski MK, Haas TJ, Otegui MS. The ESCRT-Related CHMP1A and B Proteins Mediate Multivesicular Body Sorting of Auxin Carriers in Arabidopsis and Are Required for Plant Development. Plant Cell 2009;21:749–766. [PubMed: 19304934]
- Spitzer C, Schellmann S, Sabovljevic A, Shahriari M, Keshavaiah C, Bechtold N, Herzog M, Muller S, Hanisch FG, Hulskamp M. The Arabidopsis elch mutant reveals functions of an ESCRT component in cytokinesis. Development 2006;133:4679–4689. [PubMed: 17090720]
- Stahelin RV, Long F, Diraviyam K, Bruzik KS, Murray D, Cho W. Phosphatidylinositol 3-phosphate induces the membrane penetration of the FYVE domains of Vps27p and Hrs. J Biol Chem 2002;277:26379–26388. [PubMed: 12006563]
- Stawiecka-Mirota M, Pokrzywa W, Morvan J, Zoladek T, Haguenauer-Tsapis R, Urban-Grimal D, Morsomme P. Targeting of Sna3p to the endosomal pathway depends on its interaction with Rsp5p and multivesicular body sorting on its ubiquitylation. Traffic 2007;8:1280–1296. [PubMed: 17645729]
- Steigemann P, Gerlich DW. Cytokinetic abscission: cellular dynamics at the midbody. Trends Cell Biol 2009;19:606–616. [PubMed: 19733077]
- Stuchell-Brereton M, Skalicky J, Kieffer C, Karren MA, Ghaffarian S, Sundquist WI. ESCRT-III recognition by VPS4 ATPases. Nature 2007;449:740–744. [PubMed: 17928862]
- Stuffers S, Brech A, Stenmark H. ESCRT proteins in physiology and disease. Exp Cell Res 2009;315:1619–1626. [PubMed: 19013455]
- Sundquist WI, Schubert HL, Kelly BN, Hill GC, Holton JM, Hill CP. Ubiquitin recognition by the human TSG101 protein. Mol Cell 2004;13:783–789. [PubMed: 15053872]
- Suzuki H, Kawasaki M, Inuzuka T, Okumura M, Kakiuchi T, Shibata H, Wakatsuki S, Maki M. Structural Basis for Ca2+-Dependent Formation of ALG-2/Alix Peptide Complex: Ca2+/EF3-Driven Arginine Switch Mechanism. Structure 2008;16:1562–1573. [PubMed: 18940611]
- Swanson KA, Kang RS, Stamenova SD, Hicke L, Radhakrishnan I. Solution structure of Vps27 UIMubiquitin complex important for endosomal sorting and receptor downregulation. Embo J 2003;22:4597–4606. [PubMed: 12970172]
- Takata H, Kato M, Denda K, Kitamura N. A Hrs binding protein having a Src homology 3 domain is involved in intracellular degradation of growth factors and their receptors. Genes Cells 2000;5:57– 69. [PubMed: 10651905]
- Teis D, Saksena S, Emr SD. Ordered Assembly of the ESCRT-III Complex on Endosomes Is Required to Sequester Cargo during MVB Formation. Dev Cell 2008;15:578–589. [PubMed: 18854142]
- Teis D, Saksena S, Judson BL, Emr SD. ESCRT-II coordinates the assembly of ESCRT-III filaments for cargo sorting and multivesicular body vesicle formation. Embo J 29:871–883. [PubMed: 20134403]
- Teo H, Perisic O, Gonzalez B, Williams RL. ESCRT-II, an endosome-associated complex required for protein sorting: Crystal structure and interactions with ESCRT-III and membranes. Dev Cell 2004a; 7:559–569. [PubMed: 15469844]
- Teo H, Veprintsev DB, Williams RL. Structural insights into endosomal sorting complex required for transport (ESCRT-I) recognition of ubiquitinated proteins. J Biol Chem 2004b;279:28689–28696. [PubMed: 15044434]
- Teo HL, Gill DJ, Sun J, Perisic O, Veprintsev DB, Vallis Y, Emr SD, Williams RL. ESCRT-I core and ESCRT-II GLUE domain structures reveal role for GLUE in linking to ESCRT-I and membranes. Cell 2006;125:99–111. [PubMed: 16615893]

- Theos AC, Truschel ST, Tenza D, Hurbain I, Harper DC, Berson JF, Thomas PC, Raposo G, Marks MS. A lumenal domain-dependent pathway for sorting to intralumenal vesicles of multivesicular endosomes involved in organelle morphogenesis. Dev Cell 2006;10:343–354. [PubMed: 16516837]
- Thompson BJ, Mathieu J, Sung HH, Loeser E, Rorth P, Cohen SM. Tumor suppressor properties of the ESCRT-II complex component vps25 in Drosophila. Dev Cell 2005;9:711–720. [PubMed: 16256745]
- Toyoshima M, Tanaka N, Aoki J, Tanaka Y, Murata K, Kyuuma M, Kobayashi H, Ishii N, Yaegashi N, Sugamura K. Inhibition of tumor growth and metastasis by depletion of vesicular sorting protein Hrs: Its regulatory role on E-cadherin and beta-catenin. Cancer Res 2007;67:5162–5171. [PubMed: 17545595]
- Trajkovic K, Hsu C, Chiantia S, Rajendran L, Wenzel D, Wieland F, Schwille P, Brugger B, Simons M. Ceramide triggers budding of exosome vesicles into multivesicular Endosomes. Science 2008;319:1244–1247. [PubMed: 18309083]
- Tsang HTH, Connell JW, Brown SE, Thompson A, Reid E, Sanderson CM. A systematic analysis of human CHMP protein interactions: Additional MIT domain-containing proteins bind to multiple components of the human ESCRT III complex. Genomics 2006;88:333–346. [PubMed: 16730941]
- Urbe S, Sachse M, Row PE, Preisinger C, Barr FA, Strous G, Klumperman J, Clague MJ. The UIM domain of Hrs couples receptor sorting to vesicle formation. J Cell Sci 2003;116:4169–4179. [PubMed: 12953068]
- Usami Y, Popov S, Gottlinger HG. Potent rescue of human immunodeficiency virus type 1 late domain mutants by ALIX/AIP1 depends on its CHMP4 binding site. J Virol 2007;81:6614–6622. [PubMed: 17428861]
- Usami Y, Popov S, Popova E, Gottlinger HG. Efficient and specific rescue of human immunodeficiency virus type 1 budding defects by a Nedd4-like ubiquitin ligase. J Virol 2008;82:4898–4907. [PubMed: 18321969]
- Usami Y, Popov S, Popova E, Inoue M, Weissenhorn W, Gottlinger HG. The ESCRT pathway and HIV-1 budding. Biochem Soc Trans 2009;37:181–184. [PubMed: 19143627]
- Vaccari T, Bilder D. The Drosophila tumor suppressor vps25 prevents nonautonomous overproliferation by regulating Notch trafficking. Dev Cell 2005;9:687–698. [PubMed: 16256743]
- Vardhana S, Choudhuri K, Varma R, Dustin ML. Essential Role of Ubiquitin and TSG101 Protein in Formation and Function of the Central Supramolecular Activation Cluster. Immunity 2010;32:531– 540. [PubMed: 20399684]
- Vito P, Pellegrini L, Guiet C, D'Adamio L. Cloning of AIP1, a novel protein that associates with the apoptosis-linked gene ALG-2 in a Ca2+-dependent reaction. J Biol Chem 1999;274:1533–1540. [PubMed: 9880530]
- Wang TL, Hong WJ. RILP interacts with VPS22 and VPS36 of ESCRT-II and regulates their membrane recruitment. Biochem Biophys Res Comm 2006;350:413–423. [PubMed: 17010938]
- Ward DMV, Vaughn MB, Shiflett SL, White PL, Pollock AL, Hill J, Schnegelberger R, Sundquist WI, Kaplan J. The role of LIP5 and CHMP5 in multivesicular body formation and HIV-1 budding in mammalian cells. J Biol Chem 2005;280:10548–10555. [PubMed: 15644320]
- Watson H, Bonifacino JS. Direct binding to Rsp5p regulates ubiquitination-independent vacuolar transport of Sna3p. Mol Biol Cell 2007;18:1781–1789. [PubMed: 17332499]
- Whitley P, Reaves BJ, Hashimoto M, Riley AM, Potter BVL, Holman GD. Identification of mammalian Vps24p as an effector of phosphatidylinositol 3,5-bisphosphate-dependent endosome compartmentalization. J Biol Chem 2003;278:38786–38795. [PubMed: 12878588]
- Williams RL, Urbe S. The emerging shape of the ESCRT machinery. Nat Rev Mol Cell Biol 2007;8:355–368. [PubMed: 17450176]
- Wollert T, Hurley JH. Molecular mechanism of multivesicular body biogenesis by the ESCRT complexes. Nature 2010;464:864–869. [PubMed: 20305637]
- Wollert T, Wunder C, Lippincott-Schwartz J, Hurley JH. Membrane scission by the ESCRT-III complex. Nature 2009a;458:172–177. [PubMed: 19234443]
- Wollert T, Yang D, Ren XF, Lee HH, Im YJ, Hurley JH. The ESCRT machinery at a glance. J Cell Sci 2009b;122:2163–2166. [PubMed: 19535731]

- Worsdorfer P, Maxeiner S, Markopoulos C, Kirfel G, Wulf V, Auth T, Urschel S, von Maltzahn J, Willecke K. Connexin expression and functional analysis of gap junctional communication in mouse embryonic stem cells. Stem Cells 2008;26:431–439. [PubMed: 18055446]
- Xiao J, Xia H, Zhou J, Azmi I, Davies BA, Katzmann DJ, Xu Z. Structural basis of Vta1 function in the multi-vesicular body sorting pathway. Dev Cell 2008;14:37–49. [PubMed: 18194651]
- Xiao JY, Chen XW, Davies BA, Saltiel AR, Katzmann DJ, Xu ZH. Structural Basis of Ist1 Function and Ist1-Did2 Interaction in the Multivesicular Body Pathway and Cytokinesis. Mol Biol Cell 2009;20:3514–3524. [PubMed: 19477918]
- Xiao JY, Xia HC, Yoshino-Koh K, Zhou JH, Xu ZH. Structural characterization of the ATPase reaction cycle of endosomal AAA protein Vps4. J Mol Biol 2007;374:655–670. [PubMed: 17949747]
- Xie WQ, Li LM, Cohen SN. Cell cycle-dependent subcellular localization of the TSG101 protein and mitotic and nuclear abnormalities associated with TSG101 deficiency. Proc Natl Acad Sci USA 1998;95:1595–1600. [PubMed: 9465061]
- Yamashita Y, Kojima K, Tsukahara T, Agawa H, Yamada K, Amano Y, Kurotori N, Tanaka N, Sugamura K, Takeshita T. Ubiquitin-independent binding of Hrs mediates endosomal sorting of the interleukin-2 receptor beta-chain. J Cell Sci 2008;121:1727–1738. [PubMed: 18445679]
- Yang D, Hurley JH. Structural role of the Vps4-Vta1 interface in ESCRT-III recycling. Structure. 2010 in press.
- Yang D, Rismanchi N, Renvoisé B, Lippincott-Schwartz J, Blackstone C, Hurley JH. Structural basis for midbody targeting of spastin by the ESCRT-III protein CHMP1B. Nat Struct Mol Biol 2008;15:1278–1286. [PubMed: 18997780]
- Yeo SCL, Xu LH, Ren JH, Boulton VJ, Wagle MD, Liu C, Ren G, Wong P, Zahn R, Sasajala P, et al. Vps20p and Vta1p interact with Vps4p and function in multivesicular body sorting and endosomal transport in Saccharomyces cerevisiae. J Cell Sci 2003;116:3957–3970. [PubMed: 12953057]
- Yu ZH, Gonciarz MD, Sundquist WI, Hill CP, Jensen GJ. Cryo-EM structure of dodecameric Vps4p and its 2 : 1 complex with Vta1p. J Mol Biol 2008;377:364–377. [PubMed: 18280501]
- Zamborlini A, Usami Y, Radoshitzky SR, Popova E, Palu G, Gottlinger H. Release of autoinhibition converts ESCRT-III components into potent inhibitors of HIV-1 budding. Proc Natl Acad Sci USA 2006;103:19140–19145. [PubMed: 17146056]
- Zhadina M, McClure M, Johnson MC, Bieniasz PD. Ubiquitin-dependent virus particle budding without viral protein ubiquitination. Proc Natl Acad Sci USA 2007;104:20031–20036. [PubMed: 18056634]
- Zhai Q, Fisher RD, Chung HY, Myszka DG, Sundquist WI, Hill CP. Structural and functional studies of ALIX interactions with YPXnL late domains of HIV-1 and EIAV. Nat Struct Mol Biol 2008;15:43– 49. [PubMed: 18066081]
- Zhou X, Pan S, Sun L, Corvera J, Lin SH, Kuang J. The HIV-1 p6/EIAV p9 docking site in ALIX is autoinhibited as revealed by a conformation-sensitive anti-ALIX monoclonal antibody. Biochem J 2008;414:215–220. [PubMed: 18476810]



Figure 1. Multivesicular bodies. Color versions of all figures are available on line.



Figure 2. Major cellular functions of the ESCRTs.



Figure 3.

Schematic of the organization of ESCRT-0. a) yeast and (b) human ESCRT-0 shown docked to a flat, cargo-bearing membrane.



Figure 4.

Schematic of the organization of ESCRT-I and –II. (a) yeast and (b) human ESCRT-I and –II depicted as a supercomplex assembled at a membrane neck.



Figure 5.

Speculative schematic of the organization of ESCRT-III. The assembly is depicted as a spiral on the basis of the EM images of (Hanson et al., 2008) and for simplicity. The actual assembly is more likely dome-shaped rather than flat (Fabrikant *et al.*, 2009). The order of assembly of the first four subunits (beginning with Vps20 at the outside of the spiral), and the preponderance of Snf7 subunits shown is based on the observations of (Teis et al., 2008).



Figure 6. Schematic of the organization of Vps4-Vta1.

Crit Rev Biochem Mol Biol. Author manuscript; available in PMC 2011 December 1.

NIH-PA Author Manuscript



Figure 7. Schematic of the organization of Bro1.

Crit Rev Biochem Mol Biol. Author manuscript; available in PMC 2011 December 1.

NIH-PA Author Manuscript



Figure 8.

Ubiquitin binding domains. Ubiquitin is shown in yellow with Ile44 shown with space-filling spheres. UBDs are shown in blue. The VHS domain complex is shown for the human STAM subunit of ESCRT-0 (pdb entry 3LDZ) but is representative of all of the yeast and human ESCRT VHS domains. The UIM is shown for yeast ESCRT-0 subunit Vps27 (1Q0W) and is probably also representative of other "single" or conventional UIMs in the yeast ESCRT-0 Hse1 subunit and the human ESCRT-0 STAM subunit (Fig. 3). The DUIM is shown for human ESCRT-0 subunit Hrs (2D3G). The UEV domain is shown for human ESCRT-1 (1S1Q) and is also representative of the yeast UEV-ubiquitin complex. The structure of the yeast Mvb12 C-terminal UBD in complex with ubiquitin has not been reported. The structure of the NZF of Npl4 is shown (1Q5W), as a reasonable representation of the yeast ESCRT-II NZF2-ubiquitin interaction, as no structure is available for the latter. This domain is not present in human ESCRT-II. The structure of the GLUE domain of human ESCRT-II is shown (2HTH); the yeast GLUE domain does not bind ubiquitin.



Figure 9.

Membrane binding domains. Protein surfaces are colored green (hydrophobic residues), white (uncharged polar residues), red (acidic residues), and blue (basic residues). Structures are oriented such that the plane of the membrane is beneath each structure. The FYVE domain of ESCRT-0 (1VFY) targets phosphatidylinositol 3-phosphate, shown here as the soluble headgroup inositol 1,3-bisphosphate docked onto to the unliganded Vps27 FYVE domain structure from the liganded EEA1 FYVE structure (1JOC). The GLUE domain of yeast ESCRT-II (2CAY) is shown, with the non-canonical phosphoinositide binding pocket marked by a bound sulfate ion. The α 1- α 4 membrane binding core of the ESCRT-III subunit VPS24 (2GD5) is shown, with the autoinhibitory helix omitted.



Figure 10.

MIT domain-MIM complexes. The MIT domain of yeast Vps4(cyan) is shown bound to the MIM1 of Vps2 (2V6X). The MIT domain of human VPS4A (cyan) is shown in complex with the MIM2 of human VPS20 (2K3W). The MIT domain of human spastin (blue) is shown bound to the extended MIM1 of human DID2B (3EAB). MIMs are orange.

ALIX-SNF7B



Figure 11.

The Bro1 domain. The structure of the Bro1 domain of ALIX (cyan) is shown in complex with SNF7B (orange, 3C3Q). Conserved interaction residues are highlighted in stick models, and the Src binding site is also shown.



Figure 12.

Viral late domains and ESCRTs. Top, the PTAP motif (stick model) of HIV-1 bound to the UEV domain of TSG101 (1M4Q). Botton, the YPX₃L motif (stick model) of HIV-1 bound to the V domain of ALIX (2R02). Protein surfaces are colored green (hydrophobic residues), white (uncharged polar residues), red (acidic residues), and blue (basic residues).



Figure 13.

GPPX₃Y motif targeting to midbodies. The two subunits of CEP55 are colored cyan and orange, and the GPPX₃Y peptide of ALIX is shown in a stick model.

~
~
_
_
0
-
-
~
-
<u> </u>
-
-
0
<u> </u>
_
~
<
_
ຸດາ
~
_
-
_
c n
0
-
+

NIH-PA Author Manuscript

Table 1

	domain and coiled-																			
Motifs and Domains	VHS, FYVE, UIM (yeast), DUIM (metazoan), P(S/T)XP, GAT of coil core, clathrin-binding	VHS, UIM, SH3, GAT domain and coiled-coil core	UEV, PRD, stalk, headpiece	headpiece, Vps28 CTD	basic helix, stalk, headpiece	stalk, headpiece ("UMA domain"), MAPB; UBAP1 and LOC290595 are the 3rd and 4th putative Human MVB12 isoforms	basic helix, Winged-helix	Winged-helix	Winged-helix, GLUE, NZF1 (yeast), NZF2 (yeast)	MIMI	MIM2	weak MIMI	weak MIM2		MIMI		MIM1, MIM2	MIT, AAA	MIT	Brol, V, PRD
Synonyms (metazoan)	Hrs,HRS, Hgs	STAM1, 2	TSG101			UBAP1, LOC390595	EAP30	EAP20	EAP45	CHMP2A,B	CHMP6	CHMP3	CHMP4A, B, C	CHMP5	CHMP1A, B			SKD1	LIP5	AIP1
Metazoan Protein (yeast-centric terminology, this review)	VPS27	HSEI	VPS23	VPS28	VPS37A,B,C,D	MVB12A,B	VPS22	VPS25	VPS36	VPS2A,B	VPS20	VPS24	SNF7A,B,C	VPS60	DID2A, B	CHMP7	ISTI	VPS4A,B	VTA1	ALIX
Selected synonyms (yeast)			Stp22		Srn2		Snf8			Did4 Chm2	Chm6		Vps32	Chm5	Chm1 Vps46					Vps31
east or generic otein	ps27	sel	ps23	ps28	ps37	vb12	ps22	ps25	ps36	ps2	ps20	ps24	ıf7	ps60	id2		11	ps4	tal	ro1
Complex Y _i	ESCRT-0 V ₁	H	ESCRT-I V ₁	V	V1	Μ	ESCRT-II V ₁	V	V1	ESCRT-III V ₁	V	V	Sn	V	D		Isi	Vps4-Vta1 V	V	Bro1/ALIX B1

Crit Rev Biochem Mol Biol. Author manuscript; available in PMC 2011 December 1.

Table 2

Viral late domains and ESCRT recruitment. This table is adapted and updated from one appearing in the comprehensive review of virus budding by (Chen and Lamb, 2008).

PTAP-TSG101 UEV		
Virus	Reference	notes
HIV-1	(Demirov <i>et al.</i> , 2002; Garrus <i>et al.</i> , 2001; Martin-Serrano <i>et al.</i> , 2001; VerPlank <i>et al.</i> , 2001)	
Feline immunodeficiency virus	(Luttge et al., 2008)	
Mason-Pfizer Monkey virus	(Gottwein <i>et al.</i> , 2003)	
Moloney murine leukemia virus	(Segura-Morales et al., 2005)	
HTLV-1	(Blot <i>et al.</i> , 2004; Bouann <i>et al.</i> , 2003; Dorweiler <i>et al.</i> , 2006; Heidecker <i>et al.</i> , 2004; Wang <i>et al.</i> , 2004)	Minor/cell-type dependent role.
Prototypic foamy virus	(Patton et al., 2005)	
Vesicular stomatitis virus	(Irie et al., 2004)	
Ebola	(Licata <i>et al.</i> , 2003; Martin-Serrano et al., 2001; Timmins <i>et al.</i> , 2003)	
Japanese encephalitis virus	(Chiou <i>et al.</i> , 2003)	PTAP motif not identified but TSG101 role is confirmed.
Lymphocytic choriomeningitis virus	(Perez et al., 2003)	
Lassa virus	(Perez et al., 2003; Urata <i>et al.</i> , 2006)	
Bluetongue virus	(Wirblich <i>et al.</i> , 2006)	
YPX _n L-ALIX-dependent		
Virus		
HIV-1	(Strack <i>et al.</i> , 2003; von Schwedler <i>et al.</i> , 2003)	
Equine infectious anemia virus	(Martin-Serrano <i>et</i> al., 2003; Strack et al., 2003)	
Prototypic foamy virus	(Patton et al., 2005; Stange <i>et al.</i> , 2005)	
Sendai virus	(Gosselin-Grenet et al., 2007; Irie et al., 2007; Sakaguchi et al., 2005)	Consensus that ALIX interacts with viral YLDL sequence but (Gosselin-Grenet et al., 2007) finds no effect on budding of ALIX

PTAP-18G101 UEV						
Virus	Reference	notes				
		knockdown				
Vaccinia virus	(Honeychurch <i>et al.</i> , 2007)					
Rous sarcoma virus	(Dilley et al., 2010)	secondary				
PPXY-dependent						
Marburg	(Urata <i>et al.</i> , 2007)	Unconventional interaction with TSG101 is PTAP independent but PPXY dependent				
Rous sarcoma virus	(Kikonyogo <i>et al.</i> , 2001; Medina <i>et al.</i> , 2005)					
Mason-Pfizer Monkey virus	(Gottwein et al., 2003)					
Moloney murine leukemia virus	(Segura-Morales et al., 2005)					
HTLV-1	(Blot et al., 2004; Bouamr et al., 2003; Heidecker et al., 2004; Wang et al., 2004)					
Vesicular stomatitis virus	(Harty et al., 1999)					
Rabies virus	(Harty et al., 1999)					
Ebola virus	(Harty <i>et al.</i> , 2000; Licata et al., 2003; Timmins et al., 2003)					
Lymphocytic choriomeningitis virus	(Perez et al., 2003)					
Lassa virus	(Perez et al., 2003; Urata et al., 2006)					
Bluetongue virus	(Wirblich et al., 2006)					
ESCRTs involved						
Recruitment mechanism other or unknown						
Human cytomegalovirus	(Tandon <i>et al.</i> , 2009)	Note that (Fraile- Ramos <i>et al.</i> , 2007) find no effect of ESCRT knockdown.				
Hepatitis C	(Corless et al.)					
Hepatitis B	(Lambert et al., 2007)					
Herpes simplex type 1	(Pawliczek and Crump, 2009)					
Tomato bushy stunt virus	(Barajas <i>et al.</i> , 2009)	Ubiquitinated p33 protein binds to ESCRT-I and Ali				

NIH-PA Author Manuscript