

REVIEW

Molecular Studies on Stress-Responsive Gene Expression in *Arabidopsis* and Improvement of Stress Tolerance in Crop Plants by Regulon Biotechnology

Kazuo NAKASHIMA¹ and Kazuko YAMAGUCHI-SHINOZAKI^{1,2*}

¹ Biological Resources Division, Japan International Research Center for Agricultural Sciences (JIRCAS) (Tsukuba, Ibaraki 305–8686, Japan)

² Laboratory of Plant Molecular Physiology, Graduate School of Agricultural and Life Sciences, The University of Tokyo (Bunkyo, Tokyo 113–8657, Japan)

Abstract

Molecular studies have shown that several genes with various functions are induced by environmental stresses such as drought, high-salinity and low temperature in plants. Most of the dehydration responsive genes are induced by the plant hormone abscisic acid (ABA), but others are not. Expression analyses of dehydration-responsive genes have provided at least four independent regulatory systems (regulons) for gene expression in a model plant *Arabidopsis thaliana*. The *cis*-acting elements in the promoters of some genes that have a typical stress-inducible expression profile and the transcription factors that affect the expression of these genes have been analyzed. Transcription factors that bind to a DRE/CRT (dehydration-responsive element / C-repeat) *cis*-acting element were isolated and termed DREB1/CBF (DRE-binding protein 1/ C-repeat binding factor) and DREB2 (DRE-binding protein 2). Overexpression of DREB1/CBF in transgenic *Arabidopsis* plants increased tolerance to freezing, drought and high salt concentrations. The *DREB1/CBF* genes have been successfully used to improve abiotic stress tolerance in a number of different crop plants. Studies on the other transcription factors associated with stress response are in progress. We collaborate with many research groups to improve stress tolerant crop plants utilizing regulon biotechnology. We hope the results of these collaborative studies will contribute to the sustainable food production in developing countries and help to prevent the global-scale environmental damage.

Discipline: Biotechnology

Additional key words: DREB1, environmental stress, transcription factors, transgenic plants

Introduction

As plants are sessile organisms, they are directly exposed to environmental stresses such as drought, high salinity and low temperature. Plants respond to environmental stress, and the transduced signals cause expression of numerous genes associated with stress tolerance. A number of genes have been described that respond to environmental stresses such as drought, high salinity and low temperature in plants^{4,15,33,47,48,60}.

We isolated more than 60 independent cDNAs for dehydration inducible genes using molecular techniques such as differential screening in a model plant *Arabidopsis thaliana*^{33,47,48}. Recently, 299 drought-inducible genes, 54 cold-inducible genes, and 213 high-salinity-

stress-inducible genes were identified using a cDNA microarray containing around 7,000 independent *Arabidopsis* full-length cDNA groups^{46,48}. Functions of their gene products have been predicted from sequence homology with known proteins. Genes induced during dehydration stress conditions are thought to function not only in protecting cells from dehydration by the production of important metabolic proteins (functional proteins) but also in the regulation of genes for signal transduction in the dehydration stress response (regulatory proteins). The functional proteins contain water channel proteins, chaperons, proteases, LEA (Late Embryogenesis Abundant) proteins, and enzymes for the synthesis of osmoprotectants (compatible solutes; sugars, proline, etc.). The regulatory proteins contain transcription factors, protein kinases, and enzymes for phosphoinositide (PI) turn-

*Corresponding author: fax +81–29–838–6643; e-mail kazukoys@jircas.affrc.go.jp

Received 2 March 2005; accepted 22 April 2005.

over, and enzymes for the synthesis of the plant hormone abscisic acid (ABA). So far, various kinds of functional proteins such as enzymes for the synthesis of osmoprotectants were overexpressed in plants to improve the stress tolerance^{5,59}. However, it seems that the engineering of one enzyme is not enough as many kinds of stress responses are necessary for plants to survive in severe stress conditions.

In plants, one transcription factor can control the expression of many target genes through the specific binding of the transcription factor to the *cis*-acting element in the promoters of the target genes. Such kind of a transcription unit is called a “regulon”. Northern analysis of dehydration-inducible genes revealed that there appear to be at least four independent regulons in *Arabidopsis* (Fig. 1). They are (1) DREB regulon, (2) NAC (NAM, ATAF1, 2, and CUC2) and ZF-HD (zinc-finger homeodomain) regulon, (3) AREB/ABF (ABA-responsive ele-

ment binding protein / ABA-responsive element binding factor) regulon, and (4) MYC (myelocytomatosis oncogene) and MYB (myeloblastosis oncogene) regulon. The DREB regulon and the NAC and ZF-HD regulon are ABA-independent. The AREB/ABF regulon and the MYC and MYB regulon are ABA-dependent. Regulon biotechnology, by controlling the expression of the regulon system, is expected to improve the tolerance against stresses in plants.

DREB regulon involved in ABA-independent gene expression

1. Isolation of DREB1/CBF regulon and DREB2 regulon

The promoter of an *Arabidopsis* drought-, high-salinity- and cold-inducible gene *RD29A* (*responsive to dehydration 29A*) encoding a LEA-like protein has been




Fig. 1. Regulatory network of gene expression in response to drought, high salinity and cold stresses: specificity and crosstalk of gene networks

Cis-acting elements that are involved in stress-responsive transcription are shown in boxes. Transcription factors that control stress-inducible gene expression are shown in circles or ovals. Small circles indicate the modification of transcription factors in response to stress signals for their activation, such as phosphorylation. Dotted lines indicate possible regulation. Double arrow lines indicate possible cross talk. ABF: ABRE-binding factor, ABRE: ABA-responsive element, AREB: ABRE-binding protein, CBF: C-repeat-binding factor, CRT: C-repeat, DRE: dehydration-responsive element, DREB: DRE-binding protein, *ERD*: early responsive to dehydration, ICE: inducer of CBF expression, MYBR: MYB recognition site, MYCR: MYC recognition site, NACR: NAC recognition site, *RD*: responsive to dehydration, ZF-HD: zinc finger homeodomain protein.

found to contain two major *cis*-acting elements, the ABA-responsive element (ABRE) and the dehydration-responsive element (DRE)/C-repeat (CRT), that are involved in stress-inducible gene expression⁵⁶. DRE/CRT (CCGAC) is a *cis*-acting element that functions in ABA-independent gene expression in response to abiotic stress (Fig. 1). Transcription factors belonging to the AP2/ERF (APETALA2 / ethylene-responsive element binding factor) family that bind to DRE/CRT have been isolated and termed DREB1/CBF and DREB2^{10,26,50}. The conserved DNA-binding motif of DREB1/CBF and DREB2 is A/GCCGAC⁴². The *DREB1/CBF* genes are quickly and transiently induced by cold stress, and their products activate the expression of target stress-inducible genes. The *DREB2* genes are induced by dehydration, leading to the expression of various genes that are involved in drought-stress tolerance²⁶.

2. Improved stress tolerance of transgenic plants overexpressing *DREB1/CBF*

Overexpression of *DREB1A/CBF3* in transgenic *Arabidopsis* plants showed increased tolerance to freezing, drought and high salt concentrations^{17,19,26}, suggesting that the *DREB1A/CBF3* proteins function without modification of the proteins in the development of stress tolerance. Many candidates for the *DREB1A/CBF3* target genes have been identified using microarray^{9,30,45}. Most of these target genes contain DRE- or DRE-related CCGAC core motif sequences in their promoter regions. We analyzed the expression of these candidate genes using RNA gel blot and identified more than 40 genes as the *DREB1A* downstream genes. Many of the products of these genes were proteins known to function against stress and were probably responsible for the stress tolerance of the transgenic plants. The downstream genes also included genes for transcription factors involved in further regulation of signal transduction and gene expression in response to stress.

The overexpression of the *DREB1/CBF* gene results in multiple biochemical changes associated with cold acclimation¹¹: *DREB1A/CBF3*-expressing plants had elevated levels of proline (Pro) and total soluble sugars, including sucrose, raffinose, glucose, and fructose. Plants overexpressing *DREB1A/CBF3* also had elevated *P5CS* (for delta(1)-pyrroline-5-carboxylate synthase) transcript levels suggesting that the increase in Pro levels resulted, at least in part, from increased expression of the key Pro biosynthetic enzyme *P5CS*. These results lead us to propose that *DREB1A/CBF3* integrates the activation of multiple components of the cold acclimation response.

Dwarfism is observed in transgenic *Arabidopsis* overexpressing *DREB1A/CBF3*, *DREB1B/CBF1*,

DREB1C/CBF2 or *DREB1D/CBF4*^{11,12,19,26}. The development of dwarf phenotypes was also found in transgenic tomato overexpressing *Arabidopsis DREB1B/CBF1*, and it was prevented by exogenous application of gibberellin (GA)¹⁴. These suggest that an inhibition of GA biosynthesis is a function common to the *DREB1/CBF* genes. However, microarray analysis did not detect the changes in transcript levels of GA-related genes in transgenic *Arabidopsis* overexpressing *DREB1A/CBF3*, *DREB1B/CBF1*, or *DREB1C/CBF2*⁹. Recently, *DREB1F* is reported to be involved in the regulation of GA biosynthesis and stress tolerance²⁹. It is not clear yet whether other *DREB1/CBF* proteins are related to GA synthesis or not.

In contrast to the *DREB1/CBF* genes, overexpression of *DREB2* in transgenic plants does not improve stress tolerance, suggesting that *DREB2* proteins require posttranslational activation²⁶. The *DREB2* protein is expressed under normal growth conditions and is activated in the early stage of the osmotic stress response through posttranslational modification (Fig. 1).

3. Regulation of the expression of *DREB1/CBF* regulon

The *ICE1* (*inducer of CBF expression 1*) gene was identified through the map-based cloning of the *Arabidopsis ice1* mutation, which affected the expression of the *DREB1A/CBF3* promoter-*LUC* (*luciferase*) transgene⁶. *ICE1* encodes a MYC-type bHLH (basic helix-loop-helix) transcription factor that regulates the expression of *DREB1A/CBF3* but not of other *DREB1/CBF* genes (Fig. 1). Overexpression of *ICE1* in transgenic plants resulted in improved freezing tolerance, supporting an important role for *ICE1* in the cold-stress response. Molecular analysis of the *DREB1C/CBF2* promoter has identified multiple *cis*-acting elements that are involved in cold-inducible gene expression⁵⁷ (Imura et al., unpublished data). The DNA-binding protein has been cloned and shown to be a MYC-type bHLH transcription factor that is different from *ICE1* (Imura et al., unpublished data). These results suggest the redundant involvement of MYC-type bHLH transcription factors in the up-regulation of the *DREB1/CBF* genes. A cold signal is necessary for the activation of the *ICE* proteins but the mechanism of this signal remains to be solved. Analysis of the *cbf2* mutant, in which the *DREB1C/CBF2* gene was disrupted, indicated that *DREB1C/CBF2* is a negative regulator of *DREB1A/CBF3* and *DREB1B/CBF1* expression and plays a central role in stress tolerance in *Arabidopsis*³⁵. These data suggest that the regulation of the expression of the *DREB1/CBF* genes might be more complex than previously thought.

NAC and ZF-HD regulon involved in ABA-independent gene expression

The *ERD1* (*early responsive to dehydration 1*) gene encoding a Clp (caseinolytic protease) protease regulatory subunit responds to dehydration and high salinity before the accumulation of ABA, suggesting the existence of an ABA-independent pathway in the dehydration stress response³¹. Analysis of the *ERD1* promoter identified two novel *cis*-acting elements that are involved in induction by dehydration stress⁴⁹. Base substitution analysis showed that a 14-bp *rps1*-like region (CACTAAAT-TGTCAC) and a CATGTG motif are necessary for the induction of the *ERD1* gene in dehydrated plants (Fig. 1). Recently, we isolated three cDNA clones encoding proteins that bind to the 63-bp promoter region of *ERD1*, which contains the CATGTG motif⁵² (Fig. 1). These three cDNA clones encode proteins which belong to the NAC transcription factor family including RD26. Microarray analysis of transgenic plants overexpressing the NAC genes revealed that several drought inducible genes were up-regulated in the transgenic plants, and the plants showed significantly increased drought tolerance. However, *ERD1* was not up-regulated in the transgenic plants. We recently isolated zinc-finger homeodomain (ZF-HD) transcription factors containing a homeodomain that can bind to the *rps1* site 1-like sequence using the yeast one-hybrid system. Overexpression of both NAC and ZF-HD proteins activated the expression of *ERD1* under unstressed normal growth conditions in the transgenic *Arabidopsis* plants.

AREB/ABF regulon involved in ABA-dependent gene expression

ABRE (*ABA*-responsive elements: ACGTGG/TC) is a major *cis*-acting element in ABA-responsive gene expression (Fig. 1). Two ABRE motifs are important in the ABA-responsive expression of the *Arabidopsis* gene *RD29B* encoding a LEA-like protein⁵³. The bZIP (basic leucine zipper) transcription factors ABRE-binding protein (AREB)/ABRE-binding factor (ABF) can bind to ABRE and activate ABA-dependent gene expression^{7,53}. Activation of the AREB1 and AREB2 proteins has been shown to require an ABA-mediated modification⁵³, which is probably ABA-dependent phosphorylation (Fig. 1). Overexpression of ABF3 or AREB2/ABF4 caused ABA hypersensitivity, reduced transpiration rate and enhanced drought tolerance of the transgenic plants¹⁸. The AREB1/ABF2 is reported to be an essential component of glucose signaling and its overexpression affects multiple stress tolerance including drought, salt and

heat²².

MYC and MYB regulon involved in ABA-dependent gene expression

The induction of the *Arabidopsis* drought-inducible gene *RD22* encoding a protein having a homology to an unidentified seed protein is mediated by ABA, and this gene requires protein biosynthesis for its ABA-dependent expression¹. A MYC transcription factor, AtMYC2 (*Arabidopsis thaliana* MYC 2), and a MYB transcription factor, AtMYB2 (*Arabidopsis thaliana* MYB 2), have been shown to bind *cis*-elements, MYCR (MYC-recognition site: CANNTG) and MYBR (MYB-recognition site: C/TAAACNA/G) in the *RD22* promoter and cooperatively activate *RD22*¹ (Fig. 1). These MYC and MYB proteins are synthesized after the accumulation of endogenous ABA, indicating that their role is in a late stage of the stress responses. Microarray analysis of MYC- and MYB-overexpressing transgenic plants revealed target genes for MYC and MYB, such as the alcohol dehydrogenase gene and ABA- or jasmonic-acid (JA)-inducible genes². Overexpression of both AtMYC2 and AtMYB2 not only caused an ABA-hypersensitive phenotype but also improved the osmotic-stress tolerance of the transgenic plants².

Recently, AtMYC2 transcription factors function as members of a MYC-based regulatory system conserved in dicotyledonous plants with a key role in JA-induced defense gene activation^{3,28}. These reports highlight the crosstalk between biotic stress signaling and abiotic stress signaling.

Crosstalk between the DREB regulon and the other regulons

Many drought- and cold-inducible genes contain both DRE/CRT and ABRE motifs in their promoters. These *cis*-acting elements are thought to function independently. However, precise analysis of these *cis*-acting elements in the *RD29A* gene expression revealed that DRE/CRT functions cooperatively with ABRE as a coupling element in ABA-responsive gene expression in response to drought stress³⁴. This indicates that there are interactions between the DREB regulon and the AREB/ABF regulon (Fig. 1).

Recently, an osmotic-stress inducible *CBF4/DREB1D* gene has been identified¹². Genes of the *DREB1/CBF* family are mainly induced by cold stress, but the drought-inducible gene *CBF4/DREB1D* functions to provide crosstalk between DREB2 and DREB1/CBF regulatory systems. The drought-inducible expres-

sion of *CBF4/DREB1D* is controlled by ABA-dependent pathways, suggesting that *CBF4/DREB1D* may function in the slow response to drought that relies on the accumulation of ABA (Fig. 1). Moreover, ABA induces the *DREB1/CBF* gene transcription and subsequent induction of cold-regulated genes via the DRE/CRT promoter element²⁴. A maize DRE-binding protein, DBF1, has been shown to function as a transcriptional activator of the *rab17* (*responsive to abscisic acid 17*) promoter by ABA²³. This also suggests the existence in some plants of an ABA-dependent pathway for the regulation of stress-inducible genes that involves DRE/CRT.

Gene expression in recovery process from abiotic stress in *Arabidopsis*

Microarray analysis has revealed many genes that respond to rehydration after drought stress, indicating

their involvement in the process of recovery from abiotic stress³⁶. The products of these genes are thought to function not only in recovery from stress but also in cell growth and elongation. The expression and function of the rehydration-inducible *ERD5* gene encoding a proline dehydrogenase (*ProDH*) gene has been precisely analyzed. This gene is involved in the degradation of the proline that accumulates during dehydration³². Promoter analysis of the *ProDH* gene revealed an important *cis*-acting element, ACTCAT, that is involved in rehydration-inducible gene expression⁴³. Many rehydration-inducible gene promoters contain the ACTCAT motif. Recently we showed that the ATB2 subgroup bZIP proteins functions as transcriptional activators in hypoosmolarity-responsive expression of the *ProDH* gene in *Arabidopsis*⁴⁴. The molecular information in the process of recovery from abiotic stress may allow us to improve the resilient plants.




Fig. 2. Collaboration for development of stress tolerant crops

CGIAR: Consultative Group on International Agricultural Research, CIAT: International Center for Tropical Agriculture (Colombia), CIMMYT: International Maize and Wheat Improvement Center (Mexico), ICARDA: International Center for Agricultural Research in the Dry Areas (Syrian Arab Republic), ICRISAT: International Crops Research Institute for the Semi-Arid Tropics (India), IRRI: International Rice Research Institute (The Philippines), JIRCAS: Japan International Research Center for Agricultural Sciences (Japan).

Table 1. Abiotic stress tolerance by regulon biotechnology

Transcription factor	Type	Gene source	Transgenic Species	Promoter	Tolerance	Reference
DREB1A/CBF3	AP2/ERF	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S, RD29A	Freezing, salt, and drought	11, 19, 26
			Wheat	RD29A	Drought	39
			Tobacco	35S, RD29A	Freezing and drought	20
			<i>Brassica napus</i>	35S	Freezing and drought	16, 59
DREB1B/CBF1	AP2/ERF	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Freezing	11, 17
			Tomato	35S, HVA22	Drought, chilling, and oxidative stress	13, 14, 25
			Strawberry	35S	Freezing	37
			<i>Brassica napus</i>	35S	Freezing and drought	16, 59
DREB1C/CBF2	AP2/ERF	<i>Arabidopsis</i>	<i>Brassica napus</i>	35S	Freezing and drought	16, 59
DREB1D/CBF4	AP2/ERF	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Freezing and drought	12
DREB1E/DDF1	AP2/ERF	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	High salinity	29
ZmDREB1	AP2/ERF	Maize	<i>Arabidopsis</i>	35S	Drought and freezing	40
OsDREB1A	AP2/ERF	Rice	<i>Arabidopsis</i>	35S	Drought, salt, and freezing	8
Tsi1	AP2/ERF	Tobacco	Tobacco	35S	Salt	38
JERF1	AP2/ERF	Tomato	Tobacco	35S	Salt	58
ICE1	HLH	<i>Arabidopsis</i>	<i>Arabidopsis</i>	Superpromoter*	Freezing	6
AtMYC2 & AtMYB2	MYC & MYB	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Osmotic stress	2
CpMYB10	MYB	<i>Craterostigma plantagineum</i>	<i>Arabidopsis</i>	35S	Drought and salt	55
Osmyb4	MYB	Rice	<i>Arabidopsis</i>	35S	Cold and freezing tolerance	54
AREB1/ABF2	bZIP	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought, salt, heat, and oxidative stress	22
AREB2/ABF4	bZIP	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought, salt, chilling, freezing, heat, and oxidative stress	18, 22
ABF3	bZIP	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought, salt, chilling, freezing, heat, and oxidative stress	18, 22
ABI5	bZIP	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Water stress	27
ABI3 (plus ABA)	VP1	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Freezing	51
SCOF-1	Zn finger	Soybean	<i>Arabidopsis</i>	35S	Low temperature stress	21
			Tobacco	35S	Low temperature stress	21
STZ	Zn finger	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought	41
ANAC019/ANAC	NAC	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought	52
ANAC055/ANAC3	NAC	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought	52
ANAC072/RD26	NAC	<i>Arabidopsis</i>	<i>Arabidopsis</i>	35S	Drought	52

* Superpromoter consists of three copies of the octopine synthase upstream-activating sequence in front of the manopine synthase promoter.

Application of regulon biotechnology to improve stress tolerance in crop plants

The orthologous genes of *DREB1/CBF* have been found in many crop plants such as canola, broccoli, tomato, alfalfa, wheat, barley, corn, and rice⁵⁹. These indicate that the *DREB1/CBF* regulon system is ubiquitous in the plant kingdom, and the “DREB technology” with controlling the expression of the *DREB1/CBF* regulon system is expected to improve the tolerance against stresses in crop plants. So far the *DREB1/CBF* genes of *Arabidopsis* have been successfully used to engineer abiotic stress tolerance in a number of different species (Table 1). For example, constitutive overexpression of the *Arabidopsis DREB1/CBF* genes in canola results in increased freezing tolerance¹⁶ and drought tolerance⁵⁹.

We have isolated rice orthologs for *DREB1/CBF* and *DREB2*, four *OsDREB1s* and one *OsDREB2*, in the rice genome sequence and they function in stress-inducible gene expression⁸. Overexpression of *OsDREB1A* in *Arabidopsis* revealed that this gene has a similar function to that of its *Arabidopsis* homolog in stress-responsive gene expression and stress tolerance. This indicates that similar transcription factors function in dicotyledons and monocotyledons. A novel *DREB1/CBF* transcription factor *ZmDREB1A* was also identified in *Zea mays*⁴⁰. The *ZmDREB1A* was involved in cold-responsive gene expression and overexpression of the *ZmDREB1A* gene in *Arabidopsis* which resulted in increased drought and freezing tolerance.

However, constitutive overexpression of the *DREB1/CBF* genes in plants showed dwarf phenotype^{11,26}. To overcome these problems, stress-inducible promoters that have low background expression under normal growth condition have been used in conjunction with the *DREB1/CBF* genes to achieve increased stress tolerance without the growth retardation^{19,25}. Constitutive overexpression of *Arabidopsis DREB1A/CBF3* improved drought- and low-temperature stress tolerance in tobacco²⁰. The stress-inducible *RD29A* promoter minimized the negative effects on the plant growth in tobacco. Furthermore, we detected overexpression of stress-inducible target genes of *DREB1A/CBF3* in tobacco. The *Arabidopsis DREB1A/CBF3* gene was placed under control of the *RD29A* promoter and transferred via biolistic transformation into bread wheat³⁹. Plants expressing the *DREB1A/CBF3* gene demonstrated substantial resistance to water stress in comparison through checks under experimental greenhouse conditions, manifested by a 10-day delay in wilting when water was withheld. These results indicate that a combination of the *RD29A* promoter and *DREB1A* is use-

ful for improvement of various kinds of transgenic plants that are tolerant to environmental stress.

Now we collaborate with many research groups to improve stress tolerant crop plants utilizing regulon biotechnology (Fig. 2). We hope the results of these collaborative studies will contribute to the sustainable food production in developing countries and help to prevent the global-scale environmental damage.

References

1. Abe, H. et al. (1997) Role of *Arabidopsis* MYC and MYB homologs in drought- and abscisic acid-regulated gene expression. *Plant Cell*, **9**, 1859–1868.
2. Abe, H. et al. (2003) *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *Plant Cell*, **15**, 63–78.
3. Boter, M. et al. (2004) Conserved MYC transcription factors play a key role in jasmonate signaling both in tomato and *Arabidopsis*. *Genes Dev.*, **18**, 1577–1591.
4. Bray, E. A. (1997) Plant responses to water deficit. *Trends Plant Sci.*, **2**, 48–54.
5. Chen, T. H. & Murata, N. (2002) Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Curr. Opin. Plant Biol.*, **5**, 250–257.
6. Chinnusamy, V. et al. (2003) ICE1: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*. *Genes Dev.*, **17**, 1043–1054.
7. Choi, H. -I. et al. (2000) ABFs, a family of ABA-responsive element binding factors. *J. Biol. Chem.*, **275**, 1723–1730.
8. Dubouzet, J. G. et al. (2003) *OsDREB* genes in rice, *Oryza sativa* L., encode transcription activators that function in drought-, high salt- and cold- responsive gene expression. *Plant J.*, **33**, 751–763.
9. Fowler, S. & Thomashow, M. F. (2002) *Arabidopsis* transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell*, **14**, 1675–1690.
10. Gilmour, S. J. et al. (1998) Low temperature regulation of *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced *COR* gene expression. *Plant J.*, **16**, 433–442.
11. Gilmour, S. J. et al. (2000) Overexpression of the *Arabidopsis CBF3* transcriptional activator mimics multiple biochemical changes associated with cold acclimation. *Plant Physiol.*, **124**, 1854–1865.
12. Haake, V. et al. (2002) Transcription factor CBF4 is a regulator of drought adaptation in *Arabidopsis*. *Plant Physiol.*, **130**, 639–648.
13. Hsieh, T. H. et al. (2002) Tomato plants ectopically expressing *Arabidopsis* CBF1 show enhanced resistance to water deficit stress. *Plant Physiol.*, **130**, 618–626.
14. Hsieh, T. H. et al. (2002) Heterology expression of the *Arabidopsis C-repeat/dehydration response element binding factor 1* gene confers elevated tolerance to chilling and oxidative stresses in transgenic tomato. *Plant*

- Physiol.*, **129**, 1086–1094.
15. Ingram, J. & Bartels, D. (1996) The molecular basis of dehydration tolerance in plants. *Ann. Rev. Plant Physiol. Plant Mol. Biol.*, **47**, 377–403.
16. Jaglo, K. R. et al. (2001) Components of the Arabidopsis C-repeat/dehydration-responsive element binding factor cold-response pathway are conserved in *Brassica napus* and other plant species. *Plant Physiol.*, **127**, 910–917.
17. Jaglo-Ottosen, K. R. et al. (1998) *Arabidopsis* CBF1 overexpression induces *COR* genes and enhances freezing tolerance. *Science*, **280**, 104–106.
18. Kang, J. Y. et al. (2002) Arabidopsis basic leucine zipper proteins that mediate stress-responsive abscisic acid signaling. *Plant Cell*, **14**, 343–357.
19. Kasuga, M. et al. (1999) Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nat. Biotechnol.*, **17**, 287–291.
20. Kasuga, M. et al. (2004) A combination of the *Arabidopsis* DREB1A gene and stress-inducible *rd29A* promoter improved drought- and low-temperature stress tolerance in tobacco by gene transfer. *Plant Cell Physiol.*, **45**, 346–350.
21. Kim, J. C. et al. (2001) A novel cold-inducible zinc finger protein from soybean, SCOF-1, enhances cold tolerance in transgenic plants. *Plant J.*, **25**, 247–259.
22. Kim, S. et al. (2004) ABF2, an ABRE-binding bZIP factor, is an essential component of glucose signaling and its overexpression affects multiple stress tolerance. *Plant J.*, **40**, 75–87.
23. Kizis, D. & Pages, M. (2002) Maize DRE-binding proteins DBF1 and DBF2 are involved in *rab17* regulation through the drought-responsive element in an ABA-dependent pathway. *Plant J.*, **30**, 679–689.
24. Knight, H. et al. (2004) Abscisic acid induces *CBF* gene transcription and subsequent induction of cold-regulated genes via the CRT promoter element. *Plant Physiol.*, **135**, 1710–1717.
25. Lee, J. T. et al. (2003) Expression of Arabidopsis CBF1 regulated by an ABA/stress inducible promoter in transgenic tomato confers stress tolerance without affecting yield. *Plant Cell Environ.*, **26**, 1181–1190.
26. Liu, Q. et al. (1998) Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in Arabidopsis. *Plant Cell*, **10**, 1391–1406.
27. Lopez-Molina, L., Mongrand, S. & Chua, N. -H. (2001) A postgermination developmental arrest checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in Arabidopsis. *Proc. Natl. Acad. Sci. USA*, **98**, 4782–4787.
28. Lorenzo, O. et al. (2004) *JASMONATE-INSENSITIVE1* encodes a MYC transcription factor essential to discriminate between different jasmonate-regulated defense responses in Arabidopsis. *Plant Cell*, **16**, 1938–1950.
29. Magome, H. et al. (2004) *dwarf* and *delayed-flowering 1*, a novel Arabidopsis mutant deficient in gibberellin biosynthesis because of overexpression of a putative AP2 transcription factor. *Plant J.*, **37**, 720–729.
30. Maruyama, K. et al. (2004) Identification of cold-inducible downstream genes of the Arabidopsis DREB1A/CBF3 transcriptional factor using two microarray systems. *Plant J.*, **38**, 982–993.
31. Nakashima, K. et al. (1997) A nuclear gene, *erd1*, encoding a chloroplast-targeted Clp protease regulatory subunit homolog is not only induced by water stress but also developmentally up-regulated during senescence in Arabidopsis thaliana. *Plant J.*, **12**, 851–861.
32. Nakashima, K. et al. (1998) A gene encoding proline dehydrogenase is not only induced by proline and hypoosmolarity, but is also developmentally regulated in the reproductive organs of Arabidopsis. *Plant Physiol.*, **118**, 1233–1241.
33. Nakashima, K. & Yamaguchi-Shinozaki, K. (2002) Use of β -glucuronidase (GUS) to show dehydration and high-salt gene expression. In Molecular methods of plant analysis, Vol. 22, eds. Jackson, J. F. & Linskens, H. F., Springer-Verlag Berlin, Germany, 37–61.
34. Narusaka, Y. et al. (2003) Interaction between two cis-acting elements, ABRE and DRE, in ABA-dependent expression of Arabidopsis *rd29A* gene in response to dehydration and high-salinity stresses. *Plant J.*, **34**, 137–148.
35. Novillo, F. et al. (2004) CBF2/DREB1C is a negative regulator of CBF1/DREB1B and CBF3/DREB1A expression and plays a central role in stress tolerance in Arabidopsis. *Proc. Natl. Acad. Sci. USA*, **101**, 3985–3990.
36. Oono, Y. et al. (2003) Monitoring expression profiles of Arabidopsis gene expression during rehydration process after dehydration using ca. 7000 full-length cDNA microarray. *Plant J.*, **34**, 868–887.
37. Owens, C. L. et al. (2002) CBF1 orthologs in sour cherry and strawberry and the heterologous expression of CBF1 in strawberry. *J. Am. Soc. Hortic. Sci.*, **127**, 489–494.
38. Park, J. M. et al. (2001) Overexpression of the tobacco *Tsi1* gene encoding an EREBP/AP2-type transcription factor enhances resistance against pathogen attack and osmotic stress in tobacco. *Plant Cell*, **13**, 1035–1046.
39. Pellegrineschi, A. et al. (2004) Stress-induced expression in wheat of the Arabidopsis thaliana DREB1A gene delays water stress symptoms under greenhouse conditions. *Genome*, **47**, 493–500.
40. Qin, F. et al. (2004) Cloning and functional analysis of a novel DREB1/CBF transcription factor involved in cold-responsive gene expression in Zea mays L. *Plant Cell Physiol.*, **45**, 1042–1052.
41. Sakamoto, H. et al. (2004) Arabidopsis Cys2/His2-type zinc-finger proteins function as transcription repressors under drought, cold, and high-salinity stress conditions. *Plant Physiol.*, **136**, 2734–2746.
42. Sakuma, Y. et al. (2002) DNA-binding specificity of the ERF/AP2 domain of Arabidopsis DREBs, transcription factors involved in dehydration- and cold-inducible gene expression. *Biochem. Biophys. Res. Com.*, **290**, 998–1009.
43. Satoh, R. et al. (2002) ACTCAT, a novel cis-acting element for proline- and hypoosmolarity-responsive expression of the *ProDH* gene encoding proline dehydrogenase in Arabidopsis. *Plant Physiol.*, **130**, 709–719.
44. Satoh, R. et al. (2004) A novel subgroup of bZIP proteins

- functions as transcriptional activators in hypoosmolarity-responsive expression of the *ProDH* gene in *Arabidopsis*. *Plant Cell Physiol.*, **45**, 309–317.
45. Seki, M. et al. (2001) Monitoring the expression pattern of 1300 *Arabidopsis* genes under drought and cold stresses by using a full-length cDNA microarray. *Plant Cell*, **13**, 61–72.
 46. Seki, M. et al. (2002) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold, and high-salinity stresses using a full-length cDNA microarray. *Plant J.*, **31**, 279–292.
 47. Shinozaki, K. & Yamaguchi-Shinozaki, K. (1997) Gene expression and signal transduction in water-stress response. *Plant Physiol.*, **115**, 327–334.
 48. Shinozaki, K., Yamaguchi-Shinozaki, K. & Seki, M. (2003) Regulatory network of gene expression in the drought and cold stress responses. *Curr. Opin. Plant Biol.*, **6**, 410–417.
 49. Simpson, S. D. et al. (2003) Two different novel cis-acting elements of *erd1*, a *clpA* homologous *Arabidopsis* gene function in induction by dehydration stress and dark-induced senescence. *Plant J.*, **33**, 259–270.
 50. Stockinger, E. J., Gilmour, S. J. & Thomashow, M. F. (1997) *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc. Natl. Acad. Sci. USA*, **94**, 1035–1040.
 51. Tamminen, I. et al. (2001) Ectopic expression of *ABI3* gene enhances freezing tolerance in response to abscisic acid and low temperature in *Arabidopsis thaliana*. *Plant J.*, **25**, 1–8.
 52. Tran, L. -S. P. et al. (2004) Isolation and functional analysis of *Arabidopsis* stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the *early responsive to dehydration stress 1* promoter. *Plant Cell*, **16**, 2481–2498.
 53. Uno, Y. et al. (2000) *Arabidopsis* basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proc. Natl. Acad. Sci. USA*, **97**, 11632–11637.
 54. Vannini, C. et al. (2004) Overexpression of the rice *Osmby4* gene increases chilling and freezing tolerance of *Arabidopsis thaliana* plants. *Plant J.*, **37**, 115–127.
 55. Villalobos, M. A., Bartels, D. & Iturriaga, G. (2004) Stress tolerance and glucose insensitive phenotypes in *Arabidopsis* overexpressing the *CpMYB10* transcription factor gene. *Plant Physiol.*, **135**, 309–324.
 56. Yamaguchi-Shinozaki, K. & Shinozaki, K. (1994) A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low- temperature, or high-salt stress. *Plant Cell*, **6**, 251–264.
 57. Zarka, D. G. et al. (2003) Cold induction of *Arabidopsis* *CBF* genes involves multiple ICE (inducer of *CBF* expression) promoter elements and a cold-regulatory circuit that is desensitized by low temperature. *Plant Physiol.*, **133**, 910–918.
 58. Zhang, H. et al. (2004) The ethylene-, jasmonate-, abscisic acid- and NaCl-responsive tomato transcription factor JERF1 modulates expression of GCC box-containing genes and salt tolerance in tobacco. *Planta*, **220**, 262–270.
 59. Zhang, J. Z. et al. (2004) From laboratory to field. Using information from *Arabidopsis* to engineer salt, cold, and drought tolerance in crops. *Plant Physiol.*, **135**, 615–621.
 60. Zhu, J. K. (2002) Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.*, **53**, 247–273.