China

June 2013 Vol.58 No.18: 2097–2103 doi: 10.1007/s11434-012-5624-y

Summary of phylogeny in subfamily Colobinae (Primate: Cercopithecidae)

WANG XiaoPing¹, ZHANG YaPing^{1,3} & YU Li^{1,2*}

¹ Laboratory for Conservation and Utilization of Bio-resource, Yunnan University, Kunming 650091, China;

² Key Laboratory for Animal Genetic Diversity and Evolution of High Education in Yunnan Province, Yunnan University, Kunming 650091,

³ State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China

Received March 12, 2012; accepted May 25, 2012; published online January 9, 2013

The phylogeny of the colobine monkeys has a complex evolutionary history, evolving several distinct radiations and owned a wide range of forest and woodland habits in tropical Africa and in southern and eastern Asia. Understanding the true evolutionary history of the colobine monkeys becomes an important field and has received special attention. In this article, we review phylogeny of Colobinae based on previous fossil, cytological, morphological and molecular evidence and indicate the phylogenetic controversies. This review is expected to guide the future research of Colobinae phylogeny, and also provide theoretic evidence for the conservation of these highly endangered and unique primates.

Colobinae, phylogeny, Asian presbytina, African colobinae

Citation: Wang X P, Zhang Y P, Yu L. Summary of phylogeny in subfamily Colobinae (Primate: Cercopithecidae). Chin Sci Bull, 2013, 58: 2097–2103, doi: 10.1007/s11434-012-5624-y

The Old World monkeys consist of two living subfamilies —the cheek-pouch monkeys (Cercopithecinae) and the leafeating monkeys (Colobinae) [1], both of which are the closest living relatives to the apes (Hominoidea) (Figure 1) [1], but previous studies had focused on Cercopithecine evolution as a model for human evolution history. The phylogenetic and evolution of the leaf-eating monkeys (Colo-





^{*}Corresponding author (email: yuli1220@yahoo.com.cn)

binae), on the other hand, had been a relatively neglected issue. In these years, the colobine monkeys, especially Asian colobines, are facing extinction because of the increasing deforestation and hunting rates by people. Therefore, it is necessary for us to promote the conservation of these highly endangered and unique primates.

Compared with Cercopithecinae, the colobine monkeys have a sharp-cusped cheek teeth and relatively narrow incisors [2]. Their dimensions vary from 4 kg in the African olive colobus (*Procolobus verus*) to 20 kg of the adult males of the proboscis monkey in Borneo (*Nasalis larvatus*). They are characterized by the very short and in some case degraded thumb. The hind legs are usually longer than the front legs. Most of the species have no cheek pouch outside, and have very long tail except the pig-tailed langur (*Simias concolor*) [3].

Besides the features as described above, the subfamily Colobinae has evolved a complex ruminant-like stomach and foregut in which microbes ferment these indigestible plant materials [4], leading to their unique adaptation to the

[©] The Author(s) 2013. This article is published with open access at Springerlink.com

leaf-eating feeding habits [5–8]. In view of this, the Colobinae has also been the crucial model animals for the study of the molecular mechanisms of adaptive evolution [9,10]. The pancreatic ribonuclease gene (*RNASE*) duplication has been believed to be closely related to the adaptation of Colobines to the unique leaf-feeding lifestyle [5–8]. While all of the other primates have only one *RNASE1* gene, multiple *RNASE1* genes have been intriguingly observed in Colobines, evolving rapidly under positive selection for enhanced digestive efficiencies, as an adaptive response to the increased demands for the enzyme for digesting bacterial RNA (Figure 2).

The colobines consist of 10 genera in two subtribes- the African Colobina (including the genera Colobus, Piliocolobus, and Procolobus) and the Asian Presbytina (including the genera Pygathrix, Rhinopithecus, Nasalis, Simias, Presbytis, Trachypithecus and Semnopithecus) [1,11–14]. Based on fossil records, the divergence between ancestral African colobines and Asian colobines was estimated between 10 [15] and 13 Ma [16]. The colobines not only experience recent and rapid adaptive radiations, but also are more diverse and widespread for different climate environments [12]. In view of this, the phylogeny of the colobine monkeys has a complex evolutionary history, evolving several distinct radiations and owned a wide range of forest and woodland habits in tropical Africa and in southern and eastern Asia. Understanding the true evolutionary history of the colobine monkeys, therefore, becomes an important field and has received special attention. However, due to the fact that nearly all the extant colobine genera diversify from one another within a four-million-year window [17-19], attempts to clarify relationships among these colobine genera have encountered challenges.

In this article, we review the phylogeny of the colobine monkeys based on previous fossil, morphological physiol-



Figure 2 The unique duplication of *RNASE1* gene in Colobines compared with other primates [5].

ogy, cytology and molecular evidences, and indicated the phylogenetic controversies. This review is expected to shed light on the research of Colobinae phylogeny and adaptive evolution, and also provide a theoretical foundation for the conservation of these highly endangered and unique primates.

1 Phylogeny of African Colobinae

The diversification of African colobines, which distributed across the African rainforest belt, represented a major component of African primate evolution [20,21]. According to fossil records, the earliest African colobines were found in the late Miocene about 8.5–9 Ma [22].

African colobines include three genera, i.e., *Colobus* (black-and-white colobus), *Procolobus* (olive colobus), and *Piliocolobus* (red colobus) [20,23]. *Colobus* diverged first, and followed by the progenitor of *Piliocolobus* and *Procolobus* [17,19,24–26]. Based on both morphology and genetics evidence, it was well-recognized that *Procolobus* was the sister-taxon of (and possibly congeneric to) *Piliocolobus* [12,19,24,26–28]. In other words, there was a general consensus for the more close relationship between *Piliocolobus* and *Procolobus* and *Procolobus* within African colobines.

In earlier analyses of morphological and molecular data, all supported that the African colobine forms a monophyletic clade [1,12,19], grouping Piliocolobus, Procolobus, and Colobus together [17-19,24-26,29,30] (Figure 3). Interestingly, a recent phylogenetic analyses of 83 mobile elements from Roos et al. [19] indicated a closer association of the Procolobus/Piliocolobus clade to Asian colobines than to Colobus, a relationship that was not rejected by nuclear sequence data in their study, leading them to propose African colobines paraphyly hypothesis (Figure 3(e)). This finding thus challenged the traditional view of the monophyly of African colobines. An ancient hybridization between Colobus male and Procolobus/Piliocolobus females had been proposed by them to explain their new finding. However, a most recent study of Wang et al. [31] supported the traditional view of the monophyly of the African colobines based on 44 nuclear non-coding genes and the whole mt genome (Figure 3(d) and (g)).

2 Phylogeny of Asian Presbytina

Compared with African colobinae, Asian Presbytina has more diverse members and wider geographic distribution [1,12]. The classification of Asian genera varies from 3 to 9 genera based on previous studies. Groves [32] and Grzimek [33] classified Asian colobines into three genera: *Presbytis*, *Pygathrix*, and *Nasalis*; Elliot [34] divided them into 4 genera: *Pygathrix*, *Rhinopithecus*, *Nasalis*, and *Simias*. Macdonald [35] also recognized 4 genera, but referred as *Presbytis*, *Semnopithecus*, *Pygathrix*, and *Nasalis*. In comparison,



Figure 3 Hypotheses of phylogenetic relationships among Colobine genera. Trees were reconstructed based on (a) 12 protein-coding mt genes (10 kb) [17]; (b) fragment of X-chromosome (4.3 kb) [26] and 54 nuclear genes (35 kb) [18]; (c) complete *cytb* gene (1.8 kb) [30] and 7 mt genes (6.1 kb) [29]; (d) 15 mt genes and 43 nuclear genes (42 kb) [25], 44 nuclear non-coding genes (23 kb) [31]; (e) 83 mobile elements [19]; (f) nuclear genes (13 kb) [19]; (g) complete mt genomes (16.5 kb) [31].

Napier and Napier [36], Simpson [37], and Ellerman et al. [38] grouped the Asian colobines into five genera: Presbytis, Pygathrix, Rhinopithecus, Nasalis, and Simias, while Hershkovitz [39] and Nowak [40] proposed seven genera: Presbytis, Semnopithecus, Trachypithecus, Pygathrix, Rhinopithecus, Nasalis, and Simias. Hill [41-43] classified them into 9 genera: Presbytis, Semnopithecus, Trachypithecus, Pygathrix, Rhinopithecus, Nasalis, Simias, Kasi and Prebyticus. In recent years, there was a consensus that the Asian colobines were referred as langurs and comprised of 7 genera, i.e., Semnopithecus, Trachypithecus, Presbytis, Pygathrix, Rhinopithecus, Nasalis and Simias [1,12,14, 44,45]. In addition, these 7 genera were clustered into 2 groups, i.e., the odd-nosed monkey group (including Pygathrix, Rhinopithecus, Nasalis and Simias) and the langur group (including Presbytis, Trachypithecus and Semnopithecus) [46-48].

Asian colobines represent a typical example of an evolutionary radiation with rapid diversification events that date back to Middle Miocene about 10–15 Ma [11,12]. Therefore, phylogenetic relationships within Asian colobines, including those among genera and some species had been disputed hotly in previous studies and were not well established yet [1,17–19,25,26,29–31,49–59].

2.1 Intergeneric relationships within Asian Presbytina

(i) Phylogeny of the odd-nosed monkey group. Though

the monophyly of the odd-nosed monkey group has been unambiguously established from previous studies of both morphology and geographical distribution [1,11,12,14,60] and confirmed by genetic data [17–19,26,30,31,52], there had long been controversy over the relationships among the genera within this group. It was especially the case that among *Pygathrix, Rhinopithecus* and *Nasalis* (Figure 3) [17–19,26,30,31,48].

Based on the previous morphological studies, Grove [46] and Delson [61] favored the association of *Pygathrix* and Rhinopithecus to the exclusion of Nasalis and this was congruent with molecular evidences based on the analyses of mitochondrial (mt) ND3-ND4 (2.2 kb), 12SrRNA (387 bp) genes by Wang et al. [49], and Li et al. [51], whereas Jablonski et al. [47,48,55] grouped Nasalis with Pygathrix to the exclusion of Rhinopithecus by combining fossils and 455 characters of morphological data. Zhang et al. [50] analyzed mt cytb gene fragment (424 bp) and support the sister-grouping of Nasalis and Rhinopithecus to the exclusion of Pygathrix. More recently, further efforts had been undertaken to obtain phylogenetic relationships using large datasets from mt genomes and nuclear genes. In the studies of Chatterjee et al. [29] (6.1 kb), Fabre et al. [25] (42 kb) and Meyer et al. [30] (1.8 kb), as well as Wang et al. [31] (23 kb), all of which were conducted based on mt or nuclear sequence data, a sister-taxon association between Pygathrix and *Rhinopithecus* were consistently indicated (Figure 3(c) and (d)), whereas the studies of Roos et al. [19] and Wang et al. [31] supported a close association of *Pygathrix* with *Nasalis* to the exclusion of *Rhinopithecus* based on both 83 mobile elements (Figure 1(e)) and nuclear genes (13 kb) (Figure 3(f)) as well as the whole mt genomes (16.5 kb) (Figure 3(g)). Furthermore, Sterner et al. [17], Ting et al. [26] and Perelman et al. [18] also attempted to resolve the relationships among *Pygathrix*, *Rhinopithecus*, and *Nasalis* within the odd-nosed monkeys based on mt protein-coding genes (10 kb) (Figure 3(a)), X-chromosome (4.3 kb), and nuclear genes (35 kb) (Figure 3(b)), respectively, but the three genera formed a polytomy.

In comparison, when it came to the phylogenetic position of *Simias* in the odd-nosed monkey group, the debate on its systematic placement appeared remarkably subdued because all previous morphological and molecular analyses supported it as the sister-taxon (and possible subgenus) to *Nasalis* [19,61,62], with the exception of a recent study of 7 mitochondrial (mt) gene fragments (6.1 kb) from Chatterjee et al., [29] in which *Simias* had a closer relationship with *Pygathrix/Rhinopithecus* than *Nasalis*.

(ii) Phylogeny of the langur group. Another area being in hot dispute comes from the relationships between the langur genera and the odd-nosed monkey group. In fact, there has long been debate about the determination of three genera within the langur monkey group in earlier studies. Semnopithecus, and Trachypithecus had been ever grouped into Presbytis as one genus [46,63-66]. In comparison, more evidence separated Trachypithecus from Presbytis, including those from Oate et al. [67] and Brandon-Jones et al. [14] based on ecological and morphological data, as well as Md-Zain etc. based on mt ND3, ND4L, ND4 (2.3 kb) [68] and COII (850 bp) [69] genes. Hill [43] and Hooijer [70] subdivide the genus Presbytis into Semnopithecus and Trachypithecus subgenus. In addition, Trachypithecus had also been formerly considered as a subgenus of Semno*pithecus* [16,27,71,72].

Recent studies provided contradicting relationships between these three langur genera and the odd-nosed monkey group (Figure 3). Growing studies are supporting the nonmonophyly of the langur monkey group.

Sterner et al. [17] suggested a sister-group between *Presbytis* and *Trachypithecus* within the langur group, but failed to resolve the relationships among *Presbytis/Trachypithecus, Semnopithecus*, and the odd-nosed monkey group based on 12 mt protein-coding genes (10 kb) of 6 Asian colobine genera (Figure 3(a)). Ting et al. [26] claimed that *Presbytis* diverged earliest, followed by the split between *Trachypithecus/Semnopithecus* and the odd-nosed monkey group within the Asian colobines by examining a fragment of the X-chromosome (4.3 kb) (Figure 3(b)). The results were congruent with the study by Perelman et al. [18] including ~35 kb genome sequences from 54 nuclear genes of 186 primates. In contrast, both Chatterjee et al. [29] and Meyer et al. [30] favored the earliest divergence of *Presbytis/Trachypithecus* within Asian colobi-

nae, and the close relatedness of Semnopithecus and the odd-nosed monkey group based on the analyses of 7 mt genes (6.1 kb) and a fragment of mt genes (1.8 kb) (Figure 3(c)), but they lacked significant supports. Fabre et al. [25] analyzing combined mt and nuclear genes (42 kb), and Wang et al. [31] analyzing 44 nuclear non-coding genes (23 kb), both found support for close relationships between Trachypithecus and Semnopithecus, and between Presbytis and the odd-nosed monkeys (Figure 3(d)). These same relationships were also confirmed by Roos et al. [19], who performed an analysis of 83 mobile elements (Figure 3(e)). Moreover, they also supported the former Semnopithecus/ Trachypithecus clade, but suggested Presbytis as sister to the other Asian colobines based on nuclear genome data analyses (13 kb) (Figure 3(f)). However, by analyzing the whole mt genome (16.5 kb), Wang et al. [31] proposed that Semnopithecus diverged earliest, followed by the split between Trachypithecus/Presbytis and the odd-nosed monkey group within the Asian colobines (Figure 3(g)).

The results from the above studies demonstrated that the relationships among langur monkey group and odd-nosed monkey group remain unresolved, although hybridization had been raised for a most likely explanation for some of these incongruent relationships [19,73].

2.2 Interspecies relationships within Asian Presbytina

(i) Interspecific relationships in genus *Rhinopithecus*. The snub-nosed monkey genus *Rhinopithecus* is comprised of four distinct allopatric species [1,74]: *R. brelichi* (the gray snub-nosed monkey), *R. bieti* (the black snub-nosed monkey), *R. roxellana* (the golden snub-nosed monkey) and *R. avunculus* (the Tonkin snub-nosed monkey). Except for *R. avunculus*, which is distributed in low-altitude subtropical forests in northwestern Vietnam [75], the other three *Rhinopithecus* species are endemic to temperate areas of China and inhabit six isolated high-altitude mountainous regions [76].

The phylogenetic relationships among the four Rhinopithecus species had all along been focusing issues. As seen from Figure 4, contradictory conclusions had been reached under different analyses. Zhang and Ryder [77] and Wang et al. [31] supported the sister group of *R. avunculus* and *R. bieti* to the exclusion of *R. roxellana* by analyzing mt *cytb* genes (253 bp) and 44 nuclear non-coding genes (23 kb), respectively (Figure 4(a)). Intriguingly, both Zhang and Ryder [50] and Li et al. [51] failed to resolve the phylogenetic relationships among R. bieti, R. roxellana and R. avunculus based on mt tRNA^{Thr}/cytb genes (424 bp) (Figure 4b) and mt cytb genes (402 bp) (Figure 4(c)), respectively, but R. brelichi was supported to diverge earlier than the other three species in the latter study (Figure 4(c)). Interstingly, different from Zhang et al. [50], in which mt tRNA^{Thr}/cytb genes were analyzed using MP and ML methods, Whittaker et al. [62] analyzed these two genes



Figure 4 Hypotheses of phylogenetic relationships among four species of the genus *Rhinopithecus*. Trees were reconstructed based on (a) fragment of mt *cytb* gene (253 bp) [77], 44 nuclear non-coding genes (23 kb) [31]; (b) fragment of mt tRNA^{Thr}/*cytb* gene (424 bp) [50]; (c) fragment of mt *cytb* gene (402 bp), mt 12S rRNA gene (387 bp) [51]; (d) fragment of mt tRNA^{Thr}/*cybt* gene (424 bp) [62]; (e) 15 mt genes and 43 nuclear genes (42 kb) [25], 7 mt genes (6,1 bp) [29]; (f) complete mt genomes (16.5 kb) [31].

using Bayesian method and supported the close relationship between *R. roxellana* and *R. avunculus* to the exclusion of *R. bieti* (Figure 4(d)). This relationship was also supported by Fabre et al. [25] and Chatterjee et al. [29], which utilized 15 mt and 43 nuclear genes (42 kb) (Figure 4(e)), as well as 7 mt fragment (6.1 kb) (Figure 4(e)), respectively. Interestingly, both Fabre et al. [25] and Chatterjee et al. [29] claimed the earliest divergent species among the genus *Rhinopithecus* group was *R. bieti* with significant support (Figure 4(e)), which was different from Li et al. [51] (Figure 4(c)). Recently, using the whole mt genome (16.5 kb), Wang et al. [31] supported the sister relationship between *R. roxellana* and *R. bieti* to the exclusion of *R. avunculus* (Figure 4(f)).

(ii) Interspecific relationships in genus Trachypithecus. Trachypithecus is the largest and most diverse genus of Asian Presbytina with a widest distribution, ranging from South India and Sri Lanka through mainland Southeast Asia to the Sundaland [52]. Within genus Trachypithecus, one species that had long been considered hard to determine phylogenetically is T. pileatus (the capped langur). Up to now, there were two conflicting hypotheses based upon previous analyses of mt cytb gene and two nuclear genes concerning the phylogenetic position of T. pileatus. Studies of partial mt genes clustered it with Semnopithecus [29,52,78,79]. However, nuclear gene based on Y chromosomal data and retroposon integrations analyses grouped it with Trachypithecus [52]. The remarkable discrepancy between mt and nuclear gene trees had led Osterholz et al. [52] and Karanth [80] to predict that T. pileatus might be the result of an ancient hybridization event between Semnopithecus and Trachypithecus. Additional information from mt and nuclear genomes should be added to resolve the enigmatic phylogenetic position of T. pileatus in the future studies.

Besides *T. pileatus*, the classifications and phylogenetic positions of *T. johnii* and *T. vetulus* had also been subjects of controversies in previous studies. In earlier studies, they were advised to be placed in their own genus *Kasi* due to their significant distinction from other species [42]. The subsequent analyses from the characters of cranial mor-

phology, neonatal pelage color and sexually dichromatic pubic integument all classified them within *Trachypithecus* [14,40,45], whereas the growing molecular data clustered them within *Semnopithecus* [18,29,31,50,52,62,79], leading to the proposal that they should be reclassified to be *S. johnii* and *S. vetulus*.

3 Perspectives

The phylogeny of colobine monkeys, which are characterized by rapid species radiations and short internal tree branches, has long been one of the most plaguing and challenging problems in species tree reconstruction [81–83]. Up to now, only the monophyly of the odd-nosed monkey group had been unambiguously accepted. Main controversies were concentrated on (1) the paraphyletic or monophyletic origin of African colobina, (2) the intergeneric relationships among the Asian colobines genus, (3) the interspecific relationships within the odd-nosed monkeys group, and (4) the precise placement of *T. pileatus*.

Therefore, finding valid genetic markers that offer sufficient variation to distinguish among the recently divergent Colobinae species posed a major challenge to advancing the understanding of Colobine phylogeny. Earlier investigations of Colobinae phylogeny mainly utilized analysis of portions of a single or a small number of mt genes [49–52]. Recently, more efforts had been made to obtain the phylogenetic tree using large dataset from mt genomes and multiple nuclear genes [17–19,25,26,29–31]. However, many relationships within Colobinae remained a polytomy, evidence from additional characters were thus necessary to resolve them in the future study.

As the increasing availability of genomic data of more species, phylogenetic analysis is entering a new era—Phylogenomics, which use genomic data to infer evolutionary relationships [84]. By performing phylogenomic study, we are most likely to reconstruct a reliable phylogenetic tree using much more characters than those in previous studies, including gene families, large insertion and deletion gene fragments, and gene rearrangement [85–87]. These characters might be useful for distinguishing nodes resulting from rapid radiation episodes such as the Asian colobinae speciation events.

This work was supported by National Natural Science Foundation of China (U0836603) and Program for New Century Excellent Talents in University (NCET).

- 1 Groves C P. Primate Taxonomy. Washington: Smithsonian Institution Press, 2001
- 2 Fleagle J G. Old World Monkeys. In: Fleagle J G, ed. Primate Adaptation and Evolution. New York: Academic Press, 1998. 185–218
- 3 Oates J F, Davies A G, Delson E. What are the colobines? In: Davies A G, Oates J F, eds. Colobine Monkeys—Their Ecology, Behavior, and Evolution. Cambridge: Cambridge University Press, 1994. 1–10
- 4 Chivers D J, Hladik C M. Morphology of the gastrointestinal tract in primates: Comparisons with other mammals in relation to diet. J Morphol, 1980, 166: 337–386
- 5 Zhang J, Zhang Y P, Rosenberg H F. Adaptive evolution of a duplicated pancreatic ribonuclease gene in a leaf-eating monkey. Nat Genet, 2002, 30: 411–415
- 6 Zhang J. Parallel adaptive origins of digestive RNases in Asian and African leaf monkeys. Nat Genet, 2006, 38: 819–823
- 7 Yu L, Wang X Y, Jin W, et al. Adaptive evolution of digestive *RNASE1* genes in leaf-eating monkeys revisited: New insights from ten additional colobines. Mol Biol Evol, 2010, 27: 121–131
- 8 Wang X Y, Li N Z, Yu L, et al. Duplication and functional diversification of pancreatic ribonuclease (*RNASE1*) gene. Chin Sci Bull, 2010, 55: 2–6
- 9 Wang X P, Jin W, Yu L, et al. Molecular evolution of stress-response gene *Leptin* in high-altitude Chinese snub-nosed monkeys (*Rhiopithecus genus*). Chin Sci Bull, 2010, 55: 4132–4135
- 10 Yu L, Wang X P, Ting N, et al. Mitogenomic analysis of Chinese snub-nosed monkeys: Evidence of positive selection in NADH dehydrogenase genes in high-altitude adaptation. Mitochondrion, 2011, 11: 497–503
- 11 Groves C P. A Theory of Human and Primate Evolution. Oxford: Oxford University Press, 1989
- 12 Davies A G, Oates J F. Colobine Monkeys: Their Ecology, Behavior, and Evolution. Cambridge: Cambridge University Press, 1994
- 13 Disotell T R. Primates: Phylogenetics. Encyclopedia of the Human Genome. London: Nature Publishing Group, 2003
- 14 Brandon-Jones D, Eudey A A, Geissmann T, et al. Asian Primate Classification. Int J Primatol, 2004, 25: 97–164
- 15 Stewart C B, Disotell T R. Primate evolution—in and out of Africa. Curr Biol, 1998, 8: R582–588
- 16 Delson E. Evolutionary history of the colobine monkeys in paleoenvironmental perspective. In: Davies A G, Oates J F, eds. Colobine Monkeys: Their Ecology, Behavior, and Evolution. Cambridge: Cambridge University Press, 1994. 11–44
- 17 Sterner K N, Raaum R L, Zhang Y P, et al. Mitochondrial data support an odd-nosed colobine clade. Mol Phylogenet Evol, 2006, 40: 1–7
- 18 Perelman P, Johnson W E, Roos C, et al. A molecular phylogeny of living primates. PLoS Genet, 2011, 7: e1001342
- 19 Roos C, Zinner D, Kubatko L S, et al. Nuclear versus mitochondrial DNA: Evidence for hybridization in colobine monkeys. BMC Evol Biol, 2011, 11: 77
- 20 Groves C P. The taxonomic diversity of the Colobinae of Africa. J Anthropol Sci, 2007, 85: 7–34
- 21 Grubb P, Butynski T M, Oates J F, et al. Assessment of the diversity of African primates. Int J Primatol, 2003, 24: 1301–1357
- 22 Kingston J D, Fine Jacobs B, Hill A, et al. Stratigraphy, age and environments of the late Miocene Mpesida Beds, Tugen Hills, Kenya. J Hum Evol, 2002, 42: 95–116
- 23 Groves C P. SUBFAMILY Colobinae. In: Wilson D E, Reeder D M, eds. Mammal Species of the World. 3rd ed. Baltimore: Johns Hopkins University Press, 2005. 167–178

- 24 Ting N. Mitochondrial relationships and divergence dates of the African colobines: Evidence of Miocene origins for the living colobus monkeys. J Hum Evol, 2008, 55: 312–325
- 25 Fabre P H, Rodrigues A, Douzery E J. Patterns of macroevolution among primates inferred from a supermatrix of mitochondrial and nuclear DNA. Mol Phylogenet Evol, 2009, 53: 808–825
- 26 Ting N, Tosi A J, Li Y, et al. Phylogenetic incongruence between nuclear and mitochondrial markers in the Asian colobines and the evolution of the langurs and leaf monkeys. Mol Phylogenet Evol, 2008, 46: 466–474
- 27 Strasser E, Delson E. Cladistic analysis of cercopithecid relationships. J Hum Evol, 1987, 16: 81–99
- 28 Grubb P, Butynski T M, Oates J F, et al. Assessment of the diversity of african primates. Int J Primatol, 2003, 24: 1301–1357
- 29 Chatterjee H J, Ho S Y, Barnes I, et al. Estimating the phylogeny and divergence times of primates using a supermatrix approach. BMC Evol Biol, 2009, 9: 259
- 30 Meyer D, Rinaldi I D, Ramlee H, et al. Mitochondrial phylogeny of leaf monkeys (genus *Presbytis*, Eschscholtz, 1821) with implications for taxonomy and conservation. Mol Phylogenet Evol, 2011, 59: 311–319
- 31 Wang X P, Yu L, Roos C, et al. Phylogenetic relationships among the Colobine monkeys revisited: New insight from analyses of complete mt genomes and 44 nuclear non-coding markers. PLoS One, 2012, 7: e36274
- 32 Groves C P. The forgotten leaf-eaters and the phylogeny of Colobinae. In: Napier J R, Napier P H, eds. Old World Monkeys. New York: Academic Press, 1970. 629–647
- 33 Grzimek B. Grzimek's Encyclopedia of Mammals. New York: McGraw-Hill, 1990
- 34 Elliot D G. A Review of the Primates. New York: American Museum of Natural History, 1913
- 35 Macdonald D. The Encyclopedia of Mammals. New York: Facts on File Publication, 1984. 398–411
- 36 Napier J R, Napier P H. A Handbook of Living Primates. London: Academic Press, 1967
- 37 Simpson G G. The principles of classification and a classification of mammals. Bull Amer Mus Nat Hist, 1945, 85: 1–350
- 38 Ellerman J R, Morrison-Scott T C S. Checklist of Palaearctic and Indian mammals. London: British Musseum (Natural History), 1951. 1758–1946
- 39 Hershkovitz P. Living New World Primates (Platyrrhini), with an Introduction to Primates. Chicago: University of Chicago Press, 1977
- 40 Nowak R M, Walker S. Mammals of the World. Baltimore & London: The Johns Hopkins University Press, 1991
- 41 Hill W C O. Primates: Comparative Anatomy and Taxonomy. Edinburgh: Edinburgh University Press, 1970
- 42 Hill W C O. Evolutionary Biology of the Primates. London: Academic Press, 1972
- 43 Hill W C O. A monograph on the purpled-faced leaf-monkeys (*Pithecus vetulus*). Ceylon J Sci, 1934, 9: 23–88
- 44 Groves C P. Cercopithecus and company: A primate radiation. Science, 1989, 244: 860–861
- 45 Eudey A A. Asian primate conservation—The species and the IUCN/ SSC primate specialist network. Primate Conservation, 1997, 17: 101–110
- 46 Groves C P. The forgotten leaf-eaters and the phylogeny of Colobinae. In: Napier J R, Napier P H, eds. Old World Monkeys. New York: Academic Press, 1970. 555–586
- 47 Jablonski N G. The evolution of the doucs and snub-nosed monkeys and the question of the phyletic unity of the odd-nosed colobines. In: Jablonski N G, ed. The Natural History of the Doucs and Snub-Nosed Monkeys. Singapore: World Scientific Press, 1998. 13–52
- 48 Jablonski N G, Peng Y Z. The phylogenetic relationships and classification of the doucs and snub-nosed langurs of China and Vietnam. Folia Primatol, 1993, 60: 36–55
- 49 Wang W, Forstner M R J, Zhang Y P, et al. A phylogeny of Chinese leaf monkeys using mitochondrial *ND3-ND4* gene sequences. Int J Primatol, 1997, 18: 305–320

- 50 Zhang Y P, Ryder O A. Mitochondrial cytochrome *b* gene sequences of Old World monkeys: With special reference on evolution of Asian colobines. Primates, 1998, 39: 39–49
- 51 Li M, Wei F, Huang C, et al. Phylogeny of snub-nosed monkeys inferred from Mitochondrial DNA, Cytochrome B, and 12S rRNA sequences. Int J Primatol, 2004, 25: 861–873
- 52 Osterholz M, Walter L, Roos C. Phylogenetic position of the langur genera *Semnopithecus* and *Trachypithecus* among Asian colobines, and genus affiliations of their species groups. BMC Evol Biol, 2008, 8: 58
- 53 Bigoni F, Houck M, Ryder O, et al. Chromosome painting shows that *Pygathrix nemaeus* has the most basal karyotype among Asian colobinae. Int J Primatol, 2004, 25: 679–688
- 54 Bigoni F, Stanyon R, Wimmer R, et al. Chromosome painting shows that the proboscis monkey (*Nasalis larvatus*) has a derived karyotype and is phylogenetically nested within Asian colobines. Am J Primatol, 2003, 60: 85–93
- 55 Jablonski N G. Quaternary environments and the evolution of primates in East Asia, with notes on two new specimens of fossil Cercopithecidae from China. Folia Primatol, 1993, 60: 118–132
- 56 Napier J R, Napier P H. Old World Monkeys. New York: Academic Press, 1970
- 57 Pan R. Dental variation among Asian colobines, with specific reference to the macaques on the same continent. Zool Res, 2007, 28: 569–579
- 58 Peng Y, Pan R, Ye Z, et al. Comparative study on cranioface and brain case in Asian colobines (in Chinese). Acta Anthropol Sin, 1991, 10: 346–356
- 59 Szalay F S, Delson E. Evolutionary History of the Primates. New York: Academic Press, 1979
- 60 Weitzel V, Yang C M, Groves C P. A catalogue of primate in the Singapore Zoological Reference Collection. Raffles Bull of Zool, 1988, 36: 1–166
- 61 Delson E. Evolutionary history of the Cercopithecidae. Contrib Primatol, 1975, 5: 167–217
- 62 Whittaker D J, Ting N, Melnick D J. Molecular phylogenetic affinities of the simakobu monkey (*Simias concolor*). Mol Phylogenet Evol, 2006, 39: 887–892
- 63 Li Z Y. Preliminary investigation of the habitats of *Presbytis francoisi* si and *Presbytis leucocephalus*, with notes on the activity pattern of *Presbytis leucocephalus*. Folia Primatol (Basel), 1993, 60: 83–93
- 64 Peng Y Z, Ye Z Z, Pan R L. The classification and phylogeny of snub-nosed monkeys based on gross morphological characters. Zool Res, 1988, 9: 239–247
- 65 Pocock R I. The langurs, or leaf monkeys, of British India. Part II. J Bombay Nat Hist Soc 1928, 32: 660–677
- 66 Wolfheim J H. Primates of the World. Seattle: University of Washington Press, 1983
- 67 Oates J F, Davies A G, Delson E. The diversity of living colobines. In: Davies A G, Oates J F, eds. Colobine Monkeys: Their Ecology, Behavior, and Evolution. Cambridge: Cambridge University Press, 1994. 45–74
- 68 Md-Zain B M, Morales J C, Hasan M N, et al. Is *Presbytis* a distinct monophyletic genus: Inferences from mitochondrial DNA sequences. Asian Primates J, 2008, 1: 26–36
- 69 Md-Zain B M, Mohamad M, Ernie-Muneerah M A, et al. Phylogenetic relationships of Malaysian monkeys, Cercopithecidae, based on mitochondrial cytochrome *c* sequences. Genet Mol Res, 2010, 9:

1987-1996

- 70 Hooijer D A. Quaternary Langurs and Macaques from the Malay archipelago. Zool Verhandelingen, 1962, 55: 1–64
- 71 Brandon-Jones D. Colobus and leaf monkeys. In: Macdonald D, ed. The Encyclopaedia of Mammals. London: Allen and Unwin, 1984. 398–410
- 72 Brandon-Jones D. The Asian Colobinae (Mammalia: Cercopithecidae) as indicators of Quaternary climatic change. Biol J Linn Soc, 1996, 59: 327–350
- 73 Karanth K P. Molecular systematics and conservation of the langurs and leaf monkeys of South Asia. J Genet, 2010, 89: 393–399
- 74 Jablonski N G, Peng Y Z. The phylogenetic relationships and classification of the doucs and snub-nosed langurs of China and Vietnam. Folia Primatol (Basel), 1993, 60: 36–55
- 75 Boonratana R, Le X C. Preliminary Observations of the Ecology and Behavior of the Tonkin Snub-nosed Monkey (*Rhinopithecus [Presbytiscus] avunculus*) in Northern Vietnam. In: Jablonski N G, ed. The Natural History of the Doucs and Snub-nosed Monkeys. Singapore: World Scientific Publishing, 1998
- 76 Kirkpatrick R C. Ecology and behavior in snub-nosed and douc langurs. In: Jablonski N G, ed. The Natural History of the Doucs and Snub-nosed Monkeys. Singapore: World Scientific Press, 1998. 155–190
- 77 Zhang Y, Ryder O A. Mitochondrial DNA sequence evolution and conservation relevance of snub-nosed langurs. Acta Genet Sin, 1997, 24: 116–121
- 78 Geissmann T, Groves C P, Roos C. The Tenasserim Lutung, *Trachypithecus barbei* (Blyth, 1847) (Primates: Cercopithecidae): Description, of a live specimen, and a reassessment of phylogenetic affinities, taxonomic history, and distribution. Contrib Zool, 2004, 73: 271–282
- 79 Karanth K P, Singh L, Collura R V, et al. Molecular phylogeny and biogeography of langurs and leaf monkeys of South Asia (Primates: Colobinae). Mol Phylogenet Evol, 2008, 46: 683–694
- 80 Karanth K P. Primate numts and reticulate evolution of capped and golden leaf monkeys (Primates: Colobinae). J Biosci, 2008, 33: 761– 770
- 81 Rokas A, Carroll S B. Bushes in the tree of life. PLoS Biol, 2006, 4: 1899–1904
- 82 Philippe H, Chenuil A, Adoutte A. Can the Cambrian explosion be inferred through molecular phylogeny? Development, 1994, 120: S15–S25
- 83 Saitou N, Nei M. The number of nucleotides required to determine the branching order of three species, with special reference to the human-chimpanzee-gorilla divergence. J Mol Evol, 1986, 24: 189– 204
- 84 Yu L, Zhang Y P. Phylogenomics: An attractive avenue to reconstruct "tree of life" (in Chinese). Hereditas, 2006, 28: 1445–1450
- 85 Meslin C, Brimau F, Nagnan-Le Meillour P, et al. The evolutionary history of the SAL1 gene family in eutherian mammals. BMC Evol Biol, 2011, 11: 148
- 86 Wu Y C, Rasmussen M D, Kellis M. Evolution at the subgene level: Domain rearrangements in the *Drosophila* phylogeny. Mol Biol Evol, 2011, 29: 689–705
- 87 Yu L, Luan P T, Jin W, et al. Phylogenetic utility of nuclear introns in interfamilial relationships of Caniformia (order Carnivora). Syst Biol, 2011, 60: 175–187
- **Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.