

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Perälä, Mia-Maria; von Bonsdorff, Mikaela; Männistö, Satu; K.Salonen, Minna; Simonen, Mika; Pohjolainen, Pertti; Kajantie, Eero; Rantanen, Taina; Eriksson, Johan G.

Title: The Healthy Nordic Diet and Mediterranean Diet and Incidence of Disability 10 Years Later in Home-Dwelling Old Adults

Year: 2019

Version: Accepted version (Final draft)

Copyright: © 2018 AMDA - The Society for Post-Acute and Long-Term Care Medicine.

Rights: CC BY-NC-ND 4.0

Rights url: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the original version:

Perälä, M.-M., von Bonsdorff, M., Männistö, S., K.Salonen, M., Simonen, M., Pohjolainen, P., Kajantie, E., Rantanen, T., & Eriksson, J. G. (2019). The Healthy Nordic Diet and Mediterranean Diet and Incidence of Disability 10 Years Later in Home-Dwelling Old Adults. *Journal of the American Medical Directors Association*, 20(5), 511-516.e1.
<https://doi.org/10.1016/j.jamda.2018.09.001>

The healthy Nordic diet and Mediterranean diet and incidence of disability 10 years later in home-dwelling old adults

Mia-Maria Perälä PhD^{1,2}, Mikaela B von Bonsdorff PhD^{2,3}, Satu Männistö PhD¹, Minna K Salonen PhD^{1,2}, Mika Simonen PhD⁴, Pertti Pohjolainen PhD⁵, Eero Kajantie MD, PhD^{1,6,7}, Taina Rantanen PhD³, Johan G Eriksson MD, PhD^{1,2,8,9}

¹ Department of Public Health Solutions, National Institute for Health and Welfare, Helsinki, Finland

² Folkhälsan Research Center, Helsinki, Finland

³ Gerontology Research Center, Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

⁴ Department of Social Research, University of Helsinki, Helsinki, Finland

⁵ Age Institute, Helsinki, Finland

⁶ Hospital for Children and Adolescents, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

⁷ PEDEGO Research Unit, MRC Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland

⁸ Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, Unit of General Practice, Helsinki, Finland

⁹ Vaasa Central Hospital, Vaasa, Finland

Corresponding author: Mia-Maria Perälä, Department of Public Health Solutions, National Institute for Health and Welfare, P.O. Box 30, FI-00271 Helsinki, Finland, e-mail mia.perala@thl.fi

Key words: disability, Nordic diet, Mediterranean diet, mobility limitation, ageing

Running title: Healthy diet and disability

Word, reference, tables: the abstract 257, main text 2992, references 41, tables 3, appendix table 1.

Acknowledgements

34 This work was supported by grants from Academy of Finland, Finska Läkaresällskapet, Samfundet
35 Folkhälsan, Juho Vainio Foundation, Signe and Ane Gyllenberg Foundation, Yrjö Jahnsson
36 Foundation, Diabetes Research Foundation, Finnish Foundation for Cardiovascular Research. EU
37 H2020-PHC-2014-DynaHealth grant no. 633595. The Academy of Finland supported MBvB (grant
38 no. 257239); EK (grants no. 127437, 129306, 130326, 134791 and 263924), and JGE (grants no.
39 129369, 129907, 135072, 129255 and 126775).

40

41 **Conflict of interest**

42 None.

43

44 **The healthy Nordic diet and Mediterranean diet and incidence of disability 10 years later in**
45 **home-dwelling old adults**

46

47 **Abstract**

48 **Background/Objective:** Diet has a major impact on a person's health. However, limited
49 information exists on the long-term role of the whole diet on disability. We investigated the
50 association of the healthy Nordic diet and the Mediterranean diet with incident disability 10 years
51 later.

52 **Design:** Longitudinal, with a follow-up of 10 years.

53 **Settings/Participants:** A total of 962 home-dwelling men and women from the Helsinki Birth
54 Cohort Study, mean age 61.6 y, who were free of disability at baseline.

55 **Measurements:** At baseline, 2001-2004, the Nordic diet score (NDS) and modified Mediterranean
56 diet score (mMDS) were calculated using a validated 128-item food-frequency questionnaire.
57 Higher scores indicated better adherence to the diet. Participants' incident disability was assessed
58 during 2011-2013 by a self-reported questionnaire and was based on mobility limitations and
59 difficulties to perform self-care activities. Analyses were performed using logistic regression and
60 adjusted for potential confounding factors.

61 **Results:** In total, 94 participants (9.8%) developed mobility limitations and 45 participants (4.7%)
62 developed difficulties in self-care activities during 10 year follow-up. The likelihood of having
63 mobility limitations (odds ratio (OR) 0.42, 95% confidence interval (CI) 0.22–0.80) and difficulties
64 in self-care activities (OR 0.38, 95% CI 0.15–0.94) were lower among those in the highest NDS
65 tertile than among those in the lowest NDS tertile. Greater mMDS was associated with a lower
66 disability incidence; however, the association was not statistically significant.

67 **Conclusions/Implications:** Adherence to the healthy Nordic diet predicts 10-year incidence of
68 mobility limitations and difficulties to perform self-care activities in old age and may thus be
69 protective against disability in Nordic population.

70 **Introduction**

71 World Health Organization identifies disability in old individuals as people's difficulty to
 72 participate in daily activities or activities that are necessary for independent living and integration in
 73 their environment. ⁽¹⁾ There are several domains of disability, such as mobility limitations and
 74 difficulties in activities of daily living (ADL) or instrumental activities of daily living (IADL) that
 75 are essential for independent life. Studies have shown that mobility limitation is a strong predictor
 76 of ADL and IADL disability. ⁽²⁾ It has been estimated that 8–35% of individuals 65 years old or
 77 older have mobility limitations or difficulties in performing ADL tasks. ⁽²⁻⁴⁾ Mobility limitations
 78 and difficulties to perform self-care activities increase the risk for hospitalization, nursing home
 79 admission and are strong predictors of mortality. ^(5, 6) Therefore, it is important to understand the
 80 processes that cause the progression of disability in order to develop strategies to prevent or delay
 81 disability.

82

83 Poor health and physical performance, smoking as well as cognitive impairment are well known
 84 risk factors for disability. ^(2, 7) Diet is also a potential modifiable factor affecting disability.
 85 Observational studies have shown associations between low circulating levels of carotenoids and
 86 vitamin D and B12 ^(8, 9) as well as low intakes of single antioxidant nutrients, fruits and vegetables
 87 ^(10, 11) and mobility limitations and ADL. However, despite the previously described studies
 88 concerning single dietary factors and disability, the role of the whole diet in disablement process is
 89 still mostly unknown. A few studies have shown that adherence to Healthy Eating Index ^(12, 13) and
 90 Japanese diet ⁽¹⁴⁾ are related to a lower likelihood of mobility limitations. In addition, Mediterranean
 91 diet has been shown to decrease the risk of disability and slow the decline of mobility and walking
 92 speed. ⁽¹⁵⁻¹⁸⁾ We have recently observed a positive relationship between the healthy Nordic diet and
 93 overall physical performance and muscle strength. ^(19, 20) However, no studies exist in which the
 94 association of the healthy Nordic diet with incidence of disability has been examined. In addition,
 95 previous studies have not been investigated and compared different diet scores in the same cohort
 96 with respect to incidence of disability. Investigating different diet scores may help in identifying
 97 which diet is appropriate for population concerned. Therefore, we investigated whether two diet
 98 scores, the Nordic diet and the modified Mediterranean diet scores, predicts mobility limitations and
 99 difficulties to perform self-care activities 10 years later in initially healthy home-dwelling men and
 100 women aged 61 years on average at baseline.

101

102 **Methods**

103 *Design and Study Population*

104 The subjects in this study belong to the Helsinki Birth Cohort Study which originally included 4630
 105 men and 4130 women who were born as singletons at Helsinki University Central Hospital between
 106 1934 and 1944. ⁽²¹⁾ A random sample of 2 902 individuals were invited to participate in a clinical
 107 examination conducted between 2001 and 2004. Of these, 2003 men (n = 928) and women (n =
 108 1075) participated in the examination. From 2011–2013, 1404 subjects were still traceable. Of
 109 these, 1094 participants (478 men and 616 women) attended a clinical examination between 2011
 110 and 2013. ⁽¹⁹⁾ Those who participated in the follow-up clinical examination had lower rates of
 111 mobility limitations (6.2% v. 16.6%) and difficulties in self-care activities (4.6% v. 12.1%) at
 112 baseline compared to those who did not participate. Participants who had adequate data on diet and
 113 disability, and were free of disability at baseline were included in this study (n = 962).

114
 115 This study was conducted according to the guidelines laid down in the Declaration of Helsinki and
 116 all procedures involving human subjects were approved by the Ethics Committee of the Helsinki
 117 and Uusimaa Hospital District. Written informed consent was obtained from all subjects.

118 119 *Dietary Assessment*

120 At the baseline clinical examination, diet was assessed by a validated, self-administered, 128-item
 121 food-frequency questionnaire (FFQ). ^(22, 23) Participants were asked how often, on average, they
 122 consumed each food item in the past year. The frequency responses ranged from “never or seldom”
 123 to “at least six times a day”. The portion size was fixed for each food item or mixed dish (e.g. slice
 124 and glass) based on the national Findiet Study and present the most commonly used portion sizes in
 125 Finland. At the clinic, a research nurse checked the FFQ. The final decision regarding the
 126 completeness of the FFQ was made by a nutritionist. Exclusions were made due to incompletely
 127 filled FFQ and daily energy intake cut-off points corresponding to 0.5% at both ends of the daily
 128 energy intake distributions for men and women. ⁽²⁴⁾ The average daily intakes of nutrients and foods
 129 were calculated with the national food composition database Fineli®.

130
 131 Details of the Nordic diet score (NDS), which is also known as the Baltic Sea diet score, and the
 132 modified Mediterranean diet score (mMDS) are described in detail elsewhere. ^(25, 26) Briefly, the
 133 NDS is based on 9 components. Favourable food components (Nordic vegetables (excluding
 134 potatoes), Nordic fruits, Nordic cereals, fish, a ratio of polyunsaturated fatty acids to saturated and
 135 trans-fatty acids, and low-fat milk) were assigned values of 0–3 according to ascending sex-specific
 136 quartile ranks, and unfavourable food components (red and processed meat, and intake of total fat)
 137 were assigned values of 0–3 according to descending sex-specific quartile ranks. In addition, men

consuming ≤ 20 g/day and women consuming ≤ 10 g/day of alcohol received 1 point; otherwise, a 0 point was given. The mMDS is based on 9 components. For vegetables (excluding potatoes), fruit and nuts, cereals, legumes, fish and fish products, and the ratio of monounsaturated and polyunsaturated to saturated fat, 1 point was given if intake was above the sex-specific median, 0 otherwise. For meat and dairy products, 1 point was given if intake was below the sex-specific median, 0 otherwise. For alcohol, 1 point was given if intake was between 10–50 g/day in men and 5–25 g/day in women; otherwise, a 0 point was given. The NDS ranged from 0–25 and the mMDS from 0–9.

Disability

Disability was assessed with the Finnish validated version of the RAND 36-Item Health Survey 1.0 (Short Form 36 [SF-36])⁽²⁷⁾ at baseline and follow-up clinical examination. The question on difficulties in walking 500 m was used to assess mobility limitations and difficulties to perform self-care activities was assessed with questions on difficulties in dressing or bathing. Response categories were: no difficulties; little difficulties; many difficulties. Participants were considered to have mobility limitations if they reported any difficulties in walking 500 m and difficulties to perform self-care activities if they reported any difficulties in dressing and/or bathing.

For comparing whether self-reported mobility limitations is valid and identify those with worse physical performance, participants' physical performance was tested at follow-up clinical examination with the validated Senior Fitness Test⁽²⁸⁾ and total score was calculated as described earlier.⁽¹⁹⁾

Covariates

During the baseline clinical examination, height was measured to the nearest 0.1 cm and weight to 0.1 kg. BMI was calculated as weight (in kg) divided by height (in m²). At the baseline clinical examinations, the level of educational attainment (elementary school; vocational school; senior high school and college; university degree), smoking status (never; former; current), presence of main chronic diseases (cancer; cerebrovascular disease; hypertension; diabetes) were asked using questionnaire. Energy intake was measured with an FFQ as an average intake (in MJ per day). Participants' physical activity over the previous 12 months was assessed using a validated exercise questionnaire.⁽²⁹⁾ Depressive symptoms were assessed using the Beck depression inventory (BDI)⁽³⁰⁾, and participants with score ≥ 10 were classified as having at least mild depression. Cognitive

status was assessed by Mini-Mental State Examination (MMSE) ⁽³¹⁾ score (0-30) during 2009-2011, and participants with a score <24 were classified as having impaired cognition.

Statistical Methods

There were no significant interactions for the healthy diet and sex on disability, therefore, men and women were analysed together. Descriptive characteristics were reported as mean (standard deviation), proportions, or median (interquartile ranges) values, and the differences of baseline characteristics across tertiles were tested with one-way ANOVA or Chi-square test. The association between tertiles of NDS and mMDS and likelihood of having mobility limitations or difficulties to perform self-care activities were examined by logistic regression models. The relationships between each of the NDS and mMDS components and self-reported disability were also examined using logistic regression models. Model 1 was adjusted for age and sex. To investigate whether diet was independently related to disability, the Model 2 was further adjusted for potential confounding factors related to disability or diet: BMI, physical activity, smoking status, educational attainment, energy intake, and presence of chronic diseases and depression. ^(2, 32, 33) Model 3 was further adjusted for cognitive status and included only those participants whose MMSE test result was available (n=832). All statistical analyses were done using the SPSS Statistics version 23 for Windows® (SPSS Inc., Chicago, IL, USA). Significance was defined as P<.05.

Results

The basic characteristics of the study population for men (n=439) and women (n=523) are seen in **Table 1** and daily intakes of NDS and mMDS components in **Appendix Table 1**. The correlation between NDS and mMDS was 0.56 (P<.001). Incident mobility limitation was more common than self-care dependence. During a 10-year follow-up, 94 participants (9.8%) developed mobility limitations and 45 participants (4.7%) developed difficulties to perform self-care activities. Senior Fitness Test total score was significantly smaller among those with mobility limitations (32.9 points, SD 15.1) than those with no limitations (47.9 points, SD 16.6).

After adjusting for age, BMI, educational attainment, physical activity, smoking status, energy intake, and the presence of chronic disease and depression the likelihood of having difficulties to walk 500 m were lower among those in the highest NDS tertile than among those in the lowest NDS tertile (odds ratio (OR) 0.42, 95% confidence interval (CI) 0.22– 0.80) (**Table 2**). In addition, individuals with the highest adherence to a healthy Nordic diet at baseline had lower likelihood of developing difficulties to perform self-care activities compared with those with the lowest

adherence (OR 0.38, 95% CI 0.15–0.94). In total, 832 participants had completed MMSE test results. Including cognitive status to the model, the likelihood of having disability were lower among those with high NDS, however, only the association between the NDS and mobility limitations remained significant.

After adjusting for age and sex the likelihood of having mobility limitation (OR 0.65, 95% CI 0.35–1.22) and difficulties to perform self-care activities (OR 0.59, 95% CI 0.26–1.35) was lower among those who had the highest mMDS compared to those who had the lowest mMDS, however, the associations did not reach statistical significance (**Table 2**). Adjusting for further potential confounders did not have a major effect on the results.

When investigating the associations between NDS and mMDS components and disability, we found statistical significant associations of greater intake of fat and smaller intake of cereals with increased likelihood of having difficulties to perform self-care activities (**Table 3**). Greater intake of alcohol was significantly related to lower likelihood of having mobility limitations.

Discussion

In our study of 962 home-dwelling ageing men and women, we observed that the healthy Nordic diet is a strong predictor of two states of disability; mobility limitations and difficulties to perform self-care activities, even after adjusting for potential confounding factors including physical activity, chronic diseases and depression. Adherence to the Mediterranean diet was also related to lower likelihood of being disabled, however, the association was not statistically significant. To our knowledge, there are no previous studies that have compared different healthy diets within the same cohort for disability outcomes. Our findings extend the relevance of a healthy diet to target mobility limitations and difficulties to perform self-care activities and support the importance of diet in the pathogenesis of disability in old age.

Our results support the previous findings in which the protective effect of healthy diet, including the Healthy Eating Index, Japanese diet and Mediterranean diet, on disability has been observed.^(12–16) Even though these diets vary in food and nutrient components, they all capture essential elements of a high-quality diet including high consumption of fruits, vegetables, and fish and low consumption of red and processed meat. When we investigated single food components of NDS and mMDS, we observed that only intake of fat, alcohol and cereals were related to disability. Based on our results, we suggest that the whole diet is a stronger predictor of disability in old age than single foods or

nutrients. Therefore, the findings in our study extend the previous results and suggest that healthy diet, which include high amounts of fruit, vegetables, and fish, moderate intake of alcohol, and small amounts of red meat and fat, help to prevent disability in old age.

Even though the healthy Nordic and Mediterranean diet were related to lower likelihood of disability, only the association of healthy Nordic diet with disability was significant. The healthy Nordic diet contains locally grown healthy Nordic foods whereas the widely investigated Mediterranean diet contains foods familiar in the Mediterranean area such as seeds, nuts and olive oil. Therefore, adherence to the healthy Nordic diet may be easier in Nordic population. The major difference between these diets is that dairy products are beneficial in the NDS, whereas they are detrimental in the mMDS. In addition, as these diet scores are based on study specific cut-offs for each food components, the absolute intake of each component may differ between studies even though diet scores are identical. Indeed, we observed that the median consumption of food components included in the mMDS, such as fruits and vegetables, were in part lower than observed in previous studies ^(15, 18), suggesting that the diet of our study population is not as healthy than in the Mediterranean area. Furthermore, it should be noted that there are several different versions of Mediterranean diet score, which make it difficult to compare the results. Moreover, the range of the MDS is much smaller (0-9) compared to the NDS (0-25) which may limit the statistical power. Therefore, we propose that these could partly explain why we did not observe as strong associations between Mediterranean diet and disability as with healthy Nordic diet.

Strong evidence exist that mobility limitations are the results of physical impairments, such as decreased muscle strength ^(34, 35). Several diseases and changes in body homeostasis also affect muscle strength and thus mobility. Previously, it has been demonstrated that adherence to the healthy Nordic and Mediterranean diet predicts greater muscle strength and thus better overall physical performance. ^(16, 18-20) In addition, both of these diets have been found to be associated with a lower risk of several diseases, such as diabetes and cardiovascular disease, and changes in body homeostasis including insulin resistance and inflammation. ⁽³⁶⁻³⁹⁾ Therefore, we suggest that plausible explanations for the observed associations between the healthy diet and disability are multifactorial, including beneficial effects of diet on muscle strength and on risk factors of chronic diseases. In addition, adherence to the healthy diet may also be linked with overall healthy lifestyles practices including increased physical activity, which associates with disability.

A significant strength of our study is its longitudinal study design that provides opportunity to investigate long-term influences of diet on disability. In addition, our study included a large study population consisting of both men and women. A further strength of our study is that we measured two states of disability; difficulties to perform self-care activities, which is more serious and mobility limitations, which are more common. Our study also has some limitations as discussed earlier.⁽²⁰⁾ Firstly, we adjusted for a multitude of covariates, including physical activity, socioeconomic status and chronic diseases; however, potential unmeasured and residual confounding cannot be excluded. Secondly, diet was measured by a validated FFQ^(22, 23) only at baseline; therefore changes in the diet were not captured. Thirdly, the disability status of participants was self-reported, which may cause misclassification of disability status.⁽³⁾ However, it has been shown that self-reported difficulties in walking reflect people's physical performance status and is able to identify those with worse physical performance.^(40, 41) Therefore, it is valid and has clinical significance. We also observed that those who reported mobility limitations had significantly poorer physical performance than those with no limitations. In addition, we measured only one of the ADL components, difficulty to bath and dress, which may underestimate the difficulties in self-care activities. However, difficulty to bath is the most common ADL disability and it represents the first ADL in which most persons become disabled. Therefore, we suggest that it is a good indicator of ADL. Further, as reported earlier, those individuals who participated in the follow-up clinical examination had lower rates of disability, were younger, thinner, more educated, and had healthier diet at baseline compared with those who did not participate.⁽¹⁹⁾ Those with unhealthy lifestyle habits may have died at earlier ages or developed disability and were, therefore, excluded from the study. Selective survival may thus have influenced the association between the healthy diet and disability. In addition, the incidence of disability was low during a 10-year follow-up which may limit the statistical power. Thus, participants may not be fully representative of all older people living in Finland.

Conclusions

Our study indicates that adherence to the healthy Nordic diet decreased the likelihood of mobility limitations and difficulties to perform self-care activities in old age. In addition, Mediterranean diet may also have a beneficial impact on disability. Our results may have important clinical implications. As disability is common among older age and restricts a person's independence as well as affects overall health, it is important try to prevent disability. Our results indicates that adherence to healthy diet provide one opportunity to prevent or delay disability. In addition, it is well known that adherence to healthy diet has several other health benefits as well. Therefore,

306 adhering to a healthy diet may have major beneficial effect on people's health. However, future
307 intervention studies as well as longitudinal studies, that compare different diet scores with incidence
308 of disability, are critically needed.

309

References

1. WHO (World Health Organization). International Classification of Functioning, Disability and Health (ICF). Geneva: WHO (World Health Organization); 2001.
2. Guralnik JM, Fried LP, Salive ME. Disability as a public health outcome in the aging population. *Annu Rev Public Health* 1996;17:25-46.
3. Sainio P, Koskinen S, Heliövaara M, et al. Self-reported and test-based mobility limitations in a representative sample of Finns aged 30+. *Scand J Public Health* 2006;34:378-386.
4. Shumway-Cook A, Ciol MA, Yorkston KM, et al. Mobility limitations in the Medicare population: prevalence and sociodemographic and clinical correlates. *J Am Geriatr Soc* 2005;53:1217-1221.
5. Leskinen R, Laatikainen T, Peltonen M, et al. Self-reported walking difficulty predicts late-life mortality in Finnish war veterans: results from the Veteran 1992 Project Survey. *J Am Geriatr Soc* 2015;63:118-123.
6. Wu LW, Chen WL, Peng TC, et al. All-cause mortality risk in elderly individuals with disabilities: a retrospective observational study. *BMJ Open* 2016;6:e011164.
7. Legrand D, Vaes B, Matheï C, et al. Muscle strength and physical performance as predictors of mortality, hospitalization, and disability in the oldest old. *J Am Geriatr Soc* 2014;62:1030-1038.
8. Bartali B, Semba RD, Frongillo EA, et al. Low micronutrient levels as a predictor of incident disability in older women. *Arch Intern Med* 2006;166:2335-2340.
9. Houston DK, Neiberg RH, Tooze JA, et al. Low 25-hydroxyvitamin D predicts the onset of mobility limitation and disability in community-dwelling older adults: the Health ABC Study. *J Gerontol A Biol Sci Med Sci* 2013;68:181-187.
10. Houston DK, Stevens J, Cai J, et al. Dairy, fruit, and vegetable intakes and functional limitations and disability in a biracial cohort: the Atherosclerosis Risk in Communities Study. *Am J Clin Nutr* 2005;81:515-522.
11. Tomey KM, Sowers MR, Crandall C, et al. Dietary intake related to prevalent functional limitations in midlife women. *Am J Epidemiol* 2008;167:935-943.
12. Koster A, Penninx BW, Newman AB, et al. Lifestyle factors and incident mobility limitation in obese and non-obese older adults. *Obesity (Silver Spring)* 2007;15:3122-3132.
13. Xu B, Houston D, Locher JL, et al. The association between Healthy Eating Index-2005 scores and disability among older Americans. *Age Ageing* 2012;41:365-371.

- 343 14. Tomata Y, Watanabe T, Sugawara Y, et al. Dietary patterns and incident functional
344 disability in elderly Japanese: the Ohsaki Cohort 2006 study. *J Gerontol A Biol Sci Med Sci*
345 2014;69:843-851.
- 346 15. Féart C, Pérès K, Samieri C, et al. Adherence to a Mediterranean diet and onset of
347 disability in older persons. *Eur J Epidemiol* 2011;26:747-756.
- 348 16. Milaneschi Y, Bandinelli S, Corsi AM, et al. Mediterranean diet and mobility decline in
349 older persons. *Exp Gerontol* 2011;46:303-308.
- 350 17. Shahar DR, Houston DK, Hue TF, et al. Adherence to mediterranean diet and decline in
351 walking speed over 8 years in community-dwelling older adults. *J Am Geriatr Soc*
352 2012;60:1881-1888.
- 353 18. Zbeida M, Goldsmith R, Shimony T, et al. Mediterranean diet and functional indicators
354 among older adults in non-Mediterranean and Mediterranean countries. *J Nutr Health Aging*
355 2014;18:411-418.
- 356 19. Perälä MM, von Bonsdorff M, Männistö S, et al. A healthy Nordic diet and physical
357 performance in old age: findings from the longitudinal Helsinki Birth Cohort Study. *Br J*
358 *Nutr* 2016;115:878-886.
- 359 20. Perälä MM, von Bonsdorff MB, Männistö S, et al. The healthy Nordic diet predicts
360 muscle strength 10 years later in old women, but not old men. *Age Ageing* 2017;46:588-594.
- 361 21. Barker DJ, Osmond C, Forsén TJ, et al. Trajectories of growth among children who have
362 coronary events as adults. *N Engl J Med* 2005;353:1802-1809.
- 363 22. Männistö S, Virtanen M, Mikkonen T, et al. Reproducibility and validity of a food
364 frequency questionnaire in a case-control study on breast cancer. *J Clin Epidemiol*
365 1996;49:401-409.
- 366 23. Paalanen L, Männistö S, Virtanen MJ, et al. Validity of a food frequency questionnaire
367 varied by age and body mass index. *J Clin Epidemiol* 2006;59:994-1001.
- 368 24. Meltzer HM, Brantsaeter AL, Ydersbond TA, et al. Methodological challenges when
369 monitoring the diet of pregnant women in a large study: experiences from the Norwegian
370 Mother and Child Cohort Study (MoBa). *Mat Child Nutr* 2008;4:14-27.
- 371 25. Kanerva N, Kaartinen NE, Schwab U, et al. The Baltic Sea Diet Score: a tool for
372 assessing healthy eating in Nordic countries. *Public Health Nutr* 2013;17:1697-1705.
- 373 26. Trichopoulou A, Orfanos P, Norat T, et al. Modified Mediterranean diet and survival:
374 EPIC-elderly prospective cohort study. *BMJ* 2005;330:991.
- 375 27. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I.
376 Conceptual framework and item selection. *Med Care* 1992;30:473-483.

- 377 28. Rikli RE, Jones CJ. Development and validation of a functional fitness test for
378 community-residing older adults. *J Aging Phys Act* 1999;6:127-159.
- 379 29. Lakka TA, Salonen JT. Intra-person variability of various physical activity assessments
380 in the Kuopio Ischaemic Heart Disease Risk Factor Study. *International journal of*
381 *epidemiology* 1992;21:467-472.
- 382 30. Beck AT, Steer RA, Garbin MG. Psychometric properties of the Beck Depression
383 Inventory: Twenty-five years of evaluation. *Clin Psychol Rev* 1988;8:77-100.
- 384 31. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for
385 grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-198.
- 386 32. Maynard M, Gunnell D, Ness AR, et al. What influences diet in early old age?
387 Prospective and cross-sectional analyses of the Boyd Orr cohort. *Eur J Public Health*
388 2006;16: 316-24.
- 389 33. Thiele S, Mensink GB, Beitz R. Determinants of diet quality. *Public Health Nutr* 2004;7:
390 29-37.
- 391 34. Stenholm S, Shardell M, Bandinelli S, et al. Physiological factors contributing to
392 mobility loss over 9 years of follow-up-results from the InCHIANTI study. *J Gerontol A Biol*
393 *Sci Med Sci* 2015;70:591-597.
- 394 35. Visser M, Goodpaster BH, Kritchevsky SB, et al. Muscle mass, muscle strength, and
395 muscle fat infiltration as predictors of incident mobility limitations in well-functioning older
396 persons. *J Gerontol A Biol Sci Med Sci* 2005;60:324-333.
- 397 36. Kanerva N, Kaartinen NE, Rissanen H, et al. Associations of the Baltic Sea diet with
398 cardiometabolic risk factors--a meta-analysis of three Finnish studies. *Br J Nutr*
399 2014;112:616-626.
- 400 37. Kastorini CM, Milionis HJ, Esposito K, et al. The effect of Mediterranean diet on
401 metabolic syndrome and its components: a meta-analysis of 50 studies and 534,906
402 individuals. *Journal of the American College of Cardiology* 2011;57:1299-1313.
- 403 38. Lacoppidan SA, Kyrø C, Loft S, et al. Adherence to a Healthy Nordic Food Index Is
404 Associated with a Lower Risk of Type-2 Diabetes -The Danish Diet, Cancer and Health
405 Cohort Study. *Nutrients* 2015;7:8633-8644.
- 406 39. Sofi F, Macchi C, Abbate R, et al. Mediterranean diet and health status: an updated
407 meta-analysis and a proposal for a literature-based adherence score. *Public Health Nutr*
408 2014;17:2769-2782.

- 409 40. Fried LP, Young Y, Rubin G, et al. Self-reported preclinical disability identifies older
410 women with early declines in performance and early disease. *J Clin Epidemiol* 2001;54:889-
411 901.
- 412 41. Wang CY, Hu MH, Chen HY, Li RH. Self-reported mobility and instrumental activities
413 of daily living: Test–retest reliability and criterion validity. *J Aging Phys Act* 2012;20:186-
414 197.
- 415

416 **Table 1.** Characteristics of the subjects by adherence to the diet score tertile

	Nordic diet score							modified Mediterranean diet score						
	Low (0-11)		Middle (12-15)		High (16-25)		P	Low (0-3)		Middle (4-5)		High (6-9)		P
n (men %)	366	(48.4)	308	(42.2)	288	(45.8)	.28	284	(45.8)	423	(44.7)	255	(47.1)	.83
Age (years) *	61.0	2.7	61.3	2.8	61.4	2.7	.25	61.3	2.9	61.2	2.8	61.2	2.5	.81
Body mass index (kg/m ²) *	26.9	3.9	26.9	3.8	26.9	4.0	.97	27.0	3.8	26.9	3.9	26.7	3.9	.60
Current smokers (%) *,†	26.6		16.3		10.2		<.001	25.0		17.7		11.9		.003
University degree (%) *	22.2		27.0		29.5		.28	21.8		25.1		31.8		.015
Physical activity (MET h/week) *	34.9	23.4	38.7	29.0	42.1	24.3	<.001	36.5	26.4	36.8	23.8	42.9	27.6	.004
Energy intake (MJ) *	8.5	2.9	9.4	3.3	10.0	3.1	<.001	8.1	2.7	9.2	3.1	10.5	3.4	<.001
Chronic diseases (%) *,‡	37.5		41.0		41.3		.50	38.4		40.1		40.8		.34
Depressed (%) *,§	14.9		11.4		14.4		.36	12.0		15.9		11.8		.20
Impaired cognition (%) *	2.3		3.2		1.6		.47	2.5		2.4		2.3		.99
Mobility limitations (%) ¶	12.3		10.7		5.6		.013	9.9		11.6		6.7		.11
Self-care dependence (%) ¶,¶¶	5.7		5.5		2.4		.097	6.0		4.5		3.5		.39

417 MET, metabolic equivalent.

418 Data are expressed as mean values (with standard deviations) for continuous values or percentages for categorical values. The differences
 419 between the diet score groups were tested with one-way ANOVA for continuous variables and Chi-square test for categorical variables.

420 * At baseline clinical examination.

421 † Smoking one or more cigarettes per day.

- 422 ‡ Presence of at least 1 chronic disease.
- 423 § Having at least mild depression.
- 424 || At follow-up clinical examination.
- 425 ¶ Difficulties to perform self-care activities.
- 426

Table 2. Odds ratios of mobility limitations and difficulties to perform self-care activities over 10-year follow-up for baseline adherence to Nordic and Mediterranean diet among Helsinki Birth Cohort Study participants (n=962)

	Model 1 *			Model 2 †			Model 3 ‡		
	OR	95% CI	P §	OR	95% CI	P §	OR	95% CI	P §
Mobility limitations									
NDS tertiles									
Low	Ref.			Ref.			Ref.		
Middle	0.86	0.54–1.39		0.88	0.53–1.48		0.81	0.46–1.40	
High	0.42	0.23–0.76	.005	0.42	0.22–0.80	.010	0.42	0.21–0.84	.014
mMDS tertiles									
Low	Ref.			Ref.			Ref.		
Middle	1.21	0.74–1.97		1.15	0.68–1.98		1.15	0.65–2.01	
High	0.65	0.35–1.22	.12	0.69	0.34–1.37	.26	0.61	0.28–1.30	.16
Difficulties to perform self-care activities									
NDS tertiles									
Low	Ref.			Ref.			Ref.		
Middle	0.92	0.47–1.78		0.89	0.45–1.79		0.87	0.41–1.86	
High	0.39	0.16–0.93	.039	0.38	0.15–0.94	.039	0.45	0.17–1.16	.087
mMDS tertiles									
Low	Ref.			Ref.			Ref.		
Middle	0.75	0.38–1.47		0.60	0.30–1.23		0.60	0.28–1.27	
High	0.59	0.26–1.35	.21	0.51	0.21–1.25	.17	0.45	0.17–1.21	.12

OR, odds ratio; CI, confidence interval; NDS, Nordic diet score; mMDS, modified

Mediterranean diet score; Ref., referent value.

* Model 1, adjusted for sex and age.

† Model 2, adjusted for Model 1 plus body mass index, educational attainment, smoking status, physical activity, energy intake, presence of chronic diseases and depression.

‡ Model 3, adjusted for Model 2 plus impaired cognitive function (n=832).

§ P for a linear trend.

Table 3. Association between sex specific tertiles of Nordic and Mediterranean diet score components and risk of mobility limitations and difficulties to perform self-care activities during a 10 year follow-up of old adults (n=962) *

Component (g/d)	Mobility limitations					Difficulties in self-care activities				
	1	2	3	1	2	3	1	2	3	
	OR	95% CI	OR	95% CI	OR	95 % CI	OR	95 % CI	OR	95% CI
NDS component										
Nordic vegetables	Ref.	0.95	0.53–1.70	0.76	0.39–1.47	Ref.	0.64	0.28–1.47	0.99	0.42–2.35
Nordic fruits	Ref.	0.94	0.52–1.71	0.94	0.49–1.78	Ref.	1.26	0.56–2.82	1.30	0.53–3.21
Nordic cereals	Ref.	0.62	0.34–1.12	0.77	0.42–1.44	Ref.	0.91	0.42–1.97	0.64	0.27–1.50
Fat intake (E%)	Ref.	2.12	1.11–4.05	1.93	0.94–3.97	Ref.	3.52	1.21–10.24	4.13	1.33–12.84 †
Fat ratio ‡	Ref.	1.03	0.57–1.88	1.14	0.60–2.18	Ref.	0.98	0.45–2.13	0.65	0.26–1.62
Low-fat milk	Ref.	1.80	1.02–3.20	1.03	0.54–1.94	Ref.	2.36	1.02–5.45	1.59	0.63–4.00
Fish	Ref.	1.00	0.57–1.76	0.74	0.38–1.47	Ref.	0.50	0.22–1.16	0.68	0.28–1.65
Red meat	Ref.	1.16	0.62–2.16	1.09	0.54–2.19	Ref.	2.58	1.01–6.60	1.28	0.44–3.72
Alcohol §	Ref.	0.77	0.45–1.35	0.48	0.25–0.91 †	Ref.	1.21	0.54–2.70	1.02	0.42–2.45
mMDS component										
All vegetables	Ref.	0.53	0.29–0.98	0.80	0.42–1.51	Ref.	0.46	0.18–1.17	1.82	0.79–4.19
All fruits and nuts	Ref.	0.71	0.39–1.30	0.55	0.29–1.05	Ref.	1.38	0.61–3.15	0.88	0.36–2.15
All cereals	Ref.	0.66	0.35–1.25	0.74	0.36–1.49	Ref.	0.54	0.24–1.22	0.26	0.09–0.73 †

Legumes	Ref.	0.58	0.32–1.07	0.94	0.52–1.70	Ref.	1.21	0.56–2.60	0.59	0.24–1.43
All dairy products	Ref.	0.78	0.43–1.42	0.81	0.44–1.50	Ref.	1.11	0.48–2.53	1.07	0.46–2.49
Fat ratio ‖	Ref.	0.74	0.40–1.36	0.99	0.53–1.87	Ref.	0.63	0.28–1.41	0.49	0.20–1.21
Fish ¶	Ref.	1.76	0.99–3.10	0.94	0.48–1.86	Ref.	1.10	0.50–2.41	0.91	0.36–2.25
Meat **	Ref.	1.31	0.71–2.43	1.13	0.56–2.27	Ref.	3.11	1.24–7.78	2.29	0.80–6.61
Alcohol §	Ref.	0.74	0.43–1.28	0.45	0.24–0.85 †	Ref.	1.14	0.52–2.50	0.78	0.33–1.85

440 OR, odds ratio; CI, confidence interval; NDS, Nordic diet score; mMDS, modified Mediterranean diet score; Ref., referent value.

441 * Adjusted for sex, age, body mass index, educational attainment, smoking status, physical activity, energy intake, presence of chronic diseases
442 and depression, and mutually adjusted for the diet score components.

443 † P for a linear trend <.05.

444 ‡ A ratio of polyunsaturated fatty acids to saturated and trans-fatty acids.

445 § Both diet scores included alcohol, calculated as 100% ethanol.

446 ‖ A ratio of unsaturated fatty acids to saturated fatty acids.

447 ¶ All fish and fish products.

448 ** All meat and meat products.

449

Appendix Table 1. Daily intakes of healthy Nordic diet score and modified Mediterranean diet score components by sex in 962 Helsinki Birth Cohort Study participants

Component (g/day)	Men	Women
NDS component		
Nordic vegetables	179 (121, 268)	231 (160, 331)
Nordic fruits	84 (30, 161)	116 (58, 207)
Nordic cereals	58 (31, 81)	54 (30, 75)
Fat intake (E%)	33 (29, 36)	32 (29, 36)
Fat ratio *	0.4 (0.4, 0.5)	0.4 (0.3, 0.5)
Low-fat milk	153 (13, 412)	86 (11, 221)
Fish	45 (29, 64)	33 (21, 45)
Red and processed meat	128 (87, 178)	80 (54, 119)
Alcohol †	11 (4, 19)	3 (2, 7)
mMDS component		
All vegetables	217 (149, 324)	275 (191, 392)
All fruits and nuts	206 (93, 331)	263 (147, 428)
All cereals	151 (108, 198)	131 (97, 164)
Legumes	9 (4, 14)	8 (4, 12)
All dairy products	394 (250, 695)	442 (270, 652)
Fat ratio ‡	1.4 (1.2, 1.6)	1.4 (1.1, 1.6)
Fish and fish products	53 (36, 77)	39 (25, 53)
All meat and meat products	168 (114, 222)	117 (82, 162)
Alcohol †	11 (4, 19)	3 (2, 7)

NDS, Nordic diet score; E%, percent of energy intake, mMDS, modified Mediterranean diet score. Values are medians (interquartile ranges).

* A ratio of polyunsaturated fatty acids to saturated and trans-fatty acids.

† Both diet scores included alcohol, calculated as 100 % ethanol.

‡ A ratio of unsaturated fatty acids to saturated fatty acids.