

**Bidirectional pathways between psychosocial risk factors and paranoid ideation
in a general non-clinical population**

Running head: Psychosocial risks and paranoid ideation

Aino Saarinen^{1,2,*}, Niklas Granö³, Mirka Hintsanen¹, Terho Lehtimäki⁴,

C. Robert Cloninger⁵, Liisa Keltikangas-Järvinen²

¹ Research Unit of Psychology, University of Oulu, Finland

² Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Finland

³ Helsinki University Hospital, Department of Adolescent Psychiatry, Finland

⁴ Department of Clinical Chemistry, Fimlab Laboratories and Finnish Cardiovascular Research Center-Tampere, Faculty of Medicine and Health Technology, Tampere University, Finland

⁵ Department of Psychiatry, Washington University, St. Louis, School of Medicine, and Anthropedia Foundation, United States

* **Corresponding author:** Aino Saarinen. Department of Psychology and Logopedics, Faculty of Medicine, Haartmaninkatu 3, P.O. Box 21, 00014 University of Helsinki, Finland. E-mail: aino.i.saarinen@helsinki.fi, Tel.: +35844 307 1204.

Acknowledgments

This study was supported financially by the Academy of Finland (M.H., grant number 308676). The Young Finns Study has been financially supported by the Academy of Finland: Grants 322098; 286284, 134309 (Eye), 126925, 121584, 124282, 129378 (Salve), 117797 (Gendi), and 41071 (Skidi); the Social Insurance Institution of Finland; Competitive State Research Financing of the Expert Responsibility area of Kuopio, Tampere and Turku University Hospitals (grant X51001); the Juho Vainio Foundation; the Sigrid Juselius Foundation; the Yrjö Jahnsson Foundation; the Paavo Nurmi Foundation; the Finnish Foundation of Cardiovascular Research and Finnish Cultural Foundation; the Tampere Tuberculosis Foundation; the Emil Aaltonen Foundation; and Diabetes Research Foundation of Finnish Diabetes Association.

Role of the funding source

The funding source had no role in study design, data collection, data analysis, data interpretation, writing of the report, or in the decision to submit the article for publication.

Conflict of interest

None.

Abstract

We investigated (i) whether psychosocial factors (experienced stress, anticipatory worry, social detachment, sleeping disturbances, alcohol use) predict the course of paranoid ideation between ages 24–50 years (ii) whether the predictive relationships are more likely to proceed from the psychosocial factors to paranoid ideation, or vice versa. The participants ($N=1534$ – 1553) came from the population-based Young Finns study. Paranoid ideation and psychosocial factors were assessed by reliable self-report questionnaires in 2001, 2007, and 2011/2012. The data was analyzed using growth curve and structural equation models. High experienced stress, anticipatory worry, social detachment, frequent sleeping disturbances, and frequent alcohol use predicted more paranoid ideation. More risk factors predicted increasing paranoid ideation. There were bidirectional predictive relationships of paranoid ideation with experienced stress, anticipatory worry, social detachment, and sleeping disturbances. The link between alcohol use and paranoid ideation was only correlative. In conclusion, paranoid ideation increases by reciprocal interactions with stress, worry, social detachment, and sleeping disturbances. The findings support the threat-anticipation model of paranoid ideation, providing important implications for treatment of paranoia.

Keywords

Paranoid; Stress; Social isolation; Sleep; Alcohol use

1 Introduction

Paranoid ideation refers to an unjustified suspiciousness towards others' motives (American Psychological Association, APA, 2013). Paranoid ideation exists along a continuum of symptoms in the general population, ranging from mild to severe paranoid ideation (e.g. Freeman et al., 2005; Freeman & Garety, 2014; Freeman, 2016). Mild paranoia includes, for example, temporary thoughts that there are negative rumours circulating about the self (Freeman & Garety, 2014). More severe paranoia refers to more persistent, distressing, and convincing beliefs that others aim to deliberately cause some kind of harm to the self (Freeman, 2016). Most severe paranoia refer to psychotic-level persecutory delusions (Freeman, 2016). Paranoid ideation is related to a wide variety of comorbid psychiatric disorders, such as depression, post-traumatic stress disorder, and even suicidality (Alsawy et al., 2015; Freeman et al., 2011). Importantly, most severe forms of paranoid ideation are suggested to be a crucial risk factor for conducting violent crimes (Coid et al., 2016; Heinrichs & Sam, 2012), especially mass shooting and other mass murders (Dutton et al., 2013; Knoll & Meloy, 2014). Since the severe forms of paranoia may be related to such adverse outcomes, it is necessary to investigate paranoid ideation in a much earlier phase to provide possibilities for early preventive interventions.

Along with this, there exist several psychological models about the early development of paranoid ideation. One of them is the threat-anticipation model of paranoia (Freeman, 2007, 2016). In this model, the following psychosocial factors play a crucial role: experienced stress (state-level stress related to e.g. stressful events), disposition to feel worry or anxiety, social avoidance (related to safety behavior), sleep disturbances (e.g. insomnia or disruptions of circadian rhythm), and anomalous experiences deriving from e.g. substance use (Freeman, 2007, 2016). This model has been supported by extensive prior evidence. For example, some studies have found that elevated stress levels predict increase in paranoid states (Kesting et al., 2013; Lincoln et al., 2009). Regarding social isolation,

previous studies have found that low social support and low self-confidence in social situations (Riggio & Kwon, 2009), frequent social fears (Marteinsdottir et al., 2003), high introversion (Tackett et al., 2014), and hostile interpretations of others' socioemotional cues (Bentall, 2001; Freeman, 2007; Tone & Davis, 2012) are related to higher paranoid ideation. Moreover, in line with Freeman's model, paranoid ideation is found to be related to anxiety (Freeman et al., 2008; Jylhä & Isometsä, 2006) and expectations of future negative events (Corcoran et al., 2006). Finally, cross-sectional studies have shown that alcohol use (Dawson et al., 2005; Grant et al., 2004) and sleep disturbances, particularly insomnia, are linked to higher paranoid ideation (Reeve et al., 2015; Scott et al., 2017).

Importantly, even though previous studies have supported the roles of experienced stressful states, disposition to feel worry, social avoidance, sleep disturbances, and alcohol use for paranoid ideation, longitudinal studies with prospective follow-ups of several years have been mostly lacking. Hence, several questions related to the etiology of paranoid ideation have remained unclear. To our knowledge, among previous studies, the longest follow-up of paranoid ideation has ranged 18 months in population-based studies (Freeman et al., 2012). In clinical populations, the longest follow-up has been in a study of 301 patients first admitted to a hospital with delusional psychoses and then followed up 5 to 18 years later by Retterstol and Opjordsmoen in Norway (Opjordsmoen, 1989). The study in Norway was primarily focused on the relationship of psychiatric diagnosis to outcome and provided only broad descriptions of the associated psychosocial variables.

Given that, due to the lack of long-term follow-ups of paranoid ideation, two essential topics have remained unclear. Firstly, it has remained largely unknown whether the associations of the psychosocial risk factors with paranoid ideation are stable over age or whether they change from early adulthood to middle age. There is evidence that there are age-related changes in the prevalence of paranoia-related risk factors: for example, level of anxiety proneness seems to slightly decrease over age (Zohar et al., 2019).

Secondly, more research has been recommended in order to clarify the temporal relationships of paranoid ideation with the psychosocial risk factors (i.e. experienced stressful states, disposition to feel worry, social detachment, sleep disturbances, and alcohol use). That is, whether these psychosocial factors are more likely to predict paranoid ideation or vice versa. On one hand, sleep problems and increased psychosocial stress are found to predict increase in paranoid states (Kesting et al., 2013; Scott et al., 2017; Thewissen et al., 2011). Additionally, increases in anxiety, worry, and insomnia predict the onset and persistence of paranoid episodes (Freeman et al., 2008, 2012). Further, alcohol use in the past is found to correlate with the current level of paranoid ideation (Grant et al., 2004). On the other hand, paranoid beliefs are suggested to result in experiences of anxiety and social isolation (Freeman, 2007). Additionally, paranoid ideation may commonly cause subjective suffering and stress (Freeman, 2007, 2016) that, in turn, may predispose to alcohol use as a way of stress coping.

The first aim of the present study was to investigate (i) whether experienced stress (stressful states), anticipatory worry (a more stable disposition to feel worry), social detachment, sleep disturbances, and alcohol use are associated with the course of paranoid ideation between ages 24–50 years. We also calculated a cumulative risk score of the risk factors since cumulative risk scores have been widely used previously (e.g. Copeland et al., 2009; el Bouhaddani et al., 2019) but, to our knowledge, in the research field of paranoid ideation not before. Our long follow-up provided exceptional possibilities to investigate the accumulation of the risk factors over age. The second aim was to examine whether the temporal relationships are more likely to proceed from the psychosocial factors to paranoid ideation or vice versa. We used a population-based sample with a 10-year prospective follow-up and several measurement times of paranoid ideation and the psychosocial risk factors. This rare population-based data provides a solid empirical foundation for identifying the conditions that increase the risk of paranoid ideation at both societal and individual levels.

2 Methods

2.1 Participants

We used data from the prospective Young Finns Study. The participants were selected randomly from six age cohorts (born between 1962–1977) from the population register of the Social Insurance Institution. The Social Insurance Institution covers the whole population of Finland. The original sample included 3596 participants in the baseline measurement in 1980 (when participants were aged 3–18 years). The participants have been followed since then so that the latest follow-up measurement was in 2012 (participants were aged 35–50 years). The study was carried out in accordance with the Declaration of Helsinki. Furthermore, the design of the Young Finns Study was approved by all the Finnish universities with medical schools. Before participation, all the participants or their parents (for participants aged below 12 years) provided informed consent after the nature of the procedures had been fully explained. The design of the Young Finns Study is described with more detail elsewhere (see Raitakari et al., 2008).

For this study, paranoid ideation, experienced stress, anticipatory worry, social detachment, sleep disturbances, and alcohol use were evaluated in 2001, 2007, and 2011/2012; participants' socioeconomic factors in 2011; and parents' socioeconomic factors in 1980. In the analyses, we included all the participants with data available on the study variables. The final sample included 1534–1553 participants.

2.2 Measures

2.2.1 Cumulative risk score

Psychosocial risk factors included anticipatory worry, social detachment, experienced stress, sleep disturbances, and alcohol use in 2001, 2007, and 2012. For each measurement year, we calculated the

total score of each risk factor for all the participants who had responded to at least 80% of the items. The missing values (for participants with at least 20% values missing) were imputed with the mean of each scale. We calculated both continuous and dichotomous (0=risk not present; 1=risk present) variables for all the psychosocial risk factors. A cumulative risk score was calculated as the total score of the five dichotomous psychosocial risk factors. Hence, continuous scores were used in the main analyses reported in the tables; and dichotomous variables were used only when calculating the cumulative risk score and when illustrating the findings in the figure.

Anticipatory worry was evaluated with the Anticipatory Worry scale of the Temperament and Character Inventory (TCI) (Cloninger et al., 1994). The Anticipatory Worry scale included 11 items (e.g. “I’m never worried about terrible things that could happen me in the future” (reversed); Cronbach’s $\alpha=.80-.81$ in 2001, 2007, and 2012) that were responded with a 5-point scale (1=definitely true; 5=definitely false). We used this scale to measure trait-level disposition to feel worry. The test-retest correlations of the scale were high ($r=0.661-0.751$ between different measurement times in 2001, 2007, and 2012), indicating that the scale measured a comparatively stable disposition. The items were reversed so that high scores referred to higher anticipatory worry. High anticipatory worry was defined to be present if the mean score of the scale was >3 .

Social detachment was evaluated with the Attachment scale of the TCI. The Attachment scale included 8 items (e.g. “I usually discuss my experiences and feelings with my friends openly, instead of keeping them inside my mind” (reversed); Cronbach’s $\alpha=.82-.84$ in 2001, 2007, and 2012) that were responded with a 5-point scale (1=definitely true; 5=definitely false). The items were reversed so that high scores referred to lower attachment (i.e. higher social detachment). High social detachment was defined to be present if the mean score of the reversed scale was >3 . There is previous evidence that high social detachment is related to more severe challenges in social functioning (including challenges in social activities and initiatives and aspects of avoidance tendencies) (Jetha et

al., 2013). Additionally, high social detachment correlates with higher scores of asociality (Gutiérrez et al., 2014) and higher scores of detachment (Jönsson et al., 2003).

Experienced stress was evaluated with one item (“Stress refers to such a situation where one feels tensioned up, restless, nervous, or anxious, or where one experiences sleep disturbances because things are bothering one’s mind. Do you experience this kind of stress?”) that was responded with a 5-point scale (1=not at all; 2=only a bit; 3=somewhat; 4=quite much; 5=very much). High experienced stress was defined to be present if the participant reported experiencing stress quite much or very much. Contrary to the scale of Anticipatory Worry (measuring rather trait-level disposition to feel worry), we used the item of experienced stress to measure state-level experiences of stress. In accord with this difference, the correlation between the item of experienced stress and the scale of Anticipatory Worry was comparatively low ($r=0.387$, see Supplementary Table 2). Further, the test-retest correlations were comparatively low ($r=0.373$ - 0.454 between different measurement points in 2001, 2007, and 2012), suggesting that this item measured rather state-level or situation-dependent experiences of stress. Along with this, high scores of the item of experienced stress were found to correlate with frequent stressful life events ($r=0.528$, $p<.001$) and high stress reactivity ($r=0.421$, $p<.001$) (see Supplementary Material for a more detailed description).

Sleep disturbances were assessed with two items (“Do you have difficulties to fall asleep?”; “Do you wake up at nights?”) that were responded with a 3-point scale (0=no; 1=not sure; 2=yes). The items were selected from the Vital Exhaustion Questionnaire (Appels et al., 1987) on the basis of an exploratory factor analysis of the items (see Supplementary Table 1). The reliability and validity of the Vital Exhaustion Questionnaire have been confirmed previously (e.g. Chumaeva et al., 2010; Meesters & Appels, 1996). Frequent sleep disturbances were defined to be present if the participant responded “yes” to at least one of the items, i.e. reported having difficulties in falling asleep or being awake at nights.

Alcohol use was evaluated by asking the participants the frequency of alcohol intoxication (i.e. consuming at least six portions at one time). The frequency was evaluated with a 6-point scale (1=two times per week or more often; 6=once a year or more seldom). Frequent alcohol use was defined to be present if the participant reported becoming intoxicated at least once a week.

2.2.2 Paranoid ideation

Paranoid ideation was evaluated with the Paranoid Ideation Scale of the Symptom Checklist-90 Revised (SCL-90R; Derogatis, 1986). The scale consists of 6 items (e.g. “I think that other people would take advantage of me if I let them to do that”) that are responded with a 5-point scale (1=totally disagree; 5=totally agree). We calculated the mean score of the items for each measurement year for those participants who had responded to at least 80% of the items. The Paranoid Ideation Scale has been used also previously (e.g. Saarinen et al., 2018a,b). Previous studies have confirmed good reliability for the SCL-90R and for the subscale of paranoid ideation (e.g. Olsen et al., 2004; Schmitz et al., 2000). The internal reliability of the scale was good also in our data (Cronbach’s α =.75–.79 in 2001, 2007, and 2012).

2.2.3 Covariates

Socioeconomic factors included participants’ and their parents’ level of income and educational level. Parents’ level of income was assessed with a 8-point scale (1=less than 15 000 Finnish mark per year; 8=more than 100 000 Finnish mark per year). Participants’ level of income was measured with a 13-point scale (1=less than 5 000€ per year; 13=more than 60 000€ per year). Participants’ and their parents’ educational level was classified into three categories (1=comprehensive school; 2=high school or occupational school; 3=academic level). In case mother’s and father’s educational levels were

different, we used the higher level of education.

2.3 Statistical analyses

Statistical analyses were conducted with STATA SE (version 13.0). The association of psychosocial risk factors with paranoid ideation was investigated using multilevel models for longitudinal design (growth curve models). Growth curve models estimate “fixed effects” that refer to classic regression coefficients, and “random effects” that refer to the individual-level variance in the intercept, slopes, and residual variance (i.e. within-individual variance over the follow-up). Before conducting the analyses, the scores of psychosocial risk factors and the cumulative risk score were standardized with the mean and standard deviation of year 2001 scores, in order to stabilize the growth curve trajectories over the follow-up measurements. In all the models, the dependent variable was the course of paranoid ideation (in 2001–2012). We estimated fixed effects for the continuous scores of psychosocial risk factors, age, age-squared, and the age-interactions of the psychosocial risk factors. Continuous age was set also as random effect. As the sample consisted of six age cohorts (born in 1962–1977), the age range of the participants varied between 24–50 years in 2001–2012. Age was centered to 24 years (the age of the youngest participants in 2001). We included age-squared as fixed effect since previous studies have shown that there exist curvilinear trends in the development of paranoid ideation over age (e.g. Saarinen et al., 2018). All the models were adjusted for sex and participants’ and their parents’ socioeconomic factors.

The temporal associations of psychosocial factors with paranoid ideation were investigated using structural equation models with cross-lagged panel design (MLMV estimation). Separate models were estimated for each psychosocial risk factor (i.e. anticipatory worry, social detachment, experienced stress, sleep disturbances, and alcohol use) to investigate its temporal associations with paranoid ideation. We conducted altogether 4 models: (1) a model including only

stability coefficients (the variables at each time point were predicted by the same variables at the next time points) and covariances between the psychosocial risk factor and paranoid ideation at each time point; (2) a model that included also the predictive coefficients from the psychosocial risk factor at each time point to paranoid ideation at the next time points; (3) a model that included the predictive coefficients in the opposite direction (i.e. from paranoid ideation at each time point to the psychosocial risk factor at the next time points); and (4) a model with the cross-lagged predictive coefficients in both directions between the psychosocial risk factor and paranoid ideation. The models 1-4 are illustrated in Supplementary Figure 1. In all the models, we used continuous scores of the psychosocial risk factors.

We evaluated the goodness-of-fit of the models 1–4 by investigating the values of the Comparative Fit Index (CFI), the Root-Mean-Square Error of Approximation (RMSEA), and the Bayesian Information Criterion (BIC). Previously, it has been recommended that CFI should be >0.95 and RMSEA should be <0.06 (Hu & Bentler, 1999). Moreover, lower scores of the χ^2 test of absolute model fit and BIC indicate better model-fit (Schreiber et al., 2006). The final selection of the model with best statistical fit was done on the basis of the χ^2 test score of absolute model fit.

3 Results

The descriptive statistics of the study variables are shown in Table 1. The correlations between the study variables can be found in Supplementary Table 2. Attrition analyses showed that included participants were slightly older than excluded participants (31.681 vs. 31.261, $p<.013$). Further, women were more likely to participate than men (48.090% vs. 38.095%, $p<.001$). At the baseline measurement point (in 2001), there was no attrition bias in experienced stress or social detachment. However, included participants had slightly lower levels of anticipatory worry (2.430 vs. 2.507, $p=.0019$), sleep

disturbances (0.416 vs. 0.489, $p=.014$), and alcohol use (2.474 vs. 2.683, $p<.001$). Moreover, included participants had slightly lower level of paranoid ideation (2.281 vs. 2.414, $p<.001$). Regarding socioeconomic factors, included participants had slightly higher level of income than excluded participants (7.458 vs. 7.013, $p<.011$) and were substantially more likely to have high educational level (44.559% vs. 9.173%, $p<.001$). With regard to parental socioeconomic factors, included participants' parents had slightly higher level of income (4.962 vs. 4.658, $p<.001$) and were more likely to have high educational level (26.980% vs. 23.402%, $p=.015$). Previously, it has been found that values are missing at random in the psychosocial variables of the Young Finns data (Pulkki-Råback et al., 2015).

(TABLE 1 ABOUT HERE)

The results of the growth curve models are presented in Table 2. High experienced stress, high anticipatory worry, high social detachment, frequent alcohol use, and frequent sleep disturbances had positive main effects on the trajectory of paranoid ideation. When predicting the trajectory of paranoid ideation, there were no significant age-interactions with experienced stress, anticipatory worry, sleep disturbances, or alcohol use. This indicated that the associations of these risk factors (experienced stress, anticipatory worry, sleep disturbances, and alcohol use) with paranoid ideation did not change significantly over age. However, there were significant age-interactions with social detachment, indicating that age modified the association of social detachment with paranoid ideation. All the findings were controlled for age, sex, and participants' and their parents' socioeconomic factors. The results are illustrated in Figure 1.

Additionally, the results showed that high cumulative risk score predicted higher trajectory of paranoid ideation. The accumulation of risk factors seemed to result in larger differences

in paranoid ideation than any single risk factor. That is, differences in paranoid ideation between participants with low vs. high accumulation of risk factors seemed to be greater than the differences between participants with low vs. high value of any single risk factor.

(TABLES 2A AND 2B ABOUT HERE)

(FIGURE 1 ABOUT HERE)

Next, we investigated the temporal associations of psychosocial risk factors with paranoid ideation with structural equation models, when each psychological risk factor was added to the model separately. The goodness-of-fit indexes of the structural equation models are shown in Table 3. When investigating the temporal associations of experienced stress, anticipatory worry, and sleep disturbances with paranoid ideation, model 4 (i.e. predictive relationships in both directions between the psychosocial risk factors and paranoid ideation) fit the data better than any other model (models 1–3). With regard to the temporal associations of social detachment with paranoid ideation, model 2 (i.e. predictive pathways from social detachment to paranoid ideation but not in the opposite direction) fit the data better than any other model (models 1, 3 or 4). Regarding the temporal associations of alcohol use with paranoid ideation, model 1 (i.e. no predictive relationships between alcohol use and paranoid ideation) fit the data better than any other model (models 2–4). Taken together, the results indicated that there existed bidirectional associations of experienced stress, anticipatory worry, and sleep disturbances with paranoid ideation. That is, high experienced stress, high anticipatory worry, and frequent sleep disturbances predicted increase paranoid ideation, and vice versa. Further, high social detachment was found to predict increase in paranoid ideation but not vice versa. Additionally, there were no predictive

relationships between alcohol use and paranoid ideation. Hence, the association of alcohol use with paranoid ideation appeared to be only correlative.

4 Discussion

To our knowledge, this study was the first to investigate the relationship of psychosocial risk factors with paranoid ideation over a long-term prospective follow-up. Overall, the results provide strong support for Freeman's (2007, 2016) model of paranoid ideation. High experienced stress (i.e. state-level experiences of stress), high anticipatory worry (i.e. a comparatively stable disposition to feel worry), high social detachment, frequent sleep disturbances, and frequent alcohol use predicted higher level of developmental curve of paranoid ideation from early adulthood to middle age. Additionally, the accumulation of the risk factors predicted higher trajectory of paranoid ideation. Moreover, there were bidirectional predictive relationships of experienced stress, anticipatory worry, and sleep disturbances with paranoid ideation. That is, high experienced stress, high anticipatory worry, and frequent sleep disturbances predicted increase paranoid ideation, and vice versa. Further, high social detachment was found to predict increase in paranoid ideation but not vice versa. Additionally, we obtained no predictive relationships between paranoid ideation and alcohol use, indicating that this association appears to be only correlative.

There are several explanations for the pathways from psychosocial factors to paranoid ideation. Firstly, elevated stress may increase aberrant salience of some socioemotional stimuli and result in a higher level of anxiety and, in this way, increase risk for delusional ideation (Lincoln et al., 2009; Westermann et al., 2012). Additionally, experienced stress may increase risk for making conclusions on the basis of limited evidence ("jumping to conclusions") and attributing events to one causal factor, instead of considering the wider circumstances (Moritz et al., 2015) that, in turn,

predispose to paranoid ideation (Lincoln et al., 2009; Moritz et al., 2018; Tone & Davis, 2012).

Secondly, social detachment may reduce the possibilities to receive contradictory evidence for one's current paranoid ideas (Morse & Lynch, 2004). In some cases, social detachment may also be an indicator of weak self-esteem (Hall-Lande et al., 2007) that is suggested to be a core factor in the etiology of paranoid episodes (Thewissen et al., 2011). Finally, sleep disturbances predispose to more frequent anomalous experiences such as depersonalization (Watson, 2001), elevated levels of anxiety (Reeve et al., 2015), and reduced abilities to sustain attention on non-emotional stimuli and to leave behind negative memories (Gobin et al., 2015). All these factors, in turn, are known to elevate the risk for paranoid thoughts (Freeman, 2007). Taken together, the psychosocial risk factors may act in a reciprocal and complicated interplay in the onset and maintenance of paranoid ideation.

Importantly, high paranoid ideation was also found to predict increases in sleep disturbances, experienced stress, and anticipatory worry. In the current classifications of mental disorders, it is stated that a marker of clinical significance of paranoid ideation is that the paranoid symptoms cause subjective stress and anxiety (APA, 2013). Our findings in a non-clinical population suggest that also subclinical paranoia may predict increase in experienced stress and worry. Elevated levels of stress and worry may also provide an explanation for the pathway from paranoid ideation to increased sleep disturbances. Taken together, there may exist vicious circles between paranoid ideation and psychosocial factors that increase the risk for maintenance of paranoid ideas.

Our findings suggested that the association of paranoid ideation with alcohol use is correlative. That is, even though frequent alcohol users have on average higher level of paranoid ideation, frequent alcohol use may not predict increase in paranoid ideation or vice versa. This suggests that the link between frequent alcohol use and higher paranoid ideation may be explained by third factors that predispose to both alcohol use and paranoid ideation. Such factors could include for instance the use of other substances such as cannabis or street drugs (Grant et al., 2004; D'Souza et al.,

2004; Verdoux et al., 2003). Overall, frequent alcohol use may have a more prominent role for some other psychotic symptom dimensions than paranoid ideation, for example, hallucinations (APA, 2013).

This study had some limitations that need to be taken into consideration. Firstly, the measures of sleep disturbances and stress included only single items. Nevertheless, the items of sleep disturbances were selected from the Vital Exhaustion Questionnaire that has been widely used and validated previously (Chumaeva et al., 2010; Meesters & Appels, 1996). The construct validity of the items was confirmed with an exploratory factor analysis. Regarding the stress item, we used it to measure state-level experiences of stress (contrary to the scale of anticipatory worry that was used to measure rather trait-level disposition to feel worry). The stress item had comparatively low test-retest correlations. Further, the correlation of the stress item was quite low with anticipatory worry (i.e. trait-level disposition) but higher with stressful life events. All this provided support that the item provided information about participants' state-level experiences of stress. Additionally, the face validity of the item was good: stress was first defined for the participants in order to ensure that the participants have a similar conception of experienced stress. Overall, more research is needed with more comprehensive and widely validated measures of stress and sleep disturbances.

Secondly, the level of paranoid ideation was mostly subclinical in our population-based sample. Hence, the results may not be fully generalized to clinical populations like those studied by Retterstol in Norway (Retterstol, 1966). However, findings in clinical and non-clinical samples have suggested that severe paranoid ideation exists on the same continuum with subclinical paranoid ideation (Johns & Van Os, 2001; Van Os, 2003; Verdoux & Van Os, 2002). Along with this, it has been stated that the involved underlying psychological processes appear to be quite similar in subclinical and clinical paranoia (Elahi et al., 2017). This could be further investigated by considering the temperament and character features that may mediate the effects of psychosocial stressors on the development of paranoid ideation.

With regard to practical implications, there is encouraging evidence that cognitive-behavioral interventions targeting at interpersonal sensitivity, worry, or insomnia (e.g. by means of psychoeducation, practical advice with sleeping circumstances, and relaxation techniques) effectively reduce paranoid ideation (Bell & Freeman, 2014; Freeman et al., 2015; Myers et al., 2011). Moreover, interventions that aim to increase metacognitive awareness of emotional processing or reasoning biases seem to be effective treatments for paranoid ideation or paranoia-related distress (Hepworth et al., 2011; Waller et al., 2015). Additionally, interventions aiming to develop a mature character profile could be an effective intervention for individuals with elevated risk for paranoid ideation. Specifically, mature character profiles including high self-directedness and high cooperativeness are found to act as protective factors against paranoid ideation in individuals with vulnerable temperament profiles (Saarinen et al., 2018b).

On the basis of our findings, early interventions should carefully consider that there may exist vicious circles between psychosocial risks and paranoid ideation. Additionally, these bidirectional circles between paranoid ideation and psychosocial risks could provide treatment targets for patients aged from early adulthood to middle age. A common challenge has been that paranoid individuals may not commonly seek help for their paranoid beliefs. However, paranoid ideation could be monitored when treating patients with sleep disturbances or anxiety since they appear to be crucial correlates of paranoid ideation. Additionally, our results provide encouraging evidence that reducing the cumulative number of single psychosocial risks may predict a long-term decrease in paranoid ideation, even though early interventions could not efficiently target all the psychosocial risks.

In summary, our findings suggest a diathesis-stress model for elicitation of persecutory ideation in which threats of personal and social harm or injustice precipitate paranoia, particularly in hypersensitive individuals susceptible to feeling inferior or powerless to defend themselves. This has important implications for promotion of health at both individual and societal levels. In individual

cases, it is important to use person-centered approaches in which paranoid individuals are treated with respect and compassion in order to provide the conditions for developing more trust and feelings of security (O'Connor et al, 2007). In communities and larger societies, increases in paranoid ideation may serve as a bellwether of a breakdown in civility and social unity due to increasing perception of social injustice, lack of humane concerns for others, and threats of harm or punishment. In addition to providing person-centered services to vulnerable and alienated individuals, our findings suggest that reductions in paranoid ideation in individuals and societies require community-based efforts to improve social connectedness by creating more civil and equitable conditions of living based on respect for the dignity of all people.

References

- Alsawy, S., Wood, L., Taylor, P. J., & Morrison, A. P. (2015). Psychotic experiences and PTSD: exploring associations in a population survey. *Psychological Medicine*, 45, 2849–2859.
- American Psychiatric Association (APA) (2013). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub.
- Appels, A. P. W. M., Höppener, P., & Mulder, P. (1987). A questionnaire to assess premonitory symptoms of myocardial infarction. *International Journal of Cardiology*, 17, 15-24.
- Bell, V., & Freeman, D. (2014). A pilot trial of cognitive behavioural therapy for interpersonal sensitivity in individuals with persecutory delusions. *Journal of Behavior Therapy and Experimental Psychiatry*, 45, 441-446.
- Bentall, R. P., de Sousa, P., Varese, F., Wickham, S., Sitko, K., Haarmans, M., & Read, J. (2014). From adversity to psychosis: pathways and mechanisms from specific adversities to specific symptoms. *Social Psychiatry and Psychiatric Epidemiology*, 49, 1011-1022.

- Berry, K., Bucci, S., Kinderman, P., Emsley, R., & Corcoran, R. (2015). An investigation of attributional style, theory of mind and executive functioning in acute paranoia and remission. *Psychiatry Research*, 226, 84-90.
- Chan, R. C., Li, X., Lai, M. K., Li, H., Wang, Y., Cui, J., Deng, Y., & Raine, A. (2011). Exploratory study on the base-rate of paranoid ideation in a non-clinical Chinese sample. *Psychiatry Research*, 185, 254–260.
- Chumaeva, N., Hintsanen, M., Juonala, M., Raitakari, O. T., & Keltikangas-Järvinen, L. (2010). Sex differences in the combined effect of chronic stress with impaired vascular endothelium functioning and the development of early atherosclerosis: the Cardiovascular Risk in Young Finns study. *BMC Cardiovascular Disorders*, 10, 34.
- Cloninger, C. R., Svrakic, D. M. (2017). Personality Disorders. In Sadock, B. J., Sadock, V., & Ruiz, P. (Eds.): *Kaplan and Sadock's Comprehensive Textbook of Psychiatry* (pp. 2126-2176).
- Coid, J. W., Ullrich, S., Bebbington, P., Fazel, S., & Keers, R. (2016). Paranoid ideation and violence: meta-analysis of individual subject data of 7 population surveys. *Schizophrenia Bulletin*, 42, 907-915.
- Copeland, W., Shanahan, L., Jane Costello, E., & Angold, A. (2009). Configurations of common childhood psychosocial risk factors. *Journal of Child Psychology and Psychiatry*, 50(4), 451-459.
- Dawson, D. A., Grant, B. F., Stinson, F. S., & Chou, P. S. (2005). Psychopathology associated with drinking and alcohol use disorders in the college and general adult populations. *Drug and Alcohol Dependence*, 77, 139-150.
- Derogatis, L. R. (1986). Manual for the Symptom Checklist 90 Revised (SCL-90R). *Baltimore, MD*.
- D'Souza, D. C., Perry, E., MacDougall, L., Ammerman, Y., Cooper, T., Braley, G., ... & Krystal, J. H. (2004). The psychotomimetic effects of intravenous delta-9-tetrahydrocannabinol in healthy individuals: implications for psychosis. *Neuropsychopharmacology*, 29, 1558.

- Dutton, D. G., White, K. R., & Fogarty, D. (2013). Paranoid thinking in mass shooters. *Aggression and Violent Behavior, 18*, 548-553.
- Elahi, A., Algorta, G. P., Varese, F., McIntyre, J. C., & Bentall, R. P. (2017). Do paranoid delusions exist on a continuum with subclinical paranoia? A multi-method taxometric study. *Schizophrenia Research, 190*, 77-81.
- el Bouhaddani, S., van Domburgh, L., Schaefer, B., Doreleijers, T. A., & Veling, W. (2019). Psychotic experiences and trauma predict persistence of psychosocial problems in adolescence. *European Child & Adolescent Psychiatry*, 1-10.
- Freeman, D. (2007). Suspicious minds: the psychology of persecutory delusions. *Clinical Psychology Review, 27*, 425-457.
- Freeman, D. (2016). Persecutory delusions: a cognitive perspective on understanding and treatment. *The Lancet Psychiatry, 3*, 685-692.
- Freeman, D., Dunn, G., Startup, H., Pugh, K., Cordwell, J., Mander, H., ... & Kingdon, D. (2015). Effects of cognitive behaviour therapy for worry on persecutory delusions in patients with psychosis (WIT): a parallel, single-blind, randomised controlled trial with a mediation analysis. *The Lancet Psychiatry, 2*, 305-313.
- Freeman, D., & Freeman, J. (2008). *Paranoia: the 21st century fear*. Oxford University Press, Oxford.
- Freeman, D., Freeman, J., & Garety, P. (2016). *Overcoming paranoid and suspicious thoughts: A self-help guide using cognitive behavioural techniques*. Hachette UK.
- Freeman, D., Garety, P. A., Bebbington, P. E., Smith, B., Rollinson, R., Fowler, D., Kuipers, E., Ray, K., & Dunn, G. (2005). Psychological investigation of the structure of paranoia in a non-clinical population. *The British Journal of Psychiatry, 186*, 427-435.
- Freeman, D., & Garety, P. (2014). Advances in understanding and treating persecutory delusions: a review. *Social Psychiatry and Psychiatric Epidemiology, 49*, 1179-1189.

- Freeman, D., Garety, P. A., Kuipers, E., Fowler, D., & Bebbington, P. E. (2002). A cognitive model of persecutory delusions. *British Journal of Clinical Psychology*, *41*, 331-347.
- Freeman, D., McManus, S., Brugha, T., Meltzer, H., Jenkins, R., & Bebbington, P. (2011). Concomitants of paranoia in the general population. *Psychological Medicine*, *41*, 923-936.
- Freeman, D., Pugh, K., Antley, A., Slater, M., Bebbington, P., Gittins, M., ... & Garety, P. (2008). Virtual reality study of paranoid thinking in the general population. *The British Journal of Psychiatry*, *192*, 258-263.
- Freeman, D., Stahl, D., McManus, S., Meltzer, H., Brugha, T., Wiles, N., & Bebbington, P. (2012). Insomnia, worry, anxiety and depression as predictors of the occurrence and persistence of paranoid thinking. *Social Psychiatry and Psychiatric Epidemiology*, *47*, 1195-1203.
- Frenkel, T. I., Lamy, D., Algom, D., & Bar-Haim, Y. (2009). Individual differences in perceptual sensitivity and response bias in anxiety: Evidence from emotional faces. *Cognition and Emotion*, *23*, 688-700.
- Gobin, C. M., Banks, J. B., Fins, A. I., & Tartar, J. L. (2015). Poor sleep quality is associated with a negative cognitive bias and decreased sustained attention. *Journal of Sleep Research*, *24*, 535-542.
- Grant, B. F., Hasin, D. S., Stinson, F. S., Dawson, D. A., Chou, S. P., Ruan, W., & Pickering, R. P. (2004). Prevalence, correlates, and disability of personality disorders in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Clinical Psychiatry*, *65*, 948-958.
- Grant, J. D., Scherrer, J. F., Lynskey, M. T., Lyons, M. J., Eisen, S. A., Tsuang, M. T., ... & Bucholz, K. K. (2006). Adolescent alcohol use is a risk factor for adult alcohol and drug dependence: evidence from a twin design. *Psychological Medicine*, *36*, 109-118.
- Grant, B. F., Stinson, F. S., Dawson, D. A., Chou, S. P., Ruan, W. J., & Pickering, R. P. (2004). Co-occurrence of 12-month alcohol and drug use disorders and personality disorders in the United

- States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Archives of General Psychiatry*, 61, 361-368.
- Gutiérrez, F., Vall, G., Peri, J. M., Gárriz, M., & Garrido, J. M. (2014). A hierarchical model of normal and abnormal personality up to seven factors. *Comprehensive Psychiatry*, 55, 326-335.
- Hall-Lande, J. A., Eisenberg, M. E., Christenson, S. L., & Neumark-Sztainer, D. (2007). Social isolation, psychological health, and protective factors in adolescence. *Adolescence*, 42.
- Heinrichs, R. W., & Sam, E. P. (2012). Schizophrenia and Crime: How Predictable Are Charges, Convictions and Violence. *International Journal of Mental Health and Addiction*, 10, 122–131.
- Hepworth, C., Startup, H., & Freeman, D. (2011). Developing treatments of persistent persecutory delusions: the impact of an emotional processing and metacognitive awareness intervention. *The Journal of Nervous and Mental Disease*, 199, 653-658.
- Hong, J. P., Samuels, J., BIENVENU, O. J., Hsu, F. C., Eaton, W. W., Costa, P. T., & Nestadt, G. (2005). The longitudinal relationship between personality disorder dimensions and global functioning in a community-residing population. *Psychological Medicine*, 35, 891-895.
- Howes, O. D., & Murray, R. M. (2014). Schizophrenia: an integrated sociodevelopmental-cognitive model. *The Lancet*, 383, 1677-1687.
- Jetha, M. K., Goldberg, J. O., & Schmidt, L. A. (2013). Temperament and its relation to social functioning in schizophrenia. *International Journal of Social Psychiatry*, 59, 254-263.
- Jönsson, E. G., Cichon, S., Gustavsson, J. P., Grünhage, F., Forslund, K., Mattila-Evenden, M., ... & Nöthen, M. M. (2003). Association between a promoter dopamine D2 receptor gene variant and the personality trait detachment. *Biological Psychiatry*, 53, 577-584.
- Kendler, K. S., & Hays, P. (1981). Paranoid psychosis (delusional disorder) and schizophrenia: A family history study. *Archives of General Psychiatry*, 38, 547-551.

- Kesting, M. L., Bredenpohl, M., Klenke, J., Westermann, S., & Lincoln, T. M. (2013). The impact of social stress on self-esteem and paranoid ideation. *Journal of Behavior Therapy and Experimental Psychiatry*, 44, 122-128.
- Key, F. A., Craske, M. G., & Reno, R. M. (2003). Anxiety-based cognitive-behavioral therapy for paranoid beliefs. *Behavior Therapy*, 34, 97-115.
- Kleck, G. (2009). Mass shootings in schools: The worst possible case for gun control. *American Behavioral Scientist*, 52, 1447-1464.
- Knoll, J. L., & Meloy, J. R. (2014). Mass murder and the violent paranoid spectrum. *Psychiatric Annals*, 44, 236-243.
- Lincoln, T. M., Lange, J., Burau, J., Exner, C., & Moritz, S. (2009). The effect of state anxiety on paranoid ideation and jumping to conclusions. An experimental investigation. *Schizophrenia Bulletin*, 36, 1140-1148.
- Lincoln, T. M., Peter, N., Schäfer, M., & Moritz, S. (2009). Impact of stress on paranoia: an experimental investigation of moderators and mediators. *Psychological Medicine*, 39, 1129-1139.
- Linscott, R. J., & Van Os, J. (2013). An updated and conservative systematic review and meta-analysis of epidemiological evidence on psychotic experiences in children and adults: on the pathway from proneness to persistence to dimensional expression across mental disorders. *Psychological Medicine*, 43, 1133-1149.
- McGorry, P. D. (2010). Risk syndromes, clinical staging and DSM V: new diagnostic infrastructure for early intervention in psychiatry. *Schizophrenia Research*, 120, 49-53.
- Meesters, C., & Appels, A. (1996). An interview to measure vital exhaustion. II. Reliability and validity of the interview and correlations of vital exhaustion with personality characteristics. *Psychology and Health*, 11, 573-581.

- Mohr, P., Csemy, L., Rodriguez, M., Cermak, J., Kawaciukova, R., & Seifertova, D. (2008). Prevalence of psychotic symptoms in the general population of the Czech Republic. *European Psychiatry*, 23, 132–133.
- Moorhouse, J. C., & Wanner, B. (2006). Does gun control reduce crime or does crime increase gun control. *Cato Journal*, 26, 103-124.
- Moritz, S., Bentall, R. P., Kolbeck, K., & Roesch-Ely, D. (2018). Monocausal attribution and its relationship with reasoning biases in schizophrenia. *Schizophrenia Research*, 193, 77-82.
- Moritz, S., Köther, U., Hartmann, M., & Lincoln, T. M. (2015). Stress is a bad advisor. Stress primes poor decision making in deluded psychotic patients. *European Archives of Psychiatry and Clinical Neuroscience*, 265, 461-469.
- Morse, J. Q., & Lynch, T. R. (2004). A preliminary investigation of self-reported personality disorders in late life: prevalence, predictors of depressive severity, and clinical correlates. *Aging & Mental Health*, 8, 307–315.
- Myers, E., Startup, H., & Freeman, D. (2011). Cognitive behavioural treatment of insomnia in individuals with persistent persecutory delusions: a pilot trial. *Journal of Behavior Therapy and Experimental Psychiatry*, 42, 330-336.
- Olsen, L. R., Mortensen, E. L., & Bech, P. (2004). The SCL-90 and SCL-90R versions validated by item response models in a Danish community sample. *Acta Psychiatrica Scandinavica*, 110, 225-229.
- Opjordsmoen, S. (1989). Delusional disorders: I. Comparative long-term outcome. *Acta Psychiatrica Scandinavica*, 80, 603-612.
- Opjordsmoen, S., & Retterstøl, N. (1993). Outcome in delusional disorder in different periods of time. *Psychopathology*, 26, 90-94.

- Pulkki-Råback, L., Elovainio, M., Hakulinen, C., Lipsanen, J., Hintsanen, M., Jokela, M., ... & Keltikangas-Järvinen, L. (2015). Cumulative effect of psychosocial factors in youth on ideal cardiovascular health in adulthood: the Cardiovascular Risk in Young Finns Study. *Circulation*, 131, 245-253.
- Raitakari, O. T., Juonala, M., Rönnemaa, T., Keltikangas-Järvinen, L., Räsänen, L., Pietikäinen, M., ..., Jula, A. (2008). Cohort profile: the cardiovascular risk in Young Finns Study. *Int J Epidemiol*, 37, 1220–1226.
- Reeve, S., Sheaves, B., & Freeman, D. (2015). The role of sleep dysfunction in the occurrence of delusions and hallucinations: a systematic review. *Clinical Psychology Review*, 42, 96-115.
- Retterstol, N. (1966). *Paranoid and Paranoiac Psychoses*. Charles C. Thomas Publisher, Springfield, Illinois.
- Retterstol, N. (1970). *Prognosis in Paranoid Psychoses*. Springfield, Illinois: Charles C. Thomas Publisher.
- Riggio, H. R., & Kwong, W. Y. (2009). Social skills, paranoid thinking, and social outcomes among young adults. *Personality and Individual Differences*, 47, 492-497.
- Saarinen, A., Hintsanen, M., Hakulinen, C., Pulkki-Råback, L., Lehtimäki, T., Raitakari, O., & Keltikangas-Järvinen, L. (2018a). The co-occurrence between depressive symptoms and paranoid ideation: A population-based longitudinal study. *Journal of Affective Disorders*, 229, 48-55.
- Saarinen, A., Rosenström, T., Hintsanen, M., Hakulinen, C., Pulkki-Råback, L., Lehtimäki, T., ... & Keltikangas-Järvinen, L. (2018b). Longitudinal associations of temperament and character with paranoid ideation: a population-based study. *Psychiatry Research*, 261, 137-142.
- Schmitz, N., Hartkamp, N., Kiuse, J., Franke, G. H., Reister, G., & Tress, W. (2000). The symptom check-list-90-R (SCL-90-R): a German validation study. *Quality of Life Research*, 9, 185-193.

- Scott, J., Leboyer, M., Hickie, I., Berk, M., Kapczinski, F., Frank, E., ... & McGorry, P. (2013). Clinical staging in psychiatry: a cross-cutting model of diagnosis with heuristic and practical value. *The British Journal of Psychiatry*, 20, 243-245.
- Scott, A. J., Rowse, G., & Webb, T. L. (2017). A structural equation model of the relationship between insomnia, negative affect, and paranoid thinking. *PloS one*, 12, e0186233.
- Sitko, K., Bentall, R. P., Shevlin, M., & Sellwood, W. (2014). Associations between specific psychotic symptoms and specific childhood adversities are mediated by attachment styles: an analysis of the National Comorbidity Survey. *Psychiatry Research*, 217, 202-209.
- Startup, H., Freeman, D., & Garety, P. A. (2007). Persecutory delusions and catastrophic worry in psychosis: developing the understanding of delusion distress and persistence. *Behaviour Research and Therapy*, 45, 523-537.
- Tackett, J. L., Kushner, S. C., Herzhoff, K., Smack, A. J., & Reardon, K. W. (2014). Viewing relational aggression through multiple lenses: Temperament, personality, and personality pathology. *Development and Psychopathology*, 26, 863-877.
- Thewissen, V., Bentall, R. P., Oorschot, M., à Campo, J., van Lierop, T., van Os, J., & Myin-Germeys, I. (2011). Emotions, self-esteem, and paranoid episodes: An experience sampling study. *British Journal of Clinical Psychology*, 50, 178-195.
- Tone, E. B., & Davis, J. S. (2012). Paranoid thinking, suspicion, and risk for aggression: A neurodevelopmental perspective. *Development and Psychopathology*, 24, 1031-1046.
- Van Os, J. (2003). Is there a continuum of psychotic experiences in the general population. *Epidemiology and Psychiatric Sciences*, 12, 242-252.
- Verdoux, H., Gindre, C., Sorbara, F., Tournier, M., & Swendsen, J. D. (2003). Effects of cannabis and psychosis vulnerability in daily life: an experience sampling test study. *Psychological Medicine*, 33, 23-32.

Waller, H., Emsley, R., Freeman, D., Bebbington, P., Dunn, G., Fowler, D., ... & Garety, P. (2015).

Thinking Well: a randomised controlled feasibility study of a new CBT therapy targeting reasoning biases in people with distressing persecutory delusional beliefs. *Journal of Behavior Therapy and Experimental Psychiatry*, 48, 82-89.

Watson, D. (2001). Dissociations of the night: individual differences in sleep-related experiences and their relation to dissociation and schizotypy. *Journal of Abnormal Psychology*, 110, 526.

Westermann, S., Kesting, M. L., & Lincoln, T. M. (2012). Being deluded after being excluded? How emotion regulation deficits in paranoia-prone individuals affect state paranoia during experimentally induced social stress. *Behavior Therapy*, 43, 329-340.

Zohar, A. H., Zwir, I., Wang, J., Cloninger, C. R., & Anokhin, A. P. (2019). The development of temperament and character during adolescence: the processes and phases of change. *Development and Psychopathology*, 31, 601-617.

Table 1. The means, standard deviations (SD), frequencies, and ranges of the study variables.

	Mean	SD	Measurement range	Frequency (%)
Age (1997)	31.681	5.023	24–39	
Sex (female)				881 (56.73)
Parents' educational level (1980)				
Comprehensive school				495 (31.87)
High school or occupational school				639 (41.15)
Academic level				419 (26.98)
Parents' level of income (1980)	4.962	1.908	1–8	
Participants' educational level (2011)				
Comprehensive school				38 (2.45)
High school or occupational school				823 (52.99)
Academic level				692 (44.56)
Participants' level of income (2011)	7.458	3.049	1–13	
Cumulative risk score (2001)	0.149	0.187	0–1	
Experienced stress	2.333	0.982	1–5	
Anticipatory worry	2.430	0.519	1–5	
Social detachment	2.324	0.733	1–5	

Psychosocial risks and paranoid ideation

Sleep disturbances	0.416	0.633	0–2
Alcohol use	2.474	1.347	1–6
Cumulative risk score (2007)	0.167	0.191	0–1
Experienced stress	2.344	0.949	1–5
Anticipatory worry	2.429	0.514	1–5
Social detachment	2.423	0.693	1–5
Sleep disturbances	0.500	0.682	0–2
Alcohol use	2.386	1.388	1–6
Cumulative risk score (2011/2012)	0.173	0.197	0–1
Experienced stress	2.379	0.922	1–5
Anticipatory worry	2.436	0.517	1–5
Social detachment	2.439	0.700	1–5
Sleep disturbances	0.503	0.671	0–2
Alcohol use	2.442	1.494	1–6
Paranoid ideation			
2001	2.281	0.625	1–5
2007	2.119	0.634	1–5
2012	2.123	0.659	1–5

Psychosocial risks and paranoid ideation

Table 2. Results of the multilevel models. Unstandardized estimates (B) with p-values (within brackets) of psychosocial risk factors and age, when predicting standardized scores of paranoid ideation in adulthood.

	Fixed effects						Random effects		
	Intercept	Predictor	Age	Age ²	Predictor*Age	Predictor*Age ²	Variance of intercept	Variance of age	Residual variance
Experienced stress	1.195 (<.001)	0.139 (<.001)	-0.033 (<.001)	0.00042 (.040)	0.0038 (.546)	0.000054 (.815)	0.741 (<.05)	0.023 (<.05)	0.552 (<.05)
Anticipatory worry	0.888 (<.001)	0.268 (<.001)	-0.030 (<.001)	0.00039 (.051)	0.0079 (.176)	-0.00019 (.381)	0.717 (<.05)	0.026 (<.05)	0.543 (<.05)
Social detachment	0.951 (<.001)	0.149 (<.001)	0.038 (<.001)	0.00055 (.006)	0.011 (.049)	-0.00031 (.152)	0.751 (<.05)	0.024 (<.05)	0.552 (<.05)
Sleep disturbances	1.108 (<.001)	0.148 (<.001)	-0.034 (<.001)	0.00044 (.030)	-0.0016 (.784)	0.000064 (.763)	0.742 (<.05)	0.022 (<.05)	0.557 (<.05)
Alcohol use	1.091 (<.001)	0.111 (.007)	-0.029 (<.001)	0.00035 (.092)	-0.0057 (.322)	0.00013 (.535)	0.771 (<.05)	0.023 (<.05)	0.553 (<.05)
Cumulative risk	0.985 (<.001)	0.184 (<.001)	-0.033 (<.001)	0.00038 (.070)	0.0082 (.199)	-0.00022 (.330)	0.713 (<.05)	0.023 (<.05)	0.549 (<.05)

N=1534–1553

Adjusted for age, sex, and participants' and their parents' socioeconomic factors. Continuous scores of each psychosocial risk factor (experienced stress, anticipatory worry, social detachment, sleep disturbances, frequent alcohol use) were included as predictors separately.

Note: STATA provides the p-values for random effects only at the significance level of 0.05 (not the exact p-values).









Figure 1. The trajectories of paranoid ideation in adulthood separately for participants with low/average levels (black line) and high levels (gray line) of (a) experienced stress, (b) anticipatory worry, (c) social detachment, (d) sleep disturbances, (e) frequent alcohol use, and (f) cumulative risk score. Estimated means with 95% confidence intervals. *Note:* Adjusted for age, sex, and participants' and their parents' socioeconomic factors. For this figure, we used categorized scores of psychosocial risk factors.

Table 3. The goodness-of-fit indexes of the structural equation models, when including paranoid ideation and one psychosocial factor at a time in the model.

	CFI	RMSEA	BIC	χ^2 (df)	Comparison between the models	
					χ^2 (df)	p-value
Experienced stress and paranoid ideation						
Model 1	0.940	0.056	46416.226	173.611 (30)		
Model 2	0.948	0.055	46417.665	153.007 (27)	Model 2 vs. 1: 20.604	0.00017
Model 3	0.945	0.056	46423.150	158.492 (27)		
Model 4	0.952	0.056	46426.644	139.942 (24)	Model 4 vs. 2: 13.065	0.0045
Anticipatory worry and paranoid ideation						
Model 1	0.969	0.050	40826.945	147.655 (30)		
Model 2	0.982	0.040	40796.612	95.278 (27)	Model 2 vs. 1: 52.377	<0.00001
Model 3	0.975	0.048	40824.282	122.948 (27)		
Model 4	0.986	0.039	40803.097	79.719 (24)	Model 4 vs. 1: 15.559	0.0014
Social detachment and paranoid ideation						
Model 1	0.980	0.043	42877.257	114.345 (30)		
Model 2	0.984	0.040	42687.148	94.718 (27)	Model 2 vs. 1: 19.627	<0.00001
Model 3	0.982	0.043	42697.117	104.687 (27)		
Model 4	0.985	0.041	42894.198	87.198 (24)	Model 4 vs. 2: 7.52	0.057
Alcohol use and paranoid ideation						
Model 1	0.970	0.053	49794.020	160.057 (30)		
Model 2	0.969	0.056	49814.310	158.303 (27)		
Model 3	0.970	0.056	49813.646	157.639 (27)	Model 3 vs. 1: 2.618	0.454
Model 4	0.969	0.060	49834.093	156.042 (24)	Model 4 vs. 1: 4.015	0.260
Sleep disturbances and paranoid ideation						
Model 1	0.962	0.044	44136.565	120.582 (30)		
Model 2	0.966	0.044	44146.185	108.158 (27)		
Model 3	0.972	0.040	44131.653	93.626 (27)	Model 3 vs. 1: 26.956	<0.00001
Model 4	0.975	0.040	44143.860	83.789 (24)	Model 4 vs. 3: 9.837	0.020

N=1553

Model 1: A model including only stability coefficients and covariances between the psychosocial risk factor and paranoid ideation at each time point. Model 2: A model that included also the predictive coefficients from the psychosocial risk factor at each time point to paranoid ideation at the next time points. Model 3: A model that included the cross-lagged predictive coefficients in the opposite direction. Model 4: A model with the cross-lagged predictive coefficients in both directions between the psychosocial risk factor and paranoid ideation.

Note: In all the models, we used continuous scores of psychosocial risk factors and paranoid ideation.

Supplementary Material. The correlations of the item of experienced stress with other measures of psychological stress.

To investigate the validity of the item of experienced stress, we investigated whether it correlates with stress reactivity and stressful life events. *Stressful life events* included death of romantic partner, divorce/separation, death of a child, and miscarriage (0=not occurred; 1=occurred). We calculated a total score of the number of stressful life events within the on-going year. Frequent stressful life events correlated with higher experienced stress (Pearson $r=0.528$, $p<.001$). *Stress reactivity* was evaluated with the Negative Emotionality scale of the Emotionality, Activity, and Sociability Temperament Survey (EAS) (Buss & Plomin, 1984). The scale is known to measure stress reactivity (Strelau & Angleitner, 1991). High scores of stress reactivity were found to correlated with higher experienced stress (Pearson $r=0.421$, $p<.001$).


References:

- Buss, A. H., & Plomin, R. (1984). *Temperament: Early Developing Personality Traits*. Hillsdale, New Jersey: Lawrence Erlbaum Associates.
- Strelau, J., & Angleitner, A. (Eds.). (1991). *Explorations in Temperament: International Perspectives on Theory and Measurement*. Plenum Press, New York.


Supplementary Table 1. The results of the exploratory factor analysis with Oblimin rotation of the Vital Exhaustion Questionnaire. The factor loadings >0.5 are bolded. Factor III refers to sleep disturbances.

	Factor		
	I	II	III
Item 1: Do you feel tired?	0.377	0.666	0.321
Item 2: Do you have difficulties in falling asleep?	0.263	0.198	0.808
Item 3: Do you wake up at night repeatedly?	0.281	0.262	0.775
Item 4: Do you feel powerless?	0.655	0.426	0.306
Item 5: Do you feel that you have not been able to get many things done in the recent times?	0.563	0.479	0.063
Item 6: Do you feel that you cannot get daily problems solved as well as previously?	0.665	0.514	0.059
Item 7: Do you feel like being in the dead end?	0.802	0.301	0.132
Item 8: Have you experienced feelings of insignificance or unwillingness more often recently than previously?	0.582	0.648	0.092
Item 9: Have you experienced feelings of hopelessness recently?	0.707	0.445	0.142
Item 10: Do you need more time to solve problems than, for example, for a year ago?	0.518	0.458	0.023
Item 11: Do you think that very small things have irritated you more recently than previously?	0.357	0.630	0.091
Item 12: Do you feel that you would like to stop trying?	0.730	0.225	0.201
Item 13: Do you feel good?	-0.665	-0.425	-0.195
Item 14: Do you feel that your battery is going down?	0.665	0.580	0.215
Item 15: Do you feel that it would be better to be dead?	0.681	0.017	0.239
Item 16: Do you feel that you are unable to do anything?	0.738	0.188	0.198
Item 17: Do you feel melancholic sometimes?	0.226	0.585	0.121
Item 18: Do you feel that you would like to just cry?	0.582	0.240	0.179
Item 19: Do you ever wake up in the morning with the feeling that you are totally tired or exhausted?	0.188	0.619	0.229
Item 20: Do you have increasing difficulties to concentrate on one thing at a time for a long time period?	0.612	0.405	0.151
Item 21: Do you enjoy sexual activities as much as previously?	-0.376	-0.441	-0.017


(a)




(b)



(c)



(d)

Supplementary Figure 1. Illustrations of the structural equation models: (a) Model 1, (b) Model 2, (c) Model 3, (d) Model 4. Note: For clarity, control variables (age, sex and participants' and their parents' socioeconomic factors) and the covariances between variables at each time point were excluded from this figure.

Psychosocial risks and paranoid ideation

Supplementary Table 2. The Pearson correlations between the study variables.

	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	14.
1. Age	-											
2. Sex	0.027	-										
3. Parents' educational level	-0.262***	-0.051*	-									
4. Parents' level of income	-0.022	-0.012	0.495***	-								
5. Participants' educational level	-0.160***	0.053*	0.288***	0.240***	-							
6. Participants' level of income	0.039	-0.289***	0.148***	0.182***	0.304***	-						
7. Experienced stress	0.025	0.063*	0.012	0.004	0.007	-0.033	-					
8. Anticipatory worry	0.001	0.126***	-0.079**	-0.108***	-0.143***	-0.218***	0.387***	-				
9. Social detachment	0.096***	-0.343***	-0.046	-0.096***	-0.123***	-0.012	0.083**	0.165***	-			
10. Sleep disturbances	0.031	0.094***	-0.015	-0.017	-0.049	-0.078**	0.406***	0.306***	0.083**	-		
11. Alcohol use	-0.103***	-0.371***	0.102***	-0.037	-0.037	0.143***	0.037	-0.026	0.057*	0.050	-	
12. Paranoid ideation	-0.040	-0.037	-0.050	-0.097***	-0.116***	-0.150***	0.305***	0.398***	0.260***	0.256***	0.099***	-

* $p < .05$ ** $p < .01$ *** $p < .001$

Note: When calculating the correlations between the study variables, we used the scores of the first measurement year of each variable (i.e. parents' socioeconomic factors in 1980; psychosocial risk factors and paranoid ideation in 2001).