

## ARTICLE

Received 26 Jun 2012 | Accepted 14 Dec 2012 | Published 22 Jan 2013

DOI: 10.1038/ncomms2389

# Reduced plumage and flight ability of a new Jurassic paravian theropod from China

Pascal Godefroit<sup>1,2</sup>, Helena Demuynck<sup>3</sup>, Gareth Dyke<sup>4</sup>, Dongyu Hu<sup>5,6</sup>, François Escuillié<sup>7</sup> & Philippe Claeys<sup>3</sup>

Feathered theropods were diverse in the Early Cretaceous Jehol Group of western Liaoning Province, China. Recently, anatomically distinct feathered taxa have been discovered in the older Middle-Late Jurassic Tiaojishan Formation in the same region. Phylogenetic hypotheses including these specimens have challenged the pivotal position of *Archaeopteryx* in bird phylogeny. Here we report a basal troodontid from the Tiaojishan Formation that resembles *Anchiornis*, also from Jianchang County (regarded as sister-taxa). The feathers of *Eosinopteryx* are less extensive on the limbs and tail than *Anchiornis* and other deinonychosaurs. With reduced plumage and short uncurved pedal claws, *Eosinopteryx* would have been able to run unimpeded (with large foot remiges cursorial locomotion was likely problematic for *Anchiornis*). *Eosinopteryx* increases the known diversity of small-bodied dinosaurs in the Jurassic, shows that taxa with similar body plans could occupy different niches in the same ecosystem and suggests a more complex picture for the origin of flight.

<sup>1</sup> Department of Palaeontology, Royal Belgian Institute of Natural Sciences, rue Vautier 29, 1000 Bruxelles, Belgium. <sup>2</sup> Jilin University Geological Museum, Chaoyang Campus, 6 Ximinzhu Street, Changchun 130062, China. <sup>3</sup> Earth System Science, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium. <sup>4</sup> Ocean and Earth Science, National Oceanography Centre, University of Southampton Waterfront Campus, European Way, Southampton SO 14 3ZH, UK. <sup>5</sup> Paleontological Institute, Shenyang Normal University, 253 North Huanghe Street, Shenyang 110034, China. <sup>6</sup> Key Laboratory of Vegetation Ecology, Ministry of Education, Northeast Normal University, 5268 Renmin Street, Changchun 130024, China. <sup>7</sup> Eldonia, 9 avenue des Portes Occitanes, 3800 Gannat, France. Correspondence and requests for materials should be addressed to P.G. (email: pascal.godefroit@naturalsciences.be).

Recent discoveries of feathered theropod dinosaurs in the Middle-Late Jurassic Tiaojishan Formation of western Liaoning Province, north-eastern China, have considerably challenged our understanding of the evolution and biology of the Paraves (the most-inclusive clade containing *Passer domesticus* Linnaeus, 1758 but not *Oviraptor philoceratops* Osborn, 1924; see Supplementary information)<sup>1–3</sup>. Large pennaceous feathers were already developed on the tail, forelimbs and hindlimbs, and particularly on the metatarsus and pes, of the troodontid *Anchiornis*<sup>2</sup>. It has been postulated that large pennaceous feathers first evolved distally on the hindlimbs, as on the forelimbs and tail, in basal paravian theropods and only spread proximally at a subsequent stage in theropod evolution<sup>2</sup>. Moreover, the organization of the pedal feathers into a coherent planar surface suggests that they potentially had some aerodynamic function in *Anchiornis*<sup>2</sup>. And adding the new taxa *Xiaotingia* and *Anchiornis* from the Tiaojishan Formation in a phylogenetic analysis of paravian theropods challenges the pivotal position of the iconic *Archaeopteryx*, regarded from its discovery as the most primitive bird, shifting it to the Deinonychosauria (the most-inclusive clade containing *Dromaeosaurus albertensis* Matthew and Brown, 1922 but not *P. domesticus* Linnaeus, 1758; see Supplementary information)<sup>3</sup>.

Here we describe a new troodontid theropod from the Tiaojishan Formation of western Liaoning Province. Although it resembles *Anchiornis* from an osteological point of view, this new troodontid is characterized by a reduced plumage, in comparison with other Middle-Late Jurassic feathered deinonychosaurians: large pennaceous feathers are absent from its tail, metatarsus and pes. The plumage of paravian theropods was already diversified by the Late Jurassic, obviously adapted to different ecological niches.

## Results

### Systematic palaeontology


Class Theropoda Marsh, 1881  
Order Maniraptora Gauthier, 1986  
Suborder Troodontidae Gilmore, 1924  
*Eosinopteryx brevipenna* gen. et sp. nov.

**Etymology.** *Eo*, Greek for daybreak, dawn; *sino*, Latin for Chinese; *pteryx*, Greek for feather, wing; *brevi*, Latin for short; *penna*, Latin for feather.

**Holotype.** YFGP (Yizhou Fossil & Geology Park)-T5197, a complete articulated skeleton with associated integumentary structures (Fig. 1; see Methods for provenance and authenticity of the holotype specimen).

**Locality and horizon.** Yaolugou, Jianchang, western Liaoning, China; Middle-Late Jurassic Tiaojishan Formation<sup>1–3</sup>.


**Diagnosis.** A small paravian maniraptoran theropod that possesses the following osteological autapomorphies: a short snout, about 82% the length of the orbit; a lacrimal with a long posterior process participating in about half the length of the dorsal margin of the orbit and a vestigial anterior process; a short tail, composed of 20 caudal vertebrae, 2.7 times the length of the femur; chevrons reduced to small rod-like elements below the proximal 8th or 9th caudal; an ilium with a proportionally long, low and distally tapering postacetabular process (ratio 'length/height at mid-length' = 5), and pedal unguals shorter than corresponding penultimate phalanges. The plumage of this taxon is characterized by the absence of rectrices (versus other paravians with preserved plumage) and feathers on metatarsus and pes (versus other troodontids with preserved plumage on the hindlimb).



**Figure 1 | *Eosinopteryx brevipenna* YFGP-T5197. (a)** Photograph and **(b)** line drawing. cav, caudal vertebrae; cev, cervical vertebrae; dv, dorsal vertebrae; fu, furcula; ga, gastralia; lf, left femur; lh, left humerus; lil, left ilium; lis, left ischium; lm, left manus; lp, left pes; lr, left radius; ls, left scapula; lt, left tibia; lu, left ulna; ma, mandible; rcor, right coracoid; rf, right femur; rh, right humerus; ril, right ilium; ris, right ischium; rm, right manus; rp, right pes; rr, right radius; rs, right scapula; rt, right tibia; ru, right ulna; sk, skull.

**Description.** The holotype of *E. brevipenna* is small, just 30 cm long (Fig. 1). The skull has a subtriangular lateral profile produced by a shallow snout and expanded postorbital region (Fig. 2). Unlike *Anchiornis*<sup>2</sup>, the snout of *Eosinopteryx* is distinctly shorter (about 82% of orbital length) than its orbit. Both the rostral plate and the maxillary process of the premaxilla are particularly short. The posterior process of the lacrimal is elongated and straight, participating in about half the length of the dorsal margin of the orbit, whereas the anterior process is vestigial. The postorbital process of the jugal is broader than in *Anchiornis*<sup>2</sup>, whereas the triradiate postorbital is larger than in *Archaeopteryx*<sup>4</sup> and forms a complete postorbital bar with the jugal. There is no trace of a mandibular fenestra. The dentary has a groove that widens posteriorly and contains a row of foramina as in *Archaeopteryx*, *Anchiornis*, *Xiaotingia*, troodontids and some basal dromaeosaurids<sup>2,3</sup>. The teeth of *Eosinopteryx* are unserrated and sharp in contrast to the bulbous teeth of *Xiaotingia*<sup>3</sup>. As characteristic of troodontids<sup>5</sup>, the premaxillary and the dentary teeth in the symphyseal region are more closely packed than are the teeth in the posterior part of the dentary.





**Figure 2 | Selected skeletal elements and associated feathers of *Eosinopteryx brevipenna* YFGP-T5197.** (a) Photograph of skull and mandible in right lateral view. (b) Line drawing of skull and mandible in right lateral view. (c) Photograph of pelvis in right lateral view. (d) Line drawing of right ilium in lateral view. (e) Line drawing of right ischium in lateral view. (f) Photograph of right metatarsus and pes. (g) Line drawing of right metatarsus and pes. (h) Detail of secondary remiges. (i) Detail of plumulaceous feathers under the middle part of the tail. ac, acetabulum; lpm, left premaxilla; mt, metatarsal; obt, obturator process; postpr, posterodistal process; prac, preacetabular process; ra, right angular; rdt, right dentary; rect, right ectopterygoid; rfr, right frontal; rj, right jugal; rl, right lacrimal; rmx, right maxilla; rna, right nasal; rpa, right parietal; rpal, right palatine; rpm, right premaxilla; rpo, right postorbital; rq, right quadrate; rsq, right squamosal; rsa, right surangular; scl, scleral plate.

There are seven postaxial cervical vertebrae in YFGP-T5197. The cervical ribs are as long as their corresponding centra, contrasting with the shorter cervical ribs seen in *Troodon formosus*<sup>5</sup> and the much longer ones of *Archaeopteryx*<sup>6</sup>. The trunk is about 32% the length of the hindlimb, as in *Anchiornis*<sup>2</sup>

and *Mei*<sup>7</sup>. The middle and posterior dorsals are proportionally elongate as in *Anchiornis*, dromaeosaurids and *Archaeopteryx*<sup>8</sup>. As in the latter, *Anchiornis* and basal deinonychosaurs, the dorsal centra do not have distinct pneumatic foramina just shallow depressions on their lateral surfaces<sup>3</sup>. Numerous, very slender, abdominal ribs are preserved in YFGP-T5197, as in *Archaeopteryx*<sup>6</sup>, dromaeosaurids<sup>9</sup>, the basal pygostylian bird *Confuciusornis*<sup>10</sup> and many enantiornithines (for example, *Sinornis*). The tail of *Eosinopteryx* is particularly short, 2.7 times the length of the femur (contrasting with 3.2 in *Mei*<sup>7</sup> and 3.9 in the Eichstätt *Archaeopteryx*<sup>6</sup>), and only composed of 20 caudal vertebrae (contrasting with 21–23 in *Archaeopteryx*<sup>6</sup>, ca. 30 in *Anchiornis*<sup>1</sup> and 24–26 in *Microraptor*<sup>11</sup>). The anterior caudals are proportionally short, their transverse processes longer than corresponding centra and particularly narrow. Neural spines are developed only on the anteriormost third or fourth caudals. As in *Anchiornis* and *Archaeopteryx*<sup>2,3</sup>, the middle and posterior caudal vertebrae each bear a distinct groove on the lateral surface near the junction of the centrum and neural arch. The chevrons are reduced to small rod-like elements and are only present below the proximal eighth or ninth caudal centra, contrasting with the larger hook-like proximal chevrons of *Anchiornis*<sup>1</sup> and with the vertically oriented rectangular proximal chevrons of *Archaeopteryx*<sup>6</sup>.

The subrectangular coracoid of *Eosinopteryx* bears a distinct subglenoid fossa and has a wider proximal end than *Xiaotingia*<sup>3</sup> (Supplementary Fig. S1). The scapula (scapula/femur = 0.49) is proportionally much shorter than in *Anchiornis* (0.68). The forelimb of this small paravian (0.73 times hindlimb length) is slightly shorter than that of *Anchiornis* (0.8) (ref. 2) and *Archaeopteryx* (0.9–1.0) (ref. 6) but is much longer than *Mei* (0.5) (ref. 7). The humerus of *Eosinopteryx* is also shorter (0.8 times) than its femur; the humerus is slightly longer than the femur in *Anchiornis*, distinctly longer (1.1–1.2) (ref. 6) in *Archaeopteryx*, and is only half the femur length in *Mei*. As in *Anchiornis*, the radius and ulna of *Eosinopteryx* are straight with only a narrow gap between them. The manus is slightly longer than the femur (manus/femur = 1.2), contrasting with the short hand of *Mei* (0.8) (ref. 7) and with the proportionally more elongate hands of *Anchiornis* (1.55) (ref. 2) and *Archaeopteryx* (1.4–1.6) (ref. 6). Metacarpal II is about one third the length of metacarpal III; in contrast to *Xiaotingia*<sup>3</sup>, metacarpal IV is shorter and is much more slender than metacarpal III (Supplementary Fig. S2). The phalangeal portion of the manus is also proportionally shorter, relative to the metacarpus, than in *Xiaotingia*<sup>3</sup>. As in *Anchiornis*<sup>2</sup>, the long manual phalanx II-1 of *Eosinopteryx* is as robust as the radius; this element is much more gracile in *Sinornithoides*<sup>12</sup> and *Archaeopteryx*<sup>6</sup>. Contrasting with *Xiaotingia*<sup>3</sup>, phalanx III-2 is shorter than metacarpal III, whereas the manual unguals of *Eosinopteryx*, sheathed by long keratinous claws, are less curved than in *Archaeopteryx*<sup>6</sup>, *Anchiornis*<sup>1</sup> and *Xiaotingia*<sup>3</sup>.

The ilium of *Eosinopteryx* is characterized by a long, low and distally tapering postacetabular process, with a ratio 'length/height at mid-length' of 5 (Fig. 2). The ischium is particularly short, about 28% the femur length. With its distally displaced obturator process and its long and pointed posterodistal process, this element closely resembles that of *Anchiornis*<sup>1,2</sup> (Fig. 2). The tibiotarsus of *Eosinopteryx* ( $1.4 \times$  femur length) is proportionally shorter than *Anchiornis* (1.60), whereas pes (metatarsus + phalanges) proportions ( $1.3 \times$  femur length) are similar to those of *Mei*<sup>7</sup> and shorter than *Anchiornis* (1.56) (ref. 2). The proximal end of metatarsal III is transversely compressed, suggesting a subarctometatarsalian condition. Pedal digit I lies on the medioplantar side of metatarsal II (Fig. 2), as in *Archaeopteryx*<sup>6,13,14</sup>, but contrasts with the position of pedal


digit I in *Anchiornis*, which is medial to metatarsal II<sup>1</sup>. The phalanges of pedal toes II, III and IV decrease gradually in length proximodistally, as in *Archaeopteryx* and terrestrial cursorial birds<sup>15</sup>. The pedal unguals of *Eosinopteryx* are much shorter than those of *Archaeopteryx*<sup>6</sup>, *Anchiornis*<sup>1,2</sup>, troodontids<sup>5</sup> and dromaeosaurids<sup>9</sup>, even shorter than the corresponding penultimate phalanges.

As in *Anchiornis*<sup>2</sup>, the longest remiges of *Eosinopteryx* are at least 150% the length of the humerus (Fig. 2) and are symmetrical on the forelimbs. Long, densely packed and symmetrical pennaceous feathers are present along the posterior part of the thigh and crus, consistent in distribution with the tetrapterygian condition of several other basal paravian taxa<sup>2,16</sup>. However, careful preparation shows that feathers were absent from the metatarsus and pes of YFGP-T5197. Feathers covering other regions of the body (head, neck, thorax, back and tail) are plumulaceous and ‘rachis-like’ structures described on the plumulaceous feathers at the base of the tail and on the neck in *Mei*<sup>7</sup> are absent in *Eosinopteryx*. The feathers on the tail are short, <20 mm long; all are plumulaceous, composed of a bundle of filaments that are joined together proximally and remain nearly parallel as they pass distally (Fig. 2). Pennaceous rectrices are absent.

Discussion

Preliminary phylogenetic analysis places *E. brevipenna* sister to *Anchiornis huxleyi* at the base of Troodontidae (Fig. 3) (subject to change in subsequent analyses; see Supplementary Methods for details). Both taxa share a dorsally curved ilium shaft, a pubis moderately oriented posteriorly and a straight ulna that is not bowed away from the humerus. *Archaeopteryx* and *Wellnhoferia* form an unresolved polytomy at the base of Deinonychosauria in the strict consensus tree, supporting the hypothesis that *Archaeopteryx* is a deinonychosaurian theropod instead of a primitive bird<sup>3</sup> (Fig. 3). Note, however, that this phylogeny remains only weakly supported presumably due to the numerous homoplasies widely distributed across coelurosaurian phylogeny<sup>3,17</sup> (Supplementary Fig. S3).

The almost complete plumage of *Eosinopteryx* is less developed than that of other known Middle-Late Jurassic deinonychosaurs. The complete absence of rectrices on the tail cannot be explained by taphonomic factors (that is, more fragile plumulaceous feathers are well preserved along the whole length of the tail) and does not reflect a basal phylogenetic position for *Eosinopteryx* among paravians, because long rectrices are now known to occur in at least one basal member of each of the five major known lineages in this region of the theropod tree, including




**Figure 3 | Phylogeny of Paraves.** Time-calibrated strict consensus tree of the 10 most-parsimonious trees. Tree length = 1,386; consistency index (CI) excluding uninformative characters = 0.3; retention index (RI) = 0.7. We added four feather-related characters to the original matrix<sup>3</sup> but these do not influence the topology of the strict consensus tree. Character 375: remiges on the forelimb absent (0) or present (1); character 376: pennaceous feathers on the tibia absent (0) or present (1); character 377: feathers on metatarsus and pes absent (0) or present (1); character 378: rectrices absent (0) or present (1).

basal avialans, the basal deinonychosaurs *Archaeopteryx* and *Wellnhoferia*<sup>6</sup>, the basal troodontid *Anchiornis*<sup>18</sup>, the scansoriopterygid *Epidexipteryx*<sup>19</sup> and the basal dromaeosaurid *Microraptor*<sup>20</sup>. The secondary loss of rectrices is regarded here as an unambiguous autapomorphy for *E. brevipenna* (Fig. 4). Pennaceous rectrices attached to the tibia were also a relatively ubiquitous adaptation among maniraptorans, present in *Eosinopteryx*, *Anchiornis*<sup>2</sup>, *Xiaotingia*<sup>3</sup>, *Microraptor*<sup>15</sup>, *Pedopenna*<sup>21</sup>, *Archaeopteryx* and *Wellnhoferia*<sup>6</sup>; this character is regarded here as an unambiguous synapomorphy for Deinonychosauria (Fig. 4).

In this phylogeny, the presence of long feathers on the metatarsus and pes, observed in *Anchiornis*<sup>2</sup>, *Xiaotingia*<sup>3</sup> and *Microraptor*<sup>16</sup>, is regarded as an unambiguous synapomorphy for the clade Troodontidae + Dromaeosauridae, and their secondary loss an unambiguous autapomorphy for *Eosinopteryx* (Fig. 4). It cannot be totally excluded that the absence of pennaceous feathers on the metatarsus and pes of *Eosinopteryx* is a preservational artefact, because other integumental features (for example, scales or scutes) are not preserved at the same location. However, scales and scales on the lower leg have never been identified in any feathered dinosaur or Mesozoic bird described to date. Despite its small size, the holotype of *Eosinopteryx* had reached a late ontogenetic stage (subadult or adult): neurocentral sutures are closed on all exposed vertebrae (Supplementary Fig. S4) and the suture between the astragalus–calcaneum

complex and the tibia cannot be discerned (Supplementary Fig. S5). Ontogenetic stage, therefore, also cannot account for reduced plumage, although temporary disappearance of rectrices and pennaceous feathers on the metatarsus and pes might be explained by a seasonal moult.

It is parsimonious that the plumage differences of *Eosinopteryx* can be explained by its mode-of-life. With a shorter humerus and manus and a reduced plumage, *Eosinopteryx* had a much shorter wing span than other feathered paravians including *Archaeopteryx*, *Wellnhoferia* and *Anchiornis*. The straight and closely aligned ulna–radius of *Eosinopteryx* also means that pronation/supination of the manus with respect to the upper arm would have been limited; combined with the absence of a bony sternum and weakly developed proximal humerus, these attributes suggest that *Eosinopteryx* had little or no ability to oscillate the arms to produce a wing beat. The phalanges of the third toe of *Eosinopteryx* decrease proximodistally and its ungual phalanges are not very recurved and particularly short, suggesting a ground dwelling, ‘cursorial’ mode-of-life as in many living birds<sup>15</sup>. Reduced plumage on the tail and lower legs would have facilitated this ecology and allowed *Eosinopteryx* to run easily. This new specimen also shows that the plumage of paravian theropods was already diversified by the Late Jurassic and already had different functions: insulation, flight and visual display<sup>19,20</sup>. Although *Anchiornis* and *Eosinopteryx* are clearly similar from osteological



**Figure 4 | Mapping of feather characters onto a simplified strict consensus tree.** FO, fast optimization. SO, slow optimization. If not denoted, mapping is unambiguous.



and phylogenetic points-of-view, their plumage shows that they were adapted to different ecological niches and implies a much more complex picture in terms of the origin of flight than often supposed. Indeed, it is likely that the diversification of integumental structures facilitated the sympatry of numerous closely related paravian theropods in the Late Jurassic Tiaojishan Formation and later in the Early Cretaceous Jehol Group.

## Methods

**Origin and authenticity of the specimen.** As it is the case for most of the paravian specimens known from western Liaoning Province, the holotype and only known specimen of *E. brevipenna* was acquired by the Yizhou Fossil & Geology Park from a Chinese fossil dealer, according to whom the specimen was collected at the Yaolugou locality (Jianchang county, western Liaoning, China), not far from Daxishan village where *A. huxlei* specimen LPM (Liaoning Paleontological Museum)-B00169 was discovered<sup>2</sup>. The Tiaojishan formation is widely exposed in this area<sup>22</sup>, though it also has limited outcrops of the Early Cretaceous Yixian Formation<sup>23</sup>. The *E. brevipenna* specimen is preserved in a shale slab, the sedimentary structures of which closely resemble those of the specimens of *A. huxlei* and *Xiaotingia zhengi* also described from the Tiaojishan Formation; it is in fact extremely difficult to distinguish between Tiaojishan and Yixian shale slabs on the basis of macro-sedimentary features. It is, therefore, necessary to study in detail the micro-sedimentary structures and the pollen assemblages within the shale slabs to certify the age of the specimen, as it is also the case for the *A. huxlei* and *Xiaotingia zhengi* specimens described so far<sup>3</sup>.

The specimen was only partially prepared when it was sold by the Chinese dealer. Careful preparation by the Yizhou Fossil & Geology Park staff and by authors of this paper (FE and PG) did not reveal any trace of forgery, and the probability that the specimen is a composite is accordingly low.

**Age of the Tiaojishan Formation.** The Tiaojishan Formation crops out in the West Liaoning Province and is equivalent to the Lanqi Formation of North Hebei Province<sup>24</sup>. The reason for the abundance of excellently preserved fossils is the presence of interbedded tuffs originating from recurrent volcanic events<sup>25</sup>. In western Liaoning, the Tiaojishan Formation is 130–970 m thick<sup>25</sup> and is composed of different types of rocks, including basalts, andesites, rhyolites, tuffs, tuffaceous sandstones and conglomerates<sup>24,25</sup>. Because of the complicated stratigraphic pattern in the Tiaojishan Formation, accurate dating of the formation is particularly difficult<sup>24</sup>. Ages between  $165 \pm 1.2$  Ma and  $153 \pm 2$  Ma were obtained by SHRIMP U-Th-Pb analysis of samples from the Tiaojishan (= Lanqi) Formation<sup>26</sup>. Zhang *et al.*<sup>24</sup> concluded that the upper boundary of the Tiaojishan (= Lanqi) formation dates between 156 and 153 Ma. Chang *et al.*<sup>25</sup> obtained Ar-Ar ages of  $160.7 \pm 0.4$  Ma and  $158.7 \pm 0.6$  Ma for the basal Lanqi Formation in North Hebei Province. These results indicate that the age of the Tiaojishan Formation is between the Callovian (Middle Jurassic) and the Kimmeridgian (Late Jurassic)<sup>27</sup>.

**Nomenclatural acts.** This published work and the nomenclatural acts it contains have been registered in ZooBank, the proposed online registration system for the International Code of Zoological Nomenclature. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix 'http://zoobank.org/'. The LSIDs for this publication are: urn:lsid:zoobank.org:act:068F0446-C80D-4F90-837C-921AB291880F; urn:lsid:zoobank.org:pub:11EEC3B6-1BF7-4DDE-872F-C84DE2CBB20.

## References

- Xu, X. *et al.* A new feathered maniraptoran dinosaur fossil that fills a morphological gap in avian origin. *Chin. Sci. Bull.* **54**, 430–435 (2009).
- Hu, D. -Y. *et al.* A pre-Archaeopteryx troodontid theropod with long feathers on the metatarsus. *Nature* **461**, 640–643 (2009).
- Xu, X. *et al.* An Archaeopteryx-like theropod from China and the origin of Avialae. *Nature* **475**, 465–470 (2011).
- Elzanowski, A. & Wellnhofer, P. Cranial morphology of *Archaeopteryx*: evidence from the seventh skeleton. *J. Vert. Paleontol.* **16**, 81–94 (1996).
- Makovicky, P. J. & Norell, M. A. in *The Dinosauria*. (eds Weishampel, D. B., Dodson, P. & Osmolska, H.) 2nd edn, pp 184–195 (University of California, 2004).
- Wellnhofer, P. *Archaeopteryx—Der Urvogel von Solnhofen* (Friedrich Pfeil, 2008).
- Xu, X. & Norell, M. A. A new troodontid from China with avian-like sleeping posture. *Nature* **431**, 838–841 (2004).
- Xu, X. Deinonychosaurian fossils from the Jehol Group of western Liaoning and the coelurosaurian evolution. PhD thesis, Chinese Academy of Sciences (2002).
- Norell, M. A. & Makovicky, P. J. in *The Dinosauria*. (eds Weishampel, D. B., Dodson, P. & Osmolska, H.) 2nd edn, pp 196–209 (University of California, 2004).
- Chiappe, L. M. *et al.* Anatomy and systematics of the Confuciusornithidae (Theropoda: Aves) from the late Mesozoic of Northeastern China. *Bull. Am. Mus.* **242**, 1–89 (1999).
- Hwang, S. H. *et al.* New specimens of *Microraptor zhaoianus* (Theropoda: Dromaeosauridae) from Northeastern China. *Am. Mus. Novitates* **3381**, 1–44 (2002).
- Currie, P. J. & Dong, Z.-M. New information on Cretaceous troodontids (Dinosauria, Theropoda) from the People's Republic of China. *Can. J. Earth Sc.* **38**, 1753–1766 (2001).
- O'Connor, J., Chiappe, L. M. & Bell, A. in *Pre-modern birds: Avian divergences in the Mesozoic*. (eds Dyke, G. J. & Kaiser, G.) 39–114 (John Wiley & Sons, 2011).
- Mayr, G. *et al.* The tenth skeletal specimen of *Archaeopteryx*. *Zool. J. Linn. Soc.* **149**, 97–116 (2007).
- Hopson, J. A. in *New Perspectives on the Origin and Early Evolution of Birds*. (eds Gauthier, J. & Gall, L. F.) 211–235 (Peabody Museum of Natural History, 2001).
- Xu, X. *et al.* Four-winged dinosaurs from China. *Nature* **421**, 335–340 (2003).
- Holtz, T. R. J. in *New Perspectives on the Origin and Early Evolution of Birds*. (eds Gauthier, J. A. & Gall, L. F.) 99–124 (Peabody Museum of Natural History, 2001).
- Li, Q. *et al.* Plumage color patterns of an extinct dinosaur. *Science* **327**, 1369–1372 (2010).
- Zhang, F. -C. *et al.* A bizarre Jurassic maniraptoran from China with elongate ribbon-like feathers. *Nature* **455**, 1105–1108 (2008).
- Li, Q. *et al.* Reconstruction of *Microraptor* and the evolution of iridescent plumage. *Science* **335**, 1215–1219 (2012).
- Xu, X. & Zhang, F. -C. A new maniraptoran with long metatarsalian feathers from China. *Naturwissenschaften* **92**, 173–177 (2005).
- Duan, Y. *et al.* Preliminary report on Middle Jurassic strata and fossils from Linglongta area of Jianchang, Liaoning. *Global Geol.* **28**, 143–147 (2009).
- Xu, K. *et al.* *Jurassic system in the North of China (VII): the stratigraphic region of Northeast China* (Petroleum Industry, 2003).
- Zhang, H., Wang, M. & Liu, X. Constraints on the upper boundary age of the Tiaojishan Formation volcanic rocks in West Liaoning-North Hebei by LA-ICP-MS dating. *Chin. Sci. Bull.* **53**, 3574–3584 (2008).
- Chang, S. *et al.* High-precision  $^{40}\text{Ar}/^{39}\text{Ar}$  age constraints on the basal Lanqi Formation and its implications for the origin of angiosperm plants. *Earth Planet. Sci. Lett.* **279**, 212–221 (2009).
- Yang, W. & Li, S. Geochronology and geochemistry of the Mesozoic volcanic rocks in Western Liaoning: implications for lithospheric thinning in the North China Craton. *Lithos* **102**, 88–117 (2008).
- Gradstein, F. *et al.* *A Geologic Time Scale 2004* (Cambridge University, 2004).

## Acknowledgements

We thank B. Pohl and D. Leloup for making fossils available for study, Sun Ge and A. Cau for their help and comments on the manuscript, T. Hubin for the photographs and four referees for their helpful comments. This study was supported by a grant (BL/36/62) to P.G. from the SPP Politique Scientifique (Belgium). D.H. was supported by the Key Laboratory of Evolutionary Systematics of Vertebrates (CAS2011LESV011) and by the National Natural Science Foundation of China (41172026).

## Author contributions

P.G., F.E. and P.C. designed the project. H.D., P.G., G.D. and D.H. performed the research. P.G., H.D. and G.D. wrote the manuscript.

## Additional information

**Supplementary Information** accompanies this paper at <http://www.nature.com/naturecommunications>

**Competing financial interests:** The authors declare no competing financial interests.

**Reprints and permission** information is available online at <http://npg.nature.com/reprintsandpermissions/>

**How to cite this article:** Godefroit, P. *et al.* Reduced plumage and flight ability of a new Jurassic paravian theropod from China. *Nat. Commun.* 4:1394 doi: 10.1038/ncomms2389 (2013).