

Trends and variability in the ocean carbon sink

Review Article

Author(s):

Gruber, Nicolas ; Bakker, Dorothee C.E.; DeVries, Tim; Gregor, Luke ; Hauck, Judith; Landschützer, Peter; McKinley, Galen A.; Müller, Jens Daniel 

Publication date:

2023-02

Permanent link:

<https://doi.org/10.3929/ethz-b-000595538>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Nature Reviews Earth & Environment 4, <https://doi.org/10.1038/s43017-022-00381-x>

Funding acknowledgement:

821003 - Climate-Carbon Interactions in the Coming Century (EC)

821001 - Southern Ocean Carbon and Heat Impact on Climate (EC)

820989 - Our common future ocean – quantifying coupled cycles of carbon, oxygen, and nutrients for determining and achieving safe operating spaces with respect to tipping points (EC)

TRENDS AND VARIABILITY IN THE OCEAN CARBON SINK

Nicolas Gruber,^{1,†}, Dorothee C. E. Bakker², Tim DeVries³, Luke Gregor¹, Judith Hauck⁴, Peter Landschützer^{5,6}, Galen A. McKinley⁷, Jens Daniel Müller¹.

1) *Environmental Physics, Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, Zürich, Switzerland.*

2) *Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich, UK.*

3) *Department of Geography and Earth Research Institute, University of California, Santa Barbara, CA, USA.*

4) *Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar und Meeresforschung, Bremerhaven, Germany.*

5) *Max Planck Institute for Meteorology, Hamburg, Germany.*

6) *Flanders Marine Institute (VLIZ), Ostend, Belgium*

7) *Columbia University and Lamont Doherty Earth Observatory, Palisades, NY, USA.*

[†]e-mail: nicolas.gruber@env.ethz.ch

Abstract

The ocean has absorbed $25 \pm 2\%$ of the total anthropogenic CO₂ emissions from early 1960s to the late 2010s, with rates more than tripling over this period and with a mean uptake of -2.7 ± 0.2 Pg C yr⁻¹ for the period 1990 through 2019. This growth of the ocean sink matches expectations based on the increase in atmospheric CO₂, but research has shown that the sink is more variable than long assumed. In this Review, we discuss trends and variations in the ocean carbon sink. The sink stagnated during the 1990s with rates hovering around -2 Pg C yr⁻¹, but strengthened again after ~ 2000 , $-$ uptaking over -3 Pg C yr⁻¹ for 2010-2019. The most conspicuous changes in uptake occurred in the high latitudes, especially the Southern Ocean. These variations are caused by changes in weather and climate, but a volcanic eruption-induced reduction in the atmospheric CO₂ growth rate and the associated global cooling contributed as well. Understanding the variability of the ocean carbon sink is crucial for policy making and projecting its future evolution, especially in the context of the UN Framework Convention on Climate Change stocktaking activities and the deployment of carbon dioxide removal methods. This goal will require a global-level effort to sustain and expand the current observational networks and to better integrate these observations with models.

Table of contents summary

Carbon uptake by the ocean has increased alongside rising atmospheric CO₂ concentrations, but with substantial variability. This Review examines trends in ocean CO₂ uptake and the internal and external factors driving its variability, finding an ocean uptake rate of -2.7 ± 0.2 Pg C yr⁻¹ for the period 1990 through 2019.

Key points

- i) The long-term trend in the ocean carbon sink since the early 1960s was primarily driven by the increasing uptake of anthropogenic CO₂. Although the ocean is expected to have lost a few petagrams of natural CO₂ to the atmosphere in response to ocean warming, this loss cannot be quantified conclusively with observations.
- ii) The oceanic uptake of anthropogenic CO₂ scaled proportionally with the increase in atmospheric CO₂ between the early 1960s and late 2010s, as expected given the quasi-exponential growth of atmospheric CO₂ during this period.
- iii) The average ocean uptake rate of -2.7 ± 0.2 Pg C yr⁻¹ for the period 1990 through 2019 yields a proportionality β of 1.4 ± 0.1 Pg C per ppm of atmospheric CO₂, suggesting a trend in the uptake of -0.4 ± 0.1 Pg C yr⁻¹ decade⁻¹.
- iv) The ocean carbon sink varies by about $\pm 20\%$ around this trend, primarily caused by changes in the sources and sinks of natural CO₂, with a lesser role for variations in atmospheric CO₂ growth rates impacting the uptake of anthropogenic CO₂.
- v) The net oceanic uptake rate of CO₂ will likely decrease in the future owing to several converging trends: reduced emissions of CO₂ leading to reduced atmospheric CO₂ growth rates in response to climate policy; reduced storage capacity owing to continuing ocean acidification; and enhanced outgassing of natural CO₂ owing to ocean warming and changes in ocean circulation and biology.

[H1] Introduction

Throughout the Anthropocene, the ocean has been the largest and most persistent sink for the anthropogenic CO₂ emitted into the atmosphere by the burning of fossil fuels, cement production, and land use change¹⁻⁴. This importance was recognized already by the late 19th century^{5,6}, with the chemist Arrhenius⁷ estimating that about 83% of the emitted anthropogenic CO₂ would be taken up by the ocean. Therefore, he concluded that no noticeable global warming should be expected from the emissions of anthropogenic CO₂, since the uptake by the ocean uptake leave only a small fraction of the emissions accumulating in the atmosphere. Although his estimate of the long-term capacity of ocean uptake was accurate^{8,9}, Arrhenius was not aware that it takes thousands of years for the ocean to fully realize this capacity and not decades as he implicitly assumed⁶. Arrhenius' view was widely shared, so that the scientific community was oblivious to the growing threat from the CO₂ emissions that were increasing by several percent per year for most of the early 20th century¹⁰. Revelle and Suess¹¹ realized this mistake in 1957. Thereafter, the perspective of the scientific community on the issue of human-induced climate change shifted rapidly^{12,13}, especially after Keeling confirmed in 1960 that atmospheric CO₂ was increasing much more rapidly than implied by Arrhenius¹⁴.

Much of global ocean carbon cycle research since Revelle and Suess' discovery has focused on quantifying the fraction of the CO₂ emissions taken up by the ocean, and to understand the processes that limit this uptake, preventing the ocean from reaching the huge capacity of more than 80% that Arrhenius had identified. A crucial step to address this question was the conceptualization of the net exchange of CO₂ across the air-sea interface and the change in the stock of **dissolved inorganic carbon (DIC)** [G] to consist of two components: anthropogenic CO₂ and natural CO₂ (Box 1). The anthropogenic CO₂ component, previously often referred to as excess CO₂¹⁵, can be considered the perturbation component, as it is solely a consequence of the anthropogenic increase in atmospheric CO₂. The natural CO₂ component of the flux is associated with the pre-industrial pool of DIC in the ocean (order of 37,000 Pg C (1 Pg 10¹⁵g)¹⁶ and is involved in **air-sea gas exchange** [G], uptake and release by the biological pumps, interactions with and loss to the sediments, and input by rivers (Box 1).

Under the assumption of a steady-state ocean, which is supported by the relative constancy of climate and atmospheric CO₂ for centuries prior to the onset of the industrial revolution (~1800)¹⁷, the oceanic pool of natural CO₂ remains constant and the fluxes of natural CO₂ are globally balanced. This assumption permitted researchers already in the 1970s to use models and observations to determine the oceanic uptake of anthropogenic CO₂¹⁸⁻²⁰, with subsequent work refining the methods and improving the data base^{2,4,15,21,22}.

However, it has become increasingly clear since the 2000s that the natural carbon fluxes of the ocean are changing, and that the ocean sink for carbon is more variable²³⁻³². The natural CO₂ pool is in fact highly

mobile, responding to changes in physical forcing from the atmosphere through changes in winds and in the fluxes of heat and freshwater, inducing changes in ocean circulation, temperature, salinity, and ocean biology⁹. Moreover, the anthropogenic CO₂ pool is more changeable than previously thought, responding to changes in atmospheric CO₂ growth rate or changes in ocean circulation³³.

In this Review, we assess what is currently known about the ocean sink for CO₂, and how it has responded to the rising CO₂ emissions in recent decades, relying primarily on ocean observations. We describe the variability of this sink and its drivers, which are debated. Finally, we highlight the need for increased observational capacity to support long term decision making, especially for the use of oceanic carbon dioxide removal (negative emission) approaches.

[H1] OCEAN CARBON SINK TRENDS

Since the late 1950s, the ocean has taken up a net $25 \pm 2\%$ of the total anthropogenic CO₂ emissions¹. This fractional uptake has remained relatively constant through time, meaning that the ocean sink tripled over these six decades, increasing from about $-0.9 \text{ Pg C yr}^{-1}$ in the early 1960s to more than -3 Pg C yr^{-1} in 2020^{1,25} (note that the geophysical convention of fluxes being are considered positive here, so that an uptake of CO₂ is negative). This increasing ocean carbon sink is an ecosystem service that amounts to about 2 trillion Euros worth of emission reductions per year if valued at a typical marginal abatement cost compatible with a 1.5°C target of 200 Euro per ton of CO₂³⁴. Together with the large ocean uptake of the excess heat generated from rising atmospheric CO₂³⁵, the ocean has moderated the climate change experienced so far^{36,37}. This section reviews how this ocean sink has been determined and what drives this long-term trend.

[H2] Response to rising atmospheric CO₂

The primary driver causing a long-term (> decades) change in the ocean's inventory of DIC is the rise in atmospheric CO₂, driving a flux of anthropogenic CO₂ across the air-sea interface and then from the surface ocean to depth (Box 1). The rate limiting step for the uptake of anthropogenic CO₂ by the ocean is the transport from the surface to deeper layers³⁸, as it takes decades to centuries for waters to circulate from the surface to the deeper ocean and back again^{39,40}. In contrast, CO₂ gas exchange across the air-sea interface is comparably fast (e-folding time scale of less than a year^{9,41}), so that the CO₂ concentration of the surface ocean follows the atmospheric perturbation relatively closely^{42–44}, with the magnitude of increase determined by the surface **ocean's buffer (or Revelle) factor**^{9,45,46} **[G]**. The two processes air-sea exchange and the surface-to-deep transport of CO₂ respond approximately linearly to changes in atmospheric CO₂. However, there is some the moderate non-linearity owing to the ocean's decreasing buffering capacity due to **ocean**

acidification [G] (a decrease of about 10% since preindustrial times⁴⁷) that needs to be taken into account as well^{48,49}.

When a (near)linear system like the ocean uptake of anthropogenic CO₂ (C_{ant}) is forced exponentially with a fixed growth rate (as is the case for atmospheric CO₂ since ~1970) (Fig 1a), all components of the system will increase proportionally after an initial adjustment (which is about a decade⁵⁰). This proportionality implies a linear scaling between the forcing (atmospheric CO₂) and the response (ocean accumulation of anthropogenic CO₂), which is confirmed by results from observations^{2,4}, ocean inverse models³ [G] and forward simulations [G] with ocean biogeochemical models^{25,51} [G] (Fig 1a). The slope of this relationship (the line in Fig 1a) represents the carbon concentration feedback of the ocean^{52,53} and is described as the sensitivity, where $\beta = \partial C_{\text{ant}} / \partial \Delta \text{CO}_2^{\text{a}}$, with the exact value dependent on the forcing history and especially past atmospheric growth rate. An emergent property of this relationship is that during periods of exponential growth in atmospheric CO₂, it directly determines the global oceanic uptake flux of anthropogenic CO₂ (F_{ant}(t)) from the growth rate of atmospheric CO₂: $dC_{\text{ant}}/dt = -F_{\text{ant}}(t) = \beta \cdot d\Delta \text{CO}_2^{\text{a}}/dt$, where the negative sign in front of F_{ant} reflects the convention of ocean uptake being negative.

This simple scaling relationship does not apply once the atmospheric CO₂ growth begins to deviate substantially from an exponential, as is expected if emissions start to stabilize and decrease in response to global efforts to curb climate change⁵⁴. In such cases, more complete theories building on deconvolution concepts such as pulse response functions^{3,55} or transit time distributions (TTD)^{56–58} are much better suited to describe the oceanic uptake of anthropogenic CO₂⁵⁹. Nevertheless, the high CO₂ concentration in the atmosphere would still be the main driving force for the many centuries it takes to equilibrate the entire ocean with the atmospheric perturbation.⁶⁰.

[H2] Cumulative oceanic uptake

The tight relationship between the ocean uptake for anthropogenic CO₂ and the growth in atmospheric CO₂ was recognized by the 1970s^{18,61,62}. However, until the mid-1980s, high-quality measurements of oceanic DIC were extremely scarce⁶³, making it impossible to constrain this relationship with observations. As the number of reliable DIC measurements increased in the late 1970s methods to identify the anthropogenic CO₂ signal within the substantial background variability of DIC emerged^{19,20}. Since the data were typically available only from a one-time survey, back-calculation approaches were used that implicitly assume a steady-state ocean. In such approaches, the DIC concentration in a water parcel in the ocean's interior is traced back to its origin at the surface, correcting along the way for the biological changes that incurred along this journey from the surface to depth^{15,21}. Refinement of the initial approaches led to the ΔC^* method^{64,65}, which is the most widely used back-calculation method to identify the total amount of anthropogenic CO₂ that has accumulated

in the ocean since preindustrial times²¹. A crucial enabling development for the identification of the relatively small anthropogenic CO₂ signal (see also Box 1) was the introduction of common measurement methodologies⁶⁶ and certified reference materials^{67,68} that permitted the collation of DIC measurements taken years apart and measured by different laboratories around the world into an internally coherent data set⁶⁹.

The application of the ΔC^* approach to the data collected by the Joint Global Ocean Flux Study (JGOFS)/World Ocean Circulation Experiment (WOCE) programs in the mid-1980s to mid-1990s led to the first global data-based estimate of the accumulation of anthropogenic CO₂². This approach yielded a total anthropogenic CO₂ inventory for the nominal year 1994 of 118 ± 19 Pg C (Fig 1a), i.e., reflecting the time integrated ocean uptake since ~1800. The maps in Fig 1b show the well-established spatial variations in the vertically integrated amount of anthropogenic CO₂^{38,70–72} (Fig. 1b). Strong accumulation in the North Atlantic contrasts with regions of relatively low accumulation such as the Tropical Pacific and the polar Southern Ocean. One of the most conspicuous features of the spatial variation is the band of high accumulation north of the Southern Ocean between about 30°S to 40°S. These basin-scale differences are a direct consequence of the regional effectiveness with which anthropogenic CO₂ is transported from the surface downward into the ocean's interior^{70,72–74}. The Ocean Inversion Project used such knowledge about the surface to depth transport in the form of impulse response functions, to estimate how much uptake of anthropogenic CO₂ is required in order to match the reconstructed distribution of anthropogenic CO₂ in 1994. This estimate yielded an uptake flux of -2.2 ± 0.25 Pg C yr⁻¹ for a nominal year 1995⁷².

This inventory also provided the first observation-based estimate of β of 1.47 ± 0.24 Pg C / ppm CO₂, representing the time period 1800–1994 (Supplementary Table 1). These results confirmed many prior estimates that so far had relied on models^{18,38,71} or indirect constraints such as the changes in atmospheric oxygen⁷⁵ or budgets of the stable isotope of carbon (¹³C)^{76–78}.

[H2] Decadal trends in uptake

The linear β -scaling can be used to provide a first estimate of the further evolution of the oceanic sink. Given the observed trend in the atmospheric CO₂ growth rate of 0.3 ppm yr⁻¹ decade⁻¹ between 1994 and 2007 and the inferred sensitivity β of 1.47 ± 0.24 Pg C / ppm CO₂, one would expect the steady-state ocean sink for anthropogenic CO₂ to increase (become more negative) by about -0.4 Pg C decade⁻¹ over this period, yielding an uptake in 2007 of the order of -2.6 Pg C yr⁻¹. Forward and inverse models^{3,25,70,79} have been used to assess this trend prediction (Fig 1a), but the ultimate evidence has to come from direct documentation of the increase in the ocean's DIC pool.

Direct documentation of decadal trends in anthropogenic CO₂ uptake is not straightforward, as shorter-term variations in the natural carbon pool tend to mask the slower but steadier increase in anthropogenic CO₂. This problem can be overcome for regularly sampled timeseries^{43,80,81}, but only a few sites have sufficient

observations to distinguish the anthropogenic trend from the natural variability. In most cases, the sampling rate is once per decade, as is the case for the GO-SHIP Global Repeat Hydrography Program⁸², for example. These data suffer acutely from an overprint of short-term variability in the natural carbon cycle, typically leading to a very noisy pattern of change that is difficult to interpret⁸³.

The introduction of the extended multiple linear regression (eMLR) approach⁸⁴ enabled the change in anthropogenic CO₂ to be mostly isolated^{85,86}. This method is the most widely used approach for detecting and quantifying changes in the anthropogenic CO₂ in the interior ocean based on repeat hydrography cruises^{83,87–89}. Compared to the ΔC^* approach, the eMLR approach captures both the steady-state and the non-steady-state accumulation of anthropogenic CO₂, although with limited accuracy when reconstructing the non-steady-state component⁹⁰.

A modified version of the eMLR method (eMLR(C*) method⁹⁰) was used to estimate the change in anthropogenic CO₂, ΔC_{ant} , globally⁴, using DIC and other biogeochemical data from the JGOFS/WOCE survey for the 1990s and comparing them with the measurements from the 2000s obtained during the 1st round of the GO-SHIP Repeat Hydrography Program⁸² (Fig. 1c). Global ocean carbon storage was estimated⁴ to increase by 34 ± 4 Pg C between 1994 and 2007, bringing the total inventory for anthropogenic CO₂ for 2007 to 154 ± 19 Pg C (Fig 1a). This increase in storage corresponds to a mean ocean uptake flux of anthropogenic CO₂ of -2.6 ± 0.3 Pg C yr⁻¹ over the 1994 to 2007 period, corroborating the simple scaling prediction. It also suggests a sensitivity β of 1.39 ± 0.16 Pg C / ppm CO₂, which is statistically indistinguishable from that estimated from the anthropogenic CO₂ inventory in 1994 (1.47 ± 0.24 Pg C / ppm CO₂, Supplementary Table 1). This lack of a difference provides strong support for the steady-state assumption.

Given this steady-state, the ocean interior estimate for 1994 to 2007 can be scaled to each decade over the past 30 years using β , yielding -2.1 Pg C yr⁻¹ for 1990 to 1999, -2.6 Pg C yr⁻¹ for the subsequent decade, and -3.3 Pg C yr⁻¹ for 2010 through 2019 (Table 1). Models suggest a smaller sensitivity β , lower mean uptake and smaller decadal trends (Table 1, Supplementary Table 1). However, many of the differences are not statistically significant, confirming that the ocean acts as a strong and increasing sink for anthropogenic CO₂. Overall, the steady-state assumption is useful determining the multidecadal oceanic uptake of anthropogenic CO₂. However, this assumption does not hold as well when analyzing shorter-term variations or spatial variations.

[H2] Non-steady-state uptake

A more detailed analysis of the spatial variations in the ocean interior accumulation of anthropogenic CO₂ highlights the limits of the steady-state assumption (Fig 1b,c). To first order, the increase in anthropogenic CO₂ is proportional to how much anthropogenic CO₂ was present at the beginning^{4,42,91}. The proportionality can be estimated using the β approach, yielding a value of 0.28 ± 0.02 for the inventory in 1994 and the change in inventory⁴ between 1994 and 2007 (similar approaches using a transit-time distribution (TTD) approach⁵⁷ yield comparable results). Thus, differences in the scaled spatial distribution of C_{ant} (1994) (Fig 1b) and ΔC_{ant} (2007-1994) (Fig 1c) suggest a non-steady-state contribution. Although the uncertainties in the two reconstructions are substantial, they suggest a shift in the accumulation of anthropogenic CO₂ from the North Atlantic to the South Atlantic, potentially related to decadal shifts in the overturning circulation⁹². This pattern confirms the presence of substantial decadal variability in the ocean carbon cycle identified previously along basin-wide hydrographic sections that had been occupied multiple times^{83,89}. However, the decadal nature of the repeat hydrography program limits the ability to constrain the year-to-year variability of the ocean carbon sink via the changes in the carbon storage.

[H1] OCEAN CARBON SINK VARIABILITY

Analyses of the sea-to-air fluxes of CO₂ are better suited to address this challenge, as they can be used to analyze changes at much higher temporal resolutions. In addition, they also assess the potential contribution of the non-steady-state component of the natural CO₂ fluxes, which we expect to drive most of the ocean flux variability. The ability to constrain these sea-to-air CO₂ fluxes with observations has made large strides in the last decade for at least three reasons. First was the expansion of the surface ocean **partial pressure of CO₂ [G]** (pCO₂) measurement programs that began in the 1960s⁹³, but picked up momentum in the late 1980s and 1990s^{94–96}. Second was the collation of the available surface ocean pCO₂ measurements by the Surface Ocean CO₂ Atlas (SOCAT) effort into a quality controlled and openly accessible data product^{97–99}. More than 30 million observations are in the [SOCAT](#) V2022 release, but these observations cover only a small fraction of the ocean surface. For example, at any given month in the decade of the 2010s, only 3% of all 1°x 1° grid points of the surface ocean have at least one observation. Therefore, the third notable advance was the development of approaches to inter- and extrapolate these surface ocean pCO₂ observations to obtain space-time continuous estimates of the sea-to-air CO₂ fluxes^{100–102}. Six of these reconstructions have been harmonized into a globally consistent product¹⁰³, called SeaFlux.

The long-term mean fluxes of this ensemble are characterized by strong outgassing of CO₂ in equatorial regions, most prominently in the equatorial Pacific (Fig 2). There is a strong net uptake of CO₂ at latitudes around 45° in both hemispheres. The overall pattern of the sources and sinks of CO₂ is primarily determined by the exchange of natural CO₂, responding to heating and cooling, vertical transport and mixing, and

variations in biological productivity⁹. The uptake of anthropogenic CO₂ modifies these fluxes, most strongly in the areas of large uptake of anthropogenic CO₂ such as the tropics and the high latitudes¹⁰⁴.

There is an almost doubling of the global net sea-to-air flux of CO₂ estimated by the SeaFlux ensemble from -1.5 Pg C yr⁻¹ in 1990 to -2.7 Pg C yr⁻¹ in 2018 (Fig. 3a). A loss of natural CO₂ of 0.65 ± 0.30 Pg C yr⁻¹¹⁰⁵ needs to be subtracted from the pCO₂ based estimates to compare these net fluxes with the global ocean uptake estimates here (Table 1) and also those reported by the Global Carbon Project^{1,51}. This loss is part of a natural steady-state of the global carbon cycle, and results from the difference between the carbon input by rivers and the carbon burial in marine sediments^{105–108} (see also Box 1). Based on this information, the combined fluxes of steady-state anthropogenic CO₂ and non-steady-state natural and anthropogenic CO₂ of -2.1±0.3 Pg C yr⁻¹ in the 1990s, -2.3±0.2 Pg C yr⁻¹ in the 2000s, and -3.1±0.2 Pg C yr⁻¹ in the 2010s (Table 1) (this flux is referred to as the ocean sink S_{OCEAN} in the Global Carbon Budget^{1,51}).

[H2] Interannual to decadal variability

The overall trend from the 1990s to the present of about -0.4 Pg C yr⁻¹ decade⁻¹ is close to that estimated from the steady-state model for anthropogenic CO₂ (orange dashed line in Fig 3a). The simulated fluxes from a model run with constant circulation and constant biology (CESM-ETHZ)²⁵ show the same overall trend (red dashed line), although with some more variations, largely reflecting changes in the growth rate of atmospheric CO₂³³. Thus, when analyzed over the last three decades, the surface ocean fluxes suggest an ocean carbon sink that has increased at a rate commensurate with the steady-state prediction.

However, on interannual to decadal timescales, the ocean carbon sink diagnosed from the surface pCO₂ observations deviates substantially from the steady-state prediction (Fig 3a). The strongest deviations occur on decadal timescales, with a weakening sink during the 1990s (a decadal trend of +0.3 Pg C yr⁻¹ decade⁻¹ (1990-2001)), followed by a strong reinvigoration with a decadal trend of -0.7 Pg C yr⁻¹ decade⁻¹ (2002-2018), nearly twice the rate from the steady state model. Integrated over the three decades, the ensemble mean uptake is 6±5 Pg C (11%) smaller than expected from the steady-state prediction, that is, this difference suggests a non-steady-state or climate variability induced loss of natural and anthropogenic CO₂. The estimates from the individual pCO₂-based reconstructions (shown in grey in Fig 3a) vary substantially around the SeaFlux ensemble mean, but all agree on the strong decadal modulation of the ocean carbon sink around the long-term trend.

All ocean basins contribute to the decadal variations of the ocean carbon sink, but the largest changes occur in the Pacific Ocean and the Southern Ocean, which is defined here as the ocean south of 44°S^{24,32,109,110} (Fig 3b). Both basins experienced a strong minimum in uptake around 2002 and a recovery thereafter, while the Atlantic basin north of 44°S had a more gradual increase through time. The Pacific is the only basin that

exhibits a clear interannual variability signal on top of the trend and the decadal changes. In contrast, the carbon sink of the Indian Ocean north of 44°S remained relatively constant.

Given that all these estimates rely on the same sparsely sampled ocean pCO₂ data, though, the potential for systematic errors that transcends all interpolation methods cannot be excluded¹¹¹. The reconstructions in the Southern Hemisphere are particularly concerning, as model based analyses¹¹¹ suggest that the severe undersampling could lead to an overestimating of the diagnosed decadal variability. In addition, the cool surface ocean skin effect¹¹² and uncertainties associated with the functional dependence of the gas transfer velocity on wind and other environmental factors¹¹³ add to the overall uncertainty of the flux products. Regardless, these variations—especially the weakening and strengthening periods—are seen in other, independent estimates, including from forward models²⁵ and inverse models¹¹⁴, although with generally smaller magnitudes²³.

[H2] Patterns of variability

More details about the spatio-temporal nature of the sea-to-air flux variations can be gleaned from the pCO₂ observation-based constraints that emerged in the 2010s. A Hovmoeller plot of the zonal integrals of the anomalous air-sea fluxes (Fig 4a) shows that the largest variations occur in the regions of strong absolute fluxes, that is, either in regions of strong uptake (temperate to high latitudes) or in the regions of strong outgassing of CO₂ (tropics). On top of the year-to-year variations, which are most prominent in the tropical latitudes, the long-term changes and the superimposed decadal variability clearly emerge from the data. They indicate that the extratropics (between 30° and 60° latitude) were the most important latitudes contributing to the rapid growth in the ocean carbon sink in the 2000s and 2010s, with the southern hemisphere dominating due to its larger ocean surface area.

These fluxes are the sum of the anomalies of the anthropogenic and natural CO₂ flux components. To separate them, the Ocean Inversion Project-based steady-state estimates for the uptake of anthropogenic CO₂ from the ⁷² for the year 1995 were scaled to the entire period using the β -based scaling approach used above. The zonal integral of the anomalies of this steady-state component of the anthropogenic CO₂ flux indicates that the regions of highest uptake in the Southern Ocean, the tropics and the mid latitudes of the northern hemisphere imprint large trends on the fluxes in these regions. In contrast, other regions have only a small trend in absolute terms (Fig 4b).

By removing this anthropogenic steady-state trend from the anomalous flux, the remaining anomalies reveal a clearer picture of the non-steady-state components driven by climate variability (Fig 4c). The strong interannual nature of the variations in the tropical belt emerges even more prominently. These anomalies are

correlated to the El Niño Southern Oscillation (ENSO) [G], as indicated by the negative correlation of the zonal anomalies in the tropical belt with the multivariate ENSO index¹¹⁵ ($R = -0.79$, $p < 0.05$). However, the anomalous uptake during El Niños was strong in the 1990s and weakened substantially after the turn of the millennium. At the same time, the anomalous outgassing during La Niña conditions strengthened over time. These ENSO related trends yield a distinct decadal signal in the tropics as well, characterized by an anomalous uptake during the 1990s, neutral conditions during the first decade of the new millennium, and anomalous outgassing in the 2010s.

The decadal nature of the Southern Ocean sink variability is also more discernible in these non-steady-state fluxes (Fig 4c). Over the course of the 1990s, there was a rapid change from an anomalous uptake to an anomalous outgassing peaking around 2002. This was followed by a prolonged period of anomalous outgassing until about 2008 and a recovery to normal conditions around 2010. Thus, the strong trend in the Southern Ocean toward increasing uptake in the last two decades is largely the result of the strong trend imparted by the steady-state uptake of anthropogenic CO₂, reflecting the major role of the this region in taking up anthropogenic CO₂ from the atmosphere (Fig 4b)^{72,116}.

The trend from an anomalous sink to an anomalous source during the 1990s followed by a strengthening period after 2000 is also evident across most latitudes of the northern hemisphere (Fig 4c). This co-occurrence suggests that apart from the tropics, the decadal mode of sea-to-air CO₂ flux variability has a global component, even after accounting for the steady-state trend in the uptake flux of anthropogenic CO₂.

In summary, the pCO₂ observation-based constraints on the sea-to-air CO₂ fluxes that have emerged in the last decade have reshaped our understanding of the variability of the ocean carbon sink (Figure 5). In particular, the surface flux products have suggested the presence of an important decadal mode of variability in the extra-tropics, and particularly in the Southern Ocean (Figure 5). This observation contrasts with the results of ocean biogeochemical models, whose variability tend to be, on average, smaller, and also which tend to have most of the variability focused in the tropics^{25,30,117}. Nevertheless, the models also simulate decadal variability in the extratropics^{23,25,28,29,118}), adding further evidence that the decadal variability diagnosed from the observations is a robust feature.

[H2] Mechanisms of variability

Variations in the ocean carbon sink can either be caused by processes that are internal to the climate system or can be externally forced. Internal forcing is associated with variations in weather and climate^{24,27,28,32,109,114,119} including changes associated with anthropogenic climate change¹²⁰. Externally forced variations are caused by changes outside the climate system, such as those induced by the volcanic eruption

of Mount Pinatubo in 1991³³. Such an eruption can impact the ocean carbon sink through changes in both Earth surface temperature and atmospheric CO₂ growth rate.

Interannual variations in the ocean carbon sink are driven by internal processes, as they are associated with the ENSO-related year-to-year variations in the sea-to-air fluxes in the tropical Pacific^{30,121–123}. During El Niño conditions, reduced upwelling and thermocline deepening in the Eastern Tropical Pacific strongly decrease the vertical supply of DIC to the surface. This process causes a collapse of the high pCO₂ levels that drive CO₂ out of the ocean, even though sea-surface temperatures are warmer than normal. Reduced windspeeds during El Niño conditions tend to further reduce the outgassing and thus enhance the effect of the reduced supersaturation¹²³. The resulting sea-to-air flux anomalies are sizable and impact the regional atmospheric CO₂ concentration¹²⁴. The flux variations are most likely almost entirely driven by changes in the natural CO₂, in particular its non-steady-state component (Fig 4c).

Mechanisms driving the decadal variations in the ocean carbon sink are less understood. One argument is that at least part of the variations are externally forced³³, as the eruption of Mt Pinatubo in 1991 caused both a reduced growth rate of atmospheric CO₂ during much of the 1990s^{125–127} and a global cooling trend in the surface temperature. The low growth rate reduces the ocean carbon sink directly by modifying the air-sea pCO₂ gradient. This effect would be enhanced by the upper ocean cooling and the associated enhanced ocean mixing caused by the global cooling^{128,129}. According to this argument, these two processes would have reduced the oceanic uptake during the 1990s, while the resumption of higher atmospheric CO₂ growth rates thereafter would have caused the ocean uptake to rebound³³.

An alternative line of arguments is that these decadal changes are the result of processes that are internal to the climate system. For example, a poleward contraction and intensification of the westerly wind belt around Antarctica might have caused the weakening trend of the Southern Ocean carbon sink during the 1990s²⁸, driven primarily by a trend toward a positive phase of the **Southern Annular Mode [G]** (SAM)¹³⁰. The stronger winds led to more upwelling of CO₂ and nutrient rich deep water, increasing CO₂ outgassing (albeit partly balanced by stronger biological production)^{28,118,131,132}. Then, a shift to a zonally more undulating windfield coupled with changes in sea-surface temperature caused the reinvigoration of the Southern Ocean carbon sink thereafter³². At least a part of these wind changes, and especially those of the 1990s, have been attributed to anthropogenic warming and ozone loss forcing the positive trend in the SAM¹³³. Simulations suggest that the majority of the response of the CO₂ fluxes is driven by changes in the natural CO₂ component, with the fluxes of anthropogenic CO₂ modulating the response, often in opposite directions, thus moderating the effect^{24,28,114,132}.

In contrast to the Southern Ocean, the potential mechanisms causing the reconstructed increases in the carbon sink in the northern hemisphere after 2000 are not well investigated. They most likely mechanisms involve changes in winds, changes in temperature affecting the solubility, changes in buoyancy forcing affecting

winter mixed layers¹³⁴, and large-scale gyre changes²⁷. The latter are potentially associated with changes in the northern annular mode (NAM) or associated northern hemisphere modes of variability¹⁰⁹.

The relative roles of internal versus external forcing driving the reconstructed decadal variations still need to be firmly established. Simulations with a changing atmospheric CO₂ growth rate, but no changes in climate, suggest that the effect is visible, albeit much smaller than the observed changes (the dashed red versus orange line in Fig 3a). The effect of the cooling and warming pattern associated with Mt. Pinatubo is more difficult to quantify independently, but simulations with comprehensive **ocean biogeochemical models** [G]^{128,135} suggest an effect ≤ 0.2 Pg C yr⁻¹ during peak cooling, and rapidly decreasing thereafter. However, the ocean carbon sinks changing globally relatively synchronously supports that there was an external forcing mechanism (Fig 4). Overall, external forcing (such as by volcanos) and internal changes (as by weather and climate variability) are not mutually exclusive processes, and both likely play a role in driving ocean carbon sink variability.

[H2] Merging observational constraints

Bringing together the ocean interior constraints on the evolution of the ocean sink with those provided by the surface ocean measurements can help to better understand the mechanisms driving trends and variability (Table 1). The estimates of the ocean interior accumulation of anthropogenic CO₂ suggest an ocean that globally has operated near steady-state. The extrapolation with β -scaling suggests a cumulative uptake of about 83 Pg C between 1990 and 2018. The reconstructions of the surface fluxes, which include both natural and anthropogenic CO₂ components, suggest 6 ± 5 Pg C less uptake over the same period (Table 1, Fig 3a). This reduction is mostly attributed to a non-steady-state loss of natural CO₂, as the simulation with the observed variations in atmospheric CO₂ suggested a small change in the total uptake of anthropogenic CO₂ (red versus orange dashed lines in Fig 3). This loss needs to be taken into consideration when constructing global carbon budgets with ocean interior inventory changes. Indeed, a potential loss of 5 ± 3 Pg C was considered in the global assessment of the accumulation of anthropogenic CO₂ for the period 1994 through 2007⁴. In addition to circulation driven decadal variability, a part of this loss could be caused by ocean warming, as a warming induced loss of 5 ± 1 Pg C between 1990 and 2000¹³⁶ has been suggested (Table 1). These losses and the corresponding budget adjustments are currently very tentative, and urgently require verifications through direct observations of changes in the oceanic DIC pool, for example.

While ocean interior and surface ocean constraints are becoming more consistent, new discrepancies have arisen. Most prominent is a growing difference between the ocean sink estimates based on surface ocean pCO₂ observations and those based on ocean biogeochemical models. These estimates agree well during the first decade of the millennium, but diverge thereafter, with the observation-based estimates indicating a much

larger growth in the uptake than the models^{1,25} (Table 1). This difference is also evident in these models yielding a relatively low sensitivity β of 1.11 ± 0.18 Pg C / ppm CO₂ for the period 1990 through 2018 (Fig 1a, Supplementary Table 1). One reason is that the presently used models tend to underestimate the uptake of anthropogenic CO₂, as evidenced by direct comparison with the uptake estimates stemming from the accumulation of anthropogenic CO₂ (Fig 1b,c)¹. A model-based emergent constraint approach on a different, but related set of models suggests an underestimation of about 10%¹³⁷. Adjusting the models for this bias halves the mismatch between models and observations-based estimates for the period after 2010, but opens larger discrepancies in the earlier decades. The uncertainties in the observation-based flux products stemming from the sparse observations, and issues at the tails of the observational-based time-series¹¹² might contribute to these discrepancies.

[H1] SUMMARY AND FUTURE PERSPECTIVES

The strength of the ocean carbon sink has tripled from the 1960s until the present. Thus, the ocean has maintained its key role as a sink for the CO₂ emitted into the atmosphere as a consequence of human activities, removing about $25 \pm 2\%$ of the total emissions over six decades. The strengthening of the ocean sink has been largely driven by the increasing uptake of anthropogenic CO₂ in response to the rise in atmospheric CO₂, leading to a strong proportionality between the two. In contrast, the contribution from changes in the natural carbon cycle has been small so far, consistent with the assumption that the ocean circulation and biological pump was overall in steady-state. However, new insights and observations in the past decade challenge this assumption, especially on shorter timescales, suggesting an ocean that is more variable than previously recognized. New evidence also suggests over the past three decades a loss of natural CO₂ to the atmosphere due to ocean warming and changes in ocean circulation. If confirmed, such a loss suggests an ocean carbon sink that is rather vulnerable to climate change.

An ocean sink that is more vulnerable to climate change than currently assumed in coupled carbon-climate models⁵² would imply that the ocean will take up less CO₂ from the atmosphere in the future than anticipated. This would leave a larger fraction of the emissions in the atmosphere, causing additional global warming and climate change. In other words, the ocean carbon-climate feedback could be more positive than suggested by current coupled carbon-climate models. Moreover, the finding of the ocean sink potentially being more sensitive to changes in atmospheric CO₂ growth rates than previously recognized, implies a stronger than anticipated decline of the ocean carbon sink in ambitious mitigation scenarios^{34,138}.

The implications are large and far-reaching. Any reduction in ocean carbon uptake compared to current assumptions would require even stronger investments into decarbonization strategies, making the achievement of specific global warming targets harder. It also reduces the efficacy of the negative emission

approaches that aim to curb climate change by removing CO₂ from the atmosphere using land-based^{139,140} or ocean-based¹⁴¹ approaches.

To better constrain and predict the ocean carbon sink, there are three important challenges to address: the robustness of the reconstructed changes and variations; the processes driving these changes and variations; and predictions of the future ocean uptake, in particular the response of the ocean carbon sink to future climate change, the reduction in anthropogenic CO₂ emissions, and the potential deployment of carbon dioxide removal technologies. Addressing these challenges is important both scientifically and for policy. For example, during the upcoming Global Stocktake undertaken within the U.N. Framework Convention on Climate Change (UNFCCC), reliable estimates of the ocean carbon sink will be a crucial element to close the global carbon budget. In addition, the study of ocean-based carbon dioxide removal approaches, such as ocean alkalization, nutrient fertilization, seaweed growth, and artificial upwelling, have gained momentum¹⁴¹, requiring a thorough assessment of their effectiveness and consequences.

In our view, the following measures must be taken to answer these challenges (see also Ref¹⁴²). The existing observation networks need to be improved, expanded, and put on a much better long-term funding level. The limited sampling of the ocean carbon system is currently the largest source of uncertainty in assessing the variability of the ocean carbon sink. The current sampling is sufficient to capture the long-term time mean sink, and the year-to-year variations in the tropical Pacific and a few other regions, especially in the northern hemisphere where the sampling is relatively dense. In contrast, sampling is critical in many other key regions, such as the Southern Ocean, the South Pacific and the Indian Ocean. Higher resolution observations in time and space will also help to better understand the processes leading to these variations, including those that lead to extremes in ocean acidification and/or deoxygenation¹⁴³. Ocean observing system simulation experiments can help to determine where and when the observing density has to be increased, and to suggest optimal combinations of different observing platforms^{144,145}.

To support observation, new technologies—especially those that enhance the ability to observe ocean carbon in an autonomous manner—need to be developed, improved, and strategically deployed. Improvement of analytical techniques, sensor technology and calibration methods for ocean carbon measurements is urgently required for the provision of accurate, well-calibrated ocean carbon measurements, while improving the ease and efficiency of data collection, thus increasing the scope for autonomous data collection and reducing the cost of these measurements, such as the Biogeochemical Argo program^{146–148}.

To build on expanded and improved sampling, the existing ocean carbon synthesis projects (GLODAP and SOCAT) and the downstream efforts such as the Global Carbon Budget (GCB) and SeaFlux need to be strengthened and expanded. A more rapid update of the analyses, such as on a semi-annual basis providing closer to real-time analyses of the global carbon budget, could be useful to better linking the ocean to the

Global Stocktake activities. Similarly, models and observations need to be better integrated, especially through data assimilation and interpolation approaches^{149–151}. As part of this effort, these models should be pushed to resolve smaller spatial and temporal scales, better capturing the small-scale variability that is inherent in the data that is collected and assimilated by these models. If these models can resolve both the large scales that are representative of global budgets, and the small scales that are representative of the observations, they will be able to more accurately reflect our state of knowledge and its uncertainty.

Moving beyond carbon measurements and budgets, focused process studies need to be developed to better understand critical processes. The need to improve knowledge of the sensitivity of ocean biology to changes in temperature, ocean acidification and other parameters is pressing. In addition researchers need a better understanding of the aquatic continuum¹⁰⁵—the aquatic network that connects the land aquatic systems to the ocean, delivering inorganic and organic matter to the ocean, whose fate is critical to determine the outgassing of river-derived CO₂. Although a value of 0.65 Pg C yr⁻¹ for the degassing of terrestrially-derived CO₂ was used here and in the Global Carbon Budget¹, individual estimates range between 0.2 Pg C yr⁻¹¹⁵² and 1.2 Pg C yr⁻¹¹⁵³, reflecting the large uncertainty of this estimate. An especially under-investigated area is the fate of the river-derived carbon in the ocean, and in particular, the determination of how much carbon is buried in sediments close to the river mouths, how much enters the open ocean and how fast this carbon is remineralized back to CO₂¹⁵².

The role of the ocean in taking up additional CO₂ in response to the deployment of carbon dioxide removal technologies needs to be critically evaluated. There must be a particular focus on the efficacy of these measures and their potential for negative (unintended) consequences¹⁵⁴. Historically, the ocean sink for carbon has been considered as very robust to changes, and largely tracking the increase in atmospheric CO₂. It is time to change this perspective and to recognize that the ocean carbon cycle might be more sensitive to change than previously recognized. The size of this sink, its unknown response to a reduction in anthropogenic CO₂ emissions and its relevance for past and future climates are large enough to warrant renewed efforts to observe it, to study it, and to understand it.

References

Top references

(1) Friedlingstein et al. (2022)

Most recent version of the Global Carbon Budget, an international effort led by the Global Carbon Project to synthesize all components of the global carbon cycle.

(2) Sabine et al. (2004)

First observation-based global inventory of anthropogenic CO₂ providing a key constraint for the global anthropogenic CO₂ budget.

(3) Khatiwala et al. (2009)

Reconstruction of the entire history of the oceanic uptake of anthropogenic CO₂.

(4) Gruber et al. (2019)

Inventory of anthropogenic CO₂ that provided a second time-point describing the accumulation of anthropogenic CO₂ in the ocean based on ocean interior observations.

(25) Hauck et al. (2020)

Description and assessment of the ocean biogeochemical models currently used to determine the oceanic uptake of CO₂ in the context of the Global Carbon Budget.

(30) Le Quéré et al., (2007)

First study to point out that the Southern Ocean carbon sink weakened substantially during the 1990s.

(32) Landschützer et al. (2016)

Assessment of the decadal variability of the ocean carbon sink that reveals it is driven by the extratropical latitudes in both hemispheres.

(72) Mikaloff-Fletcher et al. (2006)

Ocean inversion-based study describing the regional distribution of the air-sea fluxes of anthropogenic CO₂ and its oceanic transport.

References by number:

1. Friedlingstein, P. *et al.* Global Carbon Budget 2021. *Earth Syst. Sci. Data* **14**, 1917–2005 (2022).
2. Sabine, C. L. *et al.* The Oceanic Sink for Anthropogenic CO₂. *Science* (80-.). **305**, 367–371 (2004).
3. Khatiwala, S., Primeau, F. & Hall, T. Reconstruction of the history of anthropogenic CO₂ concentrations in the ocean. *Nature* **462**, 346–349 (2009).
4. Gruber, N. *et al.* The oceanic sink for anthropogenic CO₂ from 1994 to 2007. *Science* (80-.). **363**, 1193–1199 (2019).
5. Revelle, R. Introduction: The scientific history of carbon dioxide. in *The Global Carbon Cycle and atmospheric CO₂: Natural Variations Archean to Present* (eds. Sundquist, E. T. & Broecker, W. S.) 1–4 (AGU, 1985).
6. Heimann, M. A review of the contemporary global carbon cycle and as seen a century ago by Arrhenius and Høgbom. *Ambio* **26**, 17–24 (1997).
7. Arrhenius, S. *Lehrbuch der kosmischen Physik*. **2**, (Hirzel, 1903).
8. Archer, D., Kheshgi, H. & Maier-Reimer, E. Multiple timescales for neutralization of fossil fuel CO₂. *Geophys. Res. Lett.* **24**, 405–408 (1997).
9. Sarmiento, J. L. & Gruber, N. *Ocean Biogeochemical Dynamics*. (Princeton University Press, 2006).
10. Callendar, G. S. The artificial production of carbon dioxide and its influence on climate. *Q. J. R. Meteor. Soc.* **64**, 223–240 (1938).
11. Revelle, R. & Suess, H. E. Carbon Dioxide Exchange between Atmosphere and Ocean and the question of an increase of atmospheric {CO₂} during the past decades. *Tellus* **9**, 18–27 (1957).
12. Revelle, R., Broecker, W. S., Craig, H., Keeling, C. D. & Smagorinsky, J. *Atmospheric Carbon Dioxide*. (1965).
13. Charney, J. G. *et al.* *Carbon Dioxide and Climate: A scientific assessment*. (1979).
14. Keeling, C. D. The concentration and isotopic abundances of carbon dioxide in the atmosphere. *Tellus* **12**, 200–203 (1960).
15. W.R.Wallace, D. Chapter 6.3 Storage and transport of excess CO₂ in the oceans:The JGOFS/WOCE global CO₂ survey. in *Ocean circulation and climate* (eds. Siedler, G., Church, J. & Gould, J.) 489–L (Academic Press, 2001). doi:10.1016/S0074-6142(01)80136-4
16. Keppler, L., Landschützer, P., Gruber, N., Lauvset, S. K. & Stemmler, I. Seasonal Carbon Dynamics in the Near-Global Ocean. *Global Biogeochem. Cycles* **34**, (2020).
17. Canadell, P. G. *et al.* Global Carbon and other Biogeochemical Cycles and Feedbacks. in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* 673–816 (2021). doi:10.1017/9781009157896.007
18. Oeschger, H., Siegenthaler, U., Schotterer, U. & Gugelmann, A. A box diffusion model to study the carbon dioxide exchange in nature. *Tellus* **27**, 168–192 (1975).
19. Brewer, P. G. Direct observation of the oceanic {CO₂} increase. *Geophys. Res. Lett.* **5**, 997–1000 (1978).
20. Chen, C.-T. A. & Millero, F. J. Gradual increase of oceanic CO₂. *Nature* **277**, 205–206 (1979).
21. Sabine, C. L. & Tanhua, T. Estimation of Anthropogenic CO₂ Inventories in the Ocean. *Ann. Rev. Mar. Sci.* **2**, 175–198 (2010).
22. Olsen, A. *et al.* The Global Ocean Data Analysis Project version 2 (GLODAPv2) – an internally consistent data product for the world ocean. *Earth Syst. Sci. Data* **8**, 297–323 (2016).
23. DeVries, T. *et al.* Decadal trends in the ocean carbon sink. *Proc. Natl. Acad. Sci.* **116**, 201900371 (2019).
24. Gruber, N., Landschützer, P. & Lovenduski, N. S. The Variable Southern Ocean Carbon Sink. *Ann. Rev. Mar. Sci.* **11**, 159–186 (2019).
25. Hauck, J. *et al.* Consistency and Challenges in the Ocean Carbon Sink Estimate for the Global Carbon Budget. *Front. Mar. Sci.* **7**, 1–33 (2020).
26. Landschützer, P., Gruber, N., Bakker, D. C. E. & Schuster, U. Recent variability of the global ocean

- 611 carbon sink. *Global Biogeochem. Cycles* **28**, 927–949 (2014).
- 612 27. Landschützer, P., Gruber, N. & Bakker, D. C. E. Decadal variations and trends of the global ocean
613 carbon sink. *Global Biogeochem. Cycles* **30**, 1396–1417 (2016).
- 614 28. Le Quere, C. *et al.* Saturation of the Southern Ocean CO₂ Sink Due to Recent Climate Change.
615 *Science* (80-.). **316**, 1735–1738 (2007).
- 616 29. Lovenduski, N. S., Gruber, N., Doney, S. C. & Lima, I. D. Enhanced CO₂ outgassing in the Southern
617 Ocean from a positive phase of the Southern Annular Mode. *Global Biogeochem. Cycles* **21**, n/a-n/a
618 (2007).
- 619 30. Le Quéré, C., Orr, J. C., Monfray, P., Aumont, O. & Madec, G. Interannual variability of the oceanic
620 sink of CO₂ from 1979 through 1997. *Global Biogeochem. Cycles* **14**, 1247–1265 (2000).
- 621 31. Fay, A. R. & McKinley, G. A. Global trends in surface ocean p CO₂ from in situ data. *Global*
622 *Biogeochem. Cycles* **27**, 541–557 (2013).
- 623 32. Landschützer, P. *et al.* The reinvigoration of the Southern Ocean carbon sink. *Science* (80-.). **349**,
624 1221–1224 (2015).
- 625 33. McKinley, G. A., Fay, A. R., Eddebbar, Y. A., Gloege, L. & Lovenduski, N. S. External Forcing
626 Explains Recent Decadal Variability of the Ocean Carbon Sink. *AGU Adv.* **1**, 1–10 (2020).
- 627 34. Rogelj, J. *et al.* Mitigation Pathways Compatible with 1.5°C of Sustainable Development. in *Global*
628 *warming of 1.5°C. An IPCC Special Report [...]* 93–174 (2018).
- 629 35. Cheng, L. *et al.* Another Record: Ocean Warming Continues through 2021 despite La Niña
630 Conditions. *Adv. Atmos. Sci.* (2022). doi:10.1007/s00376-022-1461-3
- 631 36. Abram, N. *et al.* Chapter 1: Framing and Context of the Report. in *Special Report on the Ocean and*
632 *Cryosphere (SROCC)* (eds. Pörtner, H.-O. *et al.*) (2019).
- 633 37. Bindoff, N. L. *et al.* Chapter 5: Changing Ocean, Marine Ecosystems, and Dependent Communities.
634 in *Special Report on the Ocean and Cryosphere (SROCC)* (eds. Pörtner, H.-O. *et al.*) (2019).
- 635 38. Sarmiento, J. L., Orr, J. C. & Siegenthaler, U. A perturbation Simulation of CO₂ uptake in an Ocean
636 General Circulation Model. *J. Geophys. Res.* **97**, 3621–3645 (1992).
- 637 39. Matsumoto, K. Radiocarbon-based circulation age of the world oceans. *J. Geophys. Res.* **112**, 1–7
638 (2007).
- 639 40. Holzer, M. & Primeau, F. W. The path-density distribution of oceanic surface-to-surface transport. *J.*
640 *Geophys. Res. Ocean.* **113**, 1–22 (2008).
- 641 41. Dong, Y. *et al.* Update on the Temperature Corrections of Global Air-Sea CO₂ Flux Estimates.
642 *Global Biogeochem. Cycles* **26**, 21–35 (2022).
- 643 42. Matsumoto, K. & Gruber, N. How accurate is the estimation of anthropogenic carbon in the ocean?
644 An evaluation of the ΔC^* method. *Global Biogeochem. Cycles* **19**, (2005).
- 645 43. Bates, N. *et al.* A Time-Series View of Changing Ocean Chemistry Due to Ocean Uptake of
646 Anthropogenic CO₂ and Ocean Acidification. *Oceanography* **27**, 126–141 (2014).
- 647 44. Gregor, L. & Gruber, N. OceanSODA-ETHZ: a global gridded data set of the surface ocean carbonate
648 system for seasonal to decadal studies of ocean acidification. *Earth Syst. Sci. Data* **13**, 777–808
649 (2021).
- 650 45. Broecker, W. S., Takahashi, T., Simpson, H. J. & Peng, T. H. Fate of fossil fuel carbon dioxide and
651 the global carbon budget. *Science* (80-.). **206**, 409–418 (1979).
- 652 46. Egleston, E. S., Sabine, C. L. & Morel, F. M. M. Revelle revisited: Buffer factors that quantify the
653 response of ocean chemistry to changes in DIC and alkalinity. *Global Biogeochem. Cycles* **24**, 1–9
654 (2010).
- 655 47. Jiang, L. Q., Carter, B. R., Feely, R. A., Lauvset, S. K. & Olsen, A. Surface ocean pH and buffer
656 capacity: past, present and future. *Sci. Rep.* **9**, 18624 (2019).
- 657 48. Sarmiento, J. L., LeQuéré, C. & Pacala, S. W. Limiting future atmospheric carbon dioxide. *Glob.*
658 *Biogeochem. Cycles* **9**, 121–137 (1995).
- 659 49. Joos, F. *et al.* An efficient and accurate representation of complex oceanic and biospheric models of
660 anthropogenic carbon uptake. *Tellus* **48B**, 397–417 (1996).
- 661 50. Keeling, C. D. The Suess Effect: 13-Carbon and 14-Carbon Interactions. in *Environment International*

- 2, 229–300 (Pergamon, 1979).
51. Friedlingstein, P. *et al.* Global Carbon Budget 2020. *Earth Syst. Sci. Data* **12**, 3269–3340 (2020).
52. Arora, V. K. *et al.* Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models. *Biogeosciences* **17**, 4173–4222 (2020).
53. Friedlingstein, P. *et al.* Climate-carbon cycle feedback analysis: results from the {C4MIP} model intercomparison. *J. Clim.* **19**, 3337–3353 (2006).
54. Meinshausen, M. *et al.* Realization of Paris Agreement pledges may limit warming just below 2 °C. *Nature* **604**, 304–309 (2022).
55. Joos, F. *et al.* Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: A multi-model analysis. *Atmos. Chem. Phys.* **13**, 2793–2825 (2013).
56. Waugh, D. W., Hall, T. M., Mcneil, B. I., Key, R. & Matear, R. J. Anthropogenic CO₂ in the oceans estimated using transit time distributions. *Tellus B Chem. Phys. Meteorol.* **58**, 376–389 (2006).
57. Tanhua, T. *et al.* Ventilation of the Arctic Ocean: Mean ages and inventories of anthropogenic CO₂ and CFC-11. *J. Geophys. Res.* **114**, 1–11 (2009).
58. Raimondi, L., Tanhua, T., Azetsu-Scott, K., Yashayaev, I. & Wallace, D. W. R. A 30 -Year Time Series of Transient Tracer-Based Estimates of Anthropogenic Carbon in the Central Labrador Sea. *J. Geophys. Res. Ocean.* **126**, 1–19 (2021).
59. Ridge, S. M. & McKinley, G. A. Ocean carbon uptake under aggressive emission mitigation. *Biogeosciences* **18**, 2711–2725 (2021).
60. Archer, D., Kheshgi, H. & Maier-Reimer, E. Dynamics of fossil fuel {CO}₂ neutralization by marine {CaCO}₃. *Glob. Biogeochem. Cycles* **12**, 259–276 (1998).
61. Bacastow, R. B. & Keeling, C. D. Models to predict future atmospheric CO₂ concentrations. in *Workshop on the Global Effects of Carbon Dioxide from Fossil Fuels* (eds. Elliott, W. P. & Machta, L.) 72–90 (United States Department of Energy, 1979).
62. Siegenthaler, U. & Oeschger, H. Predicting future atmospheric carbon dioxide levels. *Science* **199**, 388–95 (1978).
63. Wallace, D. W. R. *Monitoring Global Ocean Carbon Inventories*. (1995).
64. Gruber, N., Sarmiento, J. L. & Stocker, T. F. An improved method for detecting anthropogenic CO₂ in the oceans. *Global Biogeochem. Cycles* **10**, 809–837 (1996).
65. Gruber, N. Anthropogenic CO₂ in the Atlantic Ocean. *Global Biogeochem. Cycles* **12**, 165–191 (1998).
66. Dickson, A. G., Goyet, C. & DOE. *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water; version 2*. (1994).
67. Dickson, A. G., Afghan, J. D. & Anderson, G. C. Reference materials for oceanic {CO}₂ analysis: a method for the certification of total alkalinity. *Mar. Chem.* **80**, 185–197 (2003).
68. Dickson, A. G. Standards for ocean measurements. *Oceanography* **23**, 34–47 (2010).
69. Key, R. M. *et al.* A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP). *Global Biogeochem. Cycles* **18**, n/a-n/a (2004).
70. DeVries, T. The oceanic anthropogenic CO₂ sink: Storage, air-sea fluxes, and transports over the industrial era. *Global Biogeochem. Cycles* **28**, 631–647 (2014).
71. Orr, J. C. *et al.* Estimates of anthropogenic carbon uptake from four three-dimensional global ocean models. *Global Biogeochem. Cycles* (2001).
72. Mikaloff Fletcher, S. E. *et al.* Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean. *Global Biogeochem. Cycles* **20**, 1–16 (2006).
73. Davila, X. *et al.* How Is the Ocean Anthropogenic Carbon Reservoir Filled? *Global Biogeochem. Cycles* **36**, 1–16 (2022).
74. Groeskamp, S., Lenton, A., Matear, R., Sloyan, B. M. & Langlais, C. Anthropogenic carbon in the ocean-Surface to interior connections. *Global Biogeochem. Cycles* **30**, 1682–1698 (2016).
75. Keeling, R. F. & Shertz, S. R. Seasonal and interannual variations in atmospheric oxygen and implications for the global carbon cycle. *Nature* **358**, 723–727 (1992).
76. Quay, P. D., Tilbrook, B. & Wong, C. S. Oceanic Uptake of Fossil Fuel CO₂ : Carbon-13 Evidence.

- 713 *Science* (80-.). **256**, 74–79 (1992).
- 714 77. Heimann, M. & Maier-Reimer, E. On the relations between the oceanic uptake of CO₂ and its carbon
715 isotopes. *Global Biogeochem. Cycles* **10**, 89–110 (1996).
- 716 78. Gruber, N. & Keeling, C. D. An improved estimate of the isotopic air-sea disequilibrium of CO₂ :
717 Implications for the oceanic uptake of anthropogenic CO₂. *Geophys. Res. Lett.* **28**, 555–558 (2001).
- 718 79. Khatiwala, S. *et al.* Global ocean storage of anthropogenic carbon. *Biogeosciences* **10**, 2169–2191
719 (2013).
- 720 80. Friedrich, T. *et al.* Detecting regional anthropogenic trends in ocean acidification against natural
721 variability. *Nature* **2**, 167–171 (2012).
- 722 81. Munro, D. R. *et al.* Recent evidence for a strengthening CO₂ sink in the Southern Ocean from
723 carbonate system measurements in the Drake Passage (2002–2015). *Geophys. Res. Lett.* **42**, 7623–
724 7630 (2015).
- 725 82. Talley, L. D. *et al.* Changes in Ocean Heat, Carbon Content, and Ventilation: A Review of the First
726 Decade of GO-SHIP Global Repeat Hydrography. *Ann. Rev. Mar. Sci.* **8**, 185–215 (2016).
- 727 83. Wanninkhof, R. *et al.* Detecting anthropogenic CO₂ changes in the interior Atlantic Ocean between
728 1989 and 2005. *J. Geophys. Res.* **115**, C11028 (2010).
- 729 84. Friis, K., Körtzinger, A., Pätsch, J. & Wallace, D. W. R. On the temporal increase of anthropogenic
730 CO₂ in the subpolar North Atlantic. *Deep. Res. Part I* **52**, 681–698 (2005).
- 731 85. Goodkin, N. F., Levine, N. M., Doney, S. C. & Wanninkhof, R. Impacts of temporal CO₂ and climate
732 trends on the detection of ocean anthropogenic CO₂ accumulation. *Global Biogeochem. Cycles* **25**, 1–
733 11 (2011).
- 734 86. Levine, N. M., Doney, S. C., Wanninkhof, R., Lindsay, K. & Fung, I. Y. Impact of ocean carbon
735 system variability on the detection of temporal increases in anthropogenic CO₂. *J. Geophys. Res.* **113**,
736 C03019 (2008).
- 737 87. Carter, B. R. *et al.* Two decades of Pacific anthropogenic carbon storage and ocean acidification along
738 Global Ocean Ship-based Hydrographic Investigations Program sections P16 and P02. *Global*
739 *Biogeochem. Cycles* **31**, 306–327 (2017).
- 740 88. Carter, B. R. *et al.* Pacific Anthropogenic Carbon Between 1991 and 2017. *Global Biogeochem.*
741 *Cycles* 2018GB006154 (2019). doi:10.1029/2018GB006154
- 742 89. Woosley, R. J., Millero, F. J. & Wanninkhof, R. Rapid anthropogenic changes in CO₂ and pH in the
743 Atlantic Ocean: 2003–2014. *Global Biogeochem. Cycles* **30**, 70–90 (2016).
- 744 90. Clement, D. & Gruber, N. The eMLR(C*) Method to Determine Decadal Changes in the Global
745 Ocean Storage of Anthropogenic CO₂. *Global Biogeochem. Cycles* **32**, 654–679 (2018).
- 746 91. Tanhua, T., Körtzinger, A., Friis, K., Waugh, D. W. & Wallace, D. W. R. An estimate of
747 anthropogenic CO₂ inventory from decadal changes in oceanic carbon content. *Proc. Natl. Acad. Sci.*
748 **104**, 3037–3042 (2007).
- 749 92. Pérez, F. F. *et al.* Atlantic Ocean CO₂ uptake reduced by weakening of the meridional overturning
750 circulation. *Nat. Geosci.* **6**, 146–152 (2013).
- 751 93. Keeling, C. D. Carbon Dioxide in Surface Ocean Waters: 4. Global Distribution. *J. Geophys. Res.* **73**,
752 4543–4553 (1968).
- 753 94. Tans, P. P., Fung, I. Y. & Takahashi, T. Observational Constrains on the Global Atmospheric Co₂
754 Budget. *Science* (80-.). **247**, 1431–1438 (1990).
- 755 95. Takahashi, T. *et al.* Global air-sea flux of CO₂ : An estimate based on measurements of sea–air pCO₂
756 difference. *Proc. Natl. Acad. Sci.* **94**, 8292–8299 (1997).
- 757 96. Takahashi, T. *et al.* Deep-Sea Research II Climatological mean and decadal change in surface ocean
758 pCO₂ , and net sea – air CO₂ flux over the global oceans. **56**, 554–577 (2009).
- 759 97. Bakker, D. C. E. *et al.* An update to the Surface Ocean CO₂ Atlas (SOCAT version 2). *Earth Syst. Sci.*
760 *Data* **6**, 69–90 (2014).
- 761 98. Pfeil, B. *et al.* A uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT). *Earth Syst. Sci.*
762 *Data* **5**, 125–143 (2013).
- 763 99. Bakker, D. C. E. *et al.* A multi-decade record of high-quality CO₂ data in version 3 of the Surface

- 764 Ocean CO₂ Atlas (SOCAT). *Earth Syst. Sci. Data* **8**, 383–413 (2016).
- 765 100. Rödenbeck, C. *et al.* Data-based estimates of the ocean carbon sink variability - First results of the
766 Surface Ocean pCO₂ Mapping intercomparison (SOCOM). *Biogeosciences* **12**, 7251–7278 (2015).
- 767 101. Landschützer, P. *et al.* A neural network-based estimate of the seasonal to inter-annual variability of
768 the Atlantic Ocean carbon sink. *Biogeosciences* **10**, 7793–7815 (2013).
- 769 102. Gregor, L., Lebehot, A. D., Kok, S. & Scheel Monteiro, P. M. A comparative assessment of the
770 uncertainties of global surface ocean CO₂ estimates using a machine-learning ensemble (CSIR-ML6
771 version 2019a)-Have we hit the wall? *Geosci. Model Dev.* **12**, 5113–5136 (2019).
- 772 103. Fay, A. R. *et al.* SeaFlux: Harmonization of air-sea CO₂ fluxes from surface pCO₂ data products
773 using a standardized approach. *Earth Syst. Sci. Data* **13**, 4693–4710 (2021).
- 774 104. Gruber, N. *et al.* Oceanic sources, sinks, and transport of atmospheric CO₂. *Global Biogeochem.*
775 *Cycles* **23**, (2009).
- 776 105. Regnier, P., Resplandy, L., Najjar, R. G. & Ciais, P. The land-to-ocean loops of the global carbon
777 cycle. *Nature* **603**, 401–410 (2022).
- 778 106. Sarmiento, J. L. & Sundquist, E. T. Revised budget for the oceanic uptake of anthropogenic carbon
779 dioxide. *Nature* **356**, 589–593 (1992).
- 780 107. Regnier, P. *et al.* Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nat. Geosci.* **6**,
781 597–607 (2013).
- 782 108. Resplandy, L. *et al.* Revision of global carbon fluxes based on a reassessment of oceanic and riverine
783 carbon transport. *Nat. Geosci.* **11**, (2018).
- 784 109. Landschützer, P., Ilyina, T. & Lovenduski, N. S. Detecting Regional Modes of Variability in
785 Observation-Based Surface Ocean pCO₂. *Geophys. Res. Lett.* **46**, 2670–2679 (2019).
- 786 110. Ritter, R. *et al.* Observation-Based Trends of the Southern Ocean Carbon Sink. *Geophys. Res. Lett.*
787 **44**, 12,339–12,348 (2017).
- 788 111. Gloege, L. *et al.* Quantifying errors in observationally-based estimates of ocean carbon sink
789 variability. *Global Biogeochem. Cycles* 1–14 (2021). doi:10.1029/2020gb006788
- 790 112. Watson, A. J. *et al.* Revised estimates of ocean-atmosphere CO₂ flux are consistent with ocean carbon
791 inventory. *Nat. Commun.* **11**, 1–6 (2020).
- 792 113. Wanninkhof, R., Asher, W. E., Ho, D. T., Sweeney, C. & McGillis, W. R. Advances in Quantifying
793 Air-Sea Gas Exchange and Environmental Forcing*. *Ann. Rev. Mar. Sci.* **1**, 213–244 (2009).
- 794 114. DeVries, T., Holzer, M. & Primeau, F. Recent increase in oceanic carbon uptake driven by weaker
795 upper-ocean overturning. *Nature* **542**, 215–218 (2017).
- 796 115. Wolter, K. & Timlin, M. S. El Niño/Southern Oscillation behaviour since 1871 as diagnosed in an
797 extended multivariate ENSO index (MEI.ext). *Int. J. Climatol.* **31**, 1074–1087 (2011).
- 798 116. Frölicher, T. L. *et al.* Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in
799 CMIP5 Models. *J. Clim.* **28**, 862–886 (2015).
- 800 117. McKinley, G. A., Rödenbeck, C., Gloor, M., Houweling, S. & Heimann, M. Pacific dominance to
801 global air-sea {CO₂} flux variability: A novel atmospheric inversion agrees with ocean models.
802 *Geophys. Res. Lett.* **31**, (2004).
- 803 118. Lenton, A. & Mearns, R. J. Role of the Southern Annular Mode (SAM) in Southern Ocean CO₂
804 uptake. *Global Biogeochem. Cycles* **21**, 1–17 (2007).
- 805 119. Keppler, L. & Landschützer, P. Regional Wind Variability Modulates the Southern Ocean Carbon
806 Sink. *Sci. Rep.* **9**, 7384 (2019).
- 807 120. Le Quéré, C., Takahashi, T., Buitenhuis, E. T., Rödenbeck, C. & Sutherland, S. C. Impact of climate
808 change and variability on the global oceanic sink of CO₂. *Global Biogeochem. Cycles* **24**, n/a-n/a
809 (2010).
- 810 121. Feely, R. A., Wanninkhof, R., Takahashi, T. & Tans, P. Influence of El Niño on the equatorial Pacific
811 contribution to atmospheric CO₂ accumulation. *Nature* **398**, 597–601 (1999).
- 812 122. Ishii, M. *et al.* Air-sea CO₂ flux in the Pacific Ocean for the period 1990–2009. *Biogeosciences* **11**,
813 709–734 (2014).
- 814 123. McKinley, G. A., Follows, M. J. & Marshall, J. Mechanisms of air-sea CO₂ flux variability in the

- equatorial Pacific and the North Atlantic. *Global Biogeochem. Cycles* **18**, n/a-n/a (2004).
124. Chatterjee, A. *et al.* Influence of El Niño on atmospheric CO₂ over the tropical Pacific Ocean: Findings from NASA’s OCO-2 mission. *Science* (80-.). **358**, (2017).
 125. Keeling, C. D., Whorf, T. P., Wahlen, M. & v. d. Plicht, J. Interannual extremes in the rate of atmospheric carbon dioxide since 1980. *Nature* **375**, 666–670 (1995).
 126. Crisp, D. *et al.* How Well Do We Understand the Land-Ocean-Atmosphere Carbon Cycle? *Rev. Geophys.* 1–64 (2022). doi:10.1029/2021rg000736
 127. Angert, A., Biraud, S., Bonfils, C., Buermann, W. & Fung, I. CO₂ seasonality indicates origins of post-Pinatubo sink. *Geophys. Res. Lett.* **31**, 1999–2002 (2004).
 128. Eddebbar, Y. A. *et al.* El Niño-like physical and biogeochemical ocean response to tropical eruptions. *J. Clim.* **32**, 2627–2649 (2019).
 129. Marshall, L. R. *et al.* Volcanic effects on climate : recent advances and future avenues. (2022).
 130. Thompson, D. W. J. & Solomon, S. Interpretation of recent Southern Hemisphere climate change. *Science* **296**, 895–899 (2002).
 131. Hauck, J. *et al.* Seasonally different carbon flux changes in the Southern Ocean in response to the southern annular mode. *Global Biogeochem. Cycles* **27**, 1236–1245 (2013).
 132. Lovenduski, N. S., Gruber, N. & Doney, S. C. Toward a mechanistic understanding of the decadal trends in the Southern Ocean carbon sink. *Global Biogeochem. Cycles* **22**, 1–9 (2008).
 133. Gillett, N. P. & Thompson, D. W. J. Simulation of recent southern hemisphere climate change. *Science* **302**, 273–5 (2003).
 134. Gruber, N., Bates, N. R. & Keeling, C. D. Interannual variability in the Northern Atlantic carbon sink. *Science* (80-.). **298**, 2374–2378 (2002).
 135. Frölicher, T. L., Joos, F., Raible, C. C. & Sarmiento, J. L. Atmospheric CO₂ response to volcanic eruptions: The role of ENSO, season, and variability. *Global Biogeochem. Cycles* **27**, 239–251 (2013).
 136. DeVries, T. Atmospheric CO₂ and Sea Surface Temperature Variability Cannot Explain Recent Decadal Variability of the Ocean CO₂ Sink . *Geophys. Res. Lett.* **49**, 1–12 (2022).
 137. Terhaar, J., Frölicher, T. L. & Joos, F. Observation-constrained estimates of the global ocean carbon sink from Earth system models. *Biogeosciences* **19**, 4431–4457 (2022).
 138. Rogelj, J., McCollum, D. L., O’Neill, B. C. & Riahi, K. 2020 emissions levels required to limit warming to below 2 °C. *Nat. Clim. Chang.* **3**, 405–412 (2012).
 139. Keller, D. P. *et al.* The Effects of Carbon Dioxide Removal on the Carbon Cycle. *Curr. Clim. Chang. Reports* **4**, 250–265 (2018).
 140. Smith, P. *et al.* Biophysical and economic limits to negative CO₂ emissions. *Nat. Clim. Chang.* **6**, 42–50 (2016).
 141. National Academies of Sciences. *A Research Strategy for Ocean-based Carbon Dioxide Removal and Sequestration*. (National Academies Press, 2022). doi:10.17226/26278
 142. Aricò, S. *et al.* *Integrated Ocean Carbon Research: A Summary of Ocean Carbon Research, and Vision of Coordinated Ocean Carbon Research and Observations for the Next Decade*. (2021). doi:10.25607/h0gj-pq41
 143. Gruber, N., Boyd, P. W., Frölicher, T. L. & Vogt, M. Biogeochemical extremes and compound events in the ocean. *Nature* **600**, 395–407 (2021).
 144. Djeutchouang, L. M., Chang, N., Gregor, L., Vichi, M. & Monteiro, P. M. S. The sensitivity of p CO₂ reconstructions to sampling scales across a Southern Ocean sub-domain: a semi-idealized ocean sampling simulation approach. *Biogeosciences* **19**, 4171–4195 (2022).
 145. Majkut, J. D. *et al.* An observing system simulation for Southern Ocean carbon dioxide uptake. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **372**, 20130046–20130046 (2014).
 146. Claustre, H., Johnson, K. S. & Takeshita, Y. Observing the Global Ocean with Biogeochemical-Argo. *Ann. Rev. Mar. Sci.* **12**, 23–48 (2020).
 147. Gray, A. R. *et al.* Autonomous Biogeochemical Floats Detect Significant Carbon Dioxide Outgassing in the High-Latitude Southern Ocean. *Geophys. Res. Lett.* **45**, 9049–9057 (2018).
 148. Bushinsky, S. M. *et al.* Reassessing Southern Ocean air-sea CO₂ flux estimates with the addition of

- biogeochemical float observations. *Global Biogeochem. Cycles* **33**, 1–19 (2019).
149. Verdy, A. & Mazloff, M. R. A data assimilating model for estimating Southern Ocean
biogeochemistry. *J. Geophys. Res. Ocean.* **122**, 6968–6988 (2017).
150. Carroll, D. *et al.* Attribution of Space-Time Variability in Global-Ocean Dissolved Inorganic Carbon.
Global Biogeochem. Cycles **36**, 1–24 (2022).
151. Bennington, V., Gloege, L. & McKinley, G. A. Variability in the Global Ocean Carbon Sink From
1959 to 2020 by Correcting Models With Observations. *Geophys. Res. Lett.* **49**, (2022).
152. Lacroix, F., Ilyina, T. & Hartmann, J. Oceanic CO₂ outgassing and biological production hotspots
induced by pre-industrial river loads of nutrients and carbon in a global modeling approach.
Biogeosciences **17**, 55–88 (2020).
153. Kwon, E. Y. *et al.* Stable Carbon Isotopes Suggest Large Terrestrial Carbon Inputs to the Global
Ocean. *Global Biogeochem. Cycles* 1–25 (2021). doi:10.1029/2020gb006684
154. GESAMP. *High level review of a wide range of proposed marine geoengineering techniques.*
GESAMP Reports and Studies (2019).
155. Dlugokencky, E. & Tans, P. Trends in atmospheric carbon dioxide. *National Oceanic&Atmospheric
Administration; Global Monitoring Laboratory (NOAA/GML)* (2022). Available at:
<http://gml.noaa.gov/ccgg/trends/>. (Accessed: 15th July 2015)
156. Kroeker, K. J. *et al.* Impacts of ocean acidification on marine organisms: quantifying sensitivities and
interaction with warming. *Glob. Chang. Biol.* **19**, 1884–1896 (2013).
157. Landschützer, P., Gruber, N., Bakker, D. C. E., Stemmler, I. & Six, K. D. Strengthening seasonal
marine CO₂ variations due to increasing atmospheric CO₂. *Nat. Clim. Chang.* **8**, 146–150 (2018).
158. Hauck, J. & Völker, C. Rising atmospheric CO₂ leads to large impact of biology on Southern Ocean
CO₂ uptake via changes of the Revelle factor. *Geophys. Res. Lett.* **42**, 1459–1464 (2015).
159. Gruber, N. & Sarmiento, J. L. Large-scale biogeochemical/physical interactions in elemental cycles.
in *THE SEA: Biological-Physical Interactions in the Oceans* (eds. Robinson, A. R., McCarthy, J. J. &
Rothschild, B. J.) **12**, 337–399 (John Wiley and Sons, 2002).

Acknowledgements

N.G., J.-D.M., L.G, and P.L. acknowledge support from the European Union’s Horizon 2020 research and innovation programme under grant agreement. No. 821003 (project 4C). N.G. also acknowledges support from the E.U. Horizon project no. 821001 (SO-CHIC). The work of D.C.E.B. was supported by the E.U. Horizon project no. 820989 (COMFORT). The work reflects only the authors’ views; the European Commission and their executive agency are not responsible for any use that may be made of the information the work contains. G.A.M. acknowledges funding from NSF through the LEAP STC (2019625) and OCE (1948624), NASA (80NSSC22K0150) and NOAA (NA20OAR4310340). J.H. received funding from the Helmholtz Young Investigator Group Marine Carbon and Ecosystem Feedbacks in the Earth System (MarESys), grant number VH-NG-1301.

Author contributions

N.G. led the conceptual design and the implementation and also wrote the first draft. J.-D.M. was responsible for the generation of Fig 1 and Table 1. P.L. generated Fig 2, L.G. generated Figs 3 and 4, and N.G. drew Fig 5. All authors contributed to the outline, discussed the content and conclusions and provided input to the manuscript during all drafting stages.

Competing interests

The authors declare not competing interests.

Peer review information

Nature Reviews Earth & Environment thanks [Referee#1 name], [Referee#2 name] and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information

Supplementary information is available for this paper at <https://doi.org/10.1038/s415XX-XXX-XXXX-X>

926

927 **Tables**

928

929

930

931

TABLE 1.

OCEAN CO₂ UPTAKE FROM 1990-2019.

Method	Reference	Components ^(*)	1990–1999 (Pg C yr ⁻¹)	2000–2009 (Pg C yr ⁻¹)	2010–2019 (Pg C yr ⁻¹)
ATMOSPHERIC CO ₂					
Change in atmospheric CO ₂ (ppm)	(¹⁵⁵)		15.0	18.7	23.6
OCEAN CO ₂ UPTAKE					
Change in interior accumulation of ant. CO ₂ (†)	(⁴)	$F_{\text{ant}}^{\text{ss}} + F_{\text{ant}}^{\text{ns}}$	-2.1±0.2	-2.6±0.3	-3.3±0.3
Ocean inverse model (Green's function)	(³)	$F_{\text{ant}}^{\text{ss}}$	-2.0±0.6	-2.3±0.6	NA
Ocean inverse model (Adjoint)	(¹³⁶)	$F_{\text{ant}}^{\text{ss}}$	-2.2±0.1	-2.5±0.1	-2.9±0.2
Ocean inverse model (Adjoint) (§)	(¹³⁶)	$F_{\text{ant}}^{\text{ss}} + F_{\text{nat}}^{\text{ns}}$	-2.0±0.1	-2.3±0.1	-2.7±0.2
Ocean forward models	(²⁵)	$F_{\text{ant}}^{\text{ss}} + F_{\text{ant}}^{\text{ns}} + F_{\text{nat}}^{\text{ns}}$	-2.0±0.2	-2.1±0.3	-2.5±0.3
Surface ocean pCO ₂ products (#)	(¹⁰³)	$F_{\text{ant}}^{\text{ss}} + F_{\text{ant}}^{\text{ns}} + F_{\text{nat}}^{\text{ns}}$	-2.1±0.4	-2.3±0.2	-3.1±0.2

(*) $F_{\text{ant}}^{\text{ss}}$: steady-state uptake flux component of anthropogenic CO₂ (part driven solely by the increase in atmospheric CO₂); $F_{\text{ant}}^{\text{ns}}$: non-steady-state uptake component of anthropogenic CO₂ (part driven by variations in ocean circulation and other physical drivers); $F_{\text{nat}}^{\text{ns}}$: non-steady-state exchange component of natural CO₂ (part driven by variations in ocean circulation and other physical drivers). (see Box 1)

(†) scaled using $\beta = 1.39$ Pg C/ppm CO₂ and the change in atmospheric pCO₂ indicated in the first line.

(§) Non-steady component is only due to SST variability (warming).

(#) Adjusted for the steady-state outgassing of river derived CO₂.

Box 1 | Key concepts in ocean carbon sink investigations

[bH1] Natural versus anthropogenic CO₂

A key concept aiding the interpretation of the ocean carbon sink has been the separation of the air-sea CO₂ fluxes and the changes in the ocean interior storage of DIC into natural and anthropogenic CO₂ components³⁸. The natural CO₂ component (C_{nat}) is the part of the ocean's DIC pool that existed in pre-industrial times. This pool is involved in many processes, namely air-sea gas exchange, uptake and release by the biological pumps, interactions with and loss to the sediments, and input by rivers (Box Figure 1a). The anthropogenic component (C_{ant}) represents the perturbation to the DIC pool, driven by the anthropogenically-driven increases in atmospheric CO₂. It is substantially smaller than the natural DIC pool (Box Figure 1b).

An important assumption that has simplified analysis is that the anthropogenic CO₂ component does not interact with the natural CO₂ component³⁸. Therefore, the only processes of importance for anthropogenic CO₂ are the uptake from the atmosphere via air-sea gas exchange and the subsequent transport to depth (Box Figure 1a). The assumption about the lack of interaction between the two pools is generally well met, but there are some exceptions. For example, the acidification induced by the oceanic accumulation of anthropogenic CO₂ can affect ocean biology¹⁵⁶ and also has been shown to modify the flux of natural CO₂^{157,158}.

[bH1] The steady state ocean

The second key concept is steady-state, which is reached if climate forcing remains constant for long enough for ocean circulation and ocean biology to become unchanging with time. In this situation, natural CO₂ fluxes across the air-sea interface balance to zero on a global scale¹⁰⁴, with the exception of steady-state outgassing of river-derived CO₂¹⁰⁵. Biological fluxes are also balanced over the annual cycle. The only variations in time come from the steady-state uptake of anthropogenic CO₂ (Box Figure 1c, e). If climate is permitted to vary, leading to a non-steady-state situation, both natural and anthropogenic CO₂ components are affected, leading to additional fluxes and changes in storage (Box Figure 1 d,f). The non-steady-state component of natural CO₂ emerges from a situation where climate is varying, but where atmospheric CO₂ is kept at its preindustrial level. The difference between this situation with one where atmospheric CO₂ is permitted to increase gives the non-steady-state component of anthropogenic CO₂ (Box Figure 1, c-f).


977 **Figures**

Fig 1. Ocean uptake and storage of anthropogenic CO_2 . **a** | Temporal progression of the total ocean inventory of anthropogenic CO_2 as a function of the atmospheric CO_2 content. Results are from an ocean inverse model³ (black line and grey shaded band indicating uncertainty) spanning the period from 1765 until 2010, the ocean biogeochemical model results used in the Global Carbon Budget²⁵ (blue line for the mean and blue shaded band representing the standard deviation), and two observation based estimates of the ocean interior accumulation of anthropogenic CO_2 ^{2,4} for 1994 and 2007. The inset shows the time history of atmospheric CO_2 and the ocean CO_2 uptake³. The bands represent the cumulative uncertainty from the start of the respective estimate. The nearly linear scaling of the ocean uptake with the atmospheric CO_2 content is

particularly evident after 1959. The ocean biogeochemical model results shown here include also the non-steady-state, component of natural CO₂ (climate variability). **b** | Column inventory of anthropogenic CO₂ in mol m⁻² for the year 1994 estimated using the ΔC* back-calculation method ². Strong regional patterning of the accumulation of anthropogenic CO₂ in the ocean was driven by regional differences in ocean circulation and mixing. **c** | Change in the water column inventory between 1994 and 2007 estimated by the eMLR(C*) method ⁴. In b and c, the hatching indicates regions that were not mapped.




Fig 2. **Climatological-mean sea-to-air CO₂ flux.** Multi-product mean estimate based on the six pCO₂-based estimates contained in the SeaFlux product¹⁰³. The mean for the period 1990 through 2020 is depicted, representing the sum of natural, anthropogenic, and steady-state riverine flux components for the nominal year of 2005. The global ocean is characterized by regions of strong sources and sinks of CO₂, reflecting primarily the exchange of natural CO₂ across the air-sea interface. This flux is regionally modified by the uptake flux of anthropogenic CO₂. The latter integrates globally to an uptake flux of about 2.6 Pg C yr⁻¹ for this nominal year. The total CO₂ flux also includes an outgassing flux of about 0.65±0.30 Pg C yr⁻¹, reflecting the steady-state outgassing of natural CO₂ associated with the imbalance between river input and burial¹⁰⁵.




Fig 3. Temporal evolution of the global ocean CO₂ sink **a** | Global ocean CO₂ sink estimated by the 6 pCO₂ observation based products contained in SeaFlux¹⁰³. The estimated net sea-to-air fluxes were adjusted by the steady-state river outgassing flux of 0.65 Pg C yr⁻¹¹⁰⁵ to obtain the ocean CO₂ sink flux that is of relevance for balancing the global sources and sinks of CO₂ (the natural flux, F_{nat}, plus the anthropogenic flux, F_{ant}). The solid black line indicates the mean estimate with the dark grey area representing the standard error across the 6 products. The dashed grey lines indicate the uncertainty of the ocean sink and include the uncertainty of ±0.30 Pg C yr⁻¹ associated with the river outgassing flux¹⁰⁵. The dashed red line represents the steady-state uptake of anthropogenic CO₂ estimated from a global ocean model (CESM-ETHZ²⁵). The dashed orange line represents the expected steady-state uptake of anthropogenic CO₂ estimated from the sensitivity β (left axis). **b** | Contribution of individual ocean basins (north of 44°S) to the global ocean CO₂ sink based on the ensemble mean of the SeaFlux products. The grey band represents the steady-state oceanic outgassing of river-derived CO₂. It was not allocated to individual basins. El Niño related variations in the Pacific basin are

1019 represented by arrows, with the grey shading indicating strength (darker arrows for stronger events).The
1020 global ocean carbon sink varies substantially in time around the long-term trend given by the steady-state
1021 uptake of anthropogenic CO₂ with a period of stagnant uptake in the 1990s followed by a period of faster than
1022 expected growth of the ocean carbon sink after the turn of the millennium.
1023




Fig 4. Zonally integrated anomalous CO₂ fluxes and its components. **a** | Hovmöller diagram of the annual mean, zonal mean anomalies of the total air-sea CO₂ fluxes as a function of time and latitude (right panels) together with the zonal mean (left panels). The anomalies were computed by subtracting the long-term mean flux from the annual mean flux for a given year using the ensemble mean data from the SeaFlux product¹⁰³. The ribbon in the left panels shows the range of the integrated fluxes relative to the zonal mean. The zonal mean dominates compared to the interannual variability. **b** | The same as a, but for the anomalous air-sea fluxes of the steady-state component of anthropogenic CO₂. This estimate was obtained by scaling the ocean inversion-based estimate⁷² for 1995 with a β of $1.4 \text{ Pg C (ppm CO}_2\text{)}^{-1}$. The anomalies were then obtained by subtracting the long-term mean flux. **c** | The same as a, but for the anomalous air-sea fluxes of the non-steady-state component of CO₂, obtained by subtracting b from a. There is strong interannual variability of the air-sea fluxes in the tropics, largely associated with El Niño/Southern Oscillation (ENSO) dynamics as

1037 indicated by the timeseries of the multivariate ENSO index (MEI)¹¹⁵ in panel c, and the strong decadal
1038 variations in the Southern Ocean, largely driven by the non-steady state components.
1039




Fig 5. **Interannual to decadal variability in the ocean carbon sink.** The global ocean sources and sinks of CO₂ are shown along the surface ocean. The ocean interior distribution of the gas-exchange component of natural CO₂^{104,159} (colors) and of the total amount of anthropogenic CO₂ for 2007^{2,4} (isolines) are shown along the depth profile. The hotspots for interannual and decadal variability are noted by the insets. Gradients in the gas-exchange component of natural CO₂ reflects the addition or removal of natural CO₂ through air-sea exchange at the surface. The turquoise arrows indicate the sea-to-air fluxes of natural CO₂ including the type and direction of variability (hat: decadal variability, waves: interannual variability). The reddish arrows indicate the oceanic uptake of anthropogenic CO₂ which is increasing everywhere (straight line). Not shown as arrows is the outgassing flux of the river-derived natural CO₂. The icons relate the variations in the dominant regions of variability (tropical Pacific, and the higher latitudes) to the underlying processes, such as El Niño-Southern Oscillation (ENSO) in the tropical Pacific, and the high latitude modes of variability, especially the Southern Annular Mode (SAM) and the Northern Annular Mode (NAM). Changes in atmospheric CO₂ growth rates affect the global uptake of anthropogenic CO₂.

Glossary

AIR-SEA GAS EXCHANGE

A diffusion-driven process governing the transfer of gases across the air-sea interface, driven by the concentration gradient of the gas across the interface and controlled by the level of turbulence at the interface.

BUFFER FACTOR (REVELLE FACTOR)

The ocean's buffer factor describes how well seawater is able to buffer an increase in surface ocean CO₂ (pCO₂) and is thus crucial for determining the amount of anthropogenic CO₂ the surface ocean can hold.

CARBON, DISSOLVED INORGANIC (DIC)

Dissolved inorganic carbon (DIC) is the sum of all dissolved inorganic carbon species in the seawater, and includes dissolved CO₂ (CO₂^{aq}), carbonic acid (H₂CO₃), bicarbonate (HCO₃⁻), and carbonate (CO₃²⁻).

CO₂, OCEANIC PARTIAL PRESSURE OF (pCO₂)

The oceanic partial pressure of CO₂, pCO₂^{oc} or often just pCO₂, is the partial pressure of CO₂ measured in the air in equilibrium with the water parcel under consideration at one atmosphere total pressure and at the in-situ temperature of the water parcel.

EL NIÑO - SOUTHERN OSCILLATION (ENSO)

The El Niño-Southern Oscillation is a quasi-periodic oscillation of the coupled ocean-atmosphere system with the majority of the action being focused on the eastern tropical Pacific; it is globally the dominant mode of climate variability.

INVERSE MODELS

Inverse models describe a class of models that fuse observations and models in order to improve our quantitative understanding of a set of processes.

FORCING, INTERNAL AND EXTERNAL

Processes leading to changes in the ocean carbon sink: Internal forcing is primarily associated with (internally generated) weather and climate variations, while external forcing is driven by processes external to the climate system, such as volcanic eruptions.

FORWARD MODELS

Forward models, such as those used for the Global Carbon Budget, are a class models that start from initial conditions and solve the governing balance equations by time-integrating them forward using a set of provided boundary conditions.

OCEAN ACIDIFICATION

Change in the ocean's seawater chemistry (pH, [CO₃²⁻], CaCO₃ saturation state, etc) as a consequence of the oceanic uptake of anthropogenic CO₂.

1102

1103 OCEAN BIOGEOCHEMICAL MODELS

1104 Ocean biogeochemical models are a class of ocean models where the most important biogeochemical
1105 processes are explicitly represented, namely air-sea gas exchange, chemical speciation, and biological
1106 processes.