

REVIEW

[View Article Online](#)
[View Journal](#) | [View Issue](#)

Cite this: *Mol. Omics*, 2021,
17, 894

Received 25th June 2021,
Accepted 20th September 2021

DOI: 10.1039/d1mo00196e

rsc.li/molomics

The state of the art in plant lipidomics

Cheka Kehelpannala,^a *^a Thusitha Rupasinghe,^b Thomas Hennessy,^c
David Bradley,^c Berit Ebert^a and Ute Roessner^a

Lipids are a group of compounds with diverse structures that perform several important functions in plants. To unravel and better understand their *in vivo* functions, plant biologists have been using various lipidomic technologies including liquid-chromatography (LC)–mass spectrometry (MS). However, there are still significant challenges in LC-MS based plant lipidomics, which need to be addressed. In this review, we provide an overview of the key developments in LC-MS based lipidomic approaches to detect and identify plant lipids with emphasis on areas that can be further improved. Given that the cellular lipidome is estimated to contain hundreds of thousands of lipids,^{1,2} many of the lipid structures remain to be discovered. Furthermore, the plant lipidome is considered to be significantly more complex compared to that of mammals. Recent technical developments in mass spectrometry have made the detection of novel lipids possible; hence, approaches that can be used for plant lipid discovery are also discussed.

Introduction

Lipids are important components of plants; they are part of cell structures,³ act as energy reserves,⁴ participate in cell signaling,⁵ mitigate stress tolerance⁶ and play a role in both symbiotic and pathogenic interactions.⁷ The entire lipid profile of an organism, a tissue or a cell, is known as the “lipidome”,⁸ and the extensive study of lipid molecules including identification, quantification and elucidation of their role in biological systems is called “lipidomics”.^{8,9} Lipids are highly diverse both in structure and in composition.^{8,10} Structural diversity arises from the differences in the chemical structures of lipids¹⁰ while the ratio of different lipids in cells, tissues, organelles, membrane leaflets, membrane sub-domains or within an organism is known as compositional diversity.¹⁰

For many years, scientists have been studying lipids and their functions in biological systems. It has been shown that even small changes in the structure and composition of lipids can drastically affect vital biological functions.¹⁰ For example, changes in membrane lipids directly affect the function of membrane proteins and the physical properties of the cell membrane including fluidity and permeability.¹¹ Therefore, to correctly interpret the *in vivo* functions of lipids they must be studied as part of an integrated system with metabolites and

enzymes.¹² It is also important to determine not only the type of lipid present but also its time-dependent local concentration.¹²

One of the major challenges in uncovering the function of lipids at the cellular level is the existence of thousands of structurally diverse lipids, some of which may be present at trace level concentrations, thus limiting their detection and identification.¹² Consequently, a robust analysis workflow is necessary from sampling to extraction and subsequently to detection and identification of lipids. In this review, we discuss the key developments in liquid chromatography–mass spectrometry-based approaches for the extraction, detection, identification, quantification and discovery of plant lipids (Fig. 1). We conducted a comprehensive and in-depth survey of over 400 journal articles published on lipidomics from 1957 to 2021 to identify studies that have played pivotal roles in advancing plant lipidomics. Our survey indicated that many of the techniques in plant lipid analysis were adapted from lipidomic approaches developed for mammalian lipid analysis, which we have discussed in this review. We have also discussed the recently developed LC-MS based lipidomic approaches to analyse lipids in animal tissues, which can be adapted for plant lipid analysis in the future. The primary goal of this review is to provide an overview of the evolution of plant lipidomics, its current state and potential future developments with special focus on LC-MS based workflows.

Plant lipids

Lipids are classified into different groups by the LIPID MAPS (LIPID Metabolites And Pathways Strategy; <https://www.lipid>

^a School of BioSciences, The University of Melbourne, Melbourne, VIC 3010, Australia. E-mail: ckehelpannala@gmail.com; Tel: +61 451821235

^b Sciex, 2 Gilda Ct, Mulgrave, VIC 3170, Australia

^c Agilent Technologies Australia Pty Ltd, 679 Springvale Road, Mulgrave, VIC 3170, Australia






Fig. 1 The workflow applied in liquid chromatography–mass spectrometry-based lipidomics.

maps.org/) classification system based on the distinct hydrophilic and hydrophobic elements that comprise the lipid.¹³ The eight main groups include fatty acyls, glycerolipids, glycerophospholipids, sphingolipids, saccharolipids, polyketides, sterol

lipids and prenol lipids, which can be distinguished by their chemically functional backbone structures (Table 1 and Fig. 2).¹³ The major plant lipids and their functions are summarized in Table 1.

Sample preparation

Given the highly diverse nature of lipid structures (Fig. 2), a comprehensive analysis of the lipidome requires firstly an efficient sample preparation process, then an effective lipid separation strategy followed by an optimal mass spectrometric detection method.³³

The first step for lipid analysis is the successful extraction of lipids from the tissue of interest, for which a rapid, simple and efficient extraction method is desirable.³⁴ The most widely used sample preparation method in lipidomics is liquid–liquid extraction.³⁵ Most of the liquid–liquid plant lipid extraction methods are derived from extraction protocols developed for animal tissues, such as the Folch³⁶ and the Bligh and Dyer method.³⁷ In addition to these two, several other protocols using diverse mixtures of organic solvents, including hexane,³⁸

Table 1 The major plant lipids and their functions

Lipid class	Core structure	Functions in plants	Ref.
Fatty acyls	Repeating series of methylene groups	Main building blocks of complex lipids. ¹³	Fahy <i>et al.</i> , 2005
Glycerolipids	Glycerol backbone	Function as signal transduction mediators. ¹⁴ Major constituents of the thylakoid membranes essential for photosynthesis. ^{15–17} Important molecules in cell communication, signal transduction and pathogen responses. ¹⁸ Major storage reserves in plants. ¹⁹	Lim <i>et al.</i> , 2017 Dormann and Benning, 2002 Holzl and Dormann, 2007 Hölzl and Dormann, 2019 Cavaco <i>et al.</i> , 2021 Yang and Benning, 2018
Glycerophospholipids	Phosphate group esterified to a glycerol backbone	Major constituents of cell membranes. ²⁰	Reszczyńska and Hanaka, 2020
Sphingolipids	Sphingoid long-chain base backbone	Act as signaling molecules. ²¹ Involved in plant defense mechanisms against pathogens. ¹⁸ Participate in cellular signaling, ²² growth and stress responses. ^{23,24} Involved in cellular trafficking ²⁵ and defense responses against microbial pathogens. ⁷	Welter <i>et al.</i> , 2007 Cavaco <i>et al.</i> , 2021 Ali <i>et al.</i> , 2018 Hou <i>et al.</i> , 2016
Sterol lipids	Fused four ring structure	Structural components of cell membranes. ²⁰ Regulate acyl chain organization whereby the domain structures of cell membranes are reinforced. ^{26,27} Participate in plant stress responses. ²⁸ Involved in resistance mechanisms against pathogens. ⁷ Precursors of steroid hormones in plants. ²⁸	Mamode Cassim <i>et al.</i> , 2021 Reszczyńska and Hanaka, 2020 Schaller, 2004 Dufourc, 2008 Rogowska and Szakiel, 2020 Siebers <i>et al.</i> , 2016 Rogowska and Szakiel, 2020 Baczewska <i>et al.</i> , 2014
Prenol lipids	C ₅ unit	Participate in protective mechanisms supporting the adaptation of plants to environmental stresses. ^{29,30}	Bajda <i>et al.</i> , 2009 Luu <i>et al.</i> , 2017
Saccharolipids	Fatty acids linked directly to a sugar backbone	Involved in plant defense responses against insects and fungal pathogen attacks. ³¹	
Polyketides	Polyketide backbone	Responsible for the antibacterial, antifungal, antiviral and antiparasitic properties of plants. ³²	Han <i>et al.</i> , 2018






Fig. 2 Representative lipid structures. (A) Fatty acid and glycerophospholipids. The fatty acyl chains are shown in brown, glycerol backbone is indicated in blue, phosphate groups are in black and the head group is in orange. (B) Glycerolipids. The glycerol backbone is indicated in blue, fatty acyl chains are in brown and the hexose group is in orange. (C) Sphingolipids. The long-chain base is denoted in purple, fatty acyl chains are in brown and the head group is in orange. (D) Sterol lipids. The sterol group is indicated in pink, fatty acyl chains are in brown and the sugar moiety is in orange. (E) Polyketides, saccharolipids and prenol lipids. The polyketide backbone is shown in black and the five-carbon groups in prenol lipids are indicated in green. The sugar backbone in saccharolipids is shown in magenta, the phosphate group in black and the fatty acyl chains in brown. R indicates a repeating series of methylene groups. Lipid structures are based on the LIPID MAPS structure database and were generated using ChemDraw 20.0 software.

butanol³⁹ and petroleum ether,⁴⁰ and different treatments such as cold⁴¹ and heat⁴² have been developed, compared and evaluated for plant lipid extraction. Studies comparing these methods concluded that the Folch or Bligh and Dyer methods and those derived from them using chloroform, methanol and water as the extraction solvents are the most efficient methods for plant lipid extraction,^{34,42–45} with the Folch method³⁶ referred to as the “gold standard”.⁴⁶ Originally, modifications to the Folch method³⁶ led to the development of the Bligh and Dyer method.³⁷ Later, significant improvements were made to the Bligh and Dyer protocol³⁷ when it was adopted for plant lipid extractions, leading to several new protocols. For example, in the study by de la Roche *et al.*⁴² wheat, *Triticum aestivum* L. cv, seeds were boiled with isopropanol to

inactivate the lipolytic enzymes and extract neutral lipids more efficiently before applying the Bligh and Dyer method.⁴² Ryu and Wang⁴⁷ simplified the protocol published by de la Roche *et al.*⁴² and introduced 0.01% butylated hydroxytoluene (BHT) to the extraction solvent to reduce lipid oxidation.⁴⁷ The Ryu and Wang⁴⁷ method was adapted and modified with additional steps by Welti *et al.*⁴⁸ and since then this method has been extensively used in plant lipidomics. Since the method by Ryu and Wang⁴⁷ was relatively labor-intensive, a shorter high-throughput single-step extraction method with a 24 h extraction period was developed by Vu *et al.*⁴⁹ Additional modifications to this single-step extraction method of Vu *et al.*⁴⁹ resulted in an extraction protocol that uses 30 parts chloroform, 25 parts isopropanol, 41.5 parts methanol, and 3.5 parts water (v/v/v/v)



with 0.01% BHT.³⁴ This protocol developed by Shiva *et al.*³⁴ has been shown to be highly efficient in extracting lipids from *Arabidopsis thaliana* and *Sorghum bicolor* leaf tissues.³⁴ To inactivate the lipolytic enzymes, fresh leaves were harvested directly into isopropanol pre-heated to 75 °C.^{42,50} However, this single step protocol³⁴ does not include steps for tissue homogenization. Tissue homogenization, however, is crucial to obtain a representative sample as it allows the extraction solvent to access lipids even in rigid plant tissues, increases the surface area in contact with the extraction solvent⁵¹ and solubilizes the lipids. While an extraction method should contain minimal sample preparation to achieve high-quality results, homogenization and metabolic quenching are vital steps.⁵² Thus, the efficiency of a lipid extraction method can be further improved through the addition of a mechanical grinding step.⁵³ A recent comparison of four popular plant lipid extraction methods undertaken by us included the single step extraction method published by Shiva *et al.*,³⁴ the multi-step extraction method of Welti *et al.*,⁴⁸ the biphasic extraction method using methyl *tert*-butyl ether (MTBE)⁴⁶ derived from the Folch method³⁶ and a rapid method employing an extraction temperature of 4 °C.⁴¹ This study revealed that overall, the single-step extraction method³⁴ is the most efficient for extracting most of the major lipid classes from *Arabidopsis* tissue. However, due to the diverse relative hydrophobicity of lipid molecules, the development of a single protocol efficient in extracting all lipid species in a plant sample is impossible.⁴⁵ For example, several studies have shown that chloroform/methanol mixtures are not suitable for the extraction of plant sphingolipids such as glycosylinositol phosphorylceramides (GIPCs).^{34,54,55} Due to the highly amphiphilic nature of sphingolipids,⁵⁵ the standard lipid extraction methods are inefficient in solubilizing them.⁵⁴ It is therefore recommended to use a mixture of isopropanol, hexane and water for the extraction of plant sphingolipids^{55,56} as described by Markham *et al.*⁵⁴

Even though chloroform and methanol are considered ideal solvents for lipid extraction, they can have detrimental effects on human health due to their lipo-solubility.⁴⁵ Given the proven carcinogenicity of chloroform, attempts have been made to replace it with methyl-*tert*-butyl ether (MTBE), which extracts lipids into the organic layer while the polar compounds remain in the aqueous layer.⁴⁶ However, this biphasic extraction method is labor-intensive, which limits its applicability in large-scale lipidomic studies. Furthermore, significant technical errors may be introduced during the separation of the organic layer from the aqueous layer, thus ultimately affecting the extraction reproducibility.⁵³ Substituting the chloroform used in the single-step extraction protocol of Shiva *et al.*³⁴ with MTBE and comparing the extraction efficiencies are worth testing to further improve the existing extraction protocols.

Recent studies have focused on developing lipid extraction methods using environmentally friendly “green solvents”, which can be organic, ionic liquids or supercritical fluids, such as carbon dioxide under supercritical conditions.⁴⁵ Supercritical fluids are substances that are not in a distinct gas or liquid phase but can be gradually compressed from low to high density.⁵⁷

By adjusting the density, its properties can be manipulated for the required process.⁵⁷ Organic green solvents are low in toxicity, readily biodegradable, easily recyclable and have high boiling points and low miscibility, while ionic liquids are non-aqueous organic salts, which are liquid at room temperature.^{45,58} The use of green solvents is widely explored for lipid extraction in biomass processing for biofuel production where large amounts of chemicals are routinely used.⁵⁹ Thus, there could be the potential to replace chloroform with one or some of these green solvents to efficiently extract plant lipids. For example, a biphasic system of cyclopentyl methyl ether (CPME):methanol:water was as efficient as the Bligh and Dyer method in extracting triacylglycerols from wet biomass of the oleaginous yeast *Lipomyces starkeyi*,⁶⁰ thus the study concluded that CPME can be used as an alternative to chloroform.⁶⁰ However, a comparative study applying the green solvents 2-methyl tetrahydrofuran (2-MeTHF) and cyclopentyl methyl ether (CPME) revealed that the Bligh and Dyer method was more effective in extracting total lipids from the microalgae *Chlorella pyrenoidosa* for bio-diesel production.⁶¹ Although the potential of green solvents such as isoamyl acetate,⁶² terpenes, ionic liquids,⁵⁸ 2-methyl-tetrahydrofuran,⁶³ supercritical and subcritical fluids^{64,65} for lipid extraction from yeast and microalgae has been evaluated, their applicability for plant lipid extraction is yet to be demonstrated. Future work will require the comparison of the traditional plant lipid extraction methods with the green solvent-based extraction methods to evaluate the applicability of green solvents in routine plant lipidomic analysis.

Even though current plant lipidomic analyses rely on manual extraction of lipids, automated workflows for lipid analysis will be developed in the future. Automation for sample preparation would be beneficial since the quality, reproducibility and efficiency would be enhanced when compared with manual execution.

Liquid chromatography–mass spectrometry (LC-MS) in lipidomics

Electrospray ionization (ESI) mass spectrometers are the predominant instruments used for lipid analysis.⁶⁶ Lipid samples can be analyzed either by directly injecting the extracts into the mass spectrometer or by first separating them through chromatographic methods before injection into the mass spectrometer.¹ In 1994, Han and Gross⁶⁷ first proposed lipidomic analysis using ESI mass spectrometry, when they analyzed human erythrocyte plasma membrane phospholipids using a triple quadrupole with syringe pump sample injection. This method was further extended to characterize and quantify membrane lipids from cells or subcellular structures by Brügger *et al.* 1997,⁶⁸ using a nano-ESI source. This is now widely known as “shotgun lipidomics” and has seen continuous improvement over the past 25 years. The strategy used by Brügger *et al.* 1997⁶⁸ for the high-throughput profiling of complex lipids from unfractionated extracts of animal cell membranes was later adapted by Welti *et al.*⁴⁸ to profile membrane lipids in *Arabidopsis* tissues.^{48,69}



Because of the relative simplicity, high sample throughput and cost-effectiveness of the shot-gun lipidomic methods³⁴ developed by Welte and colleagues over the past 20 years,^{48,49,69–71} they are still widely used in plant lipidomic analysis.^{72,73} However, the application of direct infusion methods is not without drawbacks. They show higher ion suppression effects compared to methods with chromatographic separation, consequently hindering the analysis of low-abundant lipid species.¹ The selectivity of the direct infusion approach is limited by the existence of isobaric precursor and fragment ions.⁷⁴ The predominantly targeted analysis involving these methods makes the detection of unknowns impossible.⁷⁴ Due to these factors, the shot-gun approach was rapidly followed by LC-MS for plant lipidomics.⁷⁵

The LC-MS based methods were highly sensitive with reduced ion suppression and matrix effects,⁷⁶ able to detect novel lipids,⁷⁷ able to yield reliable identities of lipids even at low levels and able to separate isomers and isobaric compounds.⁷⁸ LC-MS is now considered a comprehensive and powerful technique for the analysis of complex lipid classes such as glycerolipids, glycerophospholipids, and glycolipids and is successfully used in plant lipid research.⁷⁹ To maximize the performance of the LC-MS systems, both the LC and MS methods must be meticulously optimized.³³ An ideal LC method should possess high peak capacity, good chromatographic separation and resolution, reproducibility, selectivity and sensitivity in addition to being compatible with the data acquisition cycles in mass spectrometers.³³ In general, lipidomic methods take long chromatographic analysis times to separate the large number of lipids, which are present over a wide concentration range in biological samples, and to separate isomeric and isobaric compounds.⁸⁰ Quicker chromatographic methods can be used; however, several limitations need to be addressed such as analyte overlap, isotopic interferences and ion suppression effects.⁸⁰ To mitigate these effects, ion mobility can be introduced into the analytical workflow. For example, a separation method employing an elution time of 3.7 min per sample has resulted in a reduced number of features compared to longer chromatographic methods.⁸⁰ The introduction of ion mobility to the analytical workflow, increased the peak capacity and helped to resolve co-eluting species, thereby increasing the number of detected features.⁸⁰

The LC technologies used in modern lipidomics include reversed-phase chromatography (RPLC), normal phase chromatography (NPLC), hydrophilic interaction liquid chromatography (HILIC) and supercritical fluid chromatography (SFC).⁷⁶ RPLC is one of the most often used techniques utilizing a non-polar stationary phase and a polar mobile phase where lipid molecules are separated based on their hydrophobicity, length of fatty acyl chain, and number and position of double bonds.⁷⁶ In general, the retention time of lipids increases with increasing carbon number and decreases with increasing number of double bonds. For example, PC 34:6 elutes before PC 36:6 while PC 34:5 elutes after PC 34:6 (Fig. 3). Non-polar lipids, such as TGs are longer retained in the column while polar lipids such as lysophospholipids elute faster.

The stationary phase of RPLC consists of silica microparticles containing hydrophobic functional groups such as C₃₀, C₁₈, C₈, C₄ hydrocarbon chains or ligands, which may be cyano, phenyl or amino derivatives with C₁₈ being the most widely used.⁷⁶ The polar mobile phase generally contains water and a water-miscible organic solvent such as acetonitrile, methanol and isopropanol.⁷⁶ The development of ultrahigh performance liquid chromatography (UHPLC) columns and core-shell materials for columns significantly improved traditional HPLC techniques providing higher resolution and requiring less time.⁸¹

NPLC separates lipid classes based on polarity and consists of a polar stationary phase and a non-polar mobile phase such as hexane or heptane with a miscible organic solvent like chloroform, isopropanol or ethylacetate.⁷⁶ NPLC can only separate non-polar lipids like DGs and TGs while phospholipids cannot be separated using this method.⁷⁶ In addition, NPLC methods are lengthy and the highly non-polar organic solvent systems used in NPLC are incompatible with ESI-MS.⁸² HILIC is a more efficient type of NPLC technique that uses less apolar solvent systems compatible with ESI-MS and separates lipid classes based on their head group polarity.⁸² The polar stationary phase consists of silica or functionalized silica while the mobile phase consists of water and water-miscible organic solvents.⁷⁶ A comprehensive description of the theories behind different chromatographic approaches has been recently published by Lange *et al.* 2019.⁷⁶ HILIC and RPLC are currently the




Fig. 3 The elution order of phosphatidylcholine (PC) 34, 36 and 38 carbon lipid species in reverse-phase liquid chromatography (RPLC).



two most widely applied chromatographic separation techniques in LC-MS based lipidomics including plant lipidomics.^{78,83} A direct comparison of HILIC-MS and RPLC-MS based lipidomic workflows concluded that both can be equally used to accurately quantify LPCs, LPEs, PCs, PEs and sphingomyelins in human blood plasma;⁸³ however, no such comparison has been reported for lipids from plant tissues. An evaluation of HILIC-MS and RPLC-MS based workflows for plant lipid analysis will be of great benefit for future studies on plant lipids.

In addition to traditional one dimensional LC methods, recently two dimensional LC (2DLC) methods have also been investigated for high-throughput lipidomic analysis of various samples including human plasma⁸⁴ and plant tissues.⁸⁵ In LC, only one liquid phase separation system is applied to the analytes in a sample while 2DLC techniques utilize two orthogonal liquid phase separation systems to separate sample components.⁸⁶ A study comparing several LC and 2DLC methods coupled with high-resolution time-of-flight mass spectrometry (ToF-MS) revealed that the C18xHILIC approach was the most effective in comprehensive lipid profiling of extracts of zebrafish embryos.⁸⁷ However, the data generated by 2DLC coupled with mass spectrometry are highly complex and make an untargeted analysis challenging,⁸⁵ consequently, advanced chemometric data analysis has been proposed for such studies.⁸⁵ For example, a 2DLC setup combining RPLC and HILIC coupled to a triple quadrupole (QQQ) mass spectrometer with chemometric data analysis was able to evaluate the effects of arsenic exposure on the rice lipidome, resolving a larger number of lipids compared to the LC system.⁸⁵ However, the identification of lipids was difficult due to poor detection sensitivity, likely caused by the dilution due to two successive chromatographic separations or because it was impossible to optimize the collision energy.⁸⁵ While the application of 2DLC systems in routine lipidomic analysis may be challenging,³ where simple, rapid, user-friendly and proven lipidomic data acquisition strategies are required, future developments particularly in data processing workflows may improve its applicability. It is worthwhile to also investigate if the 2DLC approach can be used to discover novel lipids.

The separation of lipids by LC is followed by transfer to a mass spectrometer. Mass spectrometers have been used in the analysis of lipids since the 1980s⁸⁸ and over the past decades have seen substantial improvements in the instrumentation and data acquisition methodologies. Mass spectrometric analysis of complex biological samples can be performed using untargeted or targeted approaches.⁸⁹ Untargeted methods aim to comprehensively profile all detectable compounds in a sample while targeted methods focus on analyzing a pre-defined set of metabolites.⁸⁹ In general, targeted methods are more sensitive and more quantitative, however, only a limited number of compounds can be analyzed.⁸⁹ On the other hand, untargeted methods can provide an unbiased analysis of the changes in a system and allow the discovery of novel lipids.⁹⁰ While targeted methods require prior knowledge of sample matrix and LC-MS properties of the analytes, untargeted methods do not rely on prior information about the sample matrices and

are ideal for discovery-focused lipidomic approaches.⁸³ However, due to the complexity of the data retrieved using untargeted methods and the unspecific nature of the analysis, some analytes may go undetected. Furthermore, targeted MS conditions can specifically be tuned to detect individual compounds, for example through the selection of optimized collision energies, while this option is rather limited for the discovery approach.

The commonly used mass spectrometers in lipidomics are triple quadrupoles (QQQ), quadrupole ion trap (QIT), triple quadrupole-linear ion trap (QTRAP), orbitraps and quadrupole time-of-flight (Q-ToF) instruments.^{35,91} Instruments such as Q-ToF and Orbitrap provide high mass accuracy at MS and MS/MS levels and offer both accurate identification and quantification of lipids, although at a lower sensitivity.^{89,90} In contrast, QQQ mass spectrometers show low mass accuracy but offer high sensitivity when operated in the multiple reaction monitoring (MRM) mode.^{90,91} This makes them the preferred instruments for quantitative targeted studies including the analysis of low-abundant lipid species.^{90,91} However, parallel reaction monitoring also known as MRM high-resolution (MRM^{HR}) performed on a high-resolution instrument such as Q-ToF is a more comprehensive, targeted quantitative strategy where all detectable product ions of a pre-selected precursor ion are scanned in the high-resolution mode.⁹² In parallel reaction monitoring a scheduled algorithm is used, in which a targeted/predefined set of precursor ions are isolated in the quadrupole followed by fragmentation and subsequent detection of all the product ions.⁹² This yields more accurate *m/z* and narrower peak widths, and simultaneously detects up to 100 precursor ions per duty cycle; the intensities of multiple fragments can be summed for better sensitivity.⁹² Untargeted LC-MS based lipidomic analyses were found to be independent of the type of high-resolution mass spectrometer used and led to nearly identical results when lipidomic data acquisitions across nine different platforms (one single ToF, one Q/orbital ion trap, and seven Q-ToF instruments) were compared.⁹³ This suggests that the up- and down-regulation patterns of lipids in biological samples can be studied using any type of mass spectrometer.³⁵

MS-based lipidomic analyses utilizing high-resolution mass spectrometers can apply data-dependent acquisition (DDA) or data-independent acquisition (DIA) strategies.⁸⁰ With DDA methods, full scan spectra are recorded at the MS level and MS/MS spectra are recorded by automatically fragmenting the precursor ions with an intensity above a certain threshold or as specified by a user-defined precursor list,³⁵ a given mass range or by mass defect filters.⁹⁴ Since the precursor ions for fragmentation are selected based on the MS survey scan, the DDA approach is also known as information-dependent acquisition (IDA),⁹⁴ data directed acquisition or data-directed analysis.⁹⁵ Generally, the most abundant 10–20 precursor ions are selected for fragmentation using the DDA approach,⁹⁴ which is also known as the top-N data-dependent MS/MS approach⁹⁶ with N referring to the number of the most abundant precursor ions used for fragmentation. Therefore, MS/MS spectra of low-intensity ions will be excluded³⁵ and the reproducibility of the data is low.⁹⁴ To increase the coverage of low abundant



lipids using the DDA approach, an iterative exclusion is proposed where precursors previously selected for fragmentation are excluded in subsequent injections.^{97,98} The application of the iterative exclusion to DDA workflows has shown to increase lipid identification by over 50% in some cases.⁹⁸

By contrast, the DIA methods collect spectral data of all fragments without the selection of specific precursor ions, yielding highly complicated spectra.⁹⁴ DIA methods include collision energy switching MS of everything (MS^E) also known as MS^{All} or all ion fragmentation (AIF)^{35,99} and sequential window acquisition of all theoretical fragment ion spectra (SWATH).¹⁰⁰ In the MS^{All} approach, the data-dependent acquisition is performed on all product ions regardless of the precursor ion. As all precursor ions are sent to the collision cell for fragmentation, MS^{All} relies on good chromatographic separation techniques, such as ultrahigh-performance liquid chromatography (UHPLC) to obtain quality MS/MS spectra.⁹⁵

To reduce the complexity of the spectra generated using the MS^{All} approach, SWATH was introduced by Gillet *et al.*¹⁰⁰ for proteomic experiments and was quickly adapted for metabolomic and lipidomic studies.⁹⁴ However, SWATH still provided complex spectral data and special software tools were necessary to deconvolute the data.³⁵ This issue was solved when an open-source data processing software called mass spectrometry-data independent analysis (MS-DIAL) software¹⁰¹ was introduced to deconvolute the data accumulated by SWATH, which incorporated the LipidBlast¹⁰² library to identify lipids.¹⁰¹ At present, the SWATH acquisition method is immensely popular for untargeted lipidomic analyses¹⁰³ and has been used to analyze lipids in human platelets,⁹⁴ mouse plasma,¹⁰⁴ and human plasma.¹⁰⁵ More recently it was adapted for plant lipidomics to analyse lipids from diverse *Arabidopsis* tissues.¹⁰⁶

SWATH is considered to be advantageous over the DDA methods as it offers full coverage of all MS/MS fragments, yields more lipid identifications and shows high sensitivity, which is comparable to MRM reactions on a QQQ.⁹⁹ Furthermore, the SWATH approach extracts chromatograms not only at the MS level but also at the MS/MS level whereby reliable quantification can be achieved at either the ToF-MS or SWATH-MS/MS levels using precursor or product ions, respectively.¹⁰⁴ A comparison of the IDA, SWATH and MS^{All} approaches revealed that the IDA methods yield qualitatively better MS/MS spectra but a lower number of lipid annotations, while SWATH outperformed the MS^{All} approach with better quality MS/MS spectra and identical MS/MS acquisition hit rates.⁹⁵

Here, it should be noted that the method of choice for lipidomics depends on the purpose of the study⁹⁵ and available resources (Table 2). The IDA approaches such as SWATH and MS^{All} are ideal for untargeted lipidomic approaches.^{99,106,107} However, while SWATH can be used for quantitative analyses,⁹⁹ lipid identification and quantification may be far more simple with targeted MRM methods where quantification of a targeted set of lipids is the primary focus.

Lipid identification

High-throughput untargeted data acquisition strategies on high-resolution mass spectrometers generate a large amount of spectral data,¹⁰⁸ which makes the accurate annotation of the mass spectrometric features challenging. Since the launch of the LipidBlast library in 2013,¹⁰² over 25 commercial and open-source software tools have been developed to process data and identify lipids.¹⁰⁸ However, there are currently no lipid libraries developed specifically for the identification of plant lipids. A customized library for plant lipid identification will be of great value for plant lipid researchers.

Most of the developed software tools use *in silico* lipid libraries, which can be rapidly generated by computing all possible combinations of fatty acids, head groups, linkages and backbones with fragments predicted using a simple set of rules.¹⁰⁸ Commercial software tools include for example SimLipid (PREMIER Biosoft), Lipid SearchTM (Thermo Fisher Scientific),¹⁰⁹ Lipid Annotator software⁹⁷ (Agilent Technologies), and LipidzyzerTM platform¹¹⁰ (Sciex) while freely available open-access tools include lipid data analyzer 2 (LDA2),¹¹¹ LipidBlast,¹⁰² Competitive Fragmentation Modeling-ID (CFM-ID),¹¹² LIQUID,¹¹³ MS-DIAL,¹⁰¹ Greazy,¹¹⁴ lipidr,¹¹⁵ LipiDex,¹¹⁶ LipidMS¹¹⁷ and LipidMatch Flow.¹¹⁸ From these tools, MS-DIAL is the most commonly used open-source software, which is compatible with many LC-MS workflows.¹⁰⁸ The acquisition modes, data processing pipelines and features of the above-mentioned LC-MS based lipidomic software tools have recently been reviewed and compared by Züllig and Köfeler, 2020³⁵ and Züllig *et al.*, 2020.¹¹⁹ Despite all these software solutions, unambiguous structure identification is still a challenging task, mostly due to the over annotation of lipid identities that cause false-positive results and demand manual correction of the lipid annotations.^{35,120} This problem may be circumvented by developing a lipid library where the lipid identities have been manually curated and validated at least for a model organism such as *Arabidopsis*. Recently, we have compiled a manually curated database containing over 800 lipids in *Arabidopsis* tissues with information on the precursor *m/z* values, adducts, product ions, characteristic fragmentation patterns and retention times.⁵³ The information therein can be used to confirm lipid identities and develop targeted MS methods for lipid analysis.

Lipid quantification

Qualitative analyses focus on the profiling and comparison of lipids in biological samples while quantitative analyses aim to provide the concentrations of specific lipids in a biological matrix using meticulously developed quantification methods and lipid standards.⁸⁰ Qualitative comparisons between biological samples use a limited number of internal standards or no standards at all; instead, the peak intensities or area of detected lipids are compared.² This approach is generally known as lipid profiling and can provide comprehensive comparisons between





Table 2 Mass spectrometry-based workflows used for plant lipid analysis

Plant tissue	Homogenization method	Lipid extraction method	LC conditions	MS conditions	Lipid identification	Internal standards	Lipids identified	Ref.
<i>Arabidopsis thaliana</i> rosettes, leaves, flowers, flower stalks, siliques, roots, and seeds	None	Extracted with CHCl_3 : H_2O followed by a five-fold extraction with a mixture of CHCl_3 : MeOH (2:1). The solvent was evaporated under nitrogen and resuspended in CHCl_3	N/A	QQQ in positive and negative modes	MRM transitions	di14:0-PC, di24:1-PC, 13:0-LPC, 19:0-LPC, di14:0-PE, di24:1-PE, 14:0-LPE, 18:0-LPE, di14:0-PG, di24:1-PG, 14:0-LPG, 18:0-LPG, di14:0-PA, di20:0 (phytanoyl)-PA, di14:0-PS, di20:0 (phytanoyl)-PS, 16:0-18:0-PL, di18:0-PL, 16:0-18:0-MGDG, di18:0-MGDG, 16:0-18:0-DGDG and di18:0-DGDG	Phospholipids, polar glycerolipids	Welti <i>et al.</i> , 2002, Devaiah <i>et al.</i> , 2006 ^{150,151}
<i>Arabidopsis thaliana</i> leaves	The glass plunger of the DUALL tissue grinder, attached to a Ryobi D45CK power drill	Extracted with IPP: hexane: H_2O (55:20:25). The dried extract was de-esterified using methylamine, dried under nitrogen and dissolved in THF: MeOH : H_2O (2:1:2) containing 0.1% f.a.	SUPELCOSIL ABZpPlus column (150 mm \times 3 mm \times 5 μm) A: THF: MeOH : 5 mM a.f. (3:2:5) + 0.1% f.a. B: THF: MeOH : 5 mM a.f. (7:2:1) + 0.1% f.a.	QTRAP using MRM mode in positive and negative modes	RT of known standards and by comparison with other peaks	GM1, 12-GlcCer, C12-Cer, sphingosine (C17 base), and sphingosine1-phosphate (C17 base)	Sphingolipids	Markham and Jaworski, 2007 ⁵⁵
<i>Arabidopsis thaliana</i> leaves	Retsch mill	Extracted with MeOH : MTBE (1:3) followed by H_2O : MeOH (3:1). Upper organic phase was removed, dried in a speed-vac and resuspended in a buffer	C8 RP column (100 mm \times 2.1 mm \times 1.7 μm) A: 1% 1 M NH_4Ac +0.1% a.a. B: ACN: IPP (7:3) + 1% 1 M NH_4Ac +0.1% a.a.	Orbitrap with full scan and all-ion fragmentation in positive and negative modes	Matching the m/z values with previously reported data	PE 34:0 (17:0, 17:0) and PC 34:0 (17:0, 17:0)	Phospholipids, oxidized lipids, glycerolipids, sphingolipids (less comprehensive compared to other lipid classes)	Hummel <i>et al.</i> , 2011 ¹⁵²
<i>Arabidopsis thaliana</i> rosettes	Powdered using a cryogenic grinding robot	Extracted with a mixture of CHCl_3 : MeOH : H_2O (1:2.5:1), dried down in a speed vac and resuspended in IPP: hexane: H_2O (55:20:25)	C8 RP column (150 mm \times 2.1 mm \times 1.8 μm) A: 1% 1 M NH_4Ac +0.1% a.a. in H_2O B: MeOH : IPP (5:2) + 1% 1 M NH_4Ac , 0.1% a.a.	SYNAPTMS using MS^E in negative ion mode	Targeted search based on m/z values, expected RT and manual inspection of MS/MS data	Not given	Phospholipids, galactolipids	Burgos <i>et al.</i> , 2011 ⁴¹
<i>Arabidopsis thaliana</i> leaves and roots, leaves of rice (<i>Oryza sativa</i>),	Mixer mill	Bligh and Dyer method with slight modifications. Extracted with CHCl_3 : MeOH : H_2O and the lower layer evaporated by centrifugal concentration and reconstituted in CHCl_3	HILIC silica column (100 mm \times 2.1 mm \times 3 μm) A: MeOH : H_2O (95:5) + 0.2% a.f. B: ACN: MeOH : H_2O (95:2:3) + 0.2% a.f.	IT-TOF operated at scan mode in negative ion mode	m/z values and RT of authentic compounds	10:0/10:0 PC	Novel plant lipid, glucuronosyl-diacylglycerol, phospholipids, polar glycerolipids, sterol lipids, sphingolipids	Okazaki <i>et al.</i> , 2013a, Okazaki <i>et al.</i> , 2013b ^{148,153}

Table 2 (continued)

Plant tissue	Homogenization method	Lipid extraction method	LC conditions	MS conditions	Lipid identification	Internal standards	Lipids identified	Ref.
<i>Arabidopsis thaliana</i> leaves	Mixer mill cooled with liquid nitrogen	Extracted with a hot solvent mixture containing 60% IPP	ACQUITY UPLC HSS T3 column (100 mm × 1 mm × 1.8 µm) A: MeOH : 20 mM NH ₄ Ac (3 : 7) + 0.1% a.a. B: THF : MeOH : 20 mM NH ₄ Ac (6 : 3 : 1) + 0.1% a.a.	QTRAP using MRM in positive and negative modes	MRM transitions	Not given	Phospholipids, sphingolipids, sterol lipids, glycerolipids	Tarazona <i>et al.</i> , 2015 ^{7,5}
<i>Arabidopsis thaliana</i> leaves	None	Extracted with CHCl ₃ : MeOH : 300 mM NH ₄ Ac in H ₂ O (30 : 41.5 : 3.5)	N/A	QQQ in positive and negative modes	MRM transitions	A mixture of 22 internal standards	Phospholipids, glycerolipids, acylated galactolipids, sterol lipids	Shiva <i>et al.</i> , 2020 ^{7,2}
<i>Arabidopsis thaliana</i> leaves and roots. Mature new shoots of <i>Camellia sinensis</i> (tea) plants.	Lyophilized samples were homogenized in a Retschmill	Modified Matyash <i>et al.</i> , 2008 method. Extracted with MeOH : MTBE : H ₂ O (1 : 3 : 1) followed by MeOH : water (1 : 3). Upper phase was dried down in a speed vac and resuspended in ACN : IPP (7 : 3)	C8 column (100 mm × 2.1 mm × 1.7 µm) A: H ₂ O + 1% 1 M NH ₄ Ac + 0.1% a.a. B: ACN + IPP (7 : 3) + 1% 1 M NH ₄ Ac + 0.1% a.a.	Q Exactive using full scan and AIF in positive and negative modes	In-house library	None. Content of each lipid species is presented as mean intensity	Phospholipids, galactolipids and sphingolipids	Giavalisco <i>et al.</i> , 2011, Liu <i>et al.</i> , 2017 ^{15,4,15,5}
Barley (<i>Hordeum vulgare</i> L.) roots	Ground to a fine powder using a mortar and pestle in liquid nitrogen	Sphingolipid extraction in plant tissues as described by Markham <i>et al.</i> , 2006. Extracted with IPP : hexane : H ₂ O (60 : 26 : 14), evaporated under liquid nitrogen and resuspended in IPP : MeOH : H ₂ O (4 : 4 : 1)	Poroshell EC-C18 column (100 mm × 2.1 mm × 2.7 µm) A: MeOH : 20 mM NH ₄ Ac (3 : 7) B: IPP : MeOH : 20 mM NH ₄ Ac (6 : 3 : 1)	QToF using SPRM in positive and negative modes. Oxidized lipids were analyzed based on precursor-product ion transitions.	Precursor-product ion transitions	PE(12 : 0/12 : 0) and Cer(d18 : 1/12 : 0)	Oxidized lipids, phospholipids, polar and neutral glycerolipids, sterol lipids, sphingolipids	Yu <i>et al.</i> , 2018, Yu <i>et al.</i> , 2020 ^{9,2,13,6}
<i>Arabidopsis</i> seeds	Not given	Extracted with IPP : hexane : H ₂ O (55 : 20 : 25) according to the protocol by Markham <i>et al.</i> , 2006 followed by alkaline hydrolysis, dried down and reconstituted in BuOH : MeOH (1 : 1)	Ascentis Express RP Amide column (50 mm × 2.1 mm × 2.7 µm) H ₂ O : MeOH : THF (50 : 20 : 30 to 5 : 20 : 75)	QQQ using MRM transitions	MRM transitions	16 : 0-d31 SM	Ceramides, hydroxy ceramides, glucosylceramides and GIPCs	Ebert <i>et al.</i> , 2018 ^{3,6}



Table 2 (continued)

Plant tissue	Homogenization method	Lipid extraction method	LC conditions	MS conditions	Lipid identification	Internal standards	Lipids identified	Ref.
Rice (Aerial parts and roots)	Ground to a fine powder using a mortar and pestle in liquid nitrogen and lyophilized for 24 h to dryness	Modified Matyash <i>et al.</i> 2008 method extraction with pre-cooled MeOH: MTBE (1:3) followed by MeOH: water (1:3), upper-phase dried down and resuspended in MeOH: water (1:0.85)	Column 1: RP ZORBAX Eclipse XDB-C18 column (150 mm × 2.1 mm × 5 µm) A: ACN: IPP (1:2) + 0.1% f.a. B: H ₂ O + 0.1% f.a. Column 2: KINETEX HILIC column (30 mm × 3 mm × 2.6 µm) A: 5 mM NH ₄ Ac B: 5 mM NH ₄ Ac	QQQ in full scan mode followed by MRM transitions in negative and positive modes.	Matching the experimental MS/MS spectra with the reference spectra available in Metlin	17:1 LPC, 17:0 LPC, 1,3-17:0 D5 DG, 17:0 CE, 1,2,3-17:0 TG, 16:0 D31-18:1 PC, 16:0 D31-18:1 PS, N-dodecanoylsphingosine, N-dodecanoylglucosyl-sphingosine and N-dodecanoylsphingosylphosphorylcholine.	Phospholipids, glycerolipids	Navarro-Reig <i>et al.</i> , 2018 ⁸³
Halophytes: <i>Salicornia ramosissima</i> (fresh branch tips) and <i>Halimione portulacoides</i> (leaves)	Ground to a powder	Bligh and Dyer protocol with modifications. Extracted with CHCl ₃ :MeOH (1:2) followed by the addition of water. The lower phase was dried under nitrogen and resuspended in mobile phase B.	Ascentis Si column (150 mm × 1 mm, 3 µm) A: ACN: MeOH: H ₂ O (50:25:25) + 1 mM NH ₄ Ac B: ACN: MeOH (60:40) + 1 mM NH ₄ Ac	Q-Exactive with full scan MS and DDA in positive and negative modes	MS/MS spectral data	None	Phospholipids and glycosphingolipids	Maciel <i>et al.</i> , 2018 ¹⁵⁷
Wheat leaves	Cryo-homogenization (Cryomill)	Folch method with modifications. Extracted with MeOH, CHCl ₃ vacuum dried and resuspended in BuOH: MeOH (1:1)	Ascentis Express RP-Amide column (50 mm × 2.1 mm, 2.7 µm) A: 10 mM a.f. in H ₂ O: MeOH: THF (50:20:30) B: 10 mM a.f. in H ₂ O: MeOH: THF (5:20:75)	QQQ in positive and negative modes	MRM transitions	LPC(17:0), PC(34:1), PE(34:0), PG(34:1), PI(36:2), PS(34:0).	Phospholipids	Cheong <i>et al.</i> , 2019, Cheong <i>et al.</i> , 2020 ^{158,159}
Vascular plants (kale leaves and corn roots)	Cryo-homogenization (Cryomill)	Bligh and Dyer method. Samples were incubated in hot IPP prior to extraction. Extracted with MeOH and CHCl ₃ followed by the addition of H ₂ O. The bottom organic layer was dried under a nitrogen stream and reconstituted in CHCl ₃ : MeOH (1:1)	HILIC column (Luna 100 mm × 2 mm × 3 µm) to screen for any novel lipid classes A: ACN: H ₂ O (97:3) + 10 mM NH ₄ Ac B: 10 mM NH ₄ Ac in pure H ₂ O C30-RPLC (150 mm × 2 × 2.6 µm) to confirm the HILIC results A: ACN: H ₂ O (60:40) + 10 mM a.f. + 0.1% f.a. B: IPP: ACN: H ₂ O (90:10:1) + 10 mM a.f. + 0.1% f.a.	Q-extractive orbitrap using a top-20 DDA in positive and negative modes	MS/MS fragmentation patterns	SQDG 16:0/16:0, MGDG 16:3(7Z,10Z,13Z)/18:3(9Z,12Z,15Z), PA 18:1(9Z)/18:1(9Z), PG 18:0/20:4(5Z,8Z,11Z,14Z), PC 18:0/20:4(5Z,8Z,11Z,14Z), PE 18:0/20:4(5Z,8Z,11Z,14Z), diether PC O-18:0/O-18:0, plasma-logen PC P-18:0/ malogen PE P-18:0/ malogen PE P-18:0/ 20:4(5Z,8Z,11Z,14Z), MMPE 16:0/16:0 and DMPE 16:0/16:0, LPA 20:4(5Z,8Z,11Z,14Z), LPC 18:1(9Z), LPE 18:0, plasma-logen LPE P-18:0; PI 18:0/ 20:4(5Z,8Z,11Z,14Z); DLCL 18:2(9Z,12Z)/18:2(9Z,12Z), and SM d18:1/18:0.	Phospholipids, polar and neutral glycerolipids, phytosphingolipids, glycolipids. Acylphosphatidylglycerols, a modified class of acylphosphatidylglycerols and their isomeric forms in corn roots. Regioisomers of lysophospholipids in kale leaves	Pham <i>et al.</i> , 2019 ⁹⁶



Table 2 (continued)

Plant tissue	Homogenization method	Lipid extraction method	LC conditions	MS conditions	Lipid identification	Internal standards	Lipids identified	Ref.
Root, stem, leaf, and petiole tissues of Rapeseed	Intact plant tissues were used without homogenization	Plant tissue harvested directly into hot IPP and extracted with CHCl ₃ :H ₂ O as described by Welti <i>et al.</i> , 2002	Acquity UPLCTM BEH C18 column (100 mm × 2.1 mm × 2 µm) A: H ₂ O: MeOH: ACN: 300 mM NH ₄ Ac (20:20:20:1) B: IPP: MeOH: 300 mM NH ₄ Ac (180:20:3)	TripleTOF using DDA and MS/MS ^{ALL} scan in positive and negative modes	RT, accurate <i>m/z</i> , and fragmentation ion patterns	PC-12: 0/12:0, PA-14: 0/14:0, PE-12: 0/12:0, PG-20: 0/20:0, PI-16: 0/18:0, PS-20: 0/20:0, LPC-19:0, LPG-18:0, LPE-18:0, MGDG18: 0/18:0, DGDG-18: 0/18:0. TAG-17: 0/17:0 DAG-17: 0/17:0 C16 Lactosyl(β) Ceramide (d18:1/16:0)	Phospholipids, glycerolipids	Lu <i>et al.</i> , 2019 ¹²⁰
Salad (<i>Lactuca sativa</i> var. <i>capitata nidus tenerimma</i>), deep frozen spinach (<i>Spinacia oleracea</i>), raspberries (<i>Rubus idaeus</i>) and strawberries (<i>Fragaria</i>)	Salad was manually cut into small pieces and homogenized using an ultraturax. Raspberries and strawberries were homogenized using a hand blender	Extracted with IPP, <i>n</i> -hexane and H ₂ O according to the protocol by Markham <i>et al.</i> , 2006, followed by alkaline hydrolysis, dried down and reconstituted in IPP: H ₂ O (65:35)	C18 Acquity UHPLC HSS T3 column (150 mm × 2.1 mm × 1.8 µm) A: ACN: H ₂ O (3:2) + 0.1% f.a. + 10 mM a.f. B: IPP: ACN (9:1) + 0.1% f.a. + 10 mM a.f.	Q-Exactive using top-10 DDA in positive and negative modes	Lipid Data Analyzer and manual inspection of data.		GIPC	Panzenboeck <i>et al.</i> , 2020 ¹⁶⁰
Green alga, <i>Arabidopsis</i> tissues, maize, tomato and grey poplar and common beech trees	Bead beater	Extracted with precooled MeOH: MTBE (1:3) solution followed by MeOH: water (1:3) extraction. The upper phase was dried down and resuspended in ACN: IPP (70:30)	C8 column (100 mm × 2.1 mm × 1.7 µm) A: H ₂ O: IPP: ACN (446 g:130 g:300 g)+ 10 ml 1 M NH ₄ Ac+ ml a.a. B: IPP: ACN (236 g:546 g) + 10 ml 1 M NH ₄ Ac + 1 ml a.a.	Q-Exactive with full scan mode in positive and negative modes	Matching the exact <i>m/z</i> and RT to a list of potential compounds	PC 34:0	Phospholipids, glycerolipids	Lapidot-Cohen <i>et al.</i> , 2020 ¹⁶¹
Banana fruit peel	Crushing	Extracted with MeOH followed by MTBE and H ₂ O. The upper phase was dried with nitrogen stream and resuspended in IPP	C18 column (100 mm × 2.1 mm × 1.7 µm) A: ACN: H ₂ O (6:4) + 10 mM a.f. B: ACN: IPP (1:9) + 10 mM a.f.	Q-Exactive using full scan MS in positive and negative modes	LipidSearch	None. Content of each lipid species is presented as mean intensity	Phycerophospholipids, and glycerolipids	Liu <i>et al.</i> , 2020 ¹⁶²
<i>Arabidopsis thaliana</i> leaves, rosettes, internodes, roots, seedlings and siliques	Cryo-homogenization (Cryomill)	Extracted with CHCl ₃ : MeOH: 300 mM NH ₄ Ac in H ₂ O (30:41.5:3.5), dried down in a speed vac and resuspended in BuOH:MeOH (1:1)	InfinityLab Poroshell C18 column (100 mm × 2.1 mm × 2.7 µm) A: ACN:H ₂ O (6:4) + 10 mM a.f. B: IPP:ACN (9:1) + 10 mM a.f.	QToF using SWATH in positive mode	MS-DIAL, <i>m/z</i> values, RT pattern and manual inspection of the MS/MS spectra	None	Phospholipids, glycerolipids, sphingolipids	Kehepunnala <i>et al.</i> , 2020, Kehepunnala <i>et al.</i> , 2021, Schillaci <i>et al.</i> , 2021 ^{53,106,163}

RT: retention time, THF: tetrahydrofuran, a.a: acetic acid, a.f: ammonium formate, f.a: formic acid, NH₄Ac: ammonium acetate IT-TOF: Ion trap-time-of-flight mass spectrometer, QToF: quadrupole time-of-flight mass spectrometer, QQQ: triple quadrupole mass spectrometer, SPRM: scheduled parallel reaction monitoring, MRM: multiple reaction monitoring, DDA: data dependent acquisition, SWATH: sequential window acquisition of all theoretical fragment ion spectra, GIPC: glycosyl inositol phosphoryl ceramide, CHCl₃: chloroform, MeOH: methanol, BuOH: butanol ACN: acetonitrile, IPP: isopropanol, MTBE: methyl-*tert*-butyl-ether.



biological samples when combined with substantive statistical analyses.^{2,121–126}

For the absolute quantification of lipids, ideally, one stable isotope-labelled internal standard per compound should be used to account for the ion suppression effects that can arise from matrix effects and changes in mobile phase composition.³⁵ Absolute quantification will immediately indicate whether a lipid is a major or a minor species and allows the direct comparison of lipidomic results across different laboratories, which increases the confidence in lipidomic data.¹²⁷ However, using a lipid standard for each lipid species is almost impossible, particularly in large-scale lipidomic studies where several thousand lipids are present in the biological sample of interest.⁸³ Therefore, instead of one lipid standard for each lipid species, the “minimum standards” established by the Lipidomics Standards Initiative¹²⁸ require that at least one internal standard per class should be used for lipid quantification. However, it is important to note that even then only a relative quantification can be achieved where the increasing and decreasing patterns of lipids between two samples are compared.¹²⁹ While the use of a single internal standard per lipid class may account for errors in extraction efficiency and systematic errors, concentrations of individual species cannot be established.¹²⁹ This is because the instrument's response for different lipid species even within the same lipid class can differ significantly depending on various factors.^{129,130} For example, the instrument's response for phospholipids is dependent on acyl chain length, acyl chain unsaturation, the structure of the polar head group, total lipid concentration, solvent composition, and instrument settings.¹³⁰ Consequently, multiple internal standards per lipid class distributed across the entire retention time range and acyl chain composition would be required for an accurate quantitative analysis where the concentrations of individual lipids need to be recorded.^{35,129} Hence, the choice of quantification strategy depends on the scope of the study, and it is necessary to carefully consider all the different strategies before undertaking a lipidomic analysis. The principles of quantification, selection of internal standards for different analytical platforms and the factors that affect the accurate quantification of lipids are extensively reviewed by Wang *et al.*²

Standardizing lipidomic methodologies

The lipidomics standards initiative (<https://lipidomics-standards-initiative.org/>) was established in 2018 to provide guidelines and standards for lipidomics.¹²⁸ The first draft of guidelines is now provided, which covers the entire lipidomic workflow, including pre-analytics, lipid extraction, data acquisition, lipid identification, quantification, method validation, quality control and data reporting.¹²⁸ If the guidelines introduced by the Lipidomics Standards Initiative¹²⁸ are followed by the lipid research community, this could enable the cross-comparison of lipidomic data across many different laboratories. For example, we recently developed a comprehensive lipid map of *Arabidopsis* across development; this includes a catalogue of the major lipids

observed in different *Arabidopsis* tissues and provides the initial framework for a plant lipid database.¹⁰⁶ This *Arabidopsis* lipid database has great potential to be expanded, for example, by including profiles of sphingolipids, sterol lipids, oxylipins and prenol lipids detected in diverse *Arabidopsis* tissues such as leaves, stems, flowers and roots, during development. Many laboratories around the world conduct studies comparing the lipid profiles of *Arabidopsis* wild-type and mutants under standard and various stress conditions. If the data generated from all these studies can be collated to build a common platform similar to the *Arabidopsis* lipid map, this would enable an effective sharing and comparison of data and results. Following the minimum standards set by the Lipidomics Standards Initiative¹²⁸ would be a first step towards achieving such a resource that assists the lipid research community.

Furthermore, to enable the efficient cross-comparison of lipidomic data, it is advisable to follow the quality control (QC) guidelines set by the lipidomics standards initiative (<https://lipidomics-standards-initiative.org/>). QC in lipidomic analyses is an important aspect that must be used at all times to ensure data quality. QC samples are used to stabilize the analytical instrument, monitor the instrument performance¹³¹ and account for within and between batch variations in mass spectrometric responses.¹³² The occurrence of systemic variations of instrument's response within and between batches, for example, due to variations in mobile phase composition, system calibration, differences in MS detection sensitivity and resolution, is often encountered in untargeted LC-MS metabolomics.¹³² The qualitative and quantitative tools that can be used to account for batch variations have previously been evaluated and discussed by Sanchez-Illana *et al.*, 2018¹³² while various QC strategies in mass spectrometry-based lipidomics have been reviewed by Xie *et al.*, 2017.¹³¹

Identifying lipid functions

Plant scientists have studied lipids for over 50 years and their initial focus was mainly directed at understanding their biosynthesis.¹³³ The development of *Arabidopsis thaliana* as a genetic plant model, and the sequencing of its genome,¹³⁴ enabled the exploration of many lipid biosynthetic pathways including the synthesis of membrane glycerolipids, sphingolipids, triacylglycerols, cutins, waxes and suberins.^{133,135} With the advent of innovative mass spectrometric technologies in the 1990s,⁷¹ plant biologists started applying mass spectrometry to study lipid metabolic pathways involved in plant development and stress responses,^{71,136} thereby identifying lipids that act as substrates and products of enzymes¹³⁷ and detecting lipid-metabolizing enzymes.²¹ To gain a clear understanding of the *in vivo* functions of lipids in plants, often several different approaches are required including genetic, genomic¹⁹ and biochemical studies. These studies can provide novel insights into the roles of lipids in plants and the information may help in designing engineering strategies in biotechnology.¹⁹ For example, TGs have been long known to be the major energy



reserves in seeds;^{19,138} however, newer studies have shown that they are also involved in cell division and expansion,¹³⁹ stomatal opening,¹⁴⁰ membrane lipid remodelling,¹⁹ organ formation,¹⁹ stress response¹⁴¹ and pollination.¹⁴² This suggests that strategies aiming to increase the amount of TGs in plant tissues for biodiesel and industrial chemical production must be designed carefully,¹⁹ as an overaccumulation of TGs could prove detrimental to the growth of such engineered plants.¹⁹

It has been estimated that the cellular lipidome comprises hundreds of thousands of diverse lipids^{1,143} and the plant cellular sphingolipidome alone may comprise at least 500 and perhaps thousands of different species of sphingolipids.¹⁴⁴ Theoretically, possible lipid structures exceed 180 000 without even considering all the position isomers, backbone substitutions and stereochemistry. Indeed several million potential lipids could exist when all structural differences are taken into account.⁹⁸ This suggests that much of the plant lipidome is yet to be fully described and that there are likely many lipid species to be discovered. However, new species are most often present in low levels in complex biological matrices compared to other more common lipids,¹⁰⁷ which is one of the main reasons why they are yet to be discovered. Due to their low abundance, the precursor ions will not be selected for fragmentation by data-dependent acquisition methods in the LC-MS/MS approaches,¹⁰⁷ which makes it difficult to identify them. In shotgun lipidomics, the MS/MS spectra of novel species are difficult to interpret due to the presence of product ions of co-fragmented lipids and chemical noise.¹⁰⁷ Furthermore, identifying the structures of all unknown lipids in a complex biological sample is challenging since thousands of unidentified mass spectrometric features are detected,⁹² some of which may be artifacts generated from in-source fragmentation¹⁴⁵ while some peaks could also be the result of solvent masses.¹⁴⁶ Multiple adduct formation by the same lipid species and isotopic peaks may further complicate the identification of novel lipids.¹⁴⁶ Together these factors hinder the systematic discovery of novel lipids.

A strategy that can be used to detect novel lipids is to look for new peaks in the mass spectra of the treated samples, which are absent in control samples. Since plant cells often produce lipids in response to environmental stresses and during the developmental process,¹⁴⁷ there is potential to discover new lipid species through a comparison of lipid profiles from treated and control plant tissues. Given that the respective biosynthetic genes are identified first, the functions of newly discovered lipids can be studied in the corresponding mutants. For example, a new class of lipids, glucuronosyldiacylglycerols, was discovered in phosphorus-starved *Arabidopsis* plants, and further studies showed that they are essential for protection against phosphorus depletion.¹⁴⁸ The authors observed a new peak in the mass spectra of lipid extracts from the leaves of phosphorus-starved wild-type plants but not in the extracts of the plants grown with sufficient phosphorus.¹⁴⁸ Further investigations into the new peak by tandem mass spectrometry revealed the building blocks to be a glucuronosylglycerol and two fatty acids, which suggested that the structure is a glucuronosylated diacylglycerol.¹⁴⁸ The novel compound was purified

from *Arabidopsis* leaf material and its structure was established by nuclear magnetic resonance spectroscopy (NMR).¹⁴⁸ Subsequent studies revealed that the glucuronosyldiacylglycerols mitigate phosphorus depletion stress as *sqd2* mutant plants, which were unable to accumulate SQDGs and glucuronosyldiacylglycerol, showed severe growth defects under phosphorus limiting conditions.¹⁴⁸ In another study, mechanical disruption of leaf tissue caused by wounding, pathogen attack and cold stress stimulated the production of an extensive variety of oxidized and acylated galactolipids in *Arabidopsis*.¹⁴⁹ The enzyme responsible for the acylation of the galactolipid head group was purified from oat leaves and subsequently identified using tandem MS. This led to the identification of the *Arabidopsis* ortholog, which was subsequently confirmed through lipid analysis of loss-of-function mutants.¹⁴⁹ These examples highlight that by monitoring the lipid changes of plants subjected to stress conditions and by comparing the lipid profiles of loss-of-function mutants and wild-type plants, novel lipids essential for the survival of plants under challenging environmental conditions can be detected and their specific functions can be uncovered.

Conclusions

LC-MS based technology is now considered as a powerful tool to explore lipid biosynthetic pathways, identify lipid-metabolizing enzymes, and uncover lipid functions.

Plant lipid extraction methods, which were initially adapted from animal lipid extraction methods, have also vastly improved and modified to better suit the requirements for extracting lipids from plant tissue.

Shotgun lipidomics and LC-MS based lipidomics are the two lipidomic approaches widely used to effectively analyse plant lipids. For LC-MS based plant lipidomic workflows, there is a shift towards using high-resolution mass spectrometers such as QToF and orbitrap coupled with RPLC and HILIC. Most high-resolution mass spectrometers now offer high sensitivity with higher resolution that can be used for both targeted and discovery lipidomic approaches. Nevertheless, the unambiguous identification of lipids is still extremely challenging and requires the manual curation of data sets, and yet, no lipid library specifically built for plant lipid identification is available.

The development of high-resolution mass spectrometers and comprehensive data acquisition methods enable the detection of not only known lipid species but also a plethora of unidentified features. It is currently impossible to identify all unknown mass spectrometric features that can be detected and many of them may be due to artefacts. However, recent studies have shown that by considering lipids as part of an integrated system with enzymes and metabolites and in the context of plant function, new lipids essential for the survival of plants can be identified and studied.

Conflicts of interest

There are no conflicts of interest to declare.



Acknowledgements

C. K. is funded through a Research Training Program Scholarship provided by the Australian Commonwealth Government and the University of Melbourne. BE is supported by an Australian Research Council Future Fellowship [FT160100276] and the University of Melbourne Botany Foundation.

References

- 1 R. Harkewicz and E. A. Dennis, *Annu. Rev. Biochem.*, 2011, **80**, 301–325.
- 2 M. Wang, C. Wang and X. Han, *Mass Spectrom. Rev.*, 2017, **36**, 693–714.
- 3 F. Furt, F. Simon-Plas and S. Mongrand, in *The Plant Plasma Membrane*, ed. A. S. Murphy, B. Schulz and W. Peer, Springer Berlin Heidelberg, Berlin, Heidelberg, 2011, DOI: 10.1007/978-3-642-13431-9_1, pp. 3–30.
- 4 A. Cagliari, R. Margis, F. dos Santos Maraschin, A. C. Turchetto-Zolet, G. Loss and M. Margis-Pinheiro, *Int. J. Plant Biology*, 2011, **2**, e10.
- 5 W. van Leeuwen, L. Ökrész, L. Bögre and T. Munnik, *Trends Plant Sci.*, 2004, **9**, 378–384.
- 6 Y. Okazaki and K. Saito, *Plant J.*, 2014, **79**, 584–596.
- 7 M. Siebers, M. Brands, V. Wewer, Y. Duan, G. Hözl and P. Dörmann, *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids*, 2016, **1861**, 1379–1395.
- 8 E. Fahy, D. Cotter, M. Sud and S. Subramaniam, *Biochim. Biophys. Acta*, 2011, **1811**, 637–647.
- 9 S. J. Blanksby and T. W. Mitchell, *Annu. Rev. Anal. Chem.*, 2010, **3**, 433–465.
- 10 T. Harayama and H. Riezman, *Nat. Rev. Mol. Cell Biol.*, 2018, **19**, 281–296.
- 11 Q. Guo, L. Liu and B. J. Barkla, *Int. J. Mol. Sci.*, 2019, **20**, 4264.
- 12 G. van Meer, *EMBO J.*, 2005, **24**, 3159–3165.
- 13 E. Fahy, S. Subramaniam, H. A. Brown, C. K. Glass, A. H. Merrill, Jr., R. C. Murphy, C. R. Raetz, D. W. Russell, Y. Seyama, W. Shaw, T. Shimizu, F. Spener, G. van Meer, M. S. VanNieuwenhze, S. H. White, J. L. Witztum and E. A. Dennis, *J. Lipid Res.*, 2005, **46**, 839–861.
- 14 G. H. Lim, R. Singhal, A. Kachroo and P. Kachroo, *Annu. Rev. Phytopathol.*, 2017, **55**, 505–536.
- 15 P. Dormann and C. Benning, *Trends Plant Sci.*, 2002, **7**, 112–118.
- 16 G. Hözl and P. Dormann, *Prog. Lipid Res.*, 2007, **46**, 225–243.
- 17 G. Hözl and P. Dörmann, *Annu. Rev. Plant Biol.*, 2019, **70**, 51–81.
- 18 A. R. Cavaco, A. R. Matos and A. Figueiredo, *Cell. Mol. Life Sci.*, 2021, **78**, 4399–4415.
- 19 Y. Yang and C. Benning, *Curr. Opin. Biotechnol.*, 2018, **49**, 191–198.
- 20 E. Reszczyńska and A. Hanaka, *Cell Biochem. Biophys.*, 2020, **78**, 401–414.
- 21 R. Welti, J. Shah, W. Li, M. Li, J. Chen, J. J. Burke, M. L. Fauconnier, K. Chapman, M. L. Chye and X. Wang, *Front. Biosci.*, 2007, **12**, 2494–2506.
- 22 U. Ali, H. Li, X. Wang and L. Guo, *Mol. Plant*, 2018, **11**, 1328–1343.
- 23 A. Mamode Cassim, Y. Navon, Y. Gao, M. Decossas, L. Fouillen, A. Grélard, M. Nagano, O. Lambert, D. Bahammou, P. Van Delft, L. Maneta-Peyret, F. Simon-Plas, L. Heux, B. Jean, G. Fragneto, J. C. Mortimer, M. Deleu, L. Lins and S. Mongrand, *J. Biol. Chem.*, 2021, 100602, DOI: 10.1016/j.jbc.2021.100602.
- 24 Q. Hou, G. Ufer and D. Bartels, *Plant, Cell Environ.*, 2016, **39**, 1029–1048.
- 25 P. Kachroo and A. Kachroo, *Mol. Plant*, 2020, **13**, 351–353.
- 26 H. Schaller, *Plant Physiol. Biochem.*, 2004, **42**, 465–476.
- 27 E. J. Dufourc, *Plant Signaling Behav.*, 2008, **3**, 133–134.
- 28 A. Rogowska and A. Szakiel, *Phytochem. Rev.*, 2020, **19**, 1525–1538.
- 29 A. H. Baczewska, W. Dmuchowski, A. Jozwiak, D. Gozdowski, P. Bragoszewska, P. Dabrowski and E. Swiezewska, *Dendrobiology*, 2014, **72**, 177–186.
- 30 A. Bajda, D. Konopka-Postupolska, M. Krzymowska, J. Hennig, K. Skorupinska-Tudek, L. Surmacz, J. Wójcik, Z. Matysiak, T. Chojnacki, E. Skorzyńska-Polit, M. Drazkiewicz, P. Patrzybas, M. Tomaszewska, M. Kania, M. Swist, W. Danikiewicz, W. Piotrowska and E. Swiezewska, *Physiol. Plant.*, 2009, **135**, 351–364.
- 31 V. T. Luu, A. Weinhold, C. Ullah, S. Dressel, M. Schoettner, K. Gase, E. Gaquerel, S. Xu and I. T. Baldwin, *Plant Physiol.*, 2017, **174**, 370–386.
- 32 J. W. Han, G. J. Choi and B. S. Kim, *World J. Microbiol. Biotechnol.*, 2018, **34**, 163.
- 33 A. Criscuolo, M. Zeller, K. Cook, G. Angelidou and M. Fedorova, *Chem. Phys. Lipids*, 2019, **221**, 120–127.
- 34 S. Shiva, R. Enniful, M. R. Roth, P. Tamura, K. Jagadish and R. Welti, *Plant Methods*, 2018, **14**, 14.
- 35 T. Züllig and H. C. Köfeler, *Mass Spectrom. Rev.*, 2020, **n/a**.
- 36 J. Folch, M. Lees and G. H. Sloane Stanley, *J. Biol. Chem.*, 1957, **226**, 497–509.
- 37 E. G. Bligh and W. J. Dyer, *Can. J. Biochem. Physiol.*, 1959, **37**, 911–917.
- 38 D. Shearer and R. Carson, *J. Assoc. Off. Agric. Chem.*, 1958, **41**, 414–416.
- 39 T. A. Macmurray and W. R. Morrison, *J. Sci. Food Agric.*, 1970, **21**, 520–528.
- 40 C. Brown, E. Weber and C. Wilson, *Crop Science*, 1970, **10**, 488–491.
- 41 A. Burgos, J. Szymanski, B. Seiwert, T. Degenkolbe, M. A. Hannah, P. Giavalisco and L. Willmitzer, *Plant J.*, 2011, **66**, 656–668.
- 42 I. A. de la Roche and C. J. Andrews, *Plant Physiol.*, 1973, **51**, 468–473.
- 43 M. J. Fishwick and A. J. Wright, *Phytochemistry*, 1977, **16**, 1507–1510.
- 44 M. Axelsson and F. Gentili, *PLoS One*, 2014, **9**, e89643.



- 45 S. S. de Jesus and R. M. Filho, *Renewable Sustainable Energy Rev.*, 2020, **133**, 110289.
- 46 V. Matyash, G. Liebisch, T. V. Kurzchalia, A. Shevchenko and D. Schwudke, *J. Lipid Res.*, 2008, **49**, 1137–1146.
- 47 S. B. Ryu and X. Wang, *Biochim. Biophys. Acta*, 1998, **1393**, 193–202.
- 48 C. M. Buseman, P. Tamura, A. A. Sparks, E. J. Baughman, S. Maatta, J. Zhao, M. R. Roth, S. W. Esch, J. Shah, T. D. Williams and R. Welti, *Plant Physiol.*, 2006, **142**, 28–39.
- 49 H. S. Vu, S. Shiva, M. R. Roth, P. Tamura, L. Zheng, M. Li, S. Sarowar, S. Honey, D. McElhiney, P. Hinkes, L. Seib, T. D. Williams, G. Gadbury, X. Wang, J. Shah and R. Welti, *Plant J.*, 2014, **80**, 728–743.
- 50 M. Kates, in *Advances in Lipid Research*, ed. R. Paoletti and D. Kritchevsky, Elsevier, 1970, vol. 8, pp. 225–265.
- 51 A. Krastanov, *Biotechnol. Biotechnol. Equip.*, 2010, **24**, 1537–1543.
- 52 P. Yin and G. Xu, *J. Chromatogr. A*, 2014, **1374**, 1–13.
- 53 C. Kehelpannala, T. W. T. Rupasinghe, T. Hennessy, D. Bradley, B. Ebert and U. Roessner, *Plant Methods*, 2020, **16**, 155.
- 54 J. E. Markham, J. Li, E. B. Cahoon and J. G. Jaworski, *J. Biol. Chem.*, 2006, **281**, 22684–22694.
- 55 J. E. Markham and J. G. Jaworski, *Rapid Commun Mass Spectrom.*, 2007, **21**, 1304–1314.
- 56 B. Ebert, C. Rautengarten, H. E. McFarlane, T. Rupasinghe, W. Zeng, K. Ford, H. V. Scheller, A. Bacic, U. Roessner, S. Persson and J. L. Heazlewood, *Nat Plants*, 2018, **4**, 792–801.
- 57 E. Kiran, P. G. Debenedetti and C. J. Peters, *Supercritical fluids: fundamentals and applications*, Springer Science & Business Media, 2012.
- 58 S. P. J. Kumar, S. R. Prasad, R. Banerjee, D. K. Agarwal, K. S. Kulkarni and K. V. Ramesh, *Chem. Cent. J.*, 2017, **11**, 9.
- 59 S. P. Jeevan Kumar, G. Vijay Kumar, A. Dash, P. Scholz and R. Banerjee, *Algal Res.*, 2017, **21**, 138–147.
- 60 K. V. Probst, M. D. Wales, M. E. Rezac and P. V. Vadlani, *Biotechnol. Prog.*, 2017, **33**, 1096–1103.
- 61 S. S. de Jesus, G. F. Ferreira, L. S. Moreira, M. R. Wolf Maciel and R. Maciel Filho, *Renewable Energy*, 2019, **143**, 130–141.
- 62 N. Imatoukene, M. Koubaa, E. Perdrix, M. Benali and E. Vorobiev, *Process Biochem.*, 2020, **90**, 139–147.
- 63 S. S. de Jesus, G. F. Ferreira, L. V. Fregolente and R. Maciel Filho, *Algal Res.*, 2018, **35**, 292–300.
- 64 M. Cvjetko Bubalo, S. Vidović, I. Radojčić Redovniković and S. Jokić, *Food and Bioprod. Process.*, 2018, **109**, 52–73.
- 65 S. Obeid, N. Beaufils, S. Camy, H. Takache, A. Ismail and P.-Y. Pontalier, *Algal Res.*, 2018, **34**, 49–56.
- 66 J. A. Bowden, C. Z. Ulmer, C. M. Jones, J. P. Koelmel and R. A. Yost, *Metabolomics*, 2018, **14**, 53.
- 67 X. Han and R. W. Gross, *Proc. Natl. Acad. Sci. U. S. A.*, 1994, **91**, 10635–10639.
- 68 B. Brügger, G. Erben, R. Sandhoff, F. T. Wieland and W. D. Lehmann, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 2339–2344.
- 69 R. Welti, X. Wang and T. D. Williams, *Anal. Biochem.*, 2003, **314**, 149–152.
- 70 S. Narayanan, P. J. Tamura, M. R. Roth, P. V. V. Prasad and R. Welti, *Plant, Cell Environ.*, 2016, **39**, 787–803.
- 71 R. Welti and X. Wang, *Curr. Opin. Plant Biol.*, 2004, **7**, 337–344.
- 72 S. Shiva, T. Samarakoon, K. A. Lowe, C. Roach, H. S. Vu, M. Colter, H. Porras, C. Hwang, M. R. Roth, P. Tamura, M. Li, K. Schrick, J. Shah, X. Wang, H. Wang and R. Welti, *Plants*, 2020, **9**, 845.
- 73 Z. S. Zoong Lwe, R. Welti, D. Anco, S. Naveed, S. Rustgi and S. Narayanan, *Sci. Rep.*, 2020, **10**, 22163.
- 74 H. C. Köfeler, A. Fauland, G. N. Rechberger and M. Trötschmüller, *Metabolites*, 2012, **2**, 19–38.
- 75 P. Tarazona, K. Feussner and I. Feussner, *Plant J.*, 2015, **84**, 621–633.
- 76 M. Lange, Z. Ni, A. Criscuolo and M. Fedorova, *Chromatographia*, 2019, **82**, 77–100.
- 77 H. Nygren, T. Seppänen-Laakso, S. Castillo, T. Hyötyläinen and M. Orešič, in *Metabolic Profiling: Methods and Protocols*, ed. T. O. Metz, Humana Press, Totowa, NJ, 2011, DOI: 10.1007/978-1-61737-985-7_15, pp. 247–257.
- 78 T. Cajka and O. Fiehn, *TrAC, Trends Anal. Chem.*, 2014, **61**, 192–206.
- 79 T. W. T. Rupasinghe and U. Roessner, in *Plant Metabolomics: Methods and Protocols*, ed. C. António, Springer, New York, New York, NY, 2018, DOI: 10.1007/978-1-4939-7819-9_9, pp. 125–135.
- 80 A. M. King, R. D. Trengove, L. G. Mullin, P. D. Rainville, G. Isaac, R. S. Plumb, L. A. Gethings and I. D. Wilson, *J. Chromatogr. A*, 2020, **1611**, 460597.
- 81 M. Li, L. Yang, Y. Bai and H. Liu, *Anal. Chem.*, 2014, **86**, 161–175.
- 82 G. Isaac, N. Munjoma, L. A. Gethings and R. S. Plumb, 2018.
- 83 M. Lange and M. Fedorova, *Anal. Bioanal. Chem.*, 2020, **412**, 3573–3584.
- 84 M. Holčapek, M. Ovčáčíková, M. Lísa, E. Cífková and T. Hájek, *Anal. Bioanal. Chem.*, 2015, **407**, 5033–5043.
- 85 M. Navarro-Reig, J. Jaumot and R. Tauler, *J. Chromatogr. A*, 2018, **1568**, 80–90.
- 86 M. de la Guardia and S. Armenta, in *Comprehensive Analytical Chemistry*, ed. M. D. L. Guardia and S. Armenta, Elsevier, 2011, vol. 57, pp. 121–156.
- 87 M. Xu, J. Legradi and P. Leonards, *Anal. Bioanal. Chem.*, 2020, **412**, 4313–4325.
- 88 N. J. Jensen and M. L. Gross, *Mass Spectrom. Rev.*, 1987, **6**, 497–536.
- 89 A. V. Melnik, R. R. da Silva, E. R. Hyde, A. A. Aksenov, F. Vargas, A. Bouslimani, I. Protsyuk, A. K. Jarmusch, A. Tripathi, T. Alexandrov, R. Knight and P. C. Dorrestein, *Anal. Chem.*, 2017, **89**, 7549–7559.
- 90 L. Gao, A. Cazenave-Gassiot, B. Burla, M. R. Wenk and F. Torta, *Metabolomics*, 2020, **16**, 53.
- 91 T. F. Jorge, A. T. Mata and C. António, *Philos. Trans. R. Soc., A*, 2016, **374**, 20150370.



- 92 D. Yu, T. W. T. Rupasinghe, B. A. Boughton, S. H. A. Natera, C. B. Hill, P. Tarazona, I. Feussner and U. Roessner, *Anal. Chim. Acta*, 2018, **1026**, 87–100.
- 93 T. Cajka, J. T. Smilowitz and O. Fiehn, *Anal. Chem.*, 2017, **89**, 12360–12368.
- 94 J. Schlotterbeck, M. Chatterjee, M. Gawaz and M. Lämmerhofer, *Anal. Chim. Acta*, 2019, **1046**, 1–15.
- 95 X. Zhu, Y. Chen and R. Subramanian, *Anal. Chem.*, 2014, **86**, 1202–1209.
- 96 T. H. Pham, M. Zaeem, T. A. Fillier, M. Nadeem, N. P. Vidal, C. Manful, S. Cheema, M. Cheema and R. H. Thomas, *Sci. Rep.*, 2019, **9**, 5048.
- 97 J. Koelmel, M. Sartain, J. Salcedo, A. Murali, X. Li and S. Stow, *Agilent Technologies Application Note*, 2019, 5994-0775EN.
- 98 J. P. Koelmel, N. M. Kroeger, E. L. Gill, C. Z. Ulmer, J. A. Bowden, R. E. Patterson, R. A. Yost and T. J. Garrett, *J. Am. Soc. Mass Spectrom.*, 2017, **28**, 908–917.
- 99 M. Raetz, R. Bonner and G. Hopfgartner, *Metabolomics*, 2020, **16**, 71.
- 100 L. C. Gillet, P. Navarro, S. Tate, H. Röst, N. Selevsek, L. Reiter, R. Bonner and R. Aebersold, *Mol. Cell. Proteomics*, 2012, **11**, O111.016717.
- 101 H. Tsugawa, T. Cajka, T. Kind, Y. Ma, B. Higgins, K. Ikeda, M. Kanazawa, J. VanderGheynst, O. Fiehn and M. Arita, *Nat. Methods*, 2015, **12**, 523–526.
- 102 T. Kind, K. H. Liu, D. Y. Lee, B. DeFelice, J. K. Meissen and O. Fiehn, *Nat. Methods*, 2013, **10**, 755–758.
- 103 T. Xu, C. Hu, Q. Xuan and G. Xu, *Anal. Chim. Acta*, 2020, **1137**, 156–169.
- 104 B. Drotleff, J. Illison, J. Schlotterbeck, R. Lukowski and M. Lämmerhofer, *Anal. Chim. Acta*, 2019, **1086**, 90–102.
- 105 B. Drotleff, M. Hallschmid and M. Lämmerhofer, *Anal. Chim. Acta*, 2018, **1022**, 70–80.
- 106 C. Kehelpannala, T. Rupasinghe, A. Pasha, E. Esteban, T. Hennessy, D. Bradley, B. Ebert, N. J. Provart and U. Roessner, *Plant J.*, 2021, **107**, 287–302.
- 107 M. Bilgin, P. Born, F. Fezza, M. Heimes, N. Mastrangelo, N. Wagner, C. Schultz, M. Maccarrone, S. Eaton, A. Nadler, M. Wilm and A. Shevchenko, *Sci. Rep.*, 2016, **6**, 27920.
- 108 J. P. Koelmel, X. Li, S. M. Stow, M. J. Sartain, A. Murali, R. Kemperman, H. Tsugawa, M. Takahashi, V. Vasiliou, J. A. Bowden, R. A. Yost, T. J. Garrett and N. Kitagawa, *Metabolites*, 2020, **10**, 101.
- 109 R. Taguchi and M. Ishikawa, *J. Chromatogr. A*, 2010, **1217**, 4229–4239.
- 110 B. K. Ubhi, A. Conner, E. Duchoslav, A. Evans, R. Robinson, P. R. Baker and S. Watkins, *SCIEX Technical Application Note*, 2015.
- 111 J. Hartler, A. Triebel, M. Trötzmüller, A. Ziegl, G. Rechberger, F. Spener, H. Köfeler and G. Thallinger, 2015.
- 112 Y. Djoumbou-Feunang, A. Pon, N. Karu, J. Zheng, C. Li, D. Arndt, M. Gautam, F. Allen and D. S. Wishart, *Metabolites*, 2019, **9**, 72.
- 113 J. E. Kyle, K. L. Crowell, C. P. Casey, G. M. Fujimoto, S. Kim, S. E. Dautel, R. D. Smith, S. H. Payne and T. O. Metz, *Bioinformatics*, 2017, **33**, 1744–1746.
- 114 M. A. Kochen, M. C. Chambers, J. D. Holman, A. I. Nesvizhskii, S. T. Weintraub, J. T. Belisle, M. N. Islam, J. Griss and D. L. Tabb, *Anal. Chem.*, 2016, **88**, 5733–5741.
- 115 A. Mohamed, J. Molendijk and M. M. Hill, *J. Proteome Res.*, 2020, **19**, 2890–2897.
- 116 P. D. Hutchins, J. D. Russell and J. J. Coon, *Cell Systems*, 2018, **6**(621–625), e625.
- 117 M. I. Alcoriza-Balaguer, J. C. García-Cañaveras, A. López, I. Conde, O. Juan, J. Carretero and A. Lahoz, *Anal. Chem.*, 2019, **91**, 836–845.
- 118 J. P. Koelmel, N. M. Kroeger, C. Z. Ulmer, J. A. Bowden, R. E. Patterson, J. A. Cochran, C. W. W. Beecher, T. J. Garrett and R. A. Yost, *BMC Bioinf.*, 2017, **18**, 331.
- 119 T. Zullig, M. Trotschmüller and H. C. Köfeler, *Anal. Bioanal. Chem.*, 2020, **412**, 2191–2209.
- 120 S. Lu, H. Liu, C. Jin, Q. Li and L. Guo, *Plant Direct*, 2019, **3**, e00183.
- 121 J. Liu, Q. Li, J. Chen and Y. Jiang, *Foods*, 2020, **9**, 894.
- 122 A. Millner, D. Y. Lizardo and G. E. Atilla-Gökçumen, *Proteomics*, 2020, **20**, 2000013.
- 123 X. Zhang, W. Liu, J. Zan, C. Wu and W. Tan, *Sci. Rep.*, 2020, **10**, 14509.
- 124 A. Acera, X. Pereiro, B. Abad-García, Y. Rueda, N. Ruzafa, C. Santiago, I. Barbolla, J. A. Duran, B. Ochoa and E. Vecino, *Mol. Vis.*, 2019, **25**, 934–948.
- 125 S. B. Breitkopf, S. J. H. Ricoult, M. Yuan, Y. Xu, D. A. Peake, B. D. Manning and J. M. Asara, *Metabolomics*, 2017, **13**, 30.
- 126 A. Gil, W. Zhang, J. C. Wolters, H. Permentier, T. Boer, P. Horvatovich, M. R. Heiner-Fokkema, D.-J. Reijngoud and R. Bischoff, *Anal. Bioanal. Chem.*, 2018, **410**, 5859–5870.
- 127 G. Liebisch, K. Ekroos, M. Hermansson and C. S. Ejsing, *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids*, 1862, **2017**, 747–751.
- 128 G. Liebisch, R. Ahrends, M. Arita, M. Arita, J. A. Bowden, C. S. Ejsing, W. J. Griffiths, M. Holčápek, H. Köfeler, T. W. Mitchell, M. R. Wenk, K. Ekroos and C. Lipidomics Standards Initiative, *Nature Metabolism*, 2019, **1**, 745–747.
- 129 S. K. K. Mackenzie Pearson, P. Norris and C. Hunter, *SCIEX*, 2018.
- 130 M. Koivusalo, P. Haimi, L. Heikinheimo, R. Kostinen and P. Somerharju, *J. Lipid Res.*, 2001, **42**, 663–672.
- 131 Y. Xie, H. Ma, F. Wei, X. Lyu, Z. Wu, B. Wu, S. Xu, X. Dong, H. Chen and F. Huang, *Proc. - Soil Crop Sci. Soc. Fla.*, 2017, **2**, 217–224.
- 132 Á. Sánchez-Illana, J. D. Piñeiro-Ramos, J. D. Sanjuan-Herráez, M. Vento, G. Quintás and J. Kuligowski, *Anal. Chim. Acta*, 2018, **1019**, 38–48.
- 133 H. Tenenboim, A. Burgos, L. Willmitzer and Y. Brotman, *Biochimie*, 2016, **130**, 91–96.
- 134 I. The Arabidopsis Genome, *Nature*, 2000, **408**, 796–815.
- 135 Y. Li-Beisson, B. Shorrosh, F. Beisson, M. X. Andersson, V. Arondel, P. D. Bates, S. Baud, D. Bird, A. Debono, T. P. Durrett, R. B. Franke, I. A. Graham, K. Katayama, A. A. Kelly, T. Larson, J. E. Markham, M. Miquel, I. Molina, I. Nishida, O. Rowland, L. Samuels, K. M. Schmid,



- H. Wada, R. Welte, C. Xu, R. Zallot and J. Ohlrogge, *Arabidopsis Book*, 2013, **11**, e0161.
- 136 X. Wang, *Plant Physiol.*, 2005, **139**, 566–573.
- 137 X. Wang, W. Li, M. Li and R. Welte, *Physiol. Plant.*, 2006, **126**, 90–96.
- 138 J. Bewley and M. Black, 1986.
- 139 K. Takahashi, R. Morimoto, H. Tabeta, M. Asaoka, M. Ishida, M. Maeshima, H. Tsukaya and A. Ferjani, *Plant Cell Physiol.*, 2017, **58**, 668–678.
- 140 D. H. McLachlan, J. Lan, C.-M. Geilfus, A. N. Dodd, T. Larson, A. Baker, H. Hörak, H. Kollist, Z. He, I. Graham, M. V. Mickelbart and A. M. Hetherington, *Curr. Biol.*, 2016, **26**, 707–712.
- 141 S. D. Singer, J. Zou and R. J. Weselake, *Plant Sci.*, 2016, **243**, 1–9.
- 142 M. Wolters-Arts, W. M. Lush and C. Mariani, *Nature*, 1998, **392**, 818–821.
- 143 M. Wang, C. Wang, R. H. Han and X. Han, *Prog. Lipid Res.*, 2016, **61**, 83–108.
- 144 M. O. Pata, Y. A. Hannun and C. K. Y. Ng, *New Phytologist*, 2010, **185**, 611–630.
- 145 C. Hu, W. Luo, J. Xu and X. Han, *Mass Spectrom. Rev.*, 2020, DOI: 10.1002/mas.21659.
- 146 R. M. Gathungu, P. Larrea, M. J. Sniatynski, V. R. Marur, J. A. Bowden, J. P. Koelmel, P. Starke-Reed, V. S. Hubbard and B. S. Kristal, *Anal. Chem.*, 2018, **90**, 13523–13532.
- 147 E. Ruelland and O. Valentova, *Front. Plant Sci.*, 2016, **7**, 324.
- 148 Y. Okazaki, Y. Kamide, M. Y. Hirai and K. Saito, *Metabolomics*, 2013, **9**, 121–131.
- 149 A. K. Nilsson, O. N. Johansson, P. Fahlberg, M. Kommuri, M. Töpel, L. J. Bodin, P. Sikora, M. Modarres, S. Ekengren, C. T. Nguyen, E. E. Farmer, O. Olsson, M. Ellerström and M. X. Andersson, *Plant J.*, 2015, **84**, 1152–1166.
- 150 R. Welte, W. Li, M. Li, Y. Sang, H. Biesiada, H. E. Zhou, C. B. Rajashekar, T. D. Williams and X. Wang, *J. Biol. Chem.*, 2002, **277**, 31994–32002.
- 151 S. P. Devaiah, M. R. Roth, E. Baughman, M. Li, P. Tamura, R. Jeannotte, R. Welte and X. Wang, *Phytochemistry*, 2006, **67**, 1907–1924.
- 152 J. Hummel, S. Segu, Y. Li, S. Irgang, J. Jueppner and P. Giavalisco, *Front. Plant Sci.*, 2011, **2**, 54.
- 153 Y. Okazaki, H. Otsuki, T. Narisawa, M. Kobayashi, S. Sawai, Y. Kamide, M. Kusano, T. Aoki, M. Y. Hirai and K. Saito, *Nat. Commun.*, 2013, **4**, 1510.
- 154 M.-Y. Liu, A. Burgos, L. Ma, Q. Zhang, D. Tang and J. Ruan, *BMC Plant Biol.*, 2017, **17**, 165.
- 155 P. Giavalisco, Y. Li, A. Matthes, A. Eckhardt, H.-M. Hubberten, H. Hesse, S. Segu, J. Hummel, K. Köhl and L. Willmitzer, *Plant J.*, 2011, **68**, 364–376.
- 156 D. Yu, B. A. Boughton, C. B. Hill, I. Feussner, U. Roessner and T. W. T. Rupasinghe, *Front. Plant Sci.*, 2020, **11**, 1.
- 157 E. Maciel, A. Lillebø, P. Domingues, E. da Costa, R. Calado and M. R. M. Domingues, *Phytochemistry*, 2018, **153**, 94–101.
- 158 B. E. Cheong, W. W. H. Ho, B. Biddulph, X. Wallace, T. Rathjen, T. W. T. Rupasinghe, U. Roessner and R. Dolferus, *Metabolomics*, 2019, **15**, 144.
- 159 B. E. Cheong, O. Onyemaobi, W. Wing Ho Ho, T. B. Biddulph, T. W. T. Rupasinghe, U. Roessner and R. Dolferus, *Cells*, 2020, **9**, 1309.
- 160 L. Panzenboeck, N. Troppmair, S. Schlachter, G. Koellensperger, J. Hartler and E. Rampler, *Metabolites*, 2020, **10**, 375.
- 161 T. Lapidot-Cohen, L. Rosental and Y. Brotman, *Curr Protoc Plant Biol*, 2020, **5**, e20109.
- 162 J. Liu, Q. Li, J. Chen and Y. Jiang, *Foods*, 2020, **9**, 894.
- 163 M. Schillaci, C. Kehelpannala, F. Martinez-Seidel, P. M. C. Smith, B. Arsova, M. Watt and U. Roessner, *Metabolites*, 2021, **11**, 358.

