

Simple model for tuberculosis in cattle and badgers

D. R. Cox^{*†‡}, Christl A. Donnelly^{†§}, F. John Bourne[†], George Gettinby^{†||}, John P. McInerney^{†||}, W. Ivan Morrison^{†**}, and Rosie Woodroffe^{†,††}

^{*}Nuffield College, New Road, Oxford OX1 1NF, United Kingdom; [†]Independent Scientific Group on Cattle TB, c/o Department for Environment, Food and Rural Affairs, 1A Page Street, London SW1P 4PQ, United Kingdom; [§]Department of Infectious Disease Epidemiology, Faculty of Medicine, Imperial College London, St. Mary's Campus, Norfolk Place, London W2 1PG, United Kingdom; ^{||}Department of Statistics and Modelling Science, University of Strathclyde, Richmond Street, Glasgow G1 1XH, United Kingdom; ^{||}Centre for Rural Research, University of Exeter, Lafrowda House, St. German's Road, Exeter EX4 6TL, United Kingdom; ^{**}Centre for Tropical Veterinary Medicine, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Roslin, Midlothian EH25 9RG, United Kingdom; and ^{††}Department of Wildlife, Fish, and Conservation Biology, University of California, Davis, One Shields Avenue, Davis, CA 95616

Contributed by D. R. Cox, October 17, 2005

As an aid to the study of bovine tuberculosis (TB), a simple model has been developed of an epidemic involving two species, cattle and badgers. Each species may infect the other. The proportion of animals affected is assumed relatively small so that the usual nonlinear aspects of epidemic theory are avoided. The model is used to study the long-run and transient effect on cattle of culling badgers and the effect of a period without routine testing for TB, such as occurred during the 2001 epidemic of foot-and-mouth disease in Great Britain. Finally, by examining the changes in cattle TB over the last 15 years, and with some other working assumptions, it is estimated that the net reproduction number of the epidemic is ≈ 1.1 . The implications for controlling the disease are discussed.

animal ecology | epidemiology | mathematical model

Mathematical modeling of infectious disease, especially in humans, has a long history (1). Two very different kinds of dynamical model are simple outline and quasirealistic models. The former aim to capture the essence of a complex situation in a few key aspects and then to use these to explain the qualitative patterns of behavior to be expected and even to give semiquantitative predictions. Quasirealistic models, by contrast, aim to get much closer to reality by representing, almost inevitably in a model implemented by computer rather than by mathematical analysis, as many features of reality as is feasible. Despite the attractions of greater realism, a common problem with such models is the need to specify the values of a considerable number of forms of dependence and the numerical values of unknown parameters, often, as in the present instance, aspects about which little is known. The consequent need for extensive and systematic sensitivity analyses may undermine the usefulness of such models and mean that such data as are available are consistent with many different interpretations.

In the present work we consider a deterministic multispecies transmission model of the former simpler type to describe *Mycobacterium bovis* infection in cattle and badgers. After setting out a fairly general model framework allowing both within- and between-species transmission, we look at a number of special situations corresponding in particular to interventions on one or other species.

General Model Specification

We consider a reasonably homogeneous area such that at time t there are $X_1(t) = C(t)$ infected cattle per km² and $X_2(t) = B(t)$ infected badgers per km². We suppose that the proportions of infected animals are relatively small, say <0.2 , so that the nonlinear effects characteristic of epidemic theory, because of the depletion of the pool of susceptible animals, are not important. This assumption seems reasonable because reports of more than a small number of reactor cattle within individual herd breakdowns are rare. Also, we do not distinguish between infected and infectious animals, thus making the approximation

that all infected animals of a particular species are equally infectious.

A major assumption is that of spatial homogeneity over the area considered. This assumption means, in particular, that use of the conclusions at a very local level would not be justified.

Let a_{ij} be the rate per unit time (usually a year) at which one animal of species j infects animals of species i . Let animals of species i be removed at a total rate r_i and suppose that restocking introduces new infected animals at rate s_i . Note that the removal rate of cattle at time t , $r_1(t)$, includes both routine slaughter and slaughter after identification of disease. We assume that although both removal and immigration rates may be affected by policy decisions or external conditions, they are constant over defined periods.

Let \mathbf{A} denote the matrix of the a_{ij} and $\mathbf{X}(t) = \{X_1(t), X_2(t)\}^T$. Further, let $\mathbf{R} = \text{diag}(r_1, r_2)$ and $\mathbf{S} = \{s_1, s_2\}^T$. Then

$$d\mathbf{X}(t)/dt = (\mathbf{A} - \mathbf{R})\mathbf{X} + \mathbf{S}. \quad [1]$$

Therefore, if $\mathbf{B} = \mathbf{R} - \mathbf{A}$, and if initially $\mathbf{X}(0) = \mathbf{X}_0$, then we have that

$$\mathbf{X}(t) = \mathbf{B}^{-1}\mathbf{S} + e^{-\mathbf{B}t}(\mathbf{X}_0 - \mathbf{B}^{-1}\mathbf{S}),$$

where the matrix exponential is computed through the spectral decomposition $\mathbf{B} = \mathbf{H}\mathbf{\Lambda}\mathbf{H}^{-1}$ with $\mathbf{\Lambda} = \text{diag}(\lambda_1, \lambda_2)$. Then

$$\mathbf{X}(t) = \mathbf{B}^{-1}\mathbf{S} + \mathbf{H}\text{diag}(e^{-\lambda_1 t}, e^{-\lambda_2 t})\mathbf{H}^{-1}(\mathbf{X}_0 - \mathbf{B}^{-1}\mathbf{S}).$$

Note that this discussion could be applied formally to any number of species. Furthermore, four "species" could be used to represent cattle and badgers in two geographical regions. This metapopulation approach will not be explored here.

If, on the other hand, the number of infected badgers $B(t)$ at time t were externally influenced, for example by culling (whether one-off or repeated), rather than determined by Eq. 1, the resulting equation for $C(t)$ is

$$dC(t)/dt = (a_{11} - r_1)C(t) + a_{12}B(t) + s_1(t),$$

where it is convenient to allow also that the restocking rate may vary over time.

Epidemiological Setting

Cattle herds are subject to regulations set out in the Animal Health Act requiring regular tuberculosis (TB) tests. Each herd is subjected to a parish-based testing interval of between 4 yr and 1 yr, where parishes with the lowest incidence are assigned

Conflict of interest statement: No conflicts declared.

Freely available online through the PNAS open access option.

Abbreviation: TB, tuberculosis.

[†]To whom correspondence should be addressed. E-mail: david.cox@nuffield.oxford.ac.uk.

© 2005 by The National Academy of Sciences of the USA

Table 2. Ratio C_k/C_0 (Eq. 5) of infected animals to initial level after reducing badger level by factor k as a function of the badger influence ratio I_B

k	I_B					
	0.10	0.50	1.00	2.00	5.00	10.00
0.2	0.98	0.93	0.90	0.87	0.83	0.82
0.5	0.95	0.83	0.75	0.67	0.58	0.54
0.8	0.93	0.73	0.60	0.47	0.33	0.27
1	0.91	0.67	0.50	0.33	0.16	0.09

by badgers per year to the number of infections in cattle brought in from outside the system per year.

Table 2 gives C_k/C_0 (Eq. 5) for a range of values of k and I_B .

In this formulation the proportional effect of reducing, or increasing, the badger population depends neither on the removal rate, r_1 , nor on the cattle-to-cattle infection rate, a_{11} , within an area, although these do, of course, affect the actual level attained. The level of reduction achieved in the badger population would have to be maintained by repeated culling for these results to apply. The value of I_B is unknown.

In this discussion it is assumed that only the number of infected badgers is changed. If, at the other extreme, the external incidence rate s_1 also were reduced by the same factor k then the equilibrium level is reduced by that factor, too, assuming removal rate and cattle-to-cattle transmission within the area remained unchanged.

Time-Dependent Effect of Initial Culling: A Single Cull

To study the rate at which the equilibrium levels in the previous discussion are achieved and to examine some other perturbations of the system, we solve the original differential Eq. 1 in the form

$$C(t) = C_0 e^{(\lambda-r_1)t} + \int_0^t e^{(\lambda-r_1)(t-x)} \{a_{12}B(t-x) + s_1(t-x)\} dx,$$

where C_0 is the initial value of $C(t)$.

We examine a number of different situations. First, as in the previous section, suppose that at $t = 0$ the system is in equilibrium corresponding to $B(t) = B_0$ for $t < 0$, so that C_0 is given by Eq. 4 and that the number of infected badgers is instantaneously reduced to $(1-k)B_0$ and thereafter returns to its initial level, the return being represented for $t \geq 0$ by

$$B(t) = B_0(1 - ke^{-\nu t}).$$

Thus, after time $0.69/\nu$, the badger population will have returned halfway to its original level, B_0 . The possibility of maintaining the badger population permanently at its new level is represented by $\nu = 0$.

The general solution now takes the form

$$C(t) = C_0 + a_{12}B_0k(\lambda - r_1 + \nu)^{-1}(e^{-\nu t} - e^{(\lambda-r_1)t}).$$

Thus,

$$C(t)/C_0 = 1 - \frac{kI_B(1 - \lambda/r_1)}{(1 + I_B)(1 - \lambda/r_1 - \nu/r_1)}(e^{-\nu t} - e^{(\lambda-r_1)t}). \quad [6]$$

The function $C(t)$ reaches a minimum after time

$$(\nu - r_1 + \lambda)^{-1} \log\{\nu/(r_1 - \lambda)\},$$

and then returns toward its initial value C_0 . If the two timescales determined by cattle aspects, $r_1 - \lambda$, and by badger aspects ν are very different, the more rapid timescale dominates behavior.

This equation specifies prevalence as a function of time. In some ways, incidence, that is the rate of occurrence of new infections, although not directly observable, is more relevant, and this rate is

$$\begin{aligned} dC(t)/dt + r_1C(t) &= r_1C_0 - a_{12}B_0(r_1 - k\nu)(r_1 - \lambda - \nu)^{-1}e^{-\nu t} \\ &\quad + a_{12}B_0(r_1 - kr_1 + k\lambda) \\ &\quad \cdot (r_1 - \lambda - \nu)^{-1}e^{(\lambda-r_1)t}. \end{aligned}$$

When divided by the equilibrium rate r_1C_0 this value is

$$\begin{aligned} 1 - \frac{I_B(1 - k\nu/r_1)(1 - \lambda/r_1)}{(1 + I_B)(1 - \lambda/r_1 - \nu/r_1)}e^{-\nu t} \\ + \frac{I_B(1 - k + k\lambda/r_1)(1 - \lambda/r_1)}{(1 + I_B)(1 - \lambda/r_1 - \nu/r_1)}e^{(\lambda-r_1)t}. \end{aligned} \quad [7]$$

Thus, immediately after the reduction in the number of badgers, the rate of new infections is reduced from r_1C_0 by the factor

$$1 - \frac{kI_B(1 - \lambda/r_1)}{(1 + I_B)}.$$

Time-Dependent Solution: A Maintained Cull

Next suppose that at time $t = 0$ the testing frequency is changed from every d years to every d' years so that the removal rate changes from r_1 to r'_1 , B_0 and s_1 remaining unchanged. Then from Eq. 4

$$C(t) = C'_0 + (C_0 - C'_0)e^{(\lambda-r'_1)t},$$

where C'_0 is the equilibrium level of $C(t)$ at the new testing frequency. In fact

$$C'_0/C_0 = (r_1 - \lambda)/(r'_1 - \lambda). \quad [8]$$

Equilibrium requires that the removal rates exceed λ and if that state were reached, Eq. 8 shows that the equilibrium level of prevalence achieved would be approximately inversely proportional to the removal rate.

Time-Dependent Solution: Interruption of Operations

Finally, suppose that, starting from equilibrium, there is a period $(0, t_0)$ in which there is no removal or restocking, but infection by badgers and by cattle-to-cattle transmission in the area continues. Then, after time t_0 the number of infected animals is given by the solution of the full set of equations and is

$$(C_0 + a_{12}B_0/\lambda)e^{\lambda t_0} - a_{12}B_0/\lambda. \quad [9]$$

Suppose then that normal testing and restocking resumed. Then after a further time t

$$C(t_0 + t) = \{(C_0 + a_{12}B_0/\lambda)e^{\lambda t_0} - a_{12}B_0/\lambda - C_0\}e^{(\lambda-r_1)t} + C_0. \quad [10]$$

Therefore,

$$\frac{C(t_0 + t)}{C_0} = 1 + \frac{r_1(I_B + \lambda/r_1)}{\lambda(I_B + 1)}(e^{\lambda t_0} - 1)e^{-(r_1-\lambda)t}. \quad [11]$$

If λt_0 is small, this ratio is approximately

$$1 + \frac{r_1(I_B + \lambda/r_1)}{(I_B + 1)} t_0 e^{-(r_1 - \lambda)t}.$$

This situation corresponds to the restriction of livestock movements between farms and the suspension of routine tuberculin skin testing of cattle during the foot-and-mouth disease epidemic in the United Kingdom in 2001.

Conditions for Epidemic Progression

So far, much of the discussion has centered around the notion of a system in equilibrium and suffering relatively minor deviations from that equilibrium. The condition for epidemic growth, essentially that the net reproduction rate exceeds one, is that the matrix $\mathbf{B} = \mathbf{R} - \mathbf{A}$ have at least one negative eigenvalue. This negative eigenvalue would imply an exponential increase in prevalence in both species, although of course the effect might be numerically small in one of the species.

Numerical Results Based on Model

The results sketched above depend on a limited number of unknown parameters. One of the functions of simple models such as this one is in fact to highlight the features of the system most likely to have a big impact on the outcome. An example is the index I_B influencing the outcome of a permanent reduction in the number of infected badgers, and another is k , the efficiency of culling. As has already been mentioned, the amount of information available about all of the parameters is limited, and study of a range of parameter values is thus essential. We can, however, obtain approximate relevant ranges by the following arguments.

To set out the numerical results, it is helpful to use dimensionless units, although for substantive interpretation it is usually necessary to convert back to natural units. Thus, we set $r_1 t = \tau$ so that τ is a dimensionless timescale determined by the replacement rate r_1 . Further, we define $l = \lambda/r_1$ and $v = \nu/r_1$, thereby expressing the cattle-to-cattle transmission rate and the badger return rate relative to r_1 .

First, for the most part we assume λ is small compared with r_1 , i.e., that λ is small. Were λ not small, most outbreaks in a farm would be multiple and often involve whole herd breakdowns. These are, however, relatively rare.

For interpreting τ in natural time units, typically years, we have to estimate r_1 . We take a diagnostic test sensitivity of $p = 2/3$, this value corresponding broadly to reports in the literature (12, 13) to obtain the effective removal rate as $r = 1.3 \text{ yr}^{-1}$. Lower sensitivity would imply decreased r_1 , and it is clear from Eq. 4 and other formulas that this change could substantially increase $C(t)$. If the badger population returns to its equilibrium level at a rate of $\approx 20\%$ per year, then $\nu = 0.23 \text{ yr}^{-1}$. For most purposes, therefore, we show functions of time as functions of τ at a range of values of I_B and the culling fraction k .

The effect of culling on the equilibrium level of TB is shown in simple form in Table 2. The corresponding standardized prevalence and incidence curves are shown in Figs. 1 and 2 as functions of dimensionless time τ for selected values of I_B , k , and $l = \lambda/r_1$. The badger return rate $\nu = \nu/r_1$ has been fixed at 0.17. In the standardized time measure τ , a unit of time corresponds to $\approx 0.75 \text{ yr}$, i.e., $\tau = 4$ translates to 3 yr.

Both prevalence and incidence fall rapidly after culling to a level determined largely by the efficiency of culling, k , and by the index of badger activity, I_B . They then return slowly to their initial level. This result implies that repeat culling at intervals of perhaps five dimensionless units should hold prevalence close to its minimum level. Clearly a more refined calculation of the effects of repeat culling would be possible.




Fig. 1. The standardized prevalence of TB in cattle (Eq. 6) as a function of time, τ , since culling; the proportion by which the badger population was reduced, k ; the relative cattle-to-cattle transmission rate, l ; and a measure of the relative influence of badgers as compared with disease in cattle restocked into the system, I_B .

Eqs. 9–11 were considered to assess the effect of a 9-month cessation of testing during the British foot-and-mouth disease epidemic. In particular, Eq. 11 shows the relative prevalence after resumption of normal procedures. Fig. 3 shows this quantity as a function of time for selected values of I_B and $l = \lambda/r_1$. For the range of values considered an initial increase in prevalence by a factor of up to 2 is to be expected followed by a relatively rapid decrease back to the equilibrium level.

Estimation of the Basic Reproductive Number

Consider the growth of the epidemic in cattle over a reasonably large and homogeneous region. Let $C(t)$ be the number of infected cattle at time t . Suppose that they are removed by detection and routine removal at rate r_1 per year. With annual testing at $2/3$ test sensitivity and “usual” demographic properties, r_1 is $\approx 1.3 \text{ yr}^{-1}$.

Assume that the proportion of animals affected is low and that there is a new infection rate of λ per year per infected animal. Note that these new infections arise from cattle-to-cattle transmission within a herd and by infection by cattle brought in from a source with about the same infection rate. In addition, suppose




Fig. 2. The standardized incidence of TB in cattle herds (Eq. 7) as a function of time, τ , since culling; the proportion by which the badger population was reduced, k ; the relative cattle-to-cattle transmission rate, l ; and a measure of the relative influence of badgers as compared with disease in cattle restocked into the system, I_B .

old is obtained by estimating $(\lambda - r_1)$ from fitting the data, obtaining the estimate of λ , denoted $\hat{\lambda}$, by subtraction based on the r_1 value obtained from the assumed testing frequency and test sensitivity. The potential net reproduction corresponding to future alternative testing frequency d' and/or test sensitivity p' is

$$\frac{\hat{\lambda}}{1/\mu_n + d'^{-1}/\{1/2 + (1 - p')/p'\}}.$$

For example, if the true test sensitivity over the period analyzed was 66.7% and testing had been annual in the Western Region, then an increase in test sensitivity to $\approx 73\%$ would be sufficient to drop the reproduction rate to <1 . Alternatively, if the true test sensitivity over the period analyzed was 80% and testing was annual in the Western Region, then an increase in test sensitivity to $\approx 85\%$ would be sufficient. Finally, if the true test sensitivity over the period analyzed was 90% and testing was annual in the Western Region, then an increase in test sensitivity to $\approx 94\%$ would be sufficient. Alternatively, in each of these cases, an increase in testing from annual to every 10 months was sufficient to drop the reproduction rate to <1 .

These illustrative results demonstrate that although the net reproduction number is clearly >1 , it is sufficiently close to 1 that relatively modest improvements either in test performance or testing frequency would be sufficient to bring the epidemic under control, under the highly idealized assumptions made in this model.

Discussion and Further Developments

The estimates for net reproduction number for TB in cattle ranged from 1.02 to 1.11. Published estimates for the basic reproduction number for *M. bovis* infections wildlife species have varied considerably between and within species partly as a function of density: 1.2 in badgers in Great Britain (14), 1.6–2.1 in brushtail possums in New Zealand (15, 16), and 0.2–1.2 in ferrets in New Zealand (17).

For comparison, the basic reproduction number for human TB in England and Wales dropped from ≈ 3 in 1900 to 2 by 1950 and fell to <1 in ≈ 1960 , with the net reproduction number remaining at ≈ 1 from 1900 to 1950 (18).

There are many possible extensions of the above analysis. For example, spatial aspects can be represented by having a larger discrete set of intercommunicating systems, each of the above type supplemented by some possibility of transmission of disease from one site to another. There are no mathematical difficulties in analyzing such a model; we have not done so because of the absence of the data needed to specify the numerical values of the necessary parameters. Detail on variation in testing frequency in different regions also could be analyzed simultaneously in a similar set of expanded transmission equations.

By using analytical solutions to relatively simple transmission models, this analysis provides clear estimates of the net reproduction number of bovine TB in Great Britain. This number can be used to predict the impact of additional control measures such as increased testing frequency (whether on a parish or regional basis or through the imposition of additional restrictions on cattle herds with overdue skin tests) and/or increased test sensitivity (whether through the improved application of the tuberculin skin test, wider use of the so-called “severe interpretation” of the skin test, or parallel use of other diagnostic tools such as the γ -IFN test).

Although useful insight can certainly be gained from more complex, geographically specific models such as that used to analyze data on cattle movements (19) or to simulate the impact of badger-control strategies on TB incidence in cattle (20–22), the key benefit of models such as those presented here is the logical transparency and relative simplicity of the results. This approach is in keeping with the preference for simpler model formulations, in the face of parameter and other uncertainties, expressed by the Royal Society Report *Infectious Diseases of Livestock* (23). Clearly, as more data become available, the model can be extended.

1. Anderson, R. M. & May, R. M. (1992) *Infectious Diseases in Man* (Oxford Univ. Press, Oxford).
2. Department for Environment, Food and Rural Affairs (2005) *The Incidence of TB in Cattle—Great Britain*. Available at <http://statistics.defra.gov.uk/esg/statnot/tbpn.pdf>. Accessed September 21, 2005.
3. Krebs, J. R., Anderson, R. M., Clutton-Brock, T., Morrison, W. I., Young, D. & Donnelly, C. A. (1997) *Bovine Tuberculosis in Cattle and Badgers* (MAFF, London) No. PB3423.
4. Krebs, J. R., Anderson, R. M., Clutton-Brock, T., Donnelly, C. A., Frost, S., Morrison, W. I., Woodroffe, R. & Young, D. (1998) *Science* **279**, 817–818.
5. Dunnet, G. M., Jones, D. M. & McInerney, J. P. (1986) *Badgers and Bovine Tuberculosis* (HMSO, London).
6. Zuckerman, S. (1980) *Badgers, Cattle and Tuberculosis* (HMSO, London).
7. Bourne, F. J., Donnelly, C. A., Cox, D. R., Gettinby, G., McInerney, J. P., Morrison, W. I. & Woodroffe, R. (1998) *First Report of the Independent Scientific Group on Cattle TB* (MAFF, London), No. PB3881.
8. Bourne, F. J., Donnelly, C. A., Cox, D. R., Gettinby, G., McInerney, J. P., Morrison, W. I. & Woodroffe, R. (1999) *Second Report of the Independent Scientific Group on Cattle TB* (MAFF, London), No. PB4870.
9. Bourne, F. J., Donnelly, C. A., Cox, D. R., Gettinby, G., McInerney, J. P., Morrison, W. I. & Woodroffe, R. (2001) *Third Report of the Independent Scientific Group on Cattle TB* (Department for Environment, Food and Rural Affairs, London), No. PB5497.
10. Bourne, F. J., Donnelly, C. A., Cox, D. R., Gettinby, G., McInerney, J. P., Morrison, W. I. & Woodroffe, R. (2004) *Fourth Report of the Independent Scientific Group on Cattle TB* (Department for Environment, Food and Rural Affairs, London).
11. Cox, D. R. & Miller, H. D. (1965) *Theory of Stochastic Processes* (Chapman & Hall, London), p. 348.
12. Costello, E., Egan, J. W. A., Quigley, F. C. & O'Reilly, P. F. (1997) *Vet. Rec.* **141**, 222–224.
13. Morrison, W. I., Bourne, F. J., Cox, D. R., Donnelly, C. A., Gettinby, G., McInerney, J. P. & Woodroffe, R. (2000) *Vet. Rec.* **146**, 236–242.
14. Anderson, R. M. & Trewhella, W. (1985) *Philos. Trans. R. Soc. London B* **310**, 327–381.
15. Roberts, M. G. (1996) *J. Anim. Ecol.* **65**, 451–464.
16. Barlow, N. D. (2000) *J. Anim. Ecol.* **69**, 703–713.
17. Caley, P. & Hone, J. (2005) *J. Appl. Ecol.* **42**, 708–719.
18. Vynnycky, E. & Fine, P. E. M. (1998) *Epidemiol. Infect.* **121**, 309–324.
19. Gilbert, M., Mitchell, A., Bourn, D., Mawdsley, J., Clifton-Hadley, R. & Wint, W. (2005) *Nature* **435**, 491–496.
20. Smith, G. C. (2001) *Tuberculosis* **81**, 51–64.
21. Smith, G. C., Cheeseman, C. L., Clifton-Hadley, R. S. & Wilkinson, D. (2001) *J. Appl. Ecol.* **38**, 509–519.
22. Wilkinson, D., Smith, G. C., Delahay, R. J. & Cheeseman, C. L. (2004) *J. Appl. Ecol.* **41**, 492–501.
23. Royal Society (2002) *Infectious Diseases in Livestock* (Royal Society, London).