## THE MASS DONOR OF SCORPIUS X-1 REVEALED

D. Steeghs

Department of Physics and Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK; ds@astro.soton.ac.uk

AND

J. CASARES

Instituto de Astrofisica de Canarias, Via Lactea, La Laguna E-38200, Santa Cruz de Tenerife, Spain; jcv@ll.iac.es Received 2001 June 13; accepted 2001 July 18

# ABSTRACT

We present the first detection of the mass donor star in the prototype X-ray binary Scorpius X-1. Phaseresolved spectroscopy revealed narrow emission line components from the irradiated secondary star. Radial velocity fits to the Bowen blend emission are used to establish an absolute orbital ephemeris and derive an accurate value for the systemic velocity,  $\gamma = -113.8 \pm 0.6$  km s<sup>-1</sup>. Emission from the irradiated front side of the secondary leads to solid limits to the radial velocity of the mass donor of  $87 \leq K_2 \leq 148$  km s<sup>-1</sup>. In conjunction with an upper limit to the velocity of the primary  $K_1 \leq 53$  km s<sup>-1</sup>, we derive a firm limit on the mass ratio of Sco X-1 of  $q \leq 0.61$ . A likely set of system parameters satisfying the various constraints as well as the recent inclination estimate by Fomalont, Geldzahler, & Bradshaw (2001) is  $K_1 = 40$  km s<sup>-1</sup>,  $K_2 = 133$  km s<sup>-1</sup>, q = 0.30 corresponding to component masses of  $M_1 = 1.4 M_{\odot}$  and  $M_2 = 0.42 M_{\odot}$  for an orbital inclination of 38°.

Subject headings: accretion, accretion disks — stars: individual (Scorpius X-1) — X-rays: binaries

## 1. INTRODUCTION

Scorpius X-1 is the prototype low-mass X-ray binary (LMXB), consisting of a neutron star accreting matter from a Roche lobe-filling companion star. It is the brightest persistent X-ray source in the sky and has been the target of detailed analysis for the last 35 years in all energy bands, from radio to X-rays. An orbital period of 18.9 hr was reported by Gottlieb, Wright, & Liller (1975) from brightness eye estimates on 1068 plates during the interval 1890-1974. The optical *B*-band light curve shows a 0.13 mag amplitude modulation (Augusteijn et al. 1992), which is being interpreted as the visibility of the X-ray heated companion. Radial velocity curves of He II  $\lambda$ 4686 and H I confirmed the orbital period and indicated that the inferior conjunction of the emission line regions are close to the photometric minimum, and hence, they must originate near the compact object (e.g., LaSala & Thorstensen 1985). Bradshaw, Fomalont, & Geldzahler (1999) measured the trigonometric parallax of Sco X-1 using Very Long Baseline Array (VBLA) radio observations and deduce a distance of  $2.8 \pm 0.3$  kpc. Intensive radio observations were also used to measure the properties of the twin radio lobes, which are inclined at an angle of  $44^{\circ} \pm 6^{\circ}$  relative to the line of sight (Fomalont et al. 2001). Despite intensive efforts, spectroscopic signatures of the companion star have not yet been observed in optical-IR data (e.g., Schachter, Filippenko, & Kahn 1989; Bandyopadhyay et al. 1997, 1999). Here we present high-resolution spectroscopy of Sco X-1 which enables us to detect the companion star for the first time.

### 2. OBSERVATIONS AND REDUCTION

The phase-resolved spectroscopy of Sco X-1 was obtained with the 4.2 m William Hershel Telescope on La Palma in combination with the ISIS double-beam spectrograph. The R1200B grating centered at 4601 Å delivered a wavelength coverage of 4150–5050 Å at 0.45 Å per pixel on the 2048 × 4096 pixel EEV CCD detector. A 1."2 slit width resulted in a spectral resolution of 0.84 Å, corresponding to 55 km s<sup>-1</sup> at our central wavelength. The data were part of a campaign to look for circular polarization from accreting binaries, and to this end a quarter-wave plate and a Savart calcite slab were placed in the optical path. (See Tinbergen & Rutten 1992 for more details.) The slit position angle was adjusted to remain close to parallactic throughout the observations. A total of 137 spectra were obtained in the course of three consecutive nights covering 75% of the 18.9 hr orbital cycle. A journal of observations can be found in Table 1.

The insertion of the calcite slab in the light path results in the projection of two target beams on the detector. The intensity ratio then represents the fraction of circular polarization. For the analysis presented in this paper, we ignored the polarization characteristics of the light and combined the extracted spectra from both beams. Frames were first debiased using a median-bias frame obtained by combining 10-15 individual bias exposures. The overscan strip was then used to subtract any residual bias from each frame. A median of 29 Tungsten exposures was constructed at each night in order to perform flat-field correction. Each beam was then optimally extracted. Some frames were affected by transient CCD pickup noise producing variable bias structure at the level of 3-5 ADU. For those frames affected (around 8 on the second night and 10 on the third night), we introduced an additional reduction step before optimal extraction which subtracted the pickup noise using a second-order polynomial fit to each CCD row. Regular CuAr arc lamp exposures allowed us to establish an accurate wavelength scale for each spectrum. Finally, a wide slit exposure of the flux standard BD  $+28^{\circ}4211$  (Oke 1990) was used to perform flux calibration using the MOLLY data

TABLE 1Journal of Observations

Date	UT Interval	Number of Spectra	Exposure Time (s)	Orbital Phase Interval
1999 June 28	21:55-02:10	40	300	0.81-1.03
1999 June 29	20:55-02:15	51	300	0.03-0.31
1999 June 30	20:56-01:55	46	300	0.30-0.56

analysis package. The two extracted spectra for each frame were then added to produce the final set of 137 calibrated Sco X-1 spectra.

### 3. ANALYSIS

## 3.1. The Emission Line Spectrum

The optical spectrum of Sco X-1 is dominated by strong and broad emission lines from the Balmer series as well as He I /He II. However, closer inspection of our spectra revealed a large number of weaker and very narrow emission line features that showed significant Doppler motions as a function of orbital phase. In particular, the strong Bowen blend near 4640 Å was resolved into an underlying broad component, as well as several narrow components from the individual Bowen transitions. Broad and highly variable emission around 4640 Å was already noted by Sandage et al. (1966) in the paper discussing the identification of the optical counterpart of Sco X-1. A more detailed investigation of the same data by Westphal, Sandage, & Kristian (1968) did also indicate complex and strongly timedependent fine structure in the broad emission structures and some preliminary evidence for velocity changes in the strong Balmer and He II lines. More recently, Bowen emission in Sco X-1 was noted by Willis et al. (1980) while Schachter, Filippenko, & Kahn (1989) discuss the Bowen fluorescence process in Sco X-1 in detail. However, neither had the spectral resolution to resolve the individual components. We will see in the next sections that these narrow emission features can be attributed to the irradiated companion star, thus revealing a direct signature from the mass donor in Sco X-1 for the first time. In Figure 1, we present the average normalized spectrum of Sco X-1 in the rest frame of the companion star to facilitate the identification of the narrow features. Most lines were identified using the planetary nebulae emission line lists from Kaler (1976) as well as the Kurucz atomic line list. Many narrow lines from O II are present as well as a few S III, C II, and N III transitions.

#### 3.2. Bowen Emission from the Donor

In order to establish the origin of the narrow emission line components, we measured the velocity of the narrow Bowen components as a function of phase. As can be seen in Figure 1, the strongest Bowen components are the two N III components at 4634.13 Å and 4640.64 Å and the C III components at 4647.4/4650.1 Å. In Figure 2, we display the Bowen blend and He II 4686 emission in the form of a trailed spectrogram. All the individual spectra were first normalized to the continuum using a third-order polynomial fit. The two N III components and the C III component at 4647 Å clearly move in phase with each other and are extremely narrow. In the background, more diffuse emission can be seen to move roughly in antiphase with these sharp components. For comparison, the broad and weakly double-peaked He II line shows completely different line kinematics.

The velocities of the various components were measured using multiple Gaussian fits to the individual spectra. At first, we fitted the Bowen profiles using three narrow Gaussians corresponding to the three strongest Bowen components, as well as an underlying broad component. The FWHM of the Gaussians were determined from a fit to the average Bowen blend profile and fixed to 80 km s<sup>-1</sup> for the narrow components and 1550 km s<sup>-1</sup> for the broad component. The position and strength for each Gaussian was a free parameter for each individual spectrum. The derived radial velocities of the sharp components were identical within the error bars. We thus decided to reduce the number of free parameters in our fits and give the three narrow Gaussians a common radial velocity but an independent strength. The six parameters were optimized for each observed spectrum through least-square fitting to determine the common



FIG. 1.—Average normalized spectrum of Sco X-1 in the rest frame of the donor star. Three panels are on the same intensity scale with the strong He II  $\lambda$ 4686 and H $\beta$  going off-scale in order to highlight the weaker lines. Peak strength in He II is around 1.46 compared to the continuum, whereas it is 1.39 for H $\beta$ .



FIG. 2.—Trailed spectrogram of the Bowen blend and He II  $\lambda$ 4686. Three sharp components moving in phase with each other stand out from the broad Bowen blend emission. Small gaps separate the data from the three nights.

velocity of the three narrow components and its associated formal error.

Figure 3 plots the derived radial velocity curve of the Bowen components together with a least-square sinusoidal fit. The best fit gives a velocity amplitude of  $77.2 \pm 0.4$  km  $s^{-1}$  with a mean velocity of  $-113.8 \pm 0.6$  km  $s^{-1}$ . Given the extreme narrowness of the components, they cannot originate from the accretion flow or hot spot that is expected to be responsible for the broad emission line components. Such narrow emission components are, however, regularly seen in the line profiles of cataclysmic variables where they originate from the irradiated front side of the companion star (e.g., Harlaftis et al. 1999; Schwope et al. 2000). This has also been observed in the intermediate mass X-ray binary Her X-1 (Still et al. 1997), where narrow Bowen blend components from the mass donor are present. An origin on the mass donor is furthermore supported by the fact that the crossing of the emission components from blueshift to redshift occurs near minimum light and that the strength of the emission is weakest then, since we are looking at the unirradiated backside of the secondary. The inferred systemic velocity of  $\gamma = -113.8 \pm 0.6$  km s<sup>-1</sup> furthermore agrees well with previous estimates using the strong emission lines (e.g., Lasala & Thorstensen 1985). We thus have established a radial velocity curve of the mass donor in Sco X-1 for the first time. The radial velocity curve directly gives us an unbiased value for the systemic velocity of the binary and establishes the correct absolute orbital phases. From the sinusoidal fit, we established the following orbital ephemeris, which we will use throughout this paper:

$$T_0(\text{HJD}) = 2451358.568(3) + 0.787313(1)E ,$$

where we took the value for the orbital period as derived by



FIG. 3.—*Top*: radial velocity curve of the narrow Bowen blend emission components as derived from a multiple Gaussian fit. Data are repeated over two cycles. Best sinusoidal fit is over plotted as a solid line. Orbital phase of the binary is calculated using the best fit radial velocity curve to determine the zero point. *Bottom*: Radial velocity curve of the He II emission line based on a double Gaussian fit with separation 600 km s<sup>-1</sup> and FWHM of 100 km s<sup>-1</sup>. Sinusoidal fit is over plotted indicating a phase shift of 0.6 with respect to the donor star emission.

Gottlieb, Wright, & Liller (1975). Using our ephemeris, the phase of minimum light as established by Gottlieb, Wright, & Liller corresponds to an absolute phase of  $\phi = 0.04 \pm 0.04$ , where we have allowed for the uncertainties in both  $T_0$  and P. This confirms that the photometric modulation is indeed related to the X-ray heated hemisphere of the donor star. Values of  $T_0$  based on the inferior conjunction of the He II  $\lambda$ 4686 emission, on the other hand, are delayed by  $0.10 \pm 0.02$  (Cowley & Crampton 1975) and  $0.08 \pm 0.02$  (LaSala & Thorstensen 1985), indicating that the emission line regions trail the compact object. We will revisit these emission line velocities in § 3.3.

Since we have detected narrow lines from the mass donor in emission rather than absorption, they must originate from the front half of the Roche lobe. The derived radial velocity amplitude is thus biased toward lower values, depending on the detailed distribution of the line emission over the Roche lobe surface. A similar bias toward larger velocities occurs when using absorption lines that are predominantly formed near the backside of the Roche lobe (e.g., Wade & Horne 1988). The fact that the radial velocity curve is not strictly sinusoidal also suggests that the emission is not distributed uniformly. However, allowing for an ellipticity in the fit does not significantly improve the  $\chi^2$  of the fit. We can thus constrain the true velocity of the center of mass of the mass donor ( $K_2$ ) to be  $K_2 > 77.2$  km s<sup>-1</sup>. Without any knowledge of the mass ratio and the extent of disk shadowing effects in Sco X-1, it is very difficult to quantify the K-correction that needs to be applied to the Bowen blend velocities in order to define the true  $K_2$ . A more detailed modeling, in terms of both velocity and strength, of the phase dependence of the emission from the mass donor is required and will be discussed in a future paper.

## 3.3. The Radial Velocity of the Primary

Direct inspection of the trailed spectra (Fig. 2) clearly shows that the He II  $\lambda$ 4686 line is antiphased with the narrow Bower blend components, and therefore, it must come

Vol. 568

from near the compact object. Significant phase shifts between the line core and centroid, due to possible contamination by complex variable hot spots, have been reported (e.g., Cowley & Crampton 1975). Therefore, we have followed the double-Gaussian technique to estimate the velocity semi-amplitude of the compact object  $K_1$  from the line wings. (See Schneider & Young 1980 for details.) We employed a 2 Gaussian bandpass with FWHM= 100 km  $s^{-1}$  and Gaussian separations a = 200-1000 km  $s^{-1}$  in steps of 100 km s<sup>-1</sup>. Radial velocity curves of different line sections yield consistent results with velocity semi-amplitudes in the range 50–56 km s<sup>-1</sup> and blue-to-red crossing phases 0.64–0.60. Phase delays of 0.1–0.2 relative to the expected motion of the compact object are typically observed in cataclysmic variables (CVs), probably due to residual hot spot contamination. We have also attempted to extrapolate the velocity points toward phase 0.5 using the light center technique (see Marsh 1988 for details), but the method fails given the very small derived phase drift as we move from the line core into the wings. Applying the same technique to the  $H\beta$  line produces similar values for the line core; however, the phase offset increases dramatically in the line wings. We therefore choose to use the He II lines as our basis for an estimation of  $K_1$ . The bottom panel of Figure 3 presents the radial velocity curve of He II  $\lambda$ 4686 for a = 600 km s<sup>-1</sup>, before velocity points start to be corrupted by continuum noise. A formal sine wave fit provides  $K = 53 \pm 1$  km s<sup>-1</sup>, which we take as an upper limit to the true  $K_1$ .

## 3.4. Doppler Mapping

Many of the narrow lines that can be identified in Figure 1 move in phase with the secondary star and thus are also the result of reprocessed emission from the irradiated front side of the Roche lobe. Unfortunately, due to weakness of the lines, a Gaussian fit to each spectrum is not possible. Instead, we use Doppler tomography (Marsh & Horne 1988) to measure the radial velocity amplitudes of the many weak narrow emission lines that are present. By mapping the observed data onto a velocity coordinate frame, Doppler mapping makes use of all data at once and can thus be used for features that are too weak to be separated in each individual spectrum. One effectively resolves the distribution of line emission in the corotating frame of the binary system, providing an excellent tool for identifying the origin and kinematics of the various emission components. It also allows us to map the emission line distribution from the broad He I, He II, and Balmer lines. Secondary star emission is readily identified in Doppler tomograms since the solid body rotation of the Roche lobe is mapped to a corresponding Roche lobe area along the positive  $V_y$ -axis. We employed a maximum entropy implementation of Doppler tomography whereby the data are fitted under the added constraint of maximizing the entropy, i.e., image smoothness, of the tomogram (Marsh & Horne 1988). This reduces the presence of noise artifacts in the recovered tomograms and allows a simultaneous fit to a number of heavily blended lines.

To check that the method of measuring the velocity of the narrow emission lines using tomography is consistent with the method of employing (multiple) Gaussian fits, we started by producing tomograms for the Bowen blend components. The Bowen blend data were fitted simultaneously with three Doppler images corresponding to the rest wavelengths of the three strongest Bowen emission components. The tomograms revealed the secondary star emission as a sharp spot along the  $V_{y}$ -axis, and a two-dimensional Gaussian fit was used to accurately measure the position of the spot. This lead to inferred velocities of  $76 \pm 1$ ,  $76 \pm 1$ , and  $77 \pm 1$  km s<sup>-1</sup>, respectively, confirming that this method can be used to extract reliable radial velocities. We then repeated the same procedure for a range of other weak emission lines present in the spectrum of Sco X-1. The inferred velocities for emission from the mass donor spanned from 66 km s<sup>-1</sup> for Si III at 4568 Å, 78 km s<sup>-1</sup> for O II doublet at 4415 Å, to 87 km s<sup>-1</sup> for N III at 4379 Å. An example tomogram showing secondary star emission from the N III line at 4514 Å is plotted in Figure 4. The fitted velocity based on this tomogram gives 79 km s<sup>-1</sup>. Unfortunately, the width of the narrow components are of the order of our instrumental resolution and so are not well resolved. However, since only the front half of the Roche lobe can contribute to the line emission, we can confidently increase our lower limit to  $K_2 > 87$  km s<sup>-1</sup>. This technique, in conjunction with higher spectral resolution data, could be used to map the emission on the Roche lobe as a function of ionization potential, such as was done in, for example, Harlaftis (1999), and provide even more stringent constraints to the true value of  $K_2$ .

In Figure 4, we also present Doppler tomograms of the strong and broad H $\beta$ , He II  $\lambda$ 4686, and He I  $\lambda$ 4921 lines for comparison. Clearly, their line dynamics are radically different and are dominated by contributions from the accretion disk around the primary. However, significant asymmetries are present in all lines, leading to localized hot spots in the tomograms. In the case of He II at 4686 Å, we can identify a clear emission contribution from the secondary star, whereas the Balmer and He I lines do not show a secondary star emission component. If anything, the H $\beta$  tomogram indicates a depression of line flux at the location of the mass donor. The tomograms directly illustrate that the use of emission line velocities as a measure of the radial velocity amplitude of the compact object is severely affected and distorted by asymmetries in the line profiles. Indeed, our analysis in § 3.3 revealed significant phase offsets between the phasing of the secondary star components and the broad emission lines. The different phase offsets that exist between the velocities of the H $\beta$  line a versus He II are due to the fact that the He II is mainly distorted by emission from the mass donor, whereas the H $\beta$  line is distorted by a spot in the lower left quadrant of the Doppler image. The fact that our data reveal the same phase shifts as found by Lasala & Thorstensen (1985) suggests this asymmetry is a persistent feature in Sco X-1.

The complex morphology of the Doppler tomograms complicates the reliable estimation of  $K_1$ . If the emitting gas was symmetrically distributed around the primary, the Doppler image would reveal a circular distribution centered on  $V_x = 0$ ,  $V_y = -K_1$ . In order to estimate the true center of the emission line distribution, we subtracted the symmetrical component from the H $\beta$  Doppler maps and inspected the residuals. We cycled the assumed center of symmetry between -100 and +25 km s<sup>-1</sup> in steps of 2 km s<sup>-1</sup> and measured the mean of the residuals at high velocities in the upper right quadrant, which is not affected by the spot. The mean residuals show a pronounced minimum around an assumed center of symmetry of  $V_y \sim -40$  km s<sup>-1</sup>. This suggests that the true value of  $K_1$  is indeed closer to 40 km s<sup>-1</sup> for Sco X-1.



FIG. 4.—Doppler tomograms reconstructed from the H $\beta$ , He I  $\lambda$ 4921, He II  $\lambda$ 4686 and N III  $\lambda$ 4514 lines using maximum entropy optimization. Both gray scale as well as contours span the full scale of the maps. Origin corresponds to the center of mass of the binary system using  $\gamma = -113.8$  km s<sup>-1</sup>.

## 4. DISCUSSION

We have presented the first direct detection of the mass donor in Sco X-1 by resolving emission components originating from the irradiated front side of its Roche lobe. Radial velocity fits were used to establish an absolute orbital ephemeris and measure the systemic velocity to high accuracy,  $\gamma = -113.8 \pm 0.6$  km s<sup>-1</sup>. A lower limit to the radial velocity of the mass donor is given by the radial velocity amplitude of the sharp Bowen blend emission components, which leads to  $K_2 > 77$  km s<sup>-1</sup>. Detection of secondary star emission at even higher velocities in other transitions leads to a more stringent lower limit of  $K_2 > 87$  km s<sup>-1</sup>. The strong broad emission lines show orbital motions roughly in anti-phase with the secondary star. We confirm significant phase shifts between the radial velocity curves of the strong lines and the absolute ephemeris. Doppler tomography reveals that these phase shifts are due to persistent asymmetries in the emission line distribution around the compact object. The formal double Gaussian method leads to a firm upper limit to the primary radial velocity amplitude of  $K_1 < 53$  km s<sup>-1</sup>. A more reliable value is derived by searching for the optimal center of symmetry in the accretion disk emission, which leads to an estimated  $K_1$  value of  $40 \pm 5$  km s<sup>-1</sup>. Our absolute ephemeris confirms previous suggestions that the phase of optical minimum light corresponds closely to orbital phase zero, when the contribution from the heated mass donor is smallest.

The combination of our upper limit to  $K_1$  and at the same time a lower limit to  $K_2$  leads to a firm limit to the mass ratio of  $q = M_2/M_1 < 0.69$  when using the formal values of  $K_1 < 53$  and  $K_2 > 77$ , as derived from the Gaussian fits. Using the more restrictive values  $K_1 \sim 40$ ,  $K_2 > 87$ , we derive a limit of  $q \leq 0.46$ . Another constraint is given by the fact that the observed emission originates from the front half of the Roche lobe and its velocity thus lies in between that of the true  $K_2$  value and the velocity of the L1 point ( $V_{L1}$ ). Since we observe emission at velocities as low as 66 km s<sup>-1</sup>, we can thus derive an upper limit on  $K_2$  for a given mass ratio in order to make sure that the velocity of L1 is not greater than 66 km s<sup>-1</sup>. In Figure 5, we plot the various limits on  $K_2$  as a function of mass ratio. As can be seen in



FIG. 5.—Plotted are constraints to the value of  $K_2$  as a function of mass ratio. Horizontal solid line is the lower limit to  $K_2$  as derived from the observed line emission ( $K_2 > 87$ ). Dot-dashed lines denote the upper limits on  $K_2$  given that  $K_2 < K_1/q$  for  $K_1 = 53$  (upper curve) and  $K_1 = 40$  km s<sup>-1</sup> The requirement that the velocity of the L1 point may not exceed 66 km s<sup>-1</sup> leads to the dashed curve. Two dotted curves denote the values of  $K_2$  if the primary is a 1.4  $M_{\odot}$  neutron star. The two curves correspond to  $\pm 1 \sigma$  limits for the orbital inclination based on the orientation of the radio lobes  $(44 \pm 6^{\circ}).$ 

the plot, the combination of  $V_{L1} < 66$  and  $K_2 < K_1/q$  leads to  $K_2 < 148$  km s<sup>-1</sup>, regardless of the mass ratio. This upper limit proves interesting from the point of view of the recent inclination estimate based on the active radio lobes of Sco X-1 (Fomalont et al. 2001). If the orientation of the radio lobes corresponds to the rotation axis of the orbital plane, we can estimate the component masses using Kepler's law. With a typical neutron star mass of  $M_1 \sim 1.4 \, M_{\odot}$ , this inclination (44  $\pm$  6°) predicts relatively high  $K_2$  values across the allowed mass ratio range (Fig. 5). This suggests that a slightly lower orbital inclination seems more likely, unless the neutron star is atypical and significantly less massive than the Chandresekhar limit. However, given the uncertainty on the inclination, all constraints are still compatible with a 1.4  $M_{\odot}$  neutron star within 1  $\sigma$ . A likely set of parameters for Sco X-1, given the various constraints, could be

- Augusteijn, T., et al. 1992, A&A, 265, 177

- L121 Cowley, A. P., & Crampton, D. 1975, ApJ, 201, L65 Fomalont, E. B., Geldzahler, B. J., & Bradshaw, C. F. 2001, ApJ, 558, 283 Gottlieb, E. W., Wright, E. L., & Liller, W. 1975, ApJ, 195, L33 Harlaftis, E. T. 1999, A&A, 346, 73 Harlaftis, E. T., et al. 1999, MNRAS, 306, 348 Howell, S. 2001, PASJ, 53, 675 Kaler, I. B. 1976, ApJS, 31, 517 L121

- Kaler, J. B. 1976, ApJS, 31,517
- King, A. R., Kolb, U., & Burderi, L. 1996, ApJ, 464, L127
- LaSala, J., & Thorstensen, J. R. 1985, AJ, 90, 207

 $K_1 = 40 \text{ km s}^{-1}, K_2 = 133 \text{ km s}^{-1}, q = 0.30$ , corresponding to component masses of  $M_1 = 1.4~M_{\odot}, M_2 = 0.42~M_{\odot}$  for an orbital inclination of 38°. Given the orbital period of Sco X-1, the companion star must then be a significantly evolved subgiant having a radius of more than twice of the corresponding main-sequence radius of  $\sim 0.5 R_{\odot}$ . Its mean density is solely determined by the orbital period, which gives  $\rho = 0.31 \text{g cm}^{-3}$  compared to ~5.6 g cm<sup>-3</sup> for a 0.42  $M_{\odot}$ main-sequence star. We note that these parameters are also consistent with Sco X-1 being a persistent X-ray binary with an approximate mass transfer rate of  $\dot{M}_2 \sim 8 \times 10^{-10} \ M_{\odot} \ {\rm yr}^{-1}$  using the prescription from King, Kolb, & Burderi (1996). This is enough to maintain the irradiated disk hotter than 6500 K everywhere, avoiding global instabilities and, thus, outbursts.

Bandyopadhyay et al. (1997, 1999) fail to detect any features from the companion star in their IR spectra of Sco X-1 at two epochs. Most notably, the CO band head near 2.2  $\mu$ m, which is present in late-type stars, is absent. However, the appearance of the donor star will be affected by irradiation, and both epochs were taken near orbital phase 0.5 (using our derived ephemeris), when these effects would be particularly severe. In addition, the use of the CO bands for spectral type classification can be misleading, because their strength is strongly dependent on effective temperature and, furthermore, the CO abundances can be significantly nonsolar for companion stars in interactive binaries (e.g., Howell 2001). In order to derive more accurate system parameters, we need to be able to model the K-correction and use higher spectral resolution data to resolve the emission across the Roche lobe for a range of transitions using the methods described in this paper. Detecting the companion star in absorption should be possible around true orbital phase zero, when the effects of X-ray heating are minimal.

D. S. is supported by a PPARC Fellowship. Use of the MOLLY and DOPPLER software developed by T. R. Marsh is gratefully acknowledged. The William Hershel Telescope is operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias.

## REFERENCES

- Marsh, T.R. 1988, MNRAS, 231, 1117 Marsh, T. R., & Horne, K. 1988, MNRAS, 235, 269 Oke, J. B. 1990, AJ, 99, 1621 Sandage, A. R., et al. 1966, ApJ, 146, 316 Schachter, J., Filippenko, A. V., & Kahn, S. M. 1989, ApJ, 340, 1049 Schneider, D. P., & Young, P. 1980, ApJ, 238, 946 Schwope, A. D., et al. 2000, MNRAS, 313, 533 Still, M. D., et al. 1997, MNRAS, 292, 52 Tinbergen, J., & Rutten, R. 1992, La Palma ING User Manu

- Tinbergen, J., & Rutten, R. 1992, La Palma ING User Manual 21, (La Palma: Isaac Newton Group)
- Wade, R. A., & Horne, K. 1988, ApJ, 324, 411 Westphal, J. A., Sandage, A., & Kristian, J. 1968, ApJ, 154, 139
- Willis, A. J., et al. 1980, ApJ, 237, 596