

Near-horizon Bondi-Metzner-Sachs symmetry, dimensional reduction, and black hole entropy

S. Carlip **Department of Physics, University of California, Davis, California 95616, USA*

(Received 8 November 2019; accepted 8 January 2020; published 5 February 2020)

In an earlier short paper [*Phys. Rev. Lett.* **120**, 101301 (2018)], I argued that the horizon-preserving diffeomorphisms of a generic black hole are enhanced to a larger three-dimensional Bondi-Metzner-Sachs symmetry, which is powerful enough to determine the Bekenstein-Hawking entropy. Here, I provide details and extensions of that argument, including a loosening of horizon boundary conditions and a more thorough treatment of dimensional reduction and meaning of a “near-horizon symmetry.”

DOI: [10.1103/PhysRevD.101.046002](https://doi.org/10.1103/PhysRevD.101.046002)

I. INTRODUCTION

The discovery by Bekenstein [1] and Hawking [2] that black holes are thermodynamic objects has led to a host of fascinating puzzles, from the information loss problem to the question of what microscopic states are responsible for black hole entropy. Here, I focus on one particular puzzle, the “problem of universality” of black hole entropy. This paper is an expanded version of a short article published in 2018 [3]; here, I discuss details and extend some of the results.

The universality of black hole entropy has two aspects, probably related but logically distinct. The first comes from the simple form of the Bekenstein-Hawking entropy,

$$S_{\text{BH}} = \frac{A_{\text{hor}}}{4G\hbar}, \quad (1.1)$$

where A_{hor} is the horizon area. If this were merely a property of, say, uncharged static black holes, it would tell us something important about the Schwarzschild solution. But it is more. The same area law, with the same coefficient, holds for black holes with any charges, any spins, in any dimensions; it holds for black strings, black rings, black branes, and black Saturns (black holes encircled by black rings); it remains true for “dirty black holes” whose horizons are distorted by nearby matter. The only known way to change the entropy (1.1) is to change the Einstein-Hilbert action, and even then the correction will be another

universal term [4]. Black hole entropy is not, it seems, a property of specific solutions, but rather a generic characteristic of horizons.

The second aspect of universality emerges when one attempts to identify the microscopic states responsible for this entropy. We do not yet have a full description of those states; that would presumably require a complete quantum theory of gravity. We do, however, have an assortment of research programs working toward the quantization of gravity, which allow partial computations of black hole entropy. In string theory, black hole entropy can be calculated from properties of weakly coupled strings and branes, from the AdS/CFT correspondence, and (probably) from an enumeration of horizonless “fuzzball” configurations. In loop quantum gravity, entropy can be calculated from a horizon Chern-Simons theory, from an analysis of spin network reconstructions in the interior, and from conformal field theory at “punctures” of the horizon. In induced gravity—an approach in which the Einstein-Hilbert action is obtained by integrating out “heavy” fields in the path integral—entropy can be calculated from the properties of the heavy fields. In semiclassical gravity, entropy can be calculated from either a single instanton approximation or pair production. And, of course, entropy can be calculated using Hawking’s original approach, which involved only quantum field theory in a fixed black hole background.

None of these methods is complete. String theory calculations, for instance, are cleanest for near-extremal black holes, while loop quantum gravity calculations may depend on a new universal constant, the Barbero-Immirzi parameter. But although they describe very different microstates, each of these methods, within its range of validity, reproduces the standard Bekenstein-Hawking entropy. (For a review and further references, see Ref. [5]).

*carlip@physics.ucdavis.edu

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP³.

One might worry about a selection effect here: perhaps models that give the “wrong” entropy are less likely to be published. But even the elegant analysis of BPS black holes in string theory [6], the first really successful microscopic calculation of black hole entropy, illustrates the problem. Given a spacetime dimension and a set of charges and spins, one can calculate the entropy of a gas of strings and branes at weak coupling, separately calculate the horizon area of a black hole at strong coupling, and compare the results. But although the final answer always matches the Bekenstein-Hawking area law (1.1), each new choice of dimension, spins, and charges requires a new computation. Some underlying structure is clearly missing.

A first guess for this deeper structure is that the relevant degrees of freedom live on the horizon [1]. But this is not enough; while it could explain an area law for black hole entropy, there is no obvious reason why the coefficient $1/4$ should be universal. An elaboration of this idea, first suggested (I believe) in Ref. [7], is that the entropy is governed by a horizon symmetry. This is, of course, a very strong requirement; symmetries can place some restrictions on the density of states, but they are rarely strong enough to actually determine the entropy. But we know one symmetry that has the same kind of universal properties we see in black hole entropy. As Cardy first showed in 1986 [8,9], two-dimensional conformal symmetry is so restrictive that it completely fixes the asymptotic density of states in terms of a few parameters, independent of any of the fine details of the theory. More recently, it has been shown that a related symmetry, that of the three-dimensional Bondi-Metzner-Sachs group (BMS_3), exhibits the same universality [10]. The possibility of a connection with black hole entropy has obvious appeal.

This connection was first confirmed for the $(2+1)$ -dimensional BTZ black hole in 1998 [11,12]. Attempts to extend those results to higher dimensions soon followed [13,14]. These efforts, which typically involve a search for a suitable two-dimensional group of horizon symmetries, have had significant success; see Ref. [15] for a review. But they have been plagued by several problems:

- (i) The symmetries are typically located either at infinity or on a timelike “stretched horizon” just outside the actual horizon (although with occasional exceptions [16]). The physics at infinity is extremely powerful, especially for asymptotically anti-de Sitter spaces. Indeed, the BTZ black hole calculations were among the first examples of the now famous AdS/CFT correspondence. But the symmetries alone are not enough; by themselves, for instance, they cannot distinguish a black hole from a star with the same mass.

The stretched horizon more directly captures the local properties of the black hole. But the definition of the stretched horizon is not unique, and different limits can lead to different entropies [17,18]. Moreover,

while the entropy has a well-defined limit at the horizon, other parameters in the symmetry algebra typically blow up at the horizon [19–21] (again with occasional exceptions [22]).

- (ii) The approach fails in what should be the simplest case, two-dimensional dilaton gravity, where the zero-dimensional boundary of a Cauchy surface simple does not have “room” for the required central term in the conformal algebra. There are *ad hoc* fixes—lifting the theory to three dimensions [23] or artificially introducing an integral over time [24]—but none of them is very convincing.
- (iii) In higher dimensions, the relevant symmetries are those of the “ $r-t$ plane” picked out by the horizon. But to obtain a well-behaved symmetry algebra, one must introduce an extra *ad hoc* dependence on angles, with no clear physical justification. This is especially problematic for the Schwarzschild black hole, for which the angular dependence breaks spherical symmetry in a manner that seems quite arbitrary.

Here (and in a shorter form in Ref. [3]), I describe an approach that avoids these problems. A basic limitation of past work, I argue, was the attempt to force the horizon symmetry into the form of a two-dimensional conformal symmetry. This was an understandable choice; until quite recently, this was the only symmetry known to be powerful enough to determine the asymptotic density of states. But with the discovery that BMS_3 symmetry also has this universal property, the possibilities have expanded.

Starting with the intuitive idea that the relevant symmetries should lie in the $r-t$ plane picked out by the null generators of the horizon [13,19,22], I first show how to reduce the problem to an effective two-dimensional model. For spherically symmetric black holes, such a dimensional reduction already appeared in some of the earliest work on horizon symmetries [14,25], but I demonstrate that the relevant near-horizon properties are far more general. I next establish that the obvious horizon symmetries, the horizon-preserving diffeomorphisms, are enhanced by a particular shift invariance, as anticipated in Refs. [26,27]. This new symmetry may be viewed as a generalization of the global conformal symmetry found by Wall for horizon quantum field theory on a fixed background [28]; it is exact at the horizon, and, in a sense I explain, it can be made arbitrarily close to exact near the horizon. Using covariant phase space methods [19,29], I show that the generators of these symmetries can be expressed as integrals along the horizon, with no need to go to a stretched horizon. Finally, I confirm that the resulting generators satisfy a centrally extended BMS_3 algebra that determines the correct Bekenstein-Hawking entropy.

II. BMS_3 SYMMETRY

Let us start with a brief review of BMS_3 symmetry, along with a discussion of the perhaps puzzling question of how a

classical symmetry can determine the number of quantum states.

The BMS₃ algebra is described by two sets of generators L_n and M_n ($n \in \mathbb{Z}$) with Poisson brackets

$$\begin{aligned} i\{L_m, L_n\} &= (m-n)L_{m+n}, \\ i\{M_m, M_n\} &= 0, \\ i\{L_m, M_n\} &= (m-n)M_{m+n} + c_{LM}m(m^2-1)\delta_{m+n,0}, \end{aligned} \quad (2.1)$$

where the possibility of a classical central charge c_{LM} was first noted by Barnich and Compere [30].¹ This algebra can be obtained as a contraction of the usual two-dimensional conformal (Virasoro) algebra [30], and is also isomorphic to the two-dimensional Galilean Conformal Algebra [31]. While the BMS₃ algebra is not as thoroughly studied as the conformal algebra, a fair amount is understood about its properties and representations [32–36].

We shall see below that this algebra describes the classical horizon symmetry of a generic black hole. Let us assume that the same symmetry, perhaps deformed, is realized in the quantum theory. We make the usual substitutions

$$\begin{aligned} \{\bullet, \bullet\} &\rightarrow \frac{1}{i\hbar}[\bullet, \bullet], \\ \frac{1}{\hbar}L &\rightarrow \hat{L}, \quad \frac{1}{\hbar}M \rightarrow \hat{M}, \quad \frac{1}{\hbar}c \rightarrow \hat{c}, \end{aligned} \quad (2.2)$$

where the factors of \hbar in \hat{L} and \hat{M} ensure that the operators are dimensionless. (It is not always discussed explicitly in quantum mechanics textbooks, but the same substitution is used to go from the Poisson algebra of angular momentum to the Lie algebra of rotations.) We thus obtain a quantum operator algebra,

$$\begin{aligned} [\hat{L}_m, \hat{L}_n] &= (m-n)\hat{L}_{m+n}, \\ [\hat{M}_m, \hat{M}_n] &= 0, \\ [\hat{L}_m, \hat{M}_n] &= (m-n)\hat{M}_{m+n} + \hat{c}_{LM}m(m^2-1)\delta_{m+n,0}. \end{aligned} \quad (2.3)$$

Classical values of the zero modes L_0 and M_0 now become eigenvalues $h_L = L_0/\hbar$, $h_M = M_0/\hbar$ of the corresponding operators. The true quantum symmetry may be a deformation of (2.3)—other central terms may appear, for example—but differences will be suppressed by factors of \hbar .

Now, it is well known that for a theory with a two-dimensional conformal symmetry, the central charge completely fixes the asymptotic behavior of the density of states [8,9]. For the simplest case of free bosons and fermions, the Cardy formula for the density of states is just the Hardy-Ramanujan formula for partitions of an

integer [37]. For the general case, I know of no elementary explanation; for a careful but not terribly intuitive derivation, see Ref. [38]. Roughly speaking, exact conformal symmetry is powerful enough to prevent any exponential growth in the number of states, which can occur only because of the anomalous symmetry breaking characterized by the central charge c .

The BMS₃ symmetry (2.3) is not quite a conformal symmetry, but as Bagchi *et al.* have shown [10], it has its own version of the Cardy formula for the asymptotic density of states. In hindsight, this is not so surprising, since BMS₃ can be obtained as a contraction of the two-dimensional conformal algebra. The resulting entropy—the logarithm of the density of states at fixed eigenvalues h_L and h_M —has the asymptotic behavior

$$S \sim 2\pi h_L \sqrt{\frac{\hat{c}_{LM}}{2h_M}} = \frac{2\pi}{\hbar} L_0 \sqrt{\frac{c_{LM}}{2M_0}}, \quad (2.4)$$

where L_0 , M_0 , and c_{LM} in the last equality are the classical values. Note that the factors of Planck's constant combine to give an overall $1/\hbar$, an expected feature of an entropy described in terms of a classical phase space.

III. REDUCTION TO TWO DIMENSIONS

The first step in our derivation of black hole entropy will be to reduce the problem to two dimensions. To understand this process, it is helpful to start with a rather elaborate description of an ordinary Schwarzschild black hole in D spacetime dimensions. The horizon Δ of such a black hole is a Killing horizon, that is, a null $(D-1)$ -manifold whose null normal coincides with a Killing vector χ . The integral curves of χ on Δ are null geodesics, the generators of Δ . Since χ is timelike outside the horizon, it determines a preferred time coordinate, and through that a foliation of Δ by $(D-2)$ -spheres $\hat{\Delta}$ of constant time.

This structure allows us to identify two “preferred” directions at any point p on Δ : the direction of the Killing vector at p and the outward radial direction transverse to Δ and normal to the slice $\hat{\Delta}$ containing p . These determine a local “ r - v plane,” where r is a radial coordinate and v is a parameter along the null generators of the horizon. If we further choose coordinates y^μ on $\hat{\Delta}$, then (v, r, y^μ) can be extended to form a Gaussian null coordinate system near the horizon (see, for instance, Appendix A of Ref. [39]). For the Schwarzschild case, these coordinates are essentially Eddington-Finkelstein coordinates, with r shifted so that $r=0$ at the horizon. By spherical symmetry, each slice $\hat{\Delta}$ at constant (v, r) is invariant under rotations, so standard Kaluza-Klein methods can reduce the Einstein-Hilbert action to that of two-dimensional dilaton gravity [40].

¹Other central elements can also be added, but only c_{LM} is relevant in the present context.

The reason I have given such a complicated description of a relatively simple procedure is that most of the steps generalize quite broadly. Let Δ be a nonexpanding horizon [41,42], that is, a null $(D-1)$ -dimensional manifold with null normal ℓ such that:

- (1) Δ has the topology $(0,1) \times \hat{\Delta}$, where $\hat{\Delta}$ is usually taken to be compact (a sphere for a black hole, a torus for a black ring, etc.).
- (2) The expansion $\theta^{(\ell)}$ of the null normal vanishes. If the stress-energy tensor satisfies the null energy condition for ℓ , that is, $T_{AB}\ell^A\ell^B \geq 0$, then this condition further implies [42] that ℓ has vanishing shear and that

$$\mathcal{L}_\ell q_{ab} = 0, \quad (3.1)$$

where \mathcal{L} is the Lie derivative and q_{ab} is the (degenerate) induced metric on Δ .

A nonexpanding horizon generalizes the notion of a Killing horizon, dropping the requirement of a Killing vector but retaining a time translation symmetry on the horizon itself. As in the Schwarzschild case, the integral curves of ℓ^a on Δ are the null geodesic generators of Δ . In the absence of a Killing vector, there is no preferred time coordinate, but it may be shown that a generic nonexpanding horizon has a preferred foliation, a set of “good cuts,” that generalize the constant time slices of the Schwarzschild metric [43]. Hence, we can again construct an r - v plane and a Gaussian null coordinate system near Δ .

The main change from the Schwarzschild case is that the cross section $\hat{\Delta}$ of a nonexpanding horizon need not have any symmetries, so conventional Kaluza-Klein methods no longer apply. As Yoon has shown, though, there is a generalized Kaluza-Klein reduction even in the absence of symmetries [44,45]. Start by writing the metric in the general form

$$\begin{aligned} ds^2 &= g_{AB}dz^A dz^B \\ &= g_{ab}dx^a dx^b + \phi_{\mu\nu}(dy^\mu + A_a^\mu dx^a)(dy^\nu + A_b^\nu dx^b), \end{aligned} \quad (3.2)$$

where lowercase roman indices (a,b,...) run from 0 to 1, lowercase greek indices (μ, ν, \dots) run from 2 to $D-1$, and uppercase roman indices (A,B,...) run from 0 to $D-1$. The “x” coordinates label our preferred two-dimensional manifold, while the “y” coordinates are the transverse directions. Define the Kaluza-Klein-like derivatives

$$\begin{aligned} \hat{\partial}_a &= \partial_a - A_a^\mu \partial_\mu, \\ D_a \phi_{\mu\nu} &= \hat{\partial}_a \phi_{\mu\nu} - (\partial_\mu A_a^\rho) \phi_{\rho\nu} - (\partial_\nu A_a^\rho) \phi_{\rho\mu}; \end{aligned} \quad (3.3)$$

connections

$$\begin{aligned} \hat{\Gamma}_{bc}^a &= \frac{1}{2} g^{ad} (\hat{\partial}_b g_{dc} + \hat{\partial}_c g_{db} - \hat{\partial}_d g_{bc}), \\ \hat{\Gamma}_{\mu\nu}^\rho &= \frac{1}{2} \phi^{\rho\sigma} (\partial_\mu \phi_{\sigma\nu} + \partial_\nu \phi_{\sigma\mu} - \partial_\sigma \phi_{\mu\nu}); \end{aligned} \quad (3.4)$$

and curvatures

$$\begin{aligned} \hat{R}_{ab} &= \hat{\partial}_c \hat{\Gamma}_{ab}^c - \hat{\partial}_a \hat{\Gamma}_{bc}^c + \hat{\Gamma}_{ab}^c \hat{\Gamma}_{cd}^d - \hat{\Gamma}_{ad}^c \hat{\Gamma}_{bc}^d, \\ \mathcal{R}_{\mu\nu} &= \partial_\rho \hat{\Gamma}_{\mu\nu}^\rho - \partial_\mu \hat{\Gamma}_{\nu\rho}^\rho + \hat{\Gamma}_{\mu\nu}^\rho \hat{\Gamma}_{\rho\sigma}^\sigma - \hat{\Gamma}_{\mu\sigma}^\rho \hat{\Gamma}_{\nu\rho}^\sigma, \\ F_{ab}{}^\mu &= \hat{\partial}_a A_b{}^\mu - \hat{\partial}_b A_a{}^\mu, \\ \hat{R} &= g^{ab} \hat{R}_{ab}, \quad \mathcal{R} = \phi^{\mu\nu} \mathcal{R}_{\mu\nu}. \end{aligned} \quad (3.5)$$

A straightforward calculation of the Einstein-Hilbert action then gives

$$I = \int d^{D-2}y I_2 \quad (3.6)$$

with

$$\begin{aligned} I_2 &= \frac{1}{16\pi G} \int d^2x \sqrt{-g} \sqrt{\phi} \left\{ \hat{R} + \frac{1}{4} g^{ab} g^{cd} \phi_{\mu\nu} F_{ac}{}^\mu F_{bd}{}^\nu \right. \\ &\quad + \frac{1}{4} g^{ab} \phi^{\mu\nu} \phi^{\rho\sigma} (D_a \phi_{\mu\rho} D_b \phi_{\nu\sigma} - D_a \phi_{\mu\nu} D_b \phi_{\rho\sigma}) \\ &\quad \left. + \frac{1}{4} \phi^{\mu\nu} g^{ab} g^{cd} (\partial_\mu g_{ac} \partial_\nu g_{bd} - \partial_\mu g_{ab} \partial_\nu g_{cd}) + \mathcal{R} \right\}, \end{aligned} \quad (3.7)$$

where g and ϕ are the determinants of g_{ab} and $\phi_{\mu\nu}$.

The first two lines in (3.7) look like an ordinary Kaluza-Klein reduction, and can in fact be viewed as the action of a Kaluza-Klein theory whose gauge group is the group of diffeomorphisms of the transverse manifold [44]. A two-dimensional interpretation of the third line is less obvious. We can cure this, though, with a partial gauge fixing. First, we can always choose local coordinates in which

$$g_{ab} = \begin{pmatrix} -2h & 1 \\ 1 & 0 \end{pmatrix}. \quad (3.8)$$

Yoon calls this “Polyakov gauge,” after a similar choice in two-dimensional field theories [46], while from the D -dimensional point of view, it is essentially Bondi gauge [47] or Gaussian null coordinates [39]. It is easy to see that with this choice, even if h depends on the y^μ ,

$$\phi^{\mu\nu} g^{ab} g^{cd} (\partial_\mu g_{ac} \partial_\nu g_{bd} - \partial_\mu g_{ab} \partial_\nu g_{cd}) = 0. \quad (3.9)$$

For our purposes, Eq. (3.8) is too restrictive a gauge choice—it hides a piece of the symmetry we are trying to understand. But if we now allow an arbitrary

two-dimensional coordinate transformation $x \rightarrow \bar{x}(x)$, it may be checked that the term (3.9) still vanishes.

The remaining term of concern in (3.7) is the transverse curvature \mathcal{R} . In $D = 4$ dimensions, this term is essentially trivial,

$$\int d^2y \sqrt{\phi} \mathcal{R} = 4\pi\chi, \quad (3.10)$$

where χ is the Euler characteristic of the transverse manifold. The transverse curvature thus merely contributes an effective cosmological constant to the two-dimensional action. For $D \neq 4$, the situation is more complicated; the transverse curvature couples only to $\sqrt{-\bar{g}}$, but it can give a sort of position-dependent cosmological ‘‘constant.’’ This

should not affect the symmetries derived in Sec. VI, but a deeper understanding would be helpful. (Note also that \mathcal{R} involves only y derivatives, while we have chosen a gauge in which $\sqrt{-\bar{g}}$ depends only on x , so by rescaling $\phi_{\mu\nu}$ by an appropriate power of $\sqrt{-\bar{g}}$, we can actually remove the coupling).

To make it easier to compare this formalism to other work on dilaton gravity, it is convenient to separate out the determinant ϕ from the transverse metric $\phi_{\mu\nu}$, writing

$$\phi_{\mu\nu} = \phi^{\frac{1}{D-2}} \Phi_{\mu\nu}, \quad \det |\Phi_{\mu\nu}| = 1. \quad (3.11)$$

The determinant and $\Phi_{\mu\nu}$ then decouple in the kinetic term in (3.7),

$$\begin{aligned} \frac{1}{4} g^{ab} \phi^{\mu\nu} \phi^{\rho\sigma} (D_a \phi_{\mu\rho} D_b \phi_{\nu\sigma} - D_a \phi_{\mu\nu} D_b \phi_{\rho\sigma}) &= \frac{1}{4} g^{ab} (\Phi^{-1})^{\mu\nu} (\Phi^{-1})^{\rho\sigma} (D_a \Phi_{\mu\rho} D_b \Phi_{\nu\sigma} - D_a \Phi_{\mu\nu} D_b \Phi_{\rho\sigma}) \\ &\quad - \frac{D-3}{4(D-2)} \phi^{-2} g^{ab} D_a \phi D_b \phi. \end{aligned} \quad (3.12)$$

One more simplification is standard in dilaton gravity: by rescaling the metric g_{ab} , we can eliminate the kinetic term for ϕ . Specifically, if we set

$$\phi = \varphi^2 \quad (3.13)$$

and let

$$g_{ab} = \varphi^{-\frac{D-3}{D-2}} \bar{g}_{ab}, \quad (3.14)$$

the action (3.7) reduces to

$$\begin{aligned} I_2 &= \frac{1}{16\pi G} \int d^2x \sqrt{-\bar{g}} \left\{ \varphi \bar{R} + \frac{1}{4} \varphi^2 \bar{g}^{ab} \bar{g}^{cd} \Phi_{\mu\nu} F_{ac}{}^\mu F_{bd}{}^\nu \right. \\ &\quad \left. + \frac{1}{4} \varphi \bar{g}^{ab} (\Phi^{-1})^{\mu\nu} (\Phi^{-1})^{\rho\sigma} \right. \\ &\quad \left. \times (D_a \Phi_{\mu\rho} D_b \Phi_{\nu\sigma} - D_a \Phi_{\mu\nu} D_b \Phi_{\rho\sigma}) + \varphi^{\frac{1}{D-2}} \mathcal{R} \right\}, \end{aligned} \quad (3.15)$$

which may be recognized as the action for a gauge field and a nonlinear sigma model coupled to two-dimensional dilaton gravity.

I have, of course, glossed over an essential feature: the ‘‘two-dimensional’’ fields in (3.15) also depend on the transverse coordinates y , which must still be integrated over. This remnant of the higher-dimensional structure appears in two places. First, the two-dimensional curvature R_{ab} in (3.5) involves convective derivatives $\hat{\partial}_a = \partial_a - A_a{}^\mu \partial_\mu$, distinguishing it from the ordinary Ricci curvature of the two-dimensional metric \bar{g}_{ab} . Near a black hole horizon Δ , though, one can choose corotating coordinates

in which $A_a{}^\mu$ vanishes on the horizon and remains small in a neighborhood of Δ [39]. We will be interested in symmetries in a small region around the horizon. In such a region, to the order of approximation we will need, these coordinates will allow us to replace \bar{R} in (3.15) by the ordinary two-dimensional curvature scalar (see Appendix B for details).

Second, the ‘‘matter’’ fields $F_{ab}{}^\mu$ and $\Phi_{\mu\nu}$ in (3.15) also depend on the transverse coordinates. In a symmetric enough setting, we could expand these fields in modes to create a Kaluza-Klein tower of states. In general, though—for instance, for a black hole whose horizon is distorted by surrounding matter—this will not be possible. Fortunately, though, it is also not necessary. As Wall showed for quantum fields on a black hole horizon [28], the physics at different transverse positions decouples, and each null generator can be treated separately. While I do not know a rigorous generalization to the case of dynamical gravity, we shall see that the relevant symmetries act separately on each generator. Since these symmetries govern the density of states, this density is also determined independently on each generator, and the total entropy can be obtained by integrating. This is the underlying reason for an area law for entropy, although as we shall see, it gives more, fixing the exact coefficient of the area.

Note that we have not yet imposed the existence of a horizon. Even the use of Gaussian null coordinates requires only the presence of a null surface, which need not have vanishing expansion. In principle, it is possible to further restrict the metric (3.2), but the resulting expressions are complicated and unwieldy [48]. We will instead take a shortcut, identifying horizons in the two-dimensional action I_2 to obtain a more tractable formulation.

IV. DILATON GRAVITY WITH NULL DYADS

We now restrict our attention to the effective two-dimensional action (3.15) at fixed transverse position (fixed y). While this action will not reveal the full symmetries of the higher-dimensional theory, any y -independent symmetry of I_2 will also be a symmetry of the full theory.

To simplify notation, let us write

$$I = \frac{1}{16\pi G} \int_M (\varphi R + V[\varphi, \chi]) \epsilon, \quad (4.1)$$

where ϵ is the volume two-form² and χ denotes any further fields in the problem (here Φ , A , and any additional matter fields). The quantity φ is called the dilaton; as we have seen, it is essentially the volume element of the transverse metric. Note that while the potential V in (4.1) may be quite complicated, it contains no x derivatives of φ .

The equations of motion coming from varying g and φ in this action are

$$E_{ab} = \nabla_a \nabla_b \varphi - g_{ab} \square \varphi + \frac{1}{2} g_{ab} V = 8\pi G T_{ab}, \quad (4.2a)$$

$$R + \frac{dV}{d\varphi} = 0, \quad (4.2b)$$

where I have added a source stress-energy tensor. Equation (4.2b) is not independent, but follows from the divergence of (4.2a).

It is convenient to describe the geometry in terms of a null dyad (ℓ_a, n_a) , with $\ell^2 = n^2 = 0$ and $\ell \cdot n = -1$. In terms of such a dyad, the metric and volume form are

$$g_{ab} = -(\ell_a n_b + n_a \ell_b), \quad \epsilon_{ab} = (\ell_a n_b - n_a \ell_b). \quad (4.3)$$

When we later specialize to the case of a black hole spacetime with horizon Δ , we will choose a dyad for which ℓ is the null normal to Δ and n is the induced volume element.

To simplify later equations, we define derivatives

$$D = \ell^a \nabla_a, \quad \bar{D} = n^a \nabla_a. \quad (4.4)$$

D is essentially the same D as in the Newman-Penrose formalism. \bar{D} would ordinarily be denoted Δ in the Newman-Penrose formalism, but we are already using Δ to signify the horizon.

The dyad (ℓ, n) is determined only up to local Lorentz transformations,

$$\ell^a \rightarrow e^\lambda \ell^a, \quad n^a \rightarrow e^{-\lambda} n^a. \quad (4.5)$$

We can partially fix this freedom by choosing n_a to have vanishing acceleration, $n^b \nabla_b n^a = 0$. This condition implies that the integral curves of n are affinely parametrized null geodesics, which can be taken to start at the horizon Δ . The symmetry (4.5) is still not completely fixed, but the remaining transformations are restricted to those for which $\bar{D}\lambda = 0$.

With this condition on n_a , it is easy to check that

$$\begin{aligned} \nabla_a \ell_b &= -\kappa n_a \ell_b, & \nabla_a \ell^a &= \kappa, \\ \nabla_a n_b &= \kappa n_a n_b, & \nabla_a n^a &= 0, \end{aligned} \quad (4.6)$$

where κ will later be interpreted as the surface gravity at a horizon. By (4.3), ℓ_a is a conformal Killing vector. Under variation of the dyad, Eq. (4.6) will be preserved as long as

$$\begin{aligned} \bar{D}(\ell^c \delta n_c) &= (D + \kappa)(n^c \delta n_c), \\ \delta \kappa &= -D(n^c \delta \ell_c) + \kappa \ell^c \delta n_c + \bar{D}(\ell^c \delta \ell_c). \end{aligned} \quad (4.7)$$

We will later need to integrate by parts along the horizon. For this, it will be useful to take advantage of the identity

$$(df)_a = -Df n_a - \bar{D}f \ell_a \quad \text{for any function } f, \quad (4.8)$$

where I am treating n_a and ℓ_a as one-forms. Another identity will also be helpful:

$$[D, \bar{D}] = -\kappa \bar{D} \Leftrightarrow \bar{D}D = (D + \kappa)\bar{D}. \quad (4.9)$$

By considering the commutator $[\nabla_a, \nabla_b] \ell^b$ and recalling that in two dimensions $R_{ab} = \frac{1}{2} g_{ab} R$, it is straightforward to show from (4.6) that

$$R = 2\bar{D}\kappa. \quad (4.10)$$

The action (4.1) can thus be written as

$$I = \frac{1}{8\pi G} \int_M \left(-\kappa \bar{D}\varphi + \frac{1}{2} V[\varphi, \chi] \right) \epsilon. \quad (4.11)$$

V. HORIZONS

Our next task will be to characterize a generic black hole horizon in this two-dimensional setting, as a first step toward analyzing its near-horizon symmetries. Spacetimes containing black holes in two-dimensional dilaton gravity have essentially the same Penrose diagrams as those in higher dimensions [40], as illustrated in Fig. 1. As in higher dimensions, the horizons are Killing horizons [50], and form boundaries of trapped regions [51]. For simplicity, Fig. 1 shows an asymptotically flat black hole. This asymptotic behavior will be irrelevant for the main

²I am using the convention that the object one integrates over an n -manifold is an n -form [49]. For our two-dimensional manifold, the volume form is ϵ_{ab} . For a null line with null normal ℓ_a , the volume one-form is n_a , a null vector normalized so that $\ell \cdot n = -1$.




FIG. 1. Typical Penrose diagram for the exterior of a black hole.

argument of this paper, though; the analysis below will hold equally well for asymptotically de Sitter or anti-de Sitter black holes.

Let us choose our null dyad so that at the horizon ℓ coincides with the null normal to the horizon. This means ℓ is also tangent to the horizon—its inner product with the normal (itself) is zero. Indeed, in any dimension, the null normals to the horizon are the tangent vectors of the null generators of the horizon. (See the beginning of Ref. [52] for a nice review).

In two dimensions, this choice of dyad is straightforward. The lift to D dimensions, though, is ambiguous; depending on coordinate choices, ℓ_A and ℓ^A may have additional transverse components. Our philosophy here will be that our preferred two-dimensional subspace traces the generators of the horizon, that is, that each generator of the horizon Δ has constant transverse coordinates y . This means that ℓ as a tangent vector lies in our two-dimensional subspace; that is, $\ell^\mu = 0$ but ℓ_μ need not vanish. We will choose n so that $n_\mu = 0$. This ensures that, even from the higher-dimensional point of view, n_a is the volume one-form along a horizon generator. As long as we stick to the coordinates described at the end of Sec. III, in which $g_{\mu a} = 0$ at the horizon, these choices will be largely irrelevant, but they may be important in more general coordinate systems.

We now need a way to determine that Δ is indeed a horizon. This will require an appropriate generalization of the “nonexpanding horizon” criteria of Sec. III. From the D -dimensional point of view, the expansion of Δ is

$$\theta = (\delta_B^A + \ell^A n_B) \nabla_A \ell^B = (\delta_b^a + \ell^a n_b) \nabla_a \ell^b + \nabla_\mu \ell^\mu = \ell^a \Gamma_{a\mu}^\mu. \quad (5.1)$$

In general, this will be a complicated expression, involving both the transverse metric $\phi_{\mu\nu}$ and the mixed components A_a^μ . But recall that we have chosen coordinates in which $A_a^\mu \stackrel{\Delta}{=} 0$, where from now on I will use the symbol $\stackrel{\Delta}{=}$ to mean “equal on the horizon.” Hence,

$$\theta = \ell^a \Gamma_{a\mu}^\mu \stackrel{\Delta}{=} \frac{1}{2} \phi^{\mu\nu} \ell^a \partial_a \phi_{\mu\nu} = \frac{1}{\varphi} D\varphi. \quad (5.2)$$

The condition for vanishing expansion is thus $D\varphi = 0$, and we will use this as the means to locate the horizon. Near the horizon, $D\varphi$ can then serve as a small expansion parameter, indicating how far we have moved from Δ . The interpretation of this parametrization in terms of Gaussian null coordinates is described in Appendix B.

In higher dimensions, vanishing expansion is enough to ensure that the whole horizon geometry is stationary as well. In two dimensions, where the dilaton φ is now separate from the transverse metric, this is no longer the case, and we must separately require that $DR \stackrel{\Delta}{=} 0$. I will also impose one more boundary condition at the horizon, that the integration measure n_a remain fixed at Δ . This restriction appears to be needed for the covariant canonical symplectic form of Sec. VII to be well behaved, though further exploration would be interesting. In Ref. [3], the condition $\ell^a \delta \ell_a = 0$ was also imposed, but while this simplifies the symplectic structure, it is not really needed.

Our boundary conditions at Δ thus become

$$D\varphi \stackrel{\Delta}{=} 0, \quad (5.3a)$$

$$DR \stackrel{\Delta}{=} 0, \quad (5.3b)$$

$$\ell^a \delta n_a \stackrel{\Delta}{=} n^a \delta n_a \stackrel{\Delta}{=} 0 \quad (5.3c)$$

(where, again, $\stackrel{\Delta}{=}$ means “equal on Δ ”). Our task is to find the symmetries of the dilaton gravity action that are compatible with these conditions.

Before proceeding further, one slightly subtle issue of interpretation should be addressed. The approach here is not to first choose a fixed submanifold Δ and then impose (5.3a)–(5.3c). This is too strong a demand; it would forbid variations that changed $D\varphi$ on this fixed surface and would prohibit transverse diffeomorphisms at Δ . The philosophy is, rather, to use the condition $D\varphi = 0$ to determine the location of the horizon, and to impose the remaining conditions at that location. A variation that changes $D\varphi$ is then understood as changing the location of the horizon, and a suitable transverse diffeomorphism can be used to “move it back.”

VI. HORIZON SYMMETRIES

The action (4.1) is, of course, invariant under two-dimensional diffeomorphisms, including horizon “supertranslations” [53] generated by vector fields $\xi^a = \xi \ell^a$. Such diffeomorphisms fail to respect condition (5.3c), however, since $\ell^a \delta_\xi n_a \neq 0$. This is easily cured, by supplementing each diffeomorphism with a local Lorentz transformation $\delta \ell^a = (\delta\lambda) \ell^a$, $\delta n^a = -(\delta\lambda) n^a$ with $\delta\lambda = D\xi$. From the discussion after (4.5), this requires that $\bar{D}\xi \stackrel{\Delta}{=} 0$. We thus have an invariance,

$$\begin{aligned}\delta_\xi \ell^a &= 0, & \delta_\xi n^a &= -(D + \kappa)\xi n^a, \\ \delta_\xi g_{ab} &= (D + \kappa)\xi g_{ab}, \\ \delta_\xi \varphi &= \xi D\varphi, & \text{with } \bar{D}\xi &\triangleq 0.\end{aligned}\quad (6.1)$$

Note that from (4.6),

$$\delta_\xi \kappa = D(D + \kappa)\xi. \quad (6.2)$$

As pointed out some time ago [26,27], for configurations containing black holes, the action also has an *approximate* invariance under certain shifts of the dilaton near the horizon, with an approximation that can be made arbitrarily good by restricting the transformation to a small enough neighborhood of Δ . This is not quite an ordinary invariance, since it holds only for a restricted class of configurations, those with horizons. For such configurations, though, it can be made arbitrarily close to an exact symmetry (see Appendix C for more details).

Specifically, consider a variation,

$$\hat{\delta}_\eta \varphi = \nabla_a(\eta \ell^a) = (D + \kappa)\eta \quad \text{with } \bar{D}\eta \triangleq 0. \quad (6.3)$$

(The hat on $\hat{\delta}$ distinguishes this variation from a diffeomorphism.) The action transforms as

$$\begin{aligned}\hat{\delta}_\eta I &= \frac{1}{16\pi G} \int_M \left(R + \frac{dV}{d\varphi} \right) \hat{\delta}_\eta \varphi \epsilon \\ &= -\frac{1}{16\pi G} \int_M \eta \left[DR + \frac{d^2 V}{d\varphi^2} D\varphi \right] \epsilon.\end{aligned}\quad (6.4)$$

But $D\varphi$ and DR both vanish at the horizon, so the variation (6.4) can be made as small as one wishes by choosing η to fall off fast enough away from Δ .

There is one subtlety, however. While the transformation (6.3) does not directly act on the curvature, the change of φ “moves the horizon”—that is, the locus $D\varphi = 0$ may change under a shift of φ . In itself, this is not a problem, but DR may no longer vanish at the new location. The diffeomorphism needed to “move the horizon back” is determined by the condition

$$(\hat{\delta}_\eta + \delta_\zeta)(D\varphi) = \hat{\delta}_\eta(D\varphi) + \zeta^a \nabla_a(D\varphi) \triangleq 0 \quad \Rightarrow \quad \zeta^a = \bar{\zeta} n^a = -\frac{D(\hat{\delta}_\eta \varphi)}{\bar{D}D\varphi} n^a. \quad (6.5)$$

This change can be compensated with a “small” (order $D\varphi$) Weyl transformation of the metric to restore the condition $DR \triangleq 0$. Consider a transformation of the form

$$\hat{\delta}_\eta g_{ab} = \hat{\delta}\omega_\eta g_{ab} \quad \Leftrightarrow \quad \hat{\delta}_\eta \ell_a = \hat{\delta}\omega_\eta \ell_a, \quad (6.6)$$

where the second equality comes from the boundary condition that $\delta n_a \triangleq 0$. Define

$$\hat{\delta}\omega_\eta = X_\eta \frac{D\varphi}{\bar{D}D\varphi}. \quad (6.7)$$

Using identities from Sec. IV, it is not hard to see that R transforms as

$$\hat{\delta}_\eta R \triangleq 2(D + \kappa)X_\eta. \quad (6.8)$$

The condition that DR remain zero on Δ is thus

$$\bar{\zeta} \bar{D} DR + 2D(D + \kappa)X_\eta \triangleq 0. \quad (6.9)$$

On shell—or, less restrictively, whenever the constraint (4.2b) holds—a short calculation gives an explicit expression for X_η :

$$\bar{\zeta} \bar{D} DR + 2D(D + \kappa)X_\eta \triangleq \frac{d^2 V}{d\varphi^2} D(D + \kappa)\eta + 2D(D + \kappa)X_\eta \triangleq 0 \quad \Rightarrow \quad X_\eta \triangleq -\frac{1}{2} \frac{d^2 V}{d\varphi^2} \eta. \quad (6.10)$$

With this added transformation, the boundary condition $DR \triangleq 0$ is preserved. Like (6.3), the Weyl transformation (6.7) changes the action only by terms proportional to $\eta D\varphi$, which can be made arbitrarily small by choosing η to fall off fast enough away from the horizon.

We thus have two sets of transformations at the horizon, diffeomorphisms δ_ξ and shifts $\hat{\delta}_\eta$, which preserve the action (to an arbitrarily good approximation) as long as a horizon

actually exists. It is not too hard to check that these satisfy an algebra,

$$\begin{aligned}[\delta_{\xi_1}, \delta_{\xi_2}]f &\triangleq \delta_{\xi_{12}} f \quad \text{with } \xi_{12} = (\xi_1 D\xi_2 - \xi_2 D\xi_1), \\ [\hat{\delta}_{\eta_1}, \hat{\delta}_{\eta_2}]f &\triangleq 0, \\ [\delta_{\xi_1}, \hat{\delta}_{\eta_2}]f &\triangleq \hat{\delta}_{\eta_{12}} f \quad \text{with } \eta_{12} = -(\xi_1 D\eta_2 - \eta_2 D\xi_1).\end{aligned}\quad (6.11)$$

This may be recognized as a BMS_3 algebra, or equivalently a Galilean conformal algebra [31].

Given the rather atypical nature of this shift symmetry, we should also check the variation of the equations of motion (4.2a)–(4.2b). These are, of course, preserved by diffeomorphisms, so we need only consider the transformations (6.3) and (6.7). Since we are assuming that η falls off rapidly away from the horizon, it is enough to check the variations at Δ . By a straightforward computation, most of the equations of motion are preserved: up to terms that are themselves proportional to the equations of motion,

$$g^{ab}\hat{\delta}_\eta E_{ab} \triangleq 2(D+\kappa)\bar{D}\hat{\delta}_\eta\varphi + \frac{dV}{d\varphi}\hat{\delta}_\eta\varphi \triangleq \left(R + \frac{dV}{d\varphi}\right)(D+\kappa)\eta, \quad (6.12a)$$

$$n^a n^b \hat{\delta}_\eta E_{ab} \triangleq \bar{D}^2 \hat{\delta}_\eta \varphi - \bar{D}\varphi \bar{D}\hat{\delta}_\eta \omega \triangleq \frac{1}{2}\bar{D}\left(R + \frac{dV}{d\varphi}\right)\eta, \quad (6.12b)$$

$$\hat{\delta}_\eta\left(R + \frac{dV}{d\varphi}\right) \triangleq \hat{\delta}_\eta R + \frac{d^2 V}{d\varphi^2}\hat{\delta}_\eta\varphi, \quad (6.12c)$$

where I have used (4.9), (4.10), and the condition $\bar{D}\eta \triangleq 0$.

The remaining variation, $\ell^a \ell^b \hat{\delta}_\eta E_{ab}$, is not zero. But this is actually a familiar occurrence in conformal field theory. If we set $E_{ab} = 8\pi G T_{ab}$, we find that

$$\ell^a \ell^b \hat{\delta}_\eta T_{ab} \triangleq \frac{1}{8\pi G}(D-\kappa)D(D+\kappa)\eta, \quad (6.13)$$

which is essentially the usual anomaly for a conformal field theory with a central charge proportional to $1/G$ [54]. This is our first hint that the symmetry is anomalous.

One might worry that this anomaly could spoil the covariant phase space construction of Appendix A, since the closure of the symplectic current (A2) relies on the classical field equations. Fortunately, this is not a problem; the only dangerous term in the exterior derivative (A4) is proportional to $n_a n_b \delta g^{ab}$, which vanishes on Δ by virtue of the boundary conditions (5.3c).

VII. SYMPLECTIC STRUCTURE AND GENERATORS

To complete the analysis of the symmetries of Sec. VI, we should ask whether the algebra (6.11) can be realized—perhaps with a central extension—as a Poisson algebra of canonical generators of the symmetries, since this is the formulation that translates most directly into quantum mechanics. We have so far avoided introducing explicit coordinates. We will continue to do so, by employing the covariant canonical formalism reviewed in Appendix A.

The symplectic form (A3) is defined as an integral over a Cauchy surface Σ . To study horizon symmetries in the covariant phase space formalism, we should incorporate Δ as part of our Cauchy surface. Let us focus on the exterior region of an asymptotically flat black hole, with a Penrose diagram given by Fig. 1, and take Σ to be the union of the future horizon Δ and future null infinity \mathcal{I}^+ , with ends at the bifurcation point B and spacelike infinity. As noted earlier, the details of \mathcal{I}^+ will be unimportant, since we will be considering transformations that are nonvanishing only in a small neighborhood of the horizon.

Applying the general relations (A2)–(A3) to the action (4.11) for dilaton gravity and using the boundary condition $\delta n_a \triangleq 0$, it is straightforward to show that³

$$\begin{aligned} \Omega_\Delta[(\varphi, g); \delta_1(\varphi, g), \delta_2(\varphi, g)] \\ = \frac{1}{8\pi G} \int_\Delta [\delta_1 \varphi \delta_2 \kappa - \delta_1(\bar{D}\varphi) \ell^b \delta_2 \ell_b] n_a - (1 \leftrightarrow 2). \end{aligned} \quad (7.1)$$

The full symplectic form will include an additional integral along \mathcal{I}^+ , but this will be irrelevant to our consideration of near-horizon symmetries.

Two slightly tricky points remain, though, both related to the fact that a variation of φ can move the horizon, changing the locus of points $D\varphi = 0$. First, as discussed in Appendix A, the symplectic form itself is independent of the integration contour as long as the end points remain fixed. But Ω_Δ can change under variations that move the ends of the Cauchy surface. To avoid this behavior, we will require that $\delta(D\varphi) = 0$ at the bifurcation point B of Fig. 1, a condition that will be used in Sec. VIII.

Second, while typical changes in the horizon locus will not affect Ω_Δ , they will change objects such as Hamiltonians defined as integrals over Δ . We will account for this effect by adding a transverse diffeomorphism to “move the horizon back.” As in Sec. VI, such a diffeomorphism is determined by the condition that

$$(\delta + \delta_\zeta)(D\varphi) = \delta(D\varphi) + \zeta^a \nabla_a(D\varphi) \triangleq 0 \quad \Rightarrow \quad \zeta^a = \bar{\zeta} n^a = -\left(\frac{D\delta\varphi}{DD\varphi} + \frac{\bar{D}\varphi}{DD\varphi} \ell^b \delta \ell_b\right) n^a. \quad (7.2)$$

³The calculation simplifies if one notes that in differential form notation, $\kappa \bar{D}\varphi \epsilon = \bar{D}\varphi d\ell = -\kappa d\varphi \wedge n$.

Hence, for an object of the form $H = \int_{\Delta} \mathcal{H} n_a$, the full variation will be

$$\delta \int_{\Delta} \mathcal{H} n_a = \int_{\Delta} (\delta \mathcal{H} + \zeta^a \nabla_a \mathcal{H}) n_a. \quad (7.3)$$

We can now ask whether the transformations δ_{ξ} and $\hat{\delta}_{\eta}$ of the preceding section can be realized canonically as in (A8), that is, whether there exist generators that satisfy

$$\begin{aligned} \delta L[\xi] &= \frac{1}{8\pi G} \int_{\Delta} [\delta\varphi \delta_{\xi} \kappa - \delta_{\xi} \varphi \delta\kappa - \delta(\bar{D}\varphi) \ell^b \delta_{\xi} \ell_b + \delta_{\xi}(\bar{D}\varphi) \ell^b \delta\ell_b] n_a \\ &= \frac{1}{8\pi G} \int_{\Delta} [\delta\varphi D(D + \kappa)\xi - \xi D\varphi \delta\kappa + \{\xi \bar{D}D\varphi - (D + \kappa)\xi \bar{D}\varphi\} \ell^b \delta\ell_b] n_a, \end{aligned} \quad (7.4a)$$

$$\begin{aligned} \delta M[\eta] &= \frac{1}{8\pi G} \int_{\Delta} [\delta\varphi \hat{\delta}_{\eta} \kappa - \hat{\delta}_{\eta} \varphi \delta\kappa - \delta(\bar{D}\varphi) \ell^b \hat{\delta}_{\eta} \ell_b + \hat{\delta}_{\eta}(\bar{D}\varphi) \ell^b \delta\ell_b] n_a \\ &= \frac{1}{8\pi G} \int_{\Delta} [-\delta\omega_{\eta} D\delta\varphi - \delta\kappa(D + \kappa)\eta + \{\bar{D}(D + \kappa)\eta - \delta\omega_{\eta} \bar{D}\varphi\} \ell^b \delta\ell_b] n_a, \end{aligned} \quad (7.4b)$$

where in the last line I have used the fact that $\hat{\delta}_{\eta} \kappa = D\hat{\delta}\omega_{\eta}$.

It is not at all clear that such generators exist; there is no obvious reason that the near-horizon symmetry (6.3) should have a canonical realization. In fact, though, the quantities

$$L[\xi] = \frac{1}{8\pi G} \int_{\Delta} [\xi D^2 \varphi - \kappa \xi D\varphi] n_a, \quad (7.5a)$$

$$M[\eta] = \frac{1}{8\pi G} \int_{\Delta} \eta \left(D\kappa - \frac{1}{2} \kappa^2 \right) n_a \quad (7.5b)$$

do the job. [To obtain the $\delta\omega_{\eta}$ terms in (7.4b), one must use the full variation (7.3), along with equation (6.10) for X_{η} and the fact that $\bar{D}\eta \triangleq 0$; again, the covariant phase space formalism allows us to impose equations of motion after variation].

Using (A10), we can now find the Poisson brackets of these generators:

$$\{L[\xi_1], L[\xi_2]\} = L[\xi_{12}], \quad (7.6a)$$

$$\{M[\eta_1], M[\eta_2]\} \triangleq 0, \quad (7.6b)$$

$$\begin{aligned} \{L[\xi_1], M[\eta_2]\} &\triangleq -M[\eta_{12}] \\ &- \frac{1}{16\pi G} \int_{\Delta} (D\xi_1 D^2 \eta_2 - D\eta_2 D^2 \xi_1) n_a, \end{aligned} \quad (7.6c)$$

where ξ_{12} and η_{12} were defined in (6.11). The $\{L, L\}$ brackets are unchanged even if Δ is not a horizon. The $\{L, M\}$ and $\{M, M\}$ brackets do change—the shift

transformations are exact symmetries only on a horizon—but modulo equations of motion, the deviations are of order $(D\varphi)^2$. The canonical generators thus give a representation of the symmetry algebra (6.11), now with an added central term. Such central terms are well understood in classical mechanics [55]; their appearance in quantum gravity was first emphasized by Brown and Henneaux [56], and as we saw in Sec. II, they play a crucial role in determining entropy.

VIII. MODES AND ZERO MODES

As described in Sec. II, we can now use the symmetry (7.6a)–(7.6c) to determine the density of states. To do so, we will need the central charge and the zero modes. These, in turn, require a mode expansion for the parameters ξ and η .

For a black hole with constant surface gravity, the appropriate modes are well known. They take the form $e^{in\kappa v}$, where v is the advanced time along the horizon, normalized so that $\ell^a \nabla_a v = 1$. Such modes are periodic in imaginary time with period $2\pi/\kappa$, as required for non-singular Green's functions. Here, though, κ is one of our canonical variables, and we cannot simply take it to be constant. In the language of Ref. [41], we are considering nonexpanding horizons but not “isolated horizons.” While we can always perform a local Lorentz transformation to make κ constant, that would require much more restrictive boundary conditions, which would hide part of the symmetry.

Fortunately, though, the appropriate generalization is straightforward. Define a phase ψ such that

$$D\psi \triangleq \kappa, \quad \bar{D}\psi \triangleq 0 \quad \Leftrightarrow \quad d\psi \triangleq -\kappa n_a \quad \Leftrightarrow \quad \psi \triangleq - \int_{\Delta} \kappa n_a \triangleq - \int \kappa dv. \quad (8.1)$$

The modes are then

$$\zeta_n \triangleq \frac{1}{\kappa} e^{im\psi} \quad (\text{where } \zeta \text{ is either } \xi \text{ or } \eta). \quad (8.2)$$

The prefactor of $1/\kappa$ has been chosen so the modes obey the ordinary algebra of diffeomorphisms of the circle,

$$\{\zeta_m, \zeta_n\} = \zeta_m D \zeta_n - \zeta_n D \zeta_m = -i(m-n)\zeta_{m+n}. \quad (8.3)$$

Setting $L_n = L[\xi_n]$ and $M_n = M[\eta_n]$, it is easy to check that our BMS_3 algebra reduces to (2.1), with a central term

$$-\frac{1}{16\pi G} \int_{\Delta} (D\xi_m D^2\eta_n - D\eta_n D^2\xi_m) n_a = -\frac{i}{16\pi G} \int_{\Delta} (mn^2 - nm^2) e^{i(m+n)\psi} d\psi. \quad (8.4)$$

If we take the integral to be over a single period—essentially mapping the problem to a circle, as is standard in conformal field theory—we obtain a central charge:

$$c_{LM} = \frac{1}{4G}. \quad (8.5)$$

We also need the zero modes of L and M . For M , this is straightforward: from (7.5b),

$$M_0 = M[\eta_0] = -\frac{1}{16\pi G} \int_{\Delta} \kappa^2 \eta_0 n_a = \frac{1}{16\pi G} \int d\psi = \frac{1}{8G}. \quad (8.6)$$

For L , the “bulk” contribution to L_0 vanishes. But L , unlike M , has a boundary contribution. Indeed, the variation leading to (7.4a) involves integration by parts, with a boundary term

$$\delta L[\xi] = \cdots + \frac{1}{8\pi G} [\xi D\delta\varphi - (D+\kappa)\xi\delta\varphi] \Big|_{\partial\Delta}. \quad (8.7)$$

As noted in Sec. VII, the covariant phase space approach requires that we set $D\delta\varphi$ to zero at the bifurcation point B . We should certainly not hold φ itself fixed, though, since that would fix φ along the entire horizon, eliminating the shift symmetry. Instead, we should fix the conjugate variable κ at B . This requires an added boundary contribution to cancel the variation (8.7),

$$L_0^{\text{bdry}} = \frac{1}{8\pi G} \varphi(D+\kappa)\xi_0 \Big|_B = \frac{\varphi_+}{8\pi G}, \quad (8.8)$$

where φ_+ is the value of φ at B .

IX. ENTROPY

We are finally in a position to compute the entropy of our black hole. Inserting (8.5), (8.6), and (8.8) into (2.4), we obtain

$$S = \frac{\varphi_+}{4G}. \quad (9.1)$$

For a purely two-dimensional theory, this is the correct Bekenstein-Hawking entropy for a black hole [40,50]. From the D -dimensional perspective, it is the contribution of a single null generator of the horizon. But the symmetries that determine (9.1) act independently on each generator, and entropy is an extensive quantity, so we can add the individual entropies:

$$S = \frac{1}{4G} \int d^{D-2}y \varphi_+ = \frac{1}{4G} \int d^{D-2}y \sqrt{\phi_+} = \frac{A_+}{4G}, \quad (9.2)$$

where A_+ is the area of the bifurcation sphere. We have thus obtained the correct Bekenstein-Hawking entropy for the full D -dimensional theory.

X. CONCLUSIONS

As anticipated, black hole entropy is indeed determined by the symmetries of the horizon. In contrast to previous efforts to demonstrate this behavior, the derivation presented here has required no stretched horizon, no extra angular dependence, and no other *ad hoc* ingredients. The main assumption has merely been that the dimensionally reduced horizon obeys the “boundary conditions” of Sec. V.

What is the meaning of the crucial BMS_3 symmetry? It is not a gauge symmetry; physical states are singlets under gauge symmetries, while our state counting only works because the relevant states transform under high-dimensional representations. This kind of behavior is typical of an asymptotic symmetry. But our BMS_3 is also not quite a standard asymptotic symmetry; while we can view the horizon as a sort of boundary, it is a boundary that exists only for a restricted class of field configurations. Physically, we are asking a question of conditional probability—if a black hole is present, what are its properties?—and the symmetries reflect this condition. This is at least vaguely analogous to entanglement entropy, which requires a similar specification of a boundary. Indeed, it is possible that our horizon degrees of freedom might be viewed as a

remnant left behind after tracing out the state behind the horizon. For three-dimensional topological field theory, this argument can be made fairly rigorous [57]; it would be interesting to investigate it further in the present context.

There are several obvious directions for generalization. A BMS symmetry at the horizon has appeared in other settings (for instance, Refs. [53,58–62]); the relationship to the BMS₃ symmetry described here should be clarified. Perhaps most fundamentally, if this symmetry really does explain the universality of black hole entropy, it should be present—although possibly hidden—in other derivations of entropy. Hints of such a hidden symmetry have been found for loop quantum gravity [63], induced gravity [64], and near-extremal black holes in string theory [65], but none of these investigations has yet exploited the full BMS₃ symmetry.

Ideally, we might hope to do even more. Many of the fundamental questions in black hole thermodynamics involve the dynamics of Hawking radiation and its coupling to gravitational degrees of freedom. In 2 + 1 dimensions, Emparan and Sachs have succeeded in using the asymptotic conformal symmetry to couple the BTZ black hole to matter and obtain Hawking radiation [66]. Perhaps our BMS₃ symmetry will ultimately allow us to do the same in arbitrary dimensions.

ACKNOWLEDGMENTS

Acknowledgments This research was supported by the U.S. Department of Energy under Grant No. DE-FG02-91ER40674.

APPENDIX A: THE COVARIANT CANONICAL FORMALISM

The idea underlying the covariant canonical formalism is that for a theory with a well-posed initial value problem—that is, well-defined and unique time evolution—the phase space, viewed as the space of initial data, is isomorphic to the space of classical solutions [67–70]. The isomorphism is not canonical, but requires a choice of a Cauchy surface Σ . Once Σ has been chosen, though, the identification is simple: initial data on Σ determines a unique classical solution, and a classical solution restricted to Σ defines a unique set of initial data. This equivalence, which can be traced back to Lagrange (see Ref. [68]), means that we can formulate all the usual ingredients of Hamiltonian mechanics without ever having to break general covariance by choosing a particular time slicing.

Consider a theory in a D -dimensional spacetime with fields Φ^A (for us, φ , g , and ψ) and a Lagrangian density $L[\Phi]$, which we view as a D -form. Under a general variation of the fields, $L[\Phi]$ changes as

$$\delta L = E_A \delta \Phi^A + d\Theta[\Phi, \delta \Phi], \quad (\text{A1})$$

where the equations of motion are $E_A = 0$ and the last “boundary” term comes from integration by parts. We normally ignore this boundary term, but in the covariant canonical formalism, it is crucial. The symplectic current ω is defined by a second variation,

$$\omega[\Phi; \delta_1 \Phi, \delta_2 \Phi] = \delta_1 \Theta[\Phi, \delta_2 \Phi] - \delta_2 \Theta[\Phi, \delta_1 \Phi], \quad (\text{A2})$$

and the symplectic form is

$$\Omega[\Phi; \delta_1 \Phi, \delta_2 \Phi] = \int_{\Sigma} \omega[\Phi; \delta_1 \Phi, \delta_2 \Phi] = \int_{\Sigma} \omega_{AB} \delta_1 \Phi^A \wedge \delta_2 \Phi^B, \quad (\text{A3})$$

where Σ is a Cauchy surface. (More precisely, Ω is often a presymplectic form, with degenerate directions that must be factored out to obtain a true symplectic form [69]).

In keeping with the covariant phase space philosophy, $\Omega[\Phi; \delta_1 \Phi, \delta_2 \Phi]$ depends on a classical solution Φ , which fixes a point in the phase space. Ω itself is a two-form on the phase space, and the variations $\delta \Phi$ are tangent vectors to the space of classical solutions, that is, solutions of the linearized equations of motion. For a field theory in flat spacetime, it is not hard to check that when Σ is a surface of constant time, (A3) is equivalent to the ordinary symplectic form.

The symplectic current (A2) is closed,

$$\begin{aligned} d\omega[\Phi; \delta_1 \Phi, \delta_2 \Phi] &= \delta_1 d\Theta[\Phi, \delta_2 \Phi] - \delta_2 d\Theta[\Phi, \delta_1 \Phi] \\ &= -\delta_1 E_A \wedge \delta_2 \Phi^A + \delta_2 E_A \wedge \delta_1 \Phi^A = 0, \end{aligned} \quad (\text{A4})$$

since the variations satisfy the linearized equations of motion $\delta E_A = 0$. Hence, the symplectic form (A3) will depend only weakly on the choice of Cauchy surface; integrals over two surfaces Σ_1 and Σ_2 can differ only by boundary terms that might arise if $\partial \Sigma_1 \neq \partial \Sigma_2$. In particular, for a diffeomorphism-invariant theory, a diffeomorphism generated by a vector field ζ^a transverse to Σ may be viewed as a deformation of the Cauchy surface, and we have

$$\Omega[\Phi; \delta_1 \Phi, \delta_{\zeta} \Phi] = 0 \quad (\text{A5})$$

as long as ζ^a vanishes at $\partial \Sigma$. This may be checked explicitly for the symplectic form (7.1): under a diffeomorphism generated by a vector field $\zeta^a = \bar{\zeta}^a n^a$, one finds

$$\Omega_{\Delta}[(\varphi, g); \delta(\varphi, g), \delta_{\zeta}(\varphi, g)] = \frac{1}{8\pi G} \bar{\zeta}^a \delta(\bar{D}\varphi) \Big|_{\partial \Delta} + \dots, \quad (\text{A6})$$

where the omitted terms are proportional to either the equations of motion or their first variations, which are both set to zero in the covariant canonical formalism.

As in ordinary mechanics, the symplectic form determines Poisson brackets and Hamiltonians. Schematically, the Poisson bracket of two functions X and Y is

$$\{X, Y\} = \int_{\Sigma} \frac{\delta X}{\delta \Phi^A} (\omega^{-1})^{AB} \frac{\delta Y}{\delta \Phi^B}. \quad (\text{A7})$$

Given a family of transformations $\delta_{\tau} \Phi^A$ labeled by a parameter τ , the Hamiltonian $H[\tau]$ that generates the transformations is determined by the condition

$$\delta H[\tau] = \Omega[\delta \Phi, \delta_{\tau} \Phi] \quad (\text{A8})$$

for an arbitrary variation $\delta \Phi$. Using (A3), we can see that this is just a disguised form of Hamilton's equations of motion,

$$\delta_{\tau} \Phi^A = (\omega^{-1})^{AB} \frac{\delta H[\tau]}{\delta \Phi^B}. \quad (\text{A9})$$

The Poisson bracket of two such generators is

$$\{H[\tau_1], H[\tau_2]\} = \delta_{\tau_2} H[\tau_1] = -\Omega[\delta_{\tau_1} \Phi, \delta_{\tau_2} \Phi]. \quad (\text{A10})$$

APPENDIX B: GAUSSIAN NULL COORDINATES AND DIMENSIONAL REDUCTION

In Sec. III, Gaussian null coordinates were used to help reduce the D -dimensional Einstein-Hilbert action to an effective two-dimensional form. Here, I describe these coordinates in a bit more detail.

We start with the Gaussian null coordinate system⁴ described in Appendix A of Ref. [39] and in Ref. [48]. In such coordinates, a general metric takes the form

$$ds^2 = -r \cdot f dv^2 + 2drdv + 2r \cdot h_{\mu} dv dy^{\mu} + \phi_{\mu\nu} dy^{\mu} dy^{\nu}. \quad (\text{B1})$$

The surface $r = 0$ is null; here, we will take it to be the horizon Δ .

The coordinates (B1) have clear geometrical meanings. The horizon $r = 0$ is a null surface with null normal $\ell_A dz^A = dr$. (The surface gravity is $\kappa = \frac{1}{2} f|_{r=0}$). Since Δ is null, its normals are also tangent vectors; indeed, the integral curves of the tangent vectors $\ell^A \partial_A = \frac{\partial}{\partial v}$ are the null geodesic generators of Δ . The ‘‘orthogonal’’ vectors $n^A \partial_A = \frac{\partial}{\partial r}$ are null even off the horizon, and their integral curves are null geodesics transverse to the horizon. As in Sec. III, the coordinates y^{μ} parametrize a spacelike cross section $\hat{\Delta}$ of the horizon, and may be extended to a neighborhood of Δ by requiring that they be constant on both sets of null geodesics.

⁴My notation differs a bit from that of Ref. [39]; I use different index conventions, my v is their u , my ℓ^a is their k^a , and my n_a is their ℓ_a .

The coordinate r is an affine parameter along the transverse geodesics, and thus provides a natural geometric notion of ‘‘distance from the horizon.’’ For the Schwarzschild metric, in particular, Gaussian null coordinates are Eddington-Finkelstein coordinates with r shifted to vanish at the horizon. Near the horizon, $r \approx \rho^2/8m$, where ρ is the proper distance to the horizon at constant time. Appendix A of Ref. [48] gives an explicit expression for the Kerr-Newman metric in Gaussian null coordinates.

As claimed in Sec. III, the components $A_a{}^{\mu}$ of Yoon's metric (3.2)—here of the form $r \cdot h_{\mu}$ —vanish on the horizon, and are $\mathcal{O}(r)$ near Δ . In fact, a direct calculation in these coordinates shows that the ordinary two-dimensional scalar curvature R differs from the quantity \hat{R} of Eq. (3.5) by terms of order r^2 , justifying the near-horizon form (4.1) of the action.

The metric (B1) is of the general Kaluza-Klein-like form (3.2). But Gaussian null coordinates are too restrictive to exhibit the full set of available symmetries. As in Sec. III, though, we can move out of the Polyakov-like gauge by allowing an arbitrary two-dimensional coordinate transformation $x \rightarrow \bar{x}(x)$. This will restore the general structure of the metric (3.2), while still restricting the y dependence of the metric; for instance, although it will no longer be the case that $\sqrt{-\bar{g}} = 1$, it will remain true that $\partial_{\mu} \sqrt{-\bar{g}} = 0$. It may be checked that after such a transformation, R continues to differ from \hat{R} only by terms of order r^2 .

We may next ask how the D -dimensional vectors ℓ^A and n^A are related to their two-dimensional counterparts ℓ^a and n^a of Sec. IV. For ℓ^A , this is simple: we have only defined ℓ^A on the horizon, where it is the tangent field to the null generators of Δ , and thus coincides with ℓ^a . For n^A , the essential feature is that its D -dimensional integral curves are affinely parametrized null geodesics:

$$n^B \nabla_B n_A = 0 = n^B (\nabla_B n_A - \nabla_A n_B) = n^B (\partial_B n_A - \partial_A n_B), \quad (\text{B2})$$

where I have used the fact that $n_A n^A = 0$. Now, the only nonvanishing components of n in Gaussian null coordinates are n^r and n_v , and under two-dimensional coordinate transformations $x \rightarrow \bar{x}(x)$, it remains true that only two-dimensional components n^a and n_b are present. Thus, Eq. (B2) becomes

$$n^b (\partial_b n_a - \partial_a n_b) = 0 = n^{b(2)} \nabla_b n_a, \quad (\text{B3})$$

which was the defining property of n_a in Sec. IV. The D -dimensional transverse vectors n_A thus coincide with the two-dimensional vectors n_a . In particular, the affine parameter r gives a good measure of distance from the horizon in both D and two dimensions.

Note that if F is any function that vanishes at the horizon and is smooth near Δ ,

$$F = r\partial_r F + \mathcal{O}(r^2) = r\bar{D}F + \mathcal{O}(r^2). \quad (\text{B4})$$

In particular,

$$D\varphi = r\bar{D}D\varphi + \mathcal{O}(r^2), \quad (\text{B5})$$

quantifying the notion that $D\varphi$ is a measure of distance from the horizon.

APPENDIX C: SOME DETAILS OF NEAR-HORIZON SYMMETRIES

Section VI discussed a “near-horizon symmetry” that played a crucial role in counting states. Specifically, I argued that for any η satisfying $\bar{D}\eta \triangleq 0$, the transformation

$$\hat{\delta}_\eta\varphi = \nabla_a(\eta\ell^a) + \mathcal{L}_\zeta\varphi, \quad (\text{C1a})$$

$$\hat{\delta}_\eta g_{ab} = X_\eta \frac{D\varphi}{\bar{D}D\varphi} g_{ab} + \mathcal{L}_\zeta g_{ab}, \quad (\text{C1b})$$

$$\hat{\delta}_\eta\chi = \mathcal{L}_\zeta\chi, \quad (\text{C1c})$$

$$\text{with } \zeta^a = -\frac{D(D+\kappa)\eta}{\bar{D}D\varphi} n^a$$

$$\text{and } 2D(D+\kappa)X_\eta + \bar{\zeta} \bar{D}DR = 0 \quad (\text{C1d})$$

is an “approximate symmetry” of the action (4.1) with horizon boundary conditions (5.3a)–(5.3c), in the sense that the variation of the action could be made “arbitrarily small.”

As stated, this claim is a bit ambiguous. First of all, the variation $\hat{\delta}_\eta I$ will be inherently small as the support of η shrinks to a small neighborhood of the horizon, simply because the integration region becomes small. Second, for

this particular variation it may be seen from (7.6a)–(7.6c) that the symmetry algebra is unchanged under a constant rescaling $\eta \rightarrow k\eta$, while the variation $\hat{\delta}_\eta I$ of the action scales by k . It is thus not entirely clear what “small” means.

To remove these ambiguities, let us define an approximate near-horizon symmetry as one for which the quantity

$$\bar{\delta}_\eta I = \hat{\delta}_\eta I / \int |\eta| \epsilon \quad (\text{C2})$$

becomes arbitrarily small as the support of η shrinks to a small enough neighborhood of the horizon. (The absolute value in the denominator eliminates problems that could occur if $\int_\Delta \eta n_a = 0$.) This expression is invariant under rescalings of η , and the integral in the denominator compensates for the effects of a shrinking region of integration. If, as in Eq. (C5) below, η has support only in a band $r < \epsilon$, this condition is roughly equivalent to normalizing η at the horizon and then demanding that $\hat{\delta}_\eta I$ go to zero faster than ϵ .

To apply this criterion to the transformations (C1a)–(C1c), we should first check that they preserve our boundary conditions (5.3a)–(5.3c). Condition (5.3c) simply tells us that the Weyl transformation $\hat{\delta}_\eta g_{ab}$ acts only on ℓ_a and not on n_a (and therefore on n^a and not ℓ^a). Condition (5.3a) then determines the form of the transverse diffeomorphism $\bar{\zeta}$, and as shown in Sec. VI, condition (5.3b) gives the equation in (C1d) that fixes X_η .

We next examine the effect of this transformation on the action (4.1). The action is diffeomorphism invariant, so we can ignore ζ^a and consider only the shift of the dilaton and the Weyl transformation of the metric. The variation of the action is then

$$\delta_\eta I = \int \left[\frac{\delta I}{\delta\varphi} \hat{\delta}_\eta\varphi + \frac{\delta I}{\delta g_{ab}} \hat{\delta}_\eta g_{ab} \right] \epsilon = \frac{1}{16\pi G} \int \left[\nabla_a(\eta\ell^a) \left(R + \frac{dV}{d\varphi} \right) + X_\eta \frac{D\varphi}{\bar{D}D\varphi} (-\square\varphi + V) \right] \epsilon = \frac{1}{16\pi G} \int [\eta A + X_\eta B] \epsilon \quad (\text{C3})$$

with

$$A = -D \left(R + \frac{dV}{d\varphi} \right) = -DR - D\varphi \frac{d^2V}{d\varphi^2} = -r \left(\bar{D}DR + \bar{D}D\varphi \frac{d^2V}{d\varphi^2} \right) + \mathcal{O}(r^2) \quad (\text{C4a})$$

$$B = \frac{D\varphi}{\bar{D}D\varphi} (-\square\varphi + V) = \frac{D\varphi}{\bar{D}D\varphi} (2\bar{D}D\varphi + V) = r(2\bar{D}D\varphi + V) + \mathcal{O}(r^2), \quad (\text{C4b})$$

where in the last equalities I have used (B4) to write the result in Gaussian null coordinates.

We have assumed that the parameter η falls off rapidly away from the horizon—this is, after all, a “near-horizon” symmetry. Let us make this explicit by writing

$$\eta = \eta_\Delta \cdot \Psi_\epsilon(r), \quad (\text{C5})$$

where η_Δ is the restriction of η to the horizon and Ψ is a smooth bump function

$$\Psi_\varepsilon(r) = \begin{cases} 1 & r = 0 \\ \text{smooth interpolation} & 0 < r < \varepsilon \\ 0 & r > \varepsilon \end{cases}$$

$$\text{with } \partial_r \Psi_\varepsilon|_{r=0} = 0, \quad (\text{C6})$$

where the last condition ensures that $\bar{D}\eta \triangleq 0$. The variation (C3) is then

$$\delta_\eta I = \frac{\varepsilon^2}{32\pi G} \int_\Delta \left[-\eta_\Delta \left(\bar{D}DR + \bar{D}D\varphi \frac{d^2V}{d\varphi^2} \right) + X_{\eta_\Delta} (2\bar{D}D\varphi + V) \right] n_a + \mathcal{O}(\varepsilon^3). \quad (\text{C7})$$

The denominator in (C2), on the other hand, is

$$\int |\eta| \varepsilon = \varepsilon \int_\Delta |\eta_\Delta| n_a + \mathcal{O}(\varepsilon^2). \quad (\text{C8})$$

Thus, as long as the integrand in (C7) remains well behaved near the horizon, the variation $\bar{\delta}_\eta I$ is of order ε , and can be made arbitrarily small by shrinking the support of η .

Note that while η must have large ($\mathcal{O}(1/\varepsilon)$) radial derivatives, these never appear in the variation of the action. For the first term in (C7), this is obvious; for the second, it follows from the fact that the defining equation (C1d) for X_η involves no radial derivatives. Recall also from Sec. III that the corrections to the near-horizon form (4.1) of the action are at most of order r , so any additional variation of the action coming from these terms will also fall off as ε^2 .

-
- [1] J. D. Bekenstein, Black holes and entropy, *Phys. Rev. D* **7**, 2333 (1973).
 - [2] S. W. Hawking, Black hole explosions, *Nature (London)* **248**, 30 (1974).
 - [3] S. Carlip, Black Hole Entropy from Bondi-Metzner-Sachs Symmetry at the Horizon, *Phys. Rev. Lett.* **120**, 101301 (2018).
 - [4] V. Iyer and R. M. Wald, Some properties of Noether charge and a proposal for dynamical black hole entropy, *Phys. Rev. D* **50**, 846 (1994).
 - [5] S. Carlip, Black hole entropy and the problem of universality, in *Quantum Mechanics of Fundamental Systems: The Quest for Beauty and Simplicity*, edited by M. Henneaux and J. Zanelli (Springer, New York, 2009).
 - [6] A. Strominger and C. Vafa, Microscopic origin of the Bekenstein-Hawking entropy, *Phys. Lett. B* **379**, 99 (1996).
 - [7] S. Carlip, Statistical mechanics and black hole entropy, in *Field Theory, Integrable Systems and Symmetries*, edited by F. Khanna and L. Vinet (Les Publications CRM, Montreal, 1997).
 - [8] J. A. Cardy, Operator content of two-dimensional conformally invariant theories, *Nucl. Phys.* **B270**, 186 (1986).
 - [9] H. W. J. Blöte, J. A. Cardy, and M. P. Nightingale, Conformal Invariance, the Central Charge, and Universal Finite Size Amplitudes at Criticality, *Phys. Rev. Lett.* **56**, 742 (1986).
 - [10] A. Bagchi, S. Detournay, R. Fareghbal, and J. Simón, Holography of 3D Flat Cosmological Horizons, *Phys. Rev. Lett.* **110**, 141302 (2013).
 - [11] A. Strominger, Black hole entropy from near horizon microstates, *J. High Energy Phys.* **02** (1998) 009.
 - [12] D. Birmingham, I. Sachs, and S. Sen, Entropy of three-dimensional black holes in string theory, *Phys. Lett. B* **424**, 275 (1998).
 - [13] S. Carlip, Black Hole Entropy from Conformal Field Theory in any Dimension, *Phys. Rev. Lett.* **82**, 2828 (1999).
 - [14] S. N. Solodukhin, Conformal description of horizon's states, *Phys. Lett. B* **454**, 213 (1999).
 - [15] S. Carlip, Effective conformal descriptions of black hole entropy, *Entropy* **13**, 1355 (2011).
 - [16] B. R. Majhi and T. Padmanabhan, Noether current from the surface term of gravitational action, Virasoro algebra and horizon entropy, *Phys. Rev. D* **86**, 101501 (2012).
 - [17] S. Silva, Black hole entropy and thermodynamics from symmetries, *Classical Quantum Gravity* **19**, 3947 (2002).
 - [18] S. Carlip, Extremal and nonextremal Kerr/CFT correspondences, *J. High Energy Phys.* **04** (2011) 076; *J. High Energy Phys.* **01** (2012) 8.
 - [19] S. Carlip, Entropy from conformal field theory at Killing horizons, *Classical Quantum Gravity* **16**, 3327 (1999).
 - [20] O. Dreyer, A. Ghosh, and J. Wisniewski, Black hole entropy calculations based on symmetries, *Classical Quantum Gravity* **18**, 1929 (2001).

- [21] J. Koga, Asymptotic symmetries on Killing horizons, *Phys. Rev. D* **64**, 124012 (2001).
- [22] O. Dreyer, A. Ghosh, and A. Ghosh, Entropy from near-horizon geometries of Killing horizons, *Phys. Rev. D* **89**, 024035 (2014).
- [23] J. M. Izquierdo, J. Navarro-Salas, and P. Navarro, Kaluza-Klein theory, AdS/CFT correspondence and black hole entropy, *Classical Quantum Gravity* **19**, 563 (2002).
- [24] M. Cadoni and S. Mignemi, Entropy of 2-D black holes from counting microstates, *Phys. Rev. D* **59**, 081501 (1999).
- [25] A. Giacomini and N. Pinamonti, Black hole entropy from classical Liouville theory, *J. High Energy Phys.* **02** (2003) 014.
- [26] S. Carlip, Near Horizon Conformal Symmetry and Black Hole Entropy, *Phys. Rev. Lett.* **88**, 241301 (2002).
- [27] G. Kang, J.-I. Koga, and M.-I. Park, Near-horizon conformal symmetry and black hole entropy in any dimension, *Phys. Rev. D* **70**, 024005 (2004).
- [28] A. C. Wall, A proof of the generalized second law for rapidly changing fields and arbitrary horizon slices, *Phys. Rev. D* **85**, 104049 (2012).
- [29] G. Barnich and F. Brandt, Covariant theory of asymptotic symmetries, conservation laws and central charges, *Nucl. Phys.* **B633**, 3 (2002).
- [30] G. Barnich and G. Compere, Classical central extension for asymptotic symmetries at null infinity in three spacetime dimensions, *Classical Quantum Gravity* **24**, F15 (2007).
- [31] A. Bagchi, Correspondence between Asymptotically Flat Spacetimes and Nonrelativistic Conformal Field Theories, *Phys. Rev. Lett.* **105**, 171601 (2010).
- [32] G. Barnich and B. Oblak, Notes on the BMS group in three dimensions: I. Induced representations, *J. High Energy Phys.* **06** (2014) 129.
- [33] G. Barnich and B. Oblak, Notes on the BMS group in three dimensions: II. Coadjoint representation, *J. High Energy Phys.* **03** (2015) 033.
- [34] A. Campoleoni, H. A. Gonzalez, B. Oblak, and M. Riegler, BMS modules in three dimensions, *Int. J. Mod. Phys. A* **31**, 1650068 (2016).
- [35] A. Bagchi, M. Gary, and Zodinmawia, The nuts and bolts of the BMS Bootstrap, *Classical Quantum Gravity* **34**, 174002 (2017).
- [36] A. Bagchi, A. Saha, and Zodinmawia, BMS characters and modular invariance, *J. High Energy Phys.* **07** (2019) 138.
- [37] S. Ramanujan and G. H. Hardy, Asymptotic formula in combinatory analysis, *Proc. London Math. Soc. (Ser. 2)* **17**, 75 (1918).
- [38] S. Carlip, What we don't know about BTZ black hole entropy, *Classical Quantum Gravity* **15**, 3609 (1998).
- [39] H. Friedrich, I. Racz, and R. M. Wald, On the rigidity theorem for space-times with a stationary event horizon or a compact Cauchy horizon, *Commun. Math. Phys.* **204**, 691 (1999).
- [40] D. Grumiller, W. Kummer, and D. V. Vassilevich, Dilaton gravity in two-dimensions, *Phys. Rep.* **369**, 327 (2002).
- [41] A. Ashtekar, C. Beetle, and S. Fairhurst, Isolated horizons: A generalization of black hole mechanics, *Classical Quantum Gravity* **16**, L1 (1999).
- [42] J. Lewandowski and T. Pawłowski, Quasi-local rotating black holes in higher dimension: Geometry, *Classical Quantum Gravity* **22**, 1573 (2005).
- [43] M. Korzynski, J. Lewandowski, and T. Pawłowski, Mechanics of multidimensional isolated horizons, *Classical Quantum Gravity* **22**, 2001 (2005).
- [44] J.-H. Yoon, (1 + 1)-dimensional methods for general relativity, in *Directions in General Relativity*, edited by B. L. Hu and T. A. Jacobson (Cambridge University Press, Cambridge, England, 1993), Vol. 2.
- [45] J.-H. Yoon, Kaluza-Klein formalism of general space-times, *Phys. Lett. B* **451**, 296 (1999).
- [46] A. M. Polyakov, Quantum gravity in two-dimensions, *Mod. Phys. Lett. A* **02**, 893 (1987).
- [47] H. Bondi, Gravitational waves in general relativity, *Nature (London)* **186**, 535 (1960).
- [48] I. Booth, Spacetime near isolated and dynamical trapping horizons, *Phys. Rev. D* **87**, 024008 (2013).
- [49] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984), Appendix B.
- [50] D. Louis-Martinez and G. Kunstatter, Two-dimensional dilaton gravity coupled to an Abelian gauged field, *Phys. Rev. D* **52**, 3494 (1995).
- [51] R.-G. Cai and L.-M. Cao, Mechanics of apparent horizon in two dimensional Dilaton gravity, *Fundam. Theor. Phys.* **187**, 31 (2017).
- [52] E. Gourgoulhon and J. L. Jaramillo, A 3 + 1 perspective on null hypersurfaces and isolated horizons, *Phys. Rep.* **423**, 159 (2006).
- [53] L. Donnay, G. Giribet, H. A. González, and M. Pino, Supertranslations and Superrotations at the Black Hole Horizon, *Phys. Rev. Lett.* **116**, 091101 (2016).
- [54] P. Francesco, P. Mathieu, and D. Senechal, *Conformal Field Theory* (Springer, New York, 1997), Sec. V.4.
- [55] V. I. Arnold, *Mathematical Methods of Classical Mechanics* (Springer, New York, 1978), Appendix 5.
- [56] J. D. Brown and M. Henneaux, Central charges in the canonical realization of asymptotic symmetries: An example from three-dimensional gravity, *Commun. Math. Phys.* **104**, 207 (1986).
- [57] E. Witten, On holomorphic factorization of WZW and coset models, *Commun. Math. Phys.* **144**, 189 (1992).
- [58] C. Eling, On the membrane paradigm and spontaneous breaking of horizon BMS symmetries, *J. High Energy Phys.* **07** (2016) 065.
- [59] S. W. Hawking, M. J. Perry, and A. Strominger, Superrotation charge and supertranslation hair on black holes, *J. High Energy Phys.* **05** (2017) 161.
- [60] R. Fareghbal and A. Naseh, Rindler/Contracted-CFT correspondence, *J. High Energy Phys.* **06** (2014) 134.
- [61] H. Afshar, D. Grumiller, W. Merbis, A. Perez, D. Tempo, and R. Troncoso, Soft hairy horizons in three spacetime dimensions, *Phys. Rev. D* **95**, 106005 (2017).
- [62] L. Donnay and C. Marteau, Carrollian Physics at the black hole horizon, *Classical Quantum Gravity* **36**, 165002 (2019).
- [63] S. Carlip, Four-Dimensional Entropy from Three-Dimensional Gravity, *Phys. Rev. Lett.* **115**, 071302 (2015).
- [64] V. P. Frolov, D. Fursaev, and A. Zelnikov, CFT and black hole entropy in induced gravity, *J. High Energy Phys.* **03** (2003) 038.

- [65] S. Carlip, Black Hole Thermodynamics from Euclidean Horizon Constraints, *Phys. Rev. Lett.* **99**, 021301 (2007).
- [66] R. Emparan and I. Sachs, Quantization of AdS₃ Black Holes in External Fields, *Phys. Rev. Lett.* **81**, 2408 (1998).
- [67] A. Ashtekar and A. Magnon, Quantum fields in curved space-times, *Proc. R. Soc. A* **346**, 375 (1975).
- [68] A. Ashtekar, L. Bombelli, and O. Reula, The covariant phase space of asymptotically flat gravitational fields, in *Analysis, Geometry and Mechanics: 200 Years after Lagrange*, edited by M. Francaviglia (North-Holland, Amsterdam, 1991).
- [69] J. Lee and R. M. Wald, Local symmetries and constraints, *J. Math. Phys. (N.Y.)* **31**, 725 (1990).
- [70] C. Crnkovic and E. Witten, Covariant description of canonical formalism in geometrical theories, in *Three Hundred Years of Gravitation*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1987).