The boundary S-matrix and the AdS to CFT dictionary

STEVEN B. GIDDINGS[†]

Department of Physics University of California Santa Barbara, CA 93106-9530

Abstract

An S-matrix analog is defined for anti-de Sitter space by constructing "in" and "out" states that asymptote to the timelike boundary. A derivation parallel to that of the LSZ formula shows that this "boundary S-matrix" is given directly by correlation functions in the boundary conformal theory. This provides a key entry in the AdS to CFT dictionary.

[†] Email address: giddings@physics.ucsb.edu

The conjectured AdS/CFT correspondence[1] has offered a promising new window into the dynamics of string/M theory. But in order to exploit this powerful framework, we must decipher the holographic relationship between the bulk and boundary theories. Gubser, Klebanov, and Polyakov[2] and Witten[3] made important progress in this regard by providing a CFT to AdS dictionary: they show how to derive CFT correlation functions from the bulk theory in AdS. This has allowed the successful calculation of various CFT correlators.

However, in order to study bulk physics, and in particular to understand the undoubtedly profound implications of holography, a reverse dictionary is needed: we need to know which bulk quantities can be calculated, and how to calculate them, from the boundary CFT. Another important and closely related question is how to treat scattering in AdS. Due to the periodicity of particle orbits and lack of ordinary asymptotic states in AdS, a conventional S-matrix cannot be defined.¹ However, [4] outlined the definition of an AdS analog of the S-matrix in terms of scattering of states from the timelike infinity. Refs. [5,6] gave a related definition in the infinite-N limit. The purpose of the present note will be to go further and provide an intrinsic and explicit definition of this "boundary" S-matrix for arbitrary N, and to give a precise relation between it and the CFT correlators. The discussion also clarifies the relation between the framework of [5,6] and that of [4]. Other recent treatments of related aspects of the AdS/CFT dictionary include[7,8,9].

To summarize in advance, the boundary S-matrix will be defined as an overlap of certain "in" and "out" states. These will be defined so that they correspond to particles asymptotic to the timelike boundary of AdS in the past/future. An AdS analog of the LSZ formula can then be derived and relates this S-matrix to the bulk correlation functions. Finally the results of [3] are used to rewrite the boundary S-matrix in terms of the CFT correlation functions. An extremely simple relationship results: the boundary S-matrix *equals* the corresponding CFT correlator. This serves as a key entry in the AdS to CFT dictionary.

For simplicity we will consider scalar fields, with action

$$S = -\int dV \left[\frac{1}{2} (\nabla \Phi)^2 + \frac{m^2}{2} \Phi^2 + U(\Phi) \right] , \qquad (1)$$

¹ See [4] for more discussion.

where U summarizes the interaction terms. The generalization to other fields should be straightforward. We will work in global coordinates $x = (t, \rho, \Omega)$ for AdS_{d+1} ,

$$ds^{2} = R^{2} (-\sec^{2}\rho \, dt^{2} + \sec^{2}\rho \, d\rho^{2} + \tan^{2}\rho \, d\Omega_{d-1}^{2}) , \qquad (2)$$

although translation to Poincaré coordinates should also be straightforward.

Certain facts about the solutions to the free equations will be useful in the following. The effective gravitational potential of anti-de Sitter space confines particles to its interior. Solutions to the free equation

$$(\Box - m^2)\phi = 0 \tag{3}$$

therefore exist at arbitrary frequency ω , but are only normalizable (in the Klein-Gordon norm) for a discrete set of frequencies. Define the parameters h_{\pm} and ν by

$$2h_{\pm} = \frac{d}{2} \pm \nu \; ; \; \nu = \frac{1}{2}\sqrt{d^2 + 4m^2R^2} \; . \tag{4}$$

Normalizable solutions with definite angular momenta are of the form

$$\phi_{nl\vec{m}} = e^{-i\omega_{nl}t} Y_{l\vec{m}} \chi_{nl}(\rho) \tag{5}$$

and have asymptotic behavior

$$\chi_{nl}(\rho) \xrightarrow{\rho \to \pi/2} k_{nl}(\cos \rho)^{2h_+} \tag{6}$$

for some constants k_{nl} . Explicit formulas for these solutions are given in [10]. The discrete eigenfrequencies are

$$\omega_{nl} = 2h_+ + 2n + l , \ n = 0, 1, 2, \cdots .$$
(7)

Non-normalizable solutions we will write as

$$\phi_{\omega l\vec{m}} = e^{-i\omega t} Y_{l\vec{m}} \chi_{\omega l}(\rho) .$$
(8)

These have asymptotic behavior

$$\chi_{\omega l} \stackrel{\rho \to \pi/2}{\longrightarrow} (\cos \rho)^{2h_{-}} , \qquad (9)$$

where a convenient normalization convention has been chosen by fixing the overall constant.

We will also require some assumptions about the spectrum. This is classified according to the representations of SO(d, 2). A given representation is determined by its weight Δ (which corresponds to the conformal weight in the CFT), and contains states $|\Delta; n, l, \vec{m}\rangle$. Here *n* is the principal quantum number and l, \vec{m} are the standard angular quantum numbers, as above. The energy, defined with respect to global time, is given by

$$\omega_{n\ell}^{\Delta} = \Delta + 2n + l \;. \tag{10}$$

For a free field,

$$\Delta = 2h_+ \ . \tag{11}$$

We assume that the states of the interacting scalar theory consist of the vacuum, $|0\rangle$, the single particle states, $|\Delta; n, l, \vec{m}\rangle$, and multiparticle states $|\Delta_{\beta}; n, l, \vec{m}\rangle_{\beta}$ where β is an additional state label. For the interacting field Δ may be renormalized and is not necessarily given in terms of the bare mass by (11). We also assume that

$$\Delta_{\beta} > \Delta \tag{12}$$

for all multi-particle states.

Some useful properties of the bulk-boundary propagator are also needed. Suppose that we seek a solution of the free equation (3) satisfying the boundary condition

$$\phi \stackrel{\rho \to \pi/2}{\longrightarrow} (\cos \rho)^{2h_-} f(b) , \qquad (13)$$

where $b = (t, \Omega)$ denotes the boundary coordinates and f is some specified boundary value. Witten[3] defines the bulk-boundary Green function to be the kernel that provides the solution:²

$$\phi(x) = \int db f(b) G_{B\partial}(b, x) .$$
(14)

Explicit expressions for $G_{B\partial}$ can then be found[3] using the resulting condition that $G_{B\partial}$ must asymptote to a delta function at the boundary.

It is easy to derive an alternate formula for $G_{B\partial}$ in terms of the bulk Feynman propagator $G_B(x, x')$ using an AdS variant of the usual Green's theorem argument. Consider the solution ϕ with the above boundary conditions (13). Define a region V by $\rho < \bar{\rho} \approx \pi/2$. Using

$$(\Box_x - m^2)G_B(x, x') = -\delta(x, x')$$
(15)

 $^{^{2}}$ Though Witten's definition was made in the euclidean continuation of AdS, the formalism naturally extends to lorentzian signature as discussed in [11,4].

we may rewrite ϕ as

$$\phi(x') = -\int_{V} dV \phi(x) (\Box_{x} - m^{2}) G_{B}(x, x')$$
(16)

and then integrate twice by parts to find

$$\phi(x') = -\int_{\partial V} d\Sigma^{\mu} \phi(x) \overleftrightarrow{\partial_{\mu}} G_B(x, x') .$$
(17)

At the boundary the Feynman propagator scales as

$$G_B(x,x') \xrightarrow{\rho \to \pi/2} (\cos \rho)^{2h_+} G(b,x') , \qquad (18)$$

for some function G. Substituting this and the boundary behavior (13) into (17) gives

$$\phi(x') = -R^{d-1} \int db f(b) G(b, x') \lim_{\rho \to \pi/2} (\tan \rho)^{d-1} \left[(\cos \rho)^{2h_-} \overleftrightarrow{\partial_{\rho}} (\cos \rho)^{2h_+} \right] , \qquad (19)$$

with limit

$$\phi(x') = 2\nu R^{d-1} \int db f(b) G(b, x') .$$
(20)

Comparison with (14) then shows

$$G_{B\partial}(b,x') = 2\nu R^{d-1} \lim_{\rho \to \pi/2} (\cos \rho)^{-2h_+} G_B(x,x') , \qquad (21)$$

in agreement with [12].

Notice that (3), (13) do not uniquely specify the solution ϕ ; one may always add a normalizable mode without modifying the boundary behavior (13). Specifically, suppose that f(b) falls to zero in the far past and future. Then in the asymptotic past and future ϕ must be a linear combination of the normalizable modes (5). With the preceding construction of $G_{B\partial}$, (14) gives the solution that is purely positive frequency in the far future, and purely negative frequency in the far past. Other solutions can be obtained by modifying the temporal boundary conditions on the bulk Green function, *e.g.* by using retarded or advanced propagators.

Note that the non-normalizable solutions $\phi_{\omega l\vec{m}}$ can be recovered from (14); in the limit $f \to e^{-i\omega t} Y_{l\vec{m}}$, (9) coincides with (13). Therefore

$$\phi_{\omega l\vec{m}}(x) = \int db e^{-i\omega t} Y_{l\vec{m}} G_{B\partial}(b, x) \equiv G_{B\partial}(-\omega, l, -\vec{m}; x) .$$
⁽²²⁾

For a general f, we can therefore rewrite (14) in terms of the Fourier transform $f_{l\vec{m}}(\omega)$ as

$$\phi_f(x) = \int db f(b) G_{B\partial}(b, x) = \sum_{l, \vec{m}} \int \frac{d\omega}{2\pi} f_{l\vec{m}}(\omega) \phi_{\omega l\vec{m}}(x)$$
(23)

For appropriately chosen f, this function defines a solution corresponding to a wavepacket. The function f determines the packet profile.

These can be thought of as packets incident from infinity in AdS. Notice that, according to (9), they typically diverge at the boundary. There is a physical reason for this: motion in the region near the boundary is classically forbidden. Therefore the amplitude for a particle incident from infinity to reach the center of AdS is suppressed by an infinite tunneling factor. However, the amplitude for a particle to reach the center of AdS may be kept finite by rescaling the wavefunction such that the incident amplitude at infinity is infinite.

One concrete way to think of this is to imagine cutting off AdS at large but finite radius and patching the resulting AdS bubble into a spacetime with a bona-fide null infinity, as in [4]. A beam of particles from this asymptotic space can be focussed to collide with another beam in the center of the AdS region. Most of the incident flux is reflected off the potential barrier resulting from the AdS geometry, so in order for the beams to penetrate to the center the incident amplitudes must be large. The wavepacket definitions above, which are given intrinsically in AdS without reference to an auxiliary bubble picture, can be thought of as arising from the limit where the radius of the AdS bubble goes to infinity while simultaneously scaling up the incident beam amplitudes.

These wavepackets can now be used to construct operators that create "in" and "out" states.³ These asymptotic operators will be defined by

$$\alpha_f = \lim_{\bar{\rho} \to \pi/2} \int_{\Sigma} d\Sigma^{\mu} \phi_f^* \overleftrightarrow{\partial}_{\mu} \Phi , \qquad (24)$$

where $\Sigma = \partial V$ for the region V defined above and Φ is the full *interacting* field. We also define the plane wave limit of these operators,

$$\alpha_{\omega l\vec{m}} = \lim_{\bar{\rho} \to \pi/2} \int_{\Sigma} d\Sigma^{\mu} \phi^*_{\omega l\vec{m}} \overleftrightarrow{\partial_{\mu}} \Phi .$$
⁽²⁵⁾

³ Refs. [5,6] outlined the construction of such operators in the infinite-N limit. Here we will explicitly construct such operators for arbitrary N.

If Φ is replaced by the free field,

$$\phi = \sum_{n,l,\vec{m}} a_{nl\vec{m}} e^{-i\omega_{nl}t} \phi_{nl\vec{m}} + a^{\dagger}_{nl\vec{m}} e^{i\omega_{nl}t} \phi^*_{nl\vec{m}} , \qquad (26)$$

then (25) gives

$$\alpha_{\omega l\vec{m}} = -4\pi\nu R^{d-1} \sum_{n} k_{nl} \left[\delta(\omega - \omega_{nl}) a_{nl\vec{m}} + \delta(\omega + \omega_{nl}) a_{nl,-\vec{m}}^{\dagger} \right]$$
(27)

where the k_{nl} appeared in (6). This suggests that the positive and negative frequency $\alpha_{\omega l\vec{m}}$'s be thought of as annihilation and creation operators, respectively.

This is confirmed by the following critical relations, which hold for the operators constructed from the full interacting field:

$$\langle 0|\alpha_f|\Delta; n, l, \vec{m} \rangle = -2\nu R^{d-1} N(\Delta) k_{nl} f^*_{l\vec{m}}(\omega_{nl}) , \qquad (28)$$

$$\langle \Delta; n, l, \vec{m} | \alpha_f | 0 \rangle = -2\nu R^{d-1} N(\Delta) k_{nl} f^*_{l,-\vec{m}}(-\omega_{nl}) , \qquad (29)$$

and

$$\langle 0|\alpha_f|\Delta_\beta; n', l', \vec{m}'\rangle_\beta = 0 , \qquad (30)$$

where $N(\Delta)$ is another constant. Therefore α_f with positive-frequency f annihilates a particle at the boundary, and with negative-frequency f creates a particle at the boundary. Furthermore, (30) implies that the α_f 's only annihilate/create single particle states.

The first of these relations is proved by recalling that by symmetry, the full interacting field must satisfy[13]

$$\langle 0|\Phi(x)|\Delta;n,l,\vec{m}\rangle = N(\Delta)\phi_{nl\vec{m}}(x) \tag{31}$$

for some normalization factor $N(\Delta)$. Then the definition (25) and a derivation like that in (19)-(20) immediately gives (28). Note that in order for this to be true the modes in (23), (25) must be defined with the mass fixed by $2h_{+} = \Delta$, corresponding to using the renormalized physical mass of the single particle state. Analogous reasoning proves (29).

Eq. (30) is shown by noting that, again purely from the SO(d, 2) symmetry,

$$\langle 0|\Phi(x)|\Delta_{\beta}; n, l, \vec{m}\rangle_{\beta} = N_{\beta}(\Delta_{\beta})\phi_{nl\vec{m}}^{\Delta_{\beta}}(x) , \qquad (32)$$

where $\phi_{nl\vec{m}}^{\Delta_{\beta}}$ is defined using the mass parameter corresponding to the multiparticle Δ_{β} . Again, the matrix element (30) can be found from reasoning parallel to (19)-(20), but now the result contains

$$\lim_{\rho \to \pi/2} (\cos \rho)^{\Delta_{\beta} - \Delta} .$$
(33)

This vanishes by (12).

"In" and "out" states are now readily defined. For positive-frequency functions f_i , define

$$\alpha_{f_i}^{\rm in} = \alpha_{f_i} / Z_i , \qquad (34)$$

and for negative-frequency functions f'_j define

$$\alpha_{f'_j}^{\text{out}\dagger} = \alpha_{f'_j} / Z'_j \tag{35}$$

where the Z_i are wavefunction renormalization factors necessary to cancel normalization constants like those in (28),(29). The "in" and "out" states are then

$$|f_i\rangle_{\rm in} = \prod_i \alpha_{f_i}^{\rm in\dagger} |0\rangle \tag{36}$$

and

$$_{\text{out}}\langle f'_j| = \langle 0| \prod_j \alpha^{\text{out}}_{f'_j} .$$
(37)

These states in turn lead to construction of the boundary S-matrix. Suppose that the wavepackets f_i, f'_j are non-overlapping, and that the support of all the f'_j 's lies to the future of that of all the f_i 's. The boundary S-matrix is then defined as

$$S_{\partial}[f_1 \cdots f_m; f'_1 \cdots f'_n] \equiv \langle 0|T \prod_j \alpha_{f'_j}^{\text{out}} \prod_i \alpha_{f_i}^{\text{in}\dagger} |0\rangle .$$
(38)

Although the interpretation is less transparent, the same definition can be adopted for f_i and f'_i not satisfying the above conditions.

In flat space, the S-matrix is related by the LSZ formula to truncated correlation functions. A similar formula can now be derived for the boundary S-matrix, which will be given in terms of bulk correlation functions. Consider a finite region V' defined like V above, but with boundaries Σ_{-T} , Σ_T of constant time $\pm T$ lying to the far past and far future of the wavepackets' support. Gauss' theorem applied to (24) gives

$$\alpha_f^{\text{in}\dagger} = \frac{1}{Z} \lim_{\bar{\rho} \to \pi/2} \left[\int_{V'} dV \nabla^{\mu} (\phi_f \overleftrightarrow{\partial_{\mu}} \Phi) - \int_{\Sigma_{-T} + \Sigma_T} d\Sigma^{\mu} \phi_f \overleftrightarrow{\partial_{\mu}} \Phi \right]$$
(39)

where the $T \to \infty$ limit is also understood. The surface term at Σ_{-T} vanishes because ϕ_f is positive frequency and therefore vanishes in the far past. Inside (38), the time ordering

takes the surface term at Σ_T to the left. We can then insert a complete set of states $|s\rangle$ to find an expression of the form

$$\sum_{s} \int_{\Sigma_T} d\Sigma^{\mu} \phi_f \overleftarrow{\partial_{\mu}} \langle 0|\Phi|s \rangle \langle s|\psi \rangle .$$
(40)

This vanishes by (31). The bulk term is left, and after using the free equation for ϕ_f and taking $T \to \infty$ becomes

$$\alpha_f^{\text{in}\dagger} \approx \frac{1}{Z} \int dV \phi_f (\Box - m^2) \Phi \tag{41}$$

where \approx denotes equality inside (38). Similar arguments hold for α_f^{out} . The LSZ formula immediately follows:

$$S_{\partial}[f_1 \cdots f_m; f'_1 \cdots f'_n] = \int \prod_i \left[\frac{dV_i}{Z_i} \phi_{f_i}(x_i) \right] \prod_j \left[\frac{dV'_j}{Z'_j} \phi_{f'_j}(x'_j) \right] \langle 0|T \prod_i \Phi(x_i) \prod_j \Phi(x'_j)|0\rangle_T .$$

$$\tag{42}$$

Here the kinetic operator in (41) has removed the external legs, and the "T" subscript denotes the resulting truncated Green function.

The relation to correlation functions in the conformal field theory now follows trivially. Witten[3] has shown that the CFT correlators are given in terms of the truncated bulk correlators and the bulk-boundary propagator as

$$\langle T \prod_{a} \mathcal{O}(b_{a}) \rangle = \int \prod_{a} \left[dV_{a} G_{B\partial}(b_{a}, x_{a}) \right] \langle 0|T \prod_{a} \Phi(x_{a})|0\rangle_{T} .$$

$$\tag{43}$$

After substituting (23) into (42), S_{∂} is therefore given by

$$S_{\partial}[f_1 \cdots f_m; f'_1 \cdots f'_n] = \int \prod_i \left[db_i \frac{f_i(b_i)}{Z_i} \right] \int \prod_j \left[db'_j \frac{f'_j(b'_j)}{Z'_j} \right] \langle T \prod_i \mathcal{O}(b_i) \prod_j \mathcal{O}(b'_j) \rangle \quad ,$$

$$(44)$$

or in the plane wave limit $f \to e^{-i\omega t} Y_{l\vec{m}}$,

$$S_{\partial}[\{\omega_i, l_i, \vec{m}_i\}; \{\omega'_j, l'_j, \vec{m}'_j\}] = \langle T \prod_i \frac{1}{Z_i} \mathcal{O}(\omega_i, l_i, \vec{m}_i) \prod_j \frac{1}{Z'_j} \mathcal{O}(\omega'_j, l'_j, \vec{m}'_j) \rangle .$$
(45)

These strikingly simple relations tells us that the CFT correlators directly determine the boundary S-matrix, and thus provide an extremely simple dictionary between scattering amplitudes in anti-de Sitter space and the conformal field theory correlation functions.

Acknowledgments

I have greatly benefited from discussions with V. Balasubramanian and A. Lawrence in the course of writing [4]. This work was partially supported by DOE contract DE-FG-03-91ER40618.

References

- J. Maldacena, "The large-N limit of superconformal field theories and supergravity", hep-th/9711200, Adv. Theor. Math. Phys. 2 (1998) 231.
- S.S. Gubser, I. Klebanov and A. Polyakov, "Gauge theory correlators from noncritical string theory", hep-th/9802109, Phys. Lett. B428 (1998) 105.
- [3] E. Witten, "Anti-de Sitter space and holography", hep-th/9802150, Adv. Theor. Math. Phys. 2 (1998) 253.
- [4] V. Balasubramanian, S.B. Giddings, and A. Lawrence, "What do CFTs tell us about anti-de Sitter spacetimes?" hep-th/9902052.
- [5] J. Polchinski, "S-matrices from AdS spacetime", hep-th/9901076.
- [6] L. Susskind, "Holography in the flat-space limit", hep-th/9901079.
- G. Horowitz and N. Itzhaki, "Black holes, shock waves, and causality in the AdS/CFT correspondence," hep-th/9901012.
- [8] I.Ya. Aref'eva, "On the holographic S-matrix," hep-th/9902106.
- [9] J. Polchinski and L. Susskind, "Puzzles and paradoxes about holography," hepth/9902182.
- [10] V. Balasubramanian, P. Kraus and A. Lawrence, "Bulk vs. Boundary dynamics in anti-de Sitter spacetime", hep-th/9805171, Phys. Rev D59:046003 (1999).
- [11] V. Balasubramanian, P. Kraus, A. Lawrence and S. Trivedi, "Holographic probes of anti-de Sitter space-times", hep-th/9808017, Harvard preprint HUTP-98/A057, Caltech preprint CALT68-2189, Fermilab preprint Pub-98/240-T
- [12] T. Banks, M. Douglas, G. Horowitz and E. Martinec, "AdS dynamics from conformal field theory", hep-th/9808016, UCSB/ITP preprint NSF-ITP-98082, U. Chicago preprint EFI-98-30.
- [13] D.W. Düsedau and D.Z. Freedman, "Lehmann spectral representation for anti-de Sitter quantum field theory", Phys. Rev. D33 (1986) 389.