

## COMMENTARY

### Nutrition for the Brain

Commentary on the article by Isaacs *et al.* on page 308

PETRA S. HÜPPI

*Department of Pediatrics, University of Geneva, Switzerland; Department of Neurology, Harvard Medical School, Children's Hospital, Boston, Massachusetts*

The question of whether brain function is entirely genetically determined or may be influenced by the environment or by nutrition has been debated for decades. Several studies have associated breastfeeding with improved intelligence in later life (1–5). Of course many confounding factors such as socioeconomic status, perinatal, and early childhood morbidities will influence both readiness for breastfeeding as well as direct cognitive development. Effects of studies rigorously controlling for maternal intelligence did not indeed find the same major beneficial effects of breastfeeding on later intelligence (6,7). Mechanisms by which breastfeeding is supposed to exert its effects on cognitive development is mainly fatty acid (FA) composition of human milk, containing polyunsaturated FA such as omega-3 polyunsaturated fatty acids, docosahexaenoic acid (DHA), and trophic factors such as insulin-like growth factor-1 (8), both important for organ development and in particular for the brain. The Avon Longitudinal Study of Parents and Children showed that IQ increased by 3.2 points for every 100 ng/mL increase of plasma insulin-like growth factor-1 levels and this in particular for the Verbal IQ (VIQ) compared with the Performance IQ (PIQ) (9). Polyunsaturated FA, such as DHA [22:6(n-3)] are important precursors of membrane lipids and are, as such, important components of brain growth and myelination. DHA is the most abundant (n-3) fatty acid in the mammalian brain. DHA in the brain can be obtained either as DHA itself or from one of its precursors,  $\alpha$ -linolenic acid or eicosapentaenoic acid. However, the rates of conversion of  $\alpha$ -linolenic acid to DHA is low in humans (less than 1% of dietary amount). Before birth, DHA is transported through the placenta *via* pathways involving fatty acid binding proteins and  $\alpha$ -fetoprotein, before release into the fetal circulation, the rate of transfer increasing during the third trimester. High dietary intake of DHA during pregnancy results in higher maternal-to-fetal transfer (10). After birth, the infant is provided with DHA in mother's milk. However, the level of DHA can vary (from less than 0.1 to 1% of milk fatty acids) depending on the amount of DHA in the mother's diet.

For the premature infant, nutrition regularly provided by the placental transfer during the third trimester dramatically changes after birth at a time point when the organism is still dependent on nutrients transferred from the placenta. Both parental nutrition as well as breast milk of mothers after premature birth provide insufficient nutritional support to the developing brain of the premature infant and may lead

to postnatal growth restriction with the known consequences of altered hormonal status including alteration of leptin expression (11). Many recent studies have therefore assessed the effects of breastfeeding and nutritional interventions on neuro-developmental outcome of premature infants (12–16), showing advantage with early breastfeeding, FA supplementation, and higher protein intake. Preterm and low birth weight infants are often growth-restricted at hospital discharge. Feeding infants posthospital discharge with calorie- and protein-enriched formula milk might therefore facilitate “catch-up” growth, but this has not been confirmed in a recent Cochrane Database review (17). The current study by Isaacs *et al.* is one of the few studies that looks at specific brain structural effects of nutritional supplementation. Human brain growth mainly takes place during the third trimester with whole brain volume more than doubling and cortical gray matter volume increasing four-fold (18) and an increase in subcortical gray matter or basal ganglia of 70% (19,20). This is also the time period in which cortical folding and gyrification take place with an increase of brain surface and degree of sulcation index (21). Conditions such as severe prematurity and cerebral white matter injury have been shown to affect brain growth and specific structural brain development with subsequent functional consequences both at birth, in infancy, early childhood, and adolescence (22–27). Regional brain growth has been shown to be different with occipital regions growing much faster than prefrontal regions and differentially affected by conditions such as prematurity affecting growth in the central regions or brain lesions affecting both central and frontal brain regions (19,21,28). All these measures have been defined *in vivo* using advanced Magnetic Resonance Image analysis tools similar to the approach taken in the study by Isaacs *et al.* The authors in this study present data on a comparison of brain structural volumes and IQ measurements in two groups of ex-preterm infants born at a gestational age below 30 wk at adolescent age treated with a different perinatal nutritional protocol. They used an atlas-based segmentation technique to define total brain and cortical gray matter volume as well as volumes of the subcortical gray matter structures, caudate nucleus, thalamus, putamen, globus pallidum, hippocampus, and amygdala and IQ testing with Wechsler Intelligence Scale for Children defining both VIQ and PIQ. The high nutrient group ex-preterm adolescents showed significantly better performance on VIQ ex-preterm measures with no differences in PIQ measures. Structurally the two groups showed significant differences in both left- and right-sided caudate volume, with the standard nutrition group showing lower caudate volumes which further correlated with IQ scores with lower volume indicating lower VIQ. This was a gender-specific effect with mainly male preterm infants being affected by these differences in perinatal nutrition. Subcortical gray matter

Received January 2, 2008; accepted January 7, 2008.

Correspondence: Petra S. Hüppi, M.D., Department of Pediatrics, Division of Child Development & Growth, University of Geneva, 6, rue Willy-Donze, CH-1211 Geneva, Switzerland; e-mail: euro.pedres@hcuge.ch

structures have been shown to be affected by premature birth with correlations to later cognitive outcome (23,29–31) as well as in neuro-psychiatric disorders such as attention deficit hyperactivity disorder (32,33) and Depression (34). Deep nuclear gray matter volume reduction at term age has been shown in a previous study to be correlated with gestational age at birth and severity of respiratory distress syndrome, so, clearly immaturity at birth and comorbidities such as severe respiratory distress which are associated with oxidative stress lead to reduction in deep cortical gray matter volume at term. Immaturity and severity of respiratory distress syndrome on the other hand are often associated with poor nutritional status in the preterm infant, and therefore the findings of the current study by Isaacs *et al.* would suggest that some of these effects might be due to insufficient nutritional support and that some of these effects can be reversed by higher nutritional support or by additional FA such as DHA. The caudate is known to be one of the brain regions expressing high DHA content and changes can be observed after dietary depletion and repletion (35). Experimentally it is known that DHA incorporates into the membrane bilayer, and increases the degree of flexibility and direct interaction with membrane proteins. This impacts on the speed of signal transduction and neurotransmission (36). Unesterified DHA acts as ligands for brain transcription factors retinoid X receptor alpha (RXR) and peroxisome proliferator-activated receptor (PPAR), which dimerize to regulate the expression of genes involved in the control of synaptic plasticity, cytoskeleton and membrane assembly, signal transduction and ion channel formation. This gene regulation function could explain the role of DHA in many aspects of development such as neurogenesis, morphologic differentiation of catecholaminergic neurons, and activity-dependent plasticity (36,37). DHA also seems to inhibit the oxidative stress-induced induction of pro-inflammatory genes and apoptosis, and provide protection against peroxidative damage of lipids and proteins in the developing brain (37,38). Leptin regulates human eating behavior by regulating striatal brain regions (39), but is also known to regulate neuronal excitability and cognitive function in particular by influencing positively the hippocampal-dependent learning and memory (40).

Another condition by which brain development can be affected in the long term is intrauterine growth restriction (IUGR) (41). Currently the IUGR rate is the highest it has been in over 20 y and is likely to rise further due to the increasing rate of infertility treatments, multiple pregnancies, older mothers, and exposure to IUGR-inducing agents such as tobacco. All these conditions lead to poor nutritional status of the fetus and subsequent alteration of structural and functional brain development with reduction in cortical gray matter volume, reduction in striatal volume, and reduction in hippocampal volume, predominantly in boys (42–45). Children with very low birth weights have multiple rather than isolated cognitive deficits including problems with attention, memory, reading and mathematics, as well as reasoning and self regulation (46,47). These cognitive deficits are likely to have an overriding central nervous impairment with underlying brain structural changes (48). Recently epidemiologic studies assessing maternal nutrition have lead to interesting observations by which maternal consumption of seafood during pregnancy lead to higher cognitive performance in their offspring, with again the most prominent effect on VIQ (49). Fatty acid metabolism is therefore an important component of both prenatal and postnatal brain development and the current study by Isaacs *et al.* illustrates how structural and functional changes in relation to nutritional interventions can be studied. Further studies should concentrate on the combined effect of genetic background (50) and nutritional interventions on both structural and functional brain development. Understanding the effects of early antenatal, perinatal, and neonatal events on later structural and functional brain develop-

ment, aberrant or regenerative, will no doubt be essential to develop interventions and treatments for preventing developmental disabilities that have their origin in early life. Research with the aim of defining which nutrient favors adequate development of brain structure and functions during gestation and early childhood, with the ultimate goal of improving cognitive development and decreasing neuro-psychiatric disorders, will be an important task in terms of public health of the future.

## REFERENCES

- Agostoni C 1996 Early breastfeeding linked to higher intelligence quotient scores. *Acta Paediatr* 85:639
- Anderson JW, Johnstone BM, Remley DT 1999 Breast-feeding and cognitive development: a meta-analysis. *Am J Clin Nutr* 70:525–535
- Daniels MC, Adair LS 2005 Breast-feeding influences cognitive development in Filipino children. *J Nutr* 135:2589–2595
- Clark KM, Castillo M, Calatroni A, Walter T, Cayazzo M, Pino P, Lozoff B 2006 Breast-feeding and mental and motor development at 51/2 years. *Ambul Pediatr* 6:65–71
- Greene LC, Lucas A, Livingstone MB, Harland PS, Baker BA 1995 Relationship between early diet and subsequent cognitive performance during adolescence. *Biochem Soc Trans* 23:376S
- Der G, Batty GD, Deary IJ 2006 Effect of breast feeding on intelligence in children: prospective study, sibling pairs analysis, and meta-analysis. *BMJ* 333:945
- Evenhouse E, Reilly S 2005 Improved estimates of the benefits of breastfeeding using sibling comparisons to reduce selection bias. *Health Serv Res* 40:1781–1802
- Eriksson U, Duc G, Froesch ER, Zapf J 1993 Insulin-like growth factors (IGF) I and II and IGF binding proteins (IGFBPs) in human colostrum/transitory milk during the first week postpartum: comparison with neonatal and maternal serum. *Biochem Biophys Res Commun* 196:267–273
- Gunnell D, Miller LL, Rogers I, Holly JM 2005 Association of insulin-like growth factor I and insulin-like growth factor-binding protein-3 with intelligence quotient among 8- to 9-year-old children in the Avon Longitudinal Study of Parents and Children. *Pediatrics* 116:e681–e686
- Innis SM 2005 Essential fatty acid transfer and fetal development. *Placenta* 26(suppl A):S70–S75
- Patel L, Cavazzoni E, Whatmore AJ, Carney S, Wales JK, Clayton PE, Gibson AT 2007 The contributions of plasma IGF-I, IGFBP-3 and leptin to growth in extremely premature infants during the first two years. *Pediatr Res* 61:99–104
- Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C 1992 Breast milk and subsequent intelligence quotient in children born preterm. *Lancet* 339:261–264
- Lucas A, Morley R, Cole TJ 1998 Randomized trial of early diet in preterm babies and later intelligence quotient. *BMJ* 317:1481–1487
- Lucas A, Fewtrell MS, Morley R, Singhal A, Abbott RA, Isaacs E, Stephenson T, MacFadyen UM, Clements H 2001 Randomized trial of nutrient-enriched formula versus standard formula for postdischarge preterm infants. *Pediatrics* 108:703–711
- O'Connor DL, Hall R, Adamkin D, Auestad N, Castillo M, Connor WE, Connor SL, Fitzgerald K, Groh-Wargo S, Hartmann EE, Jacobs J, Janowsky J, Lucas A, Margeson D, Mena P, Neuringer M, Nesin M, Singer L, Stephenson T, Szabo J, Zemon V 2001 Growth and development in preterm infants fed long-chain polyunsaturated fatty acids: a prospective, randomized controlled trial. *Pediatrics* 108:359–371
- Fewtrell MS, Morley R, Abbott RA, Singhal A, Isaacs EB, Stephenson T, MacFadyen U, Lucas A 2002 Double-blind, randomized trial of long-chain polyunsaturated fatty acid supplementation in formula fed to preterm infants. *Pediatrics* 110:73–82
- Henderson G, Fahey T, McGuire W 2005 Calorie and protein-enriched formula versus standard term formula for improving growth and development in preterm or low birth weight infants following hospital discharge. *Cochrane Database Syst Rev* CD004696
- Hüppi PS, Warfield S, Kikinis R, Barnes P, Zientara GP, Jolesz FA, Tsuji MK, Volpe JJ 1998 Quantitative magnetic resonance imaging of brain development in premature and mature newborns. *Ann Neurol* 43:224–235
- Mewes AU, Hüppi PS, Als H, Rybicki FJ, Inder TE, McNulty GB, Mulkern RV, Robertson RL, Rivkin MJ, Warfield SK 2006 Regional brain development in serial magnetic resonance imaging of low-risk preterm infants. *Pediatrics* 118:23–33
- Zacharia A, Zimine S, Lovblad KO, Warfield S, Thoeny H, Ozdoba C, Bossi E, Kreis R, Boesch C, Schroth G, Hüppi PS 2006 Early assessment of brain maturation by MR imaging segmentation in neonates and premature infants. *AJNR Am J Neuroradiol* 27:972–977
- Dubois J, Benders M, Cachia A, Lazeyras F, Ha-Vinh LR, Sizonenko SV, Borradori-Tolsa C, Mangin JF, Hüppi PS 2007 Mapping the early cortical folding process in the preterm newborn brain. *Cereb Cortex* doi:10.1093/cercor/bhm180. Epub ahead of print.
- Allin M, Matsumoto H, Santhouse AM, Nosarti C, AlAsady MH, Stewart AL, Rifkin L, Murray RM 2001 Cognitive and motor function and the size of the cerebellum in adolescents born very pre-term. *Brain* 124:60–66
- Inder TE, Warfield SK, Wang H, Hüppi PS, Volpe JJ 2005 Abnormal cerebral structure is present at term in premature infants. *Pediatrics* 115:286–294
- Limperopoulos C, Soul JS, Haidar H, Hüppi PS, Bassan H, Warfield SK, Robertson RL, Moore M, Akins P, Volpe JJ, du Plessis AJ 2005 Impaired trophic interactions

- between the cerebellum and the cerebrum among preterm infants. *Pediatrics* 116:844–850
25. Lodygensky GA, Rademaker K, Zimine S, Gex-Fabry M, Liefink AF, Lazeyras F, Groenendaal F, de Vries LS, Hüppi PS 2005 Structural and functional brain development after hydrocortisone treatment for neonatal chronic lung disease. *Pediatrics* 116:1–7
  26. Martinussen M, Fischl B, Larsson HB, Skranes J, Kulseng S, Vangberg TR, Vik T, Brubakk AM, Haraldseth O, Dale AM 2005 Cerebral cortex thickness in 15-year-old adolescents with low birth weight measured by an automated MRI-based method. *Brain* 128:2588–2596
  27. Sowell ER, Thompson PM, Leonard CM, Welcome SE, Kan E, Toga AW 2004 Longitudinal mapping of cortical thickness and brain growth in normal children. *J Neurosci* 24:8223–8231
  28. Gilmore JH, Lin W, Prastawa MW, Looney CB, Vetsa YS, Knickmeyer RC, Evans DD, Smith JK, Hamer RM, Lieberman JA, Gerig G 2007 Regional gray matter growth, sexual dimorphism, and cerebral asymmetry in the neonatal brain. *J Neurosci* 27:1255–1260
  29. Abernethy LJ, Cooke RW, Foulder-Hughes L 2004 Caudate and hippocampal volumes, intelligence, and motor impairment in 7-year-old children who were born preterm. *Pediatr Res* 55:884–893
  30. Srinivasan L, Dutta R, Counsell SJ, Allsop JM, Boardman JP, Rutherford MA, Edwards AD 2007 Quantification of deep gray matter in preterm infants at term-equivalent age using manual volumetry of 3-tesla magnetic resonance images. *Pediatrics* 119:759–765
  31. Boardman JP, Counsell SJ, Rueckert D, Kapellou O, Bhatia KK, Aljabar P, Hajnal J, Allsop JM, Rutherford MA, Edwards AD 2006 Abnormal deep grey matter development following preterm birth detected using deformation-based morphometry. *Neuroimage* 32:70–78
  32. Casey BJ, Epstein JN, Buhle J, Liston C, Davidson MC, Tonev ST, Spicer J, Niogi S, Millner AJ, Reiss A, Garrett A, Hinshaw SP, Greenhill LL, Shafritz KM, Vitolo A, Kotler LA, Jarrett MA, Glover G 2007 Frontostriatal connectivity and its role in cognitive control in parent-child dyads with ADHD. *Am J Psychiatry* 164:1729–1736
  33. Castellanos FX, Sharp WS, Gottesman RF, Greenstein DK, Giedd JN, Rapoport JL 2003 Anatomic brain abnormalities in monozygotic twins discordant for attention deficit hyperactivity disorder. *Am J Psychiatry* 160:1693–1696
  34. Caligiuri MP, Brown GG, Meloy MJ, Ebersson S, Niculescu AB, Lohr JB 2006 Striatopallidal regulation of affect in bipolar disorder. *J Affect Disord* 91:235–242
  35. Xiao Y, Huang Y, Chen ZY 2005 Distribution, depletion and recovery of docosahexaenoic acid are region-specific in rat brain. *Br J Nutr* 94:544–550
  36. Grossfield A, Feller SE, Pitman MC 2006 A role for direct interactions in the modulation of rhodopsin by omega-3 polyunsaturated lipids. *Proc Natl Acad Sci USA* 103:4888–4893
  37. Green P, Glozman S, Weiner L, Yavin E 2001 Enhanced free radical scavenging and decreased lipid peroxidation in the rat fetal brain after treatment with ethyl docosahexaenoate. *Biochim Biophys Acta* 1532:203–212
  38. Bazan NG 2006 Cell survival matters: docosahexaenoic acid signaling, neuroprotection and photoreceptors. *Trends Neurosci* 29:263–271
  39. Farooqi IS, Bullmore E, Keogh J, Gillard J, O'Rahilly S, Fletcher PC 2007 Leptin regulates striatal regions and human eating behavior. *Science* 317:1355
  40. Harvey J 2007 Leptin regulation of neuronal excitability and cognitive function. *Curr Opin Pharmacol* 7:643–647
  41. Eide MG, Oyen N, Skjaerven R, Bjerkedal T 2007 Associations of birth size, gestational age, and adult size with intellectual performance: evidence from a cohort of Norwegian men. *Pediatr Res* 62:636–642
  42. Low JA, Handley-Derry MH, Burke SO, Peters RD, Pater EA, Killen HL, Derrick EJ 1992 Association of intrauterine fetal growth retardation and learning deficits at age 9 to 11 years. *Am J Obstet Gynecol* 167:1499–1505
  43. Tolsa CB, Zimine S, Warfield SK, Freschi M, Sancho RA, Lazeyras F, Hanquinet S, Pfizenmaier M, Hüppi PS 2004 Early alteration of structural and functional brain development in premature infants born with intrauterine growth restriction 1. *Pediatr Res* 56:132–138
  44. Toft PB, Leth H, Ring PB, Peitersen B, Lou HC, Henriksen O 1995 Volumetric analysis of the normal infant brain and in intrauterine growth retardation. *Early Hum Dev* 43:15–29
  45. Lodygensky GA, Seghier ML, Warfield S, Borradori-Tolsa C, Sizonenko S, Hüppi PS 2004 Intrauterine growth restriction affects the preterm infant's hippocampus. *Pediatr Res*, in press
  46. Isaacs EB, Edmonds CJ, Lucas A, Gadian DG 2001 Calculation difficulties in children of very low birth weight: a neural correlate. *Brain* 124:1701–1707
  47. Anderson PJ, Doyle LW 2004 Executive functioning in school-aged children who were born very preterm or with extremely low birth weight in the 1990s. *Pediatrics* 114:50–57
  48. Toft PB 1999 Prenatal and perinatal striatal injury: a hypothetical cause of attention-deficit-hyperactivity disorder? *Pediatr Neurol* 21:602–610
  49. Hibbeln JR, Davis JM, Steer C, Emmett P, Rogers I, Williams C, Golding J 2007 Maternal seafood consumption in pregnancy and neurodevelopmental outcomes in childhood (ALSPAC study): an observational cohort study. *Lancet* 369:578–585
  50. Caspi A, Williams B, Kim-Cohen J, Craig IW, Milne BJ, Poulton R, Schalkwyk LC, Taylor A, Werts H, Moffitt TE 2007 Moderation of breastfeeding effects on the IQ by genetic variation in fatty acid metabolism. *Proc Natl Acad Sci USA* 104:18860–18865