

Conservation of an intricate circuit for crucial modifications of the tRNA^{Phe} anticodon loop in eukaryotes

MICHAEL P. GUY and ERIC M. PHIZICKY

Department of Biochemistry and Biophysics, Center for RNA Biology, University of Rochester School of Medicine, Rochester, New York 14642, USA

ABSTRACT

Post-transcriptional tRNA modifications are critical for efficient and accurate translation, and have multiple different roles. Lack of modifications often leads to different biological consequences in different organisms, and in humans is frequently associated with neurological disorders. We investigate here the conservation of a unique circuitry for anticodon loop modification required for healthy growth in the yeast *Saccharomyces cerevisiae*. *S. cerevisiae* Trm7 interacts separately with Trm732 and Trm734 to 2'-O-methylate three substrate tRNAs at anticodon loop residues C₃₂ and N₃₄, and these modifications are required for efficient wybutosine formation at m¹G₃₇ of tRNA^{Phe}. Moreover, *trm7Δ* and *trm732Δ trm734Δ* mutants grow poorly due to lack of functional tRNA^{Phe}. It is unknown if this circuitry is conserved and important for tRNA^{Phe} modification in other eukaryotes, but a likely human *TRM7* ortholog is implicated in nonsyndromic X-linked intellectual disability. We find that the distantly related yeast *Schizosaccharomyces pombe* has retained this circuitry for anticodon loop modification, that *S. pombe trm7Δ* and *trm734Δ* mutants have more severe phenotypes than the *S. cerevisiae* mutants, and that tRNA^{Phe} is the major biological target. Furthermore, we provide evidence that Trm7 and Trm732 function is widely conserved throughout eukaryotes, since human *FTSJ1* and *THADA*, respectively, complement growth defects of *S. cerevisiae trm7Δ* and *trm732Δ trm734Δ* mutants by modifying C₃₂ of tRNA^{Phe}, each working with the corresponding *S. cerevisiae* partner protein. These results suggest widespread importance of 2'-O-methylation of the tRNA anticodon loop, implicate tRNA^{Phe} as the crucial substrate, and suggest that this modification circuitry is important for human neuronal development.

Keywords: FTSJ1; TRM7; TRM732; TRM734; THADA; tRNA^{Phe}

INTRODUCTION

Post-transcriptional modification of tRNA is universally required for accurate and efficient translation. Modifications are found in all characterized tRNA species (Machnicka et al. 2013), and are highly conserved within each domain of life (Grosjean 2009). Modifications have a number of different roles, with well documented examples including modulating the efficiency and specificity of charging (Muramatsu et al. 1988; Pütz et al. 1994), altering the specificity of decoding (Johansson et al. 2008), maintaining the frame for decoding (Urbonavicius et al. 2001), and preventing decay of pre-tRNA (Kadaba et al. 2004) and mature tRNA (Alexandrov et al. 2006; Chernyakov et al. 2008).

Many tRNA modifications have a similar biological impact on different organisms. For example, the genes responsible for modification of residue A₃₄ to I₃₄ (inosine) in the wobble position of tRNAs are essential in the bacterium *Escherichia coli* (Wolf et al. 2002), in the yeast *Saccharomyces cerevisiae* (Gerber and Keller 1999), and in the distantly related yeast

Schizosaccharomyces pombe (Kim et al. 2010), and RNAi against a putative homolog results in 29% embryonic lethality in the nematode *Caenorhabditis elegans* (Fernandez et al. 2005). Similarly, lack of the terminal methyl group of mcm⁵U₃₄ (5-methoxycarbonylmethyluridine) in *S. cerevisiae* due to mutation of *TRM9* results in sensitivity to aminoglycosides (Kalhor and Clarke 2003) and DNA damaging agents (Begley et al. 2007) but no other growth defects, depletion of the human protein results in DNA damage sensitivity but no other growth defects (Fu et al. 2010), and mice lacking *TRM9* (*Alkbh8*^{-/-}) appear normal (Songe-Møller et al. 2010).

In contrast, many modifications often do not have precisely the same biological impact on different organisms. Thus, for example, the genes that specify the m¹A₅₈ (1-methyladenosine) modification found in many eukaryotic tRNAs are essential in *S. cerevisiae* (Anderson et al. 1998) due to turnover of pre-tRNA_i^{Met} by the nuclear surveillance pathway (Kadaba et al. 2004), but, the corresponding gene knockouts are each

Corresponding author: eric_phizicky@urmc.rochester.edu

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.047639.114>.

© 2014 Guy and Phizicky This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

viable in *S. pombe* (Kim et al. 2010), although one knockout strain that was examined grows slowly and is more sensitive to oxidative stress (Zuin et al. 2008). Similarly, the genes required for t^6A_{37} (N^6 -threonlycarbamoyleadenosine) formation are essential in *E. coli* and *Haloferax volcanii*, but not in *S. cerevisiae*, although mutants grow poorly (El Yacoubi et al. 2011; Srinivasan et al. 2011; Naor et al. 2012).

Several modifications with more modest phenotypes in *S. cerevisiae* also appear to have a markedly different biological impact on different organisms. Thus, for example, lack of i^6A_{37} (N^6 -isopentenyladenosine) due to mutation of *MOD5* in *S. cerevisiae*, results in reduced nonsense suppression (Laten et al. 1978; Dihanich et al. 1987), but no obvious growth defect, whereas *S. pombe mod5Δ* (*Sp tit1Δ*) mutants grow slowly on glycerol or rapamycin (Lamichhane et al. 2013), *C. elegans mod-5* mutants (*gro-1*) have slowed embryogenesis and development and an increased life span (Lemieux et al. 2001), and mutations in human *MOD5* (*TRIT1*) have been linked to encephalopathy and epilepsy due to mitochondrial defects (Yarham et al. 2014).

Although only a subset of tRNA substrates of a modification enzyme are often responsible for the known phenotypes of a mutation in the corresponding modification gene (Phizicky and Alfonzo 2010; Guy et al. 2012), it is unclear if the same subset of tRNA species are equally important in each organism. For example, in *S. cerevisiae* all phenotypes associated with mutations in the ELP complex (which forms the cm^5U moiety found in mcm^5U_{34} , ncm^5U_{34} , [5-carbamoylmethyluridine], and $mcm^5s^2U_{34}$ [5-methoxycarbonylmethyl-2-thiouridine]) are due to lack of functional tRNA^{Lys(UUU)}, tRNA^{Gln(UUG)}, and tRNA^{Glu(UUC)}, although 11 tRNAs are targets of the ELP complex (Esberg et al. 2006; Johansson et al. 2008; Chen et al. 2011b); in contrast, it appears that only overexpression of tRNA^{Lys(UUU)} is required in *S. pombe* to suppress most of the sensitivity to oxidative stress in an *Sp elp3* mutant (which lacks both the cm^5 moiety and the s^2 moiety of mcm^5s^2U), although tRNA^{Gln(UUG)} and tRNA^{Glu(UUC)} have the same mcm^5s^2U modification (Fernández-Vázquez et al. 2013).

The roles and biological effects of 2'-O-methylation of the anticodon loop are of particular interest because of prior work in *S. cerevisiae* showing a severe phenotype of mutants, specificity of that phenotype for modification of only one substrate, and an intricate circuitry for modification of tRNA substrates (Pintard et al. 2002; Guy et al. 2012). *S. cerevisiae* Trm7 is required for 2'-O-methylation of the anticodon loop of tRNA^{Phe}, tRNA^{Trp}, and tRNA^{Leu(UAA)} at C₃₂ (forming 2'-O-methylcytidine, Cm₃₂) and at N₃₄ (forming Gm₃₄, Cm₃₄, and ncm^5Um_{34} , respectively); and *Sc trm7Δ* mutants have a severe growth defect (Pintard et al. 2002). Interestingly, *Sc* Trm7 interacts with *Sc* Trm732 to form Cm₃₂, and separately interacts with *Sc* Trm734 to form Nm₃₄, and both modifications are required to efficiently drive formation of yW₃₇ (wybutoxine) at m¹G₃₇ on tRNA^{Phe} (Guy et al. 2012) by *Sc* Tyw1-Tyw4 (Fig. 1A; Noma et al. 2006).

Furthermore, overexpression of only tRNA^{Phe} fully suppresses the growth defect and the aminoglycoside sensitivity of *Sc trm7Δ* mutants. In addition, the growth defect of an *Sc trm7Δ* mutant requires loss of both the Cm₃₂ and Gm₃₄ modifications of tRNA^{Phe} (and the accompanying loss of yW₃₇) since neither an *Sc trm732Δ* nor an *Sc trm734Δ* single mutant has any observable growth defect, whereas the double mutant is as sick as an *Sc trm7Δ* mutant and is equally suppressed by overexpression of tRNA^{Phe} (Guy et al. 2012).

The goal of the work described here is to determine if this intricate circuitry, the biologically significant tRNA target, and the importance of Trm7 modifications are conserved, focusing on *S. pombe*, whose lineage diverged from that of *S. cerevisiae* ~1.1 billion yr ago (Hedges 2002). Available evidence is equivocal for conservation of all these features among eukaryotes. Although Trm7 is highly conserved in each of 25 divergent eukaryotic genomes examined (Fig. 1B) (see Materials and Methods), Trm732 homologs can only be identified in 22 genomes, and Trm734 homologs only in 14 genomes, and both protein families have little overall sequence similarity, (~23% and 21% overall identity between the *Sc* proteins and their predicted human orthologs THADA [Trm732] [Fig. 1C] and WDR6 [Trm734], respectively [Shi et al. 2011]). Moreover, *S. cerevisiae* Trm734 has been implicated in regulation of Ty1 transposition (Nyswander et al. 2008) and endoplasmic recycling (Shi et al. 2011), suggesting that Trm734 family proteins from other organisms might have other roles in addition to, or instead of, tRNA modification.

It also appears that *TRM7* has an important but differing biological impact on different organisms, since a high throughput screen in *S. pombe* indicated that the putative *TRM7* gene was essential (Kim et al. 2010), and since mutations in a putative human *TRM7* homolog (*FTSJ1*) are associated with nonsyndromic X-linked intellectual disability (NSXLID) (Freude et al. 2004; Ramser et al. 2004; Froyen et al. 2007; Takano et al. 2008).

We report here that the circuitry and biologically important substrate for 2'-O-methylation of the tRNA anticodon loop are conserved in *S. pombe*, and that *S. pombe trm7Δ* and *trm734Δ* mutants are viable, but with more severe growth defects than in *S. cerevisiae*. Furthermore, we provide evidence that this circuitry is retained in other eukaryotes including humans, suggesting that defective 2'-O-methylation of tRNA is linked to NSXLID.

RESULTS

The putative *S. pombe trm7⁺* gene is required for Cm and Gm formation on tRNA^{Phe} and mutants are barely viable

To determine if *S. cerevisiae* Trm7 function is conserved in eukaryotes, we generated and examined *S. pombe* strains lacking the likely *trm7⁺* gene (*SPAC4F10.03c*). Consistent with




FIGURE 1. Trm7 modification machinery in eukaryotes. (A) Schematic of the anticodon loop of tRNA^{Phe} and tRNA^{Leu(UAA)} from *S. cerevisiae*. *S. cerevisiae* Trm7 (Sc Trm7) requires Sc Trm732 and Sc Trm734 to form Cm₃₂ and Nm₃₄, respectively, on tRNA^{Phe} and tRNA^{Leu(UAA)}. Cm₃₂ and Gm₃₄ modification then drive yW formation from m¹G on tRNA^{Phe}. Wider arrow for Gm₃₄ indicates that yW formation is more dependent on this modification than on Cm₃₂. Predicted human homologs of Sc Trm7, Sc Trm732, and Sc Trm734 are in brackets. (B) Amino acid sequence alignment of Sc Trm7 with putative homologs from *S. pombe* and *H. sapiens*. (C) Schematic representation of Sc Trm732 aligned with putative homologs from *S. pombe* and *H. sapiens*. Inset box is an amino acid alignment of the DUF2428 domain found in these proteins.

previous high throughput results (Kim et al. 2010), we were unable to generate *S. pombe* *trm7Δ* mutant haploids from a heterozygous diploid (relevant genotype: *spac4F10.03cΔ::kanMX/SPAC4F10.03c*) by sporulation and selection for the mutant haploid (data not shown). However, by first introducing a *ura4⁺* plasmid expressing *S. pombe* *trm7⁺* under control of the *P_{nmt1}* promoter (no message in thiamine), we could recover an *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *ura4⁺*] haploid after sporulation on Edinburgh minimal medium (EMM, permissive conditions) in the presence of selective drug; moreover, this *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *ura4⁺*] haploid was viable (albeit barely) when plated to EMM containing 5-fluoroorotic acid (5-FOA) to select against the *ura4⁺* plasmid. Indeed, after streaking the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *ura4⁺*] strain to EMM containing 0.5 mg/L 5-FOA at 30°C, it took 11 d of growth to attain visible colonies, compared with 3 d for wild-type cells (data not shown). Control experiments demonstrated that the growth defect was due to the *trm7Δ* mutation of the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *ura4⁺*] strain, since introduction of a [*P_{nmt1}* *Sp trm7⁺* *LEU2*] plasmid (expressing *trm7⁺* under control of the low strength no message in thiamine promot-

er), resulted in healthy growth after plating to EMM containing 5-FOA (Fig. 2A). The growth defect of *Sp trm7Δ* mutants thus appears to be more severe than that observed in the *S. cerevisiae* *trm7Δ* mutant, which is sick but viable (Pintard et al. 2002; Guy et al. 2012). Under repressive conditions in liquid YES medium, the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain (although much healthier than the *Sp trm7Δ* haploid) had a generation time of 320 min, compared with 148 min for the wild-type strain (a difference of 2.16-fold) (Table 1), which was still more than the difference of 1.94-fold in generation times observed for an *S. cerevisiae* *trm7Δ* mutant compared with its wild-type control. Furthermore, this difference was even greater in selective EMM-Leu + thiamine, with a generation time of 603 min for the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain, compared with 238 min for the wild-type control strain (a difference of 2.53-fold), whereas the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain grew nearly as well as the wild-type strain in permissive conditions (EMM-Leu), with a difference in generation times of only 1.05-fold (Table 1).

Because it was extremely difficult to obtain and grow an *Sp trm7Δ* strain for analysis of modifications, we instead examined the effect of reduced levels of *Sp* Trm7 on modification of tRNA^{Phe}

by growing the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain in EMM-Leu containing thiamine to repress *Sp* Trm7 expression, followed by purification of tRNA^{Phe} and analysis of its nucleoside content by HPLC. Under these conditions, Cm levels of tRNA^{Phe} were substantially reduced compared with those of a wild-type control strain grown in the same medium (0.13 versus 0.93 moles/mole) (Table 2; Fig. 2B), or to those when the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain was grown under permissive (EMM-Leu) conditions (0.90 moles/mole). Similarly, Gm levels were reduced in tRNA^{Phe} from the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain grown in repressive conditions (EMM-Leu containing thiamine) relative to those from the wild-type strain grown under the same conditions (0.04 versus 0.99 moles/mole), or to those when the strain was grown under permissive (EMM-Leu) conditions (0.90 moles/mole). In contrast, the control modification Ψ (pseudouridine) showed little variation in these strains (Table 2).

We also found evidence that yW modification was reduced upon repression of *trm7⁺* expression in the *Sp trm7Δ* [*P_{nmt1}* *Sp trm7⁺* *LEU2*] strain by growth in thiamine-containing




FIGURE 2. *S. pombe* *trm7Δ* mutants are barely viable, and *trm7Δ* [P_{nmt1}** *Sp trm7*⁺] strains grown in repressive conditions have reduced Cm₃₂, Gm₃₄, and yW₃₇ modification of tRNA^{Phe}. (A) *S. pombe* *trm7Δ* mutants are barely viable. *S. pombe* strains with the indicated plasmids were grown in EMM overnight, diluted to OD₆₀₀ of ~0.5 in H₂O, and serially diluted 10-fold in H₂O, and then 2 μL was spotted onto indicated media, followed by incubation for 4 d at 30°C. (B) HPLC traces of tRNA^{Phe} from an *S. pombe* *trm7Δ* [P_{nmt1}** *Sp trm7*⁺] strain. tRNA^{Phe} from indicated strains grown in EMM-Leu with thiamine (+T, repressive conditions) or without thiamine (permissive conditions) was digested to nucleosides and analyzed by HPLC as described in Materials and Methods. (C) Levels of yW on tRNA^{Phe} substantially decrease in an *S. pombe* *trm7Δ* [P_{nmt1}** *Sp trm7*⁺] strain grown under repressive conditions. tRNA^{Phe} from indicated strains grown in EMM-Leu with thiamine (+T) was analyzed as described in panel B under conditions optimized for detecting yW as described in Materials and Methods.

medium. Under these conditions, m¹G levels of tRNA^{Phe} increased substantially compared with those from the wild-type strain (1.80 versus 1.16 moles/mole), suggesting that yW formation from m¹G was reduced (Table 2; Fig. 2B). Direct measurement by HPLC (Noma et al. 2006) confirmed that levels of yW were reduced in tRNA^{Phe} from the *Sp trm7Δ* [P_{nmt1}** *Sp trm7*⁺ LEU2] strain grown under repressive conditions, to 20% of those from tRNA^{Phe} of wild-type cells (Fig. 2C). Thus, just as in *S. cerevisiae* (Guy et al. 2012), *S. pombe* mutants lacking Cm₃₂ and Gm₃₄ due to loss of *trm7*⁺ also appear to have reduced synthesis of yW from m¹G in tRNA^{Phe}.

Curiously, levels of the control modifications m²G (N²-methylguanosine) and m⁷G (7-methylguanosine) were unexpectedly reduced in tRNA^{Phe} purified from different strains

grown in EMM. m⁷G was reproducibly and significantly reduced whenever thiamine was missing from the medium, from 0.55 to 0.28 moles/mole for the wild-type strain, and from 0.46 to 0.25 moles/mole for the *Sp trm7Δ* [P_{nmt1}** *Sp trm7*⁺ LEU2] strain (Table 2; Fig. 2B). m²G levels in tRNA^{Phe} were reproducibly different in different strains, with high levels in the *Sp trm7Δ* [P_{nmt1}** *Sp trm7*⁺ LEU2] strain in the presence of thiamine (0.75 moles/mole), and reduced levels when thiamine was absent (0.12 moles/mole), and with significantly lower levels in the wild-type strain (0.29 moles/mole), further reduced in the absence of thiamine (0.11 moles/mole).

Overexpression of tRNA^{Phe} suppresses the growth defect of *S. pombe* *trm7Δ* mutants

We also found that tRNA^{Phe} was the important Trm7 substrate in *S. pombe*, since overexpression of this tRNA restored healthy growth to *Sp trm7Δ* mutants on 5-FOA medium, whereas overexpression of tRNA^{Trp} or tRNA^{Leu(UAA)}, which were not 2'-O-methylated in *S. pombe* (data not shown), did not (Fig. 3A). Indeed, the *Sp trm7Δ* mutant overexpressing tRNA^{Phe} grew nearly as well as the wild-type control strain on EMM at 25°C and 30°C (and a bit more poorly on YES medium), but was slightly temperature sensitive at 33°C and 37°C (Fig. 3B). As expected for a strain lacking Trm7, tRNA^{Phe} from the *Sp trm7Δ* [tRNA^{Phe}] strain lacked detectable Cm and Gm (Table 2), had increased m¹G levels compared with those from the wild-type strain overexpressing tRNA^{Phe} (2.18 versus 1.62 moles/mole), and had undetectable levels of yW (Fig. 3C). Thus, the important Trm7 substrate in *S. pombe* is tRNA^{Phe}, just as in *S. cerevisiae* (Guy et al. 2012).

TABLE 1. Comparison of generation times for *trm7Δ*, *trm732Δ*, and *trm734Δ* mutant strains from *S. pombe* and *S. cerevisiae* at 30°C

Organism	Strain ^{a,b}	Growth medium	Generation time (min)
<i>S. pombe</i>	Wild type	YES	148 ± 5
<i>S. pombe</i>	<i>trm7Δ</i> [P _{nmt1} ** <i>trm7</i> ⁺]	YES	320 ± 2
<i>S. pombe</i>	<i>trm732Δ</i>	YES	143 ± 11
<i>S. pombe</i>	<i>trm734Δ</i>	YES	258 ± 3
<i>S. pombe</i>	Wild type [vec]	EMM-Leu	315 ± 9
<i>S. pombe</i>	Wild type [vec]	EMM-Leu + thiamine	238 ± 3
<i>S. pombe</i>	<i>trm7Δ</i> [P _{nmt1} ** <i>trm7</i> ⁺]	EMM-Leu	330 ± 10
<i>S. pombe</i>	<i>trm7Δ</i> [P _{nmt1} ** <i>trm7</i> ⁺]	EMM-Leu + thiamine	603 ± 15
<i>S. cerevisiae</i>	Wild type	YPD	66 ± 2
<i>S. cerevisiae</i>	<i>trm7Δ</i>	YPD	128 ± 5
<i>S. cerevisiae</i>	<i>trm732Δ</i>	YPD	69 ± 6
<i>S. cerevisiae</i>	<i>trm734Δ</i>	YPD	70 ± 2

^aMean and standard deviation based on growth from three separate colonies.

^bP_{nmt1}** low strength no message in thiamine promoter.

TABLE 2. HPLC analysis of tRNA^{Phe} nucleoside content from an *S. pombe* *trm7*Δ strain

Strain ^{a,b}	Medium	Cm	Gm	m ¹ G	Ψ	m ² G	m ² C
Moles expected		1	1	1	4	1	1
Wild type [vec]	EMM-Leu	0.92 ± 0.02	0.90 ± 0.04	1.28 ± 0.05	3.85 ± 0.03	0.28 ± 0.02	0.11 ± 0.02
Wild type [vec]	EMM-Leu + thiamine	0.93 ± 0.03	0.99 ± 0.01	1.16 ± 0.01	3.90 ± 0.02	0.55 ± 0.03	0.29 ± 0.03
<i>trm7</i> Δ [P _{trm7} ⁺ <i>trm7</i> ⁺]	EMM-Leu	0.90 ± 0.04	0.90 ± 0.03	1.27 ± 0.03	3.79 ± 0.14	0.25 ± 0.03	0.12 ± 0.04
<i>trm7</i> Δ [P _{trm7} ⁺ <i>trm7</i> ⁺]	EMM-Leu + thiamine	0.13 ± 0.02	0.04 ± 0.01	1.80 ± 0.05	3.80 ± 0.06	0.46 ± 0.03	0.75 ± 0.04
Wild type [tRNA ^{Phe}]	EMM-Leu	0.80 ± 0.02	0.70 ± 0.02	1.62 ± 0.11	3.83 ± 0.02	0.19 ± 0.04	0.10 ± 0.02
<i>trm7</i> Δ [tRNA ^{Phe}]	EMM-Leu	<0.01	<0.03	2.18 ± 0.04	3.90 ± 0.03	0.23 ± 0.01	0.33 ± 0.04

^aMean and standard deviation based on three individual growths and RNA preparations.^bP_{trm7}⁺: low strength no message in thiamine promoter.

The putative *S. pombe* *trm732*⁺ and *trm734*⁺ genes are required for Cm and Gm formation, respectively, on tRNA^{Phe}

To further determine if the modification circuitry for the anticodon loop of tRNA^{Phe} is conserved between *S. cerevisiae* and *S. pombe*, we analyzed the putative *S. pombe* *trm732*Δ (SPCC1494.07) and *trm734*Δ (SPBC1306.02) strains. An *Sp* *trm732*Δ haploid mutant obtained by sporulation of the heterozygous diploid lacked detectable Cm in its tRNA^{Phe} and had normal levels of Gm compared with that from wild type (0.88 versus 0.90 moles/mole) (Table 3; Fig. 4A) and did not have any obvious growth defect on plates or in liquid YES medium (Table 1; Fig. 4B), similar to the lack of an obvious growth defect observed for the *Sc* *trm732*Δ mutant (Table 1; Guy et al. 2012).

However, an *Sp* *trm734*Δ [*Sp* *trm734*⁺ *ura4*⁺] haploid obtained by sporulation of the heterozygous diploid containing the plasmid grew poorly when plated on EMM containing 5-FOA (Fig. 4C), and the resulting *Sp* *trm734*Δ strain lacked Gm₃₄ in its tRNA^{Phe}, and had Cm levels comparable to those of wild type (0.86 versus 0.91 moles/mole) (Table 3; Fig. 4A). The *Sp* *trm734*Δ mutant had a significant growth defect on YES medium on plates or in liquid (generation time of 258 min, compared with 148 min for wild type) (Table 1; Fig. 4D), whereas the *Sc* *trm734*Δ mutant lacked any obvious growth defect on plates or in liquid medium (generation time of 70 min, compared with 66 min for wild type) (Table 1; Guy et al. 2012), demonstrating that Trm734 has a more prominent role in *S. pombe* than in *S. cerevisiae*. Furthermore, since overexpression of tRNA^{Phe} nearly completely suppressed the slow growth phenotype of the *Sp* *trm734*Δ strain over a range of temperatures (Fig. 4E), we conclude that lack of Gm₃₄ of tRNA^{Phe} is the cause of the *Sp* *trm734*Δ defect.

In addition, our analysis showed that, as in *S. cerevisiae* (Guy et al. 2012), yW formation was impaired in the *Sp* *trm734*Δ mutant (44% of wild-type levels) and to a lesser extent in the *Sp* *trm732*Δ mutant (73%) (Fig. 4F), consistent with the increased m¹G levels (Table 3; Fig. 4A). We also note that m²G levels on tRNA^{Phe} are substantially higher in the *Sp* *trm734*Δ mutant compared with those from a wild-type strain when grown in EMM (0.61 versus 0.19 moles/mole).




FIGURE 3. The biologically important *Sp* Trm7 tRNA substrate is tRNA^{Phe}. (A) Overexpression of tRNA^{Phe} suppresses the near lethality of *S. pombe* *trm7*Δ mutants. Strains with plasmids as indicated were grown overnight in EMM-Leu and analyzed as in Figure 2A, after incubation for 3 d at 30°C. (B) Overexpression of tRNA^{Phe} in *S. pombe* *trm7*Δ mutants does not completely restore wild-type growth at 37°C. Strains as indicated were purified on 5-FOA to ensure loss of the [P_{trm7}⁺ *Sp* *trm7*⁺ *ura4*⁺] plasmid, grown overnight in EMM-Leu, and then growth was analyzed as indicated in Figure 2A. (C) tRNA^{Phe} from the *S. pombe* *trm7*Δ [tRNA^{Phe}] strain lacks yW. tRNA^{Phe} purified from the indicated strains after growth in EMM-Leu was analyzed as described in Figure 2C.

TABLE 3. HPLC analysis of tRNA ^{Phe} nucleoside content from <i>S. pombe</i> <i>trm732</i> Δ and <i>S. pombe</i> <i>trm734</i> Δ strains							
Strain ^a	Medium	Cm	Gm	m ¹ G	Ψ	m ⁷ G	m ² G
Moles expected		1	1	1	4	1	1
Wild type	YES	0.88 ± 0.05	0.90 ± 0.03	0.98 ± 0.10	3.99 ± 0.08	0.32 ± 0.01	0.62 ± 0.05
<i>trm732</i> Δ	YES	<0.01	0.88 ± 0.04	1.19 ± 0.07	3.81 ± 0.17	0.31 ± 0.03	0.64 ± 0.07
Wild type [vec]	EMM-Ura	0.90 ± 0.04	0.92 ± 0.02	1.20 ± 0.21	3.94 ± 0.10	0.26 ± 0.12	0.14 ± 0.02
<i>trm734</i> Δ [<i>trm734</i> ⁺]	EMM-Ura	0.75 ± 0.18	0.85 ± 0.06	1.24 ± 0.13	3.84 ± 0.07	0.23 ± 0.05	0.16 ± 0.05
Wild type	EMM	0.91 ± 0.05	0.89 ± 0.06	1.14 ± 0.10	3.98 ± 0.04	0.21 ± 0.03	0.19 ± 0.12
<i>trm734</i> Δ	EMM	0.86 ± 0.04	<0.03	1.57 ± 0.02	4.01 ± 0.06	0.31 ± 0.05	0.61 ± 0.11

^aMean and standard deviation based on three individual growths and RNA preparations.




FIGURE 4. *S. pombe* *trm734*Δ mutants are sick due to hypomodified tRNA^{Phe}. (A) HPLC traces of tRNA^{Phe} from *S. pombe* *trm732*Δ and *S. pombe* *trm734*Δ mutants. tRNA^{Phe} was purified from the indicated strains grown in YES medium and then digested to nucleosides and analyzed by HPLC. (B) *S. pombe* *trm732*Δ mutants are healthy. Indicated strains were grown in YES medium overnight, and analyzed on YES plates as described in Figure 2A. (C) *S. pombe* *trm734*Δ mutants have a growth defect on EMM. Indicated strains were grown overnight in EMM, diluted, and plated on EMM as indicated. (D) *S. pombe* *trm734*Δ mutants have a substantial growth defect on rich medium at various temperatures. Indicated strains were transformed with indicated *ura4*⁺ plasmids, grown in EMM-Ura, diluted, and plated on YES medium at the indicated temperatures after 3 d. (E) Overexpression of tRNA^{Phe} suppresses the slow growth of *S. pombe* *trm734*Δ mutants. Strains with plasmids as indicated were grown in EMM-Leu overnight, analyzed as in Figure 2A, after incubation for 3 d at 30°C. Strains were then purified on medium containing 5-FOA, grown overnight in EMM-Leu, diluted, spotted, and analyzed at the indicated temperatures on YES medium. (F) Levels of yW in tRNA^{Phe} are decreased in *S. pombe* *trm732*Δ and *S. pombe* *trm734*Δ mutants. tRNA^{Phe} was purified from the indicated strains after growth in YES medium and analyzed as described in Figure 2C.

FTSJ1 is the human Trm7 ortholog and requires *S. cerevisiae* Trm732 to catalyze Cm₃₂ modification on tRNA^{Phe} in *S. cerevisiae*

To investigate the conservation of function of metazoan genes involved in 2'-O-methylation of the tRNA anticodon loop, we examined complementation of the corresponding *S. cerevisiae* mutants by introduction of appropriate constructs expressing metazoan TRM7 and/or its metazoan TRM732 and TRM734 partners. To identify TRM7 homologs, we examined *Sp trm7*⁺ and predicted TRM7 orthologs from *D. melanogaster* and humans by introduction of a high copy [2μ *LEU2* P_{GAL} TRM7] plasmid expressing the corresponding ortholog under galactose control to an *Sc trm7Δ* [*Sc TRM7* URA3] strain, followed by analysis of growth after plating on medium containing 5-FOA and galactose. We found that the slow growth phenotype of *Sc trm7Δ* mutants was efficiently suppressed by expression of *Sp Trm7* (~61% identical to *Sc Trm7*), *D. melanogaster* ORF CG5220 (*Dm TRM7*, ~43% identical to *Sc Trm7*), and human FTSJ1 (~50% identical to *Sc Trm7*) but not human FTSJ2 (~34% identical to *Sc Trm7*) (Fig. 5A), after selection on medium containing 5-FOA. Furthermore, the resulting *Sc trm7Δ* [2μ *LEU2* P_{GAL} FTSJ1] strain grew nearly as well as the wild-type *S. cerevisiae* strain (Fig. 5A, right panel), strongly suggesting that FTSJ1 is the human ortholog of Trm7.

The ability of human FTSJ1 and the other tested TRM7 orthologs to complement the growth defect of an *S. cerevisiae* *trm7Δ* mutant strain was surprising because *Sc Trm7* and *Sp Trm7* have a strict requirement for their corresponding Trm732 and Trm734 partner proteins to generate Cm₃₂ and Nm₃₄, respectively (Guy et al. 2012). Because tRNA^{Phe} is the *Sc Trm7* substrate that must be modified for healthy growth in *S. cerevisiae* (Guy et al. 2012), we examined 2'-O-methylation of tRNA^{Phe} from the *Sc trm7Δ* [2μ *LEU2* P_{GAL} FTSJ1] strain. We found that tRNA^{Phe} purified from this strain had high levels of Cm (0.89 versus 0.90 moles/mole in wild type), but no detectable Gm (Table 4), suggesting that human FTSJ1 is working in concert with *Sc Trm732* to modify tRNA^{Phe}. The healthy growth of this *Sc trm7Δ* [2μ *LEU2* P_{GAL} FTSJ1] strain is consistent with our previous observation that the Cm₃₂ modification of tRNA^{Phe} is sufficient for healthy growth in *S. cerevisiae* (Guy et al. 2012).

The occurrence of Cm in tRNA^{Phe} from an *Sc trm7Δ* [2μ *LEU2* P_{GAL} FTSJ1] strain suggested either that FTSJ1 does not require an interacting partner for Cm₃₂ catalysis, or that it works with *S. cerevisiae* Trm732. Since the slow growth phenotype of the *Sc trm7Δ trm732Δ* strain was not suppressed by introduction of the [2μ *LEU2* P_{GAL} FTSJ1] plasmid and no Cm formation was observed on tRNA^{Phe} purified from this strain (Table 4; Fig. 5B), we infer that FTSJ1 works with *Sc Trm732* for formation of Cm₃₂ on tRNA^{Phe}. Since tRNA^{Leu(UAA)} from an *Sc trm7Δ* [2μ *LEU2* P_{GAL} FTSJ1] strain lacked both Cm and ncm⁵Um (Table 5; Fig. 1A), the activity of FTSJ1 with Trm732 in *S. cerevisiae* does not extend to all *Sc Trm7* substrates.




FIGURE 5. FTSJ1 is the human ortholog of *S. cerevisiae* Trm7, and its function in *S. cerevisiae* requires Trm732 for Cm₃₂ modification. (A) Expression of *S. pombe trm7*⁺, *D. melanogaster TRM7*, or human FTSJ1 suppresses the slow growth *S. cerevisiae trm7Δ* mutants. Strains with plasmids as indicated were grown overnight in S-Leu medium containing raffinose and galactose, diluted to OD₆₀₀ of ~0.5 in H₂O, and serially diluted 10-fold in H₂O, and then 2 μL was spotted onto S medium containing 5-FOA, raffinose, and galactose, followed by incubation for 3 d at 30°C. (Right panel) Selected strains were then purified on medium containing 5-FOA, grown overnight in S-Leu medium containing raffinose and galactose, and analyzed by spotting to YP medium containing raffinose and galactose. (B) FTSJ1 requires *Sc TRM732* to suppress the slow growth of *Sc trm7Δ* mutants. Strains were analyzed as in A.

We note that because FTSJ1 is expressed from a high copy P_{GAL} plasmid in these experiments, it is formally possible that FTSJ1 is not the true Trm7 homolog. However, this seems unlikely due to the high sequence similarity between FTSJ1 and Trm7, and the requirement of *Sc Trm732* function for FTSJ1 activity in *S. cerevisiae*.

THADA is the human homolog of Trm732 and works with *Sc Trm7* for Cm₃₂ modification

Although Trm732 homologs have only ~20% sequence identity, we find that the putative human Trm732 homolog THADA functions efficiently to replace *Sc Trm732*. Thus, the slow growth of an *Sc trm7Δ trm732Δ* strain was

TABLE 4. HPLC analysis of tRNA^{Phe} nucleoside content from *S. cerevisiae* *trm7Δ* [*FTSJ1*] and *S. cerevisiae* *trm7Δ trm732Δ* [*FTSJ1 THADA*] strains

Strain ^a	Cm	Gm	m ¹ G	Ψ	m ⁵ C	m ² G
Moles expected	1	1	0	2	2	1
Wild type [vec]	0.90 ± 0.16	0.81 ± 0.03	<0.03	2.07 ± 0.12	1.75 ± 0.09	0.86 ± 0.05
<i>trm7Δ</i> [vec]	<0.01	<0.03	0.99 ± 0.09	2.11 ± 0.16	1.83 ± 0.06	0.98 ± 0.08
<i>trm7Δ</i> [<i>TRM7</i>]	1.0 ± 0.18	0.86 ± 0.08	<0.03	2.13 ± 0.05	1.77 ± 0.10	0.96 ± 0.08
<i>trm7Δ</i> [<i>FTSJ1</i>]	0.89 ± 0.22	<0.03	0.47 ± 0.13	2.13 ± 0.14	1.85 ± 0.09	0.95 ± 0.05
<i>trm7Δ trm732Δ</i> [vec]	<0.01	<0.03	0.91 ± 0.15	2.11 ± 0.07	1.82 ± 0.05	0.89 ± 0.21
<i>trm7Δ trm732Δ</i> [<i>FTSJ1</i>]	<0.01	<0.03	0.93 ± 0.06	2.10 ± 0.04	1.82 ± 0.02	1.01 ± 0.03
<i>trm7Δ trm732Δ</i> [<i>FTSJ1</i> + <i>THADA</i>]	0.90 ± 0.20	<0.03	0.63 ± 0.06	2.11 ± 0.05	1.86 ± 0.04	0.99 ± 0.07
<i>trm732Δ trm734Δ</i> [vec]	<0.01	<0.03	1.03 ± 0.03	2.13 ± 0.08	1.86 ± 0.06	0.98 ± 0.04
<i>trm732Δ trm734Δ</i> [<i>THADA</i>]	0.76 ± 0.13	<0.03	0.66 ± 0.19	2.08 ± 0.05	1.76 ± 0.09	0.89 ± 0.16

^aMean and standard deviation based on three individual growths and RNA preparations.

suppressed by expression of *FTSJ1* and *THADA*, but not by *FTSJ1* alone (Fig. 6A). Since tRNA^{Phe} from this *Sc trm7Δ trm732Δ* [*2μ P_{GAL} FTSJ1 P_{GAL} THADA*] strain had identical Cm levels to that from the wild-type strain (0.90 moles/mole) (Table 4) and no detectable Gm, and since tRNA^{Leu(UAA)} from this strain had substantial Cm modification (0.36 moles/mole) and no ncm⁵Um (Table 5), we conclude that *THADA* is the human *Trm732* homolog and functionally interacts with *FTSJ1*. The ability of *FTSJ1* to modify C₃₂ of tRNA^{Leu(UAA)} in combination with *THADA* but not with *Sc Trm732* may be due to higher levels of *THADA* relative to endogenous levels of yeast *Trm732*, and/or more efficient partnering of *FTSJ1* and *THADA*.

We also find evidence that *S. cerevisiae* *Trm7* can interact with human *THADA* for Cm₃₂ formation, since expression of *THADA* suppressed the slow growth of an *Sc trm732Δ trm734Δ* strain (Fig. 6A), resulting in tRNA^{Phe} with 0.76 moles/mole Cm (Table 4). Thus, our data demonstrate that human (*FTSJ1*) or *S. cerevisiae* *Trm7* can each utilize either human (*THADA*) or *S. cerevisiae* *Trm732* to catalyze formation of Cm₃₂ on tRNA^{Phe} in *S. cerevisiae*. *S. cerevisiae* *Trm7* also appears to work with *S. pombe* *Trm732*, because expres-

sion of *Sp trm732⁺* suppressed the slow growth of the *Sc trm732Δ trm734Δ* strain (Fig. 6B).

DISCUSSION

Our results demonstrate that the entire circuitry for tRNA^{Phe} anticodon loop modification established for *S. cerevisiae* has been retained in the yeast *S. pombe*, requiring *Trm7* to act with *Trm732* to 2'-O-methylate C₃₂, and with *Trm734* to 2'-O-methylate G₃₄, leading to efficient conversion of m¹G₃₇ to yW. Furthermore, we have provided strong evidence that *FTSJ1* and *THADA* are the human *Trm7* and *Trm732* orthologs, and that the human and *S. cerevisiae* proteins can act interchangeably together to catalyze 2'-O-methylation of C₃₂ of tRNA^{Phe} (Table 4). *THADA* and *S. pombe trm732⁺* each complemented the *Trm732* defect of *Sc trm732Δ trm734Δ* mutants, but have little overall conservation (~20% identity, mostly clustered in the small DUF2428 region) (Fig. 1C), suggesting that there may be substantial structural homology in less conserved regions, such as in the predicted armadillo repeats (Tewari et al. 2010).

TABLE 5. HPLC analysis of tRNA^{Leu(UAA)} nucleoside content from an *S. cerevisiae* *trm7Δ* [*FTSJ1*] strain

Modification	Moles expected	wt [vec]	<i>trm7Δ</i> [vec]	<i>trm7Δ</i> [<i>TRM7</i>]	<i>trm7Δ</i> [<i>FTSJ1</i>]	<i>trm7Δ trm732Δ</i> [vec]	<i>trm7Δ trm732Δ</i> [<i>FTSJ1</i>]	<i>trm7Δ trm732Δ</i> [<i>FTSJ1 THADA</i>]
Cm	1	0.58	0.02	0.57	0.03	0.02	0.02	0.36
ncm ⁵ Um	1	1.52	0.03	1.43	0.03	0.01	<0.01	0.02
ncm ⁵ U	0	<0.01	0.94	<0.01	1.16	1.13	1.0	0.97
m ¹ G + Gm	2	1.8	1.86	1.97	1.8	1.96	1.52	1.48
Ψ	2	2.16	2.17	2.13	2.14	2.3	2.14	2.0
m ⁵ C	1	0.82	0.83	0.92	0.74	1.02	0.77	0.97
ac ⁴ C	1	0.71	0.81	0.83	0.74	0.71	0.35	0.35
m ² G	1	0.84	0.95	1.02	0.93	1.04	0.78	0.92
m ^{2,2} G	1	0.88	0.88	0.89	0.89	0.75	0.96	0.83

Since the *S. cerevisiae* and *S. pombe* lineages diverged from one another ~1.1 billion yr ago, and since humans and fungal lineages diverged ~1.6 billion yr ago (Hedges 2002), it seems likely that this anticodon loop modification circuitry is widely conserved among eukaryotes. The absence of identifiable Trm732 and Trm734 orthologs in some eukaryotes may indeed be due to lack of universal conservation of the corresponding genes in some organisms, perhaps because of increased amounts of tRNA^{Phe} in those organisms. Alternatively the lack of identifiable Trm732 and Trm734 orthologs may reflect the poor sequence conservation of these proteins, or may be due to alternative 2'-O-methylation pathways, such as by a phylogenetically distinct anticodon loop 2'-O-methyltransferase family (Tkaczuk et al. 2007), or by a Box C/D guide RNA (Joardar et al. 2011).

We note that experiments to test the function of WDR6, the predicted human Trm734 homolog, by complementation of the appropriate *S. cerevisiae* mutants have been inconclusive. Coexpression of human WDR6 with *FTSJ1* does not suppress the slow growth of a *Sc trm7Δ trm732Δ* strain (data not shown), but WDR6 was not expressed well in *S. cerevisiae* (data not shown). Nonetheless, because Trm734 function is conserved in *S. pombe* and *S. cerevisiae*, it seems plausible that WDR6 and the other Trm734 family members will be required for Gm₃₄ formation on tRNA^{Phe} in their corresponding organisms.

Further evidence that this tRNA^{Phe} anticodon loop modification circuitry is conserved in humans derives from the observation that Ehrlich ascites tumors and neuroblastoma cells lacking O₂yW₃₇ (peroxywybutosine) on tRNA^{Phe} also lack Cm₃₂ and Gm₃₄ modifications on tRNA^{Phe} (Kuchino et al.

1982), consistent with the requirement for 2'-O-methylation of the anticodon loop of tRNA^{Phe} as a prerequisite for yW₃₇ formation, and fueling speculation that these tRNA defects arise from defective FTSJ1 function.

Our data also provide further evidence indicating that levels of tRNA modifications are regulated by cellular growth conditions. Thus, we found that m⁷G and m²G levels on tRNA^{Phe} are decreased in *S. pombe* cells grown in EMM in the absence of thiamine (Table 2), similar to the changes in tRNA modification levels observed in *S. cerevisiae* cells grown under cellular stress conditions or in cells that have undergone growth arrest (Chan et al. 2010; Chan et al. 2012; Preston et al. 2013). Furthermore, the finding that m²G levels of tRNA^{Phe} are near normal in the *Sptrm7Δ* and *Sptrm734Δ* mutants (compared with those from wild-type cells grown in EMM) implies the existence of a compensatory modification mechanism, similar to that observed in *trm9Δ* mutants, which have acquired ncm⁵U (and ncm⁵s²U) in the absence of mcm⁵U₃₄ (and mcm⁵s²U) (Chen et al. 2011a), but in this case the m²G is likely on a different residue.

Our finding that tRNA^{Phe} is the biologically important Trm7 substrate in both *S. cerevisiae* and *S. pombe* (Fig. 3; Guy et al. 2012) suggests that tRNA^{Phe} may be the important substrate throughout eukaryotes. This is consistent with our finding that human FTSJ1 and THADA readily modified C₃₂ of tRNA^{Phe} in *S. cerevisiae*, and that tRNA^{Phe} from 16 of 17 eukaryotes examined contains Cm₃₂ and Gm₃₄ (Machnicka et al. 2013). However, we note that overexpression of tRNA^{Phe} did not completely suppress the growth defect of *Sp trm7Δ* mutants, particularly at high temperature (Fig. 3). This could occur if there are other tRNA species that require modification for full function at this temperature, if the tRNA^{Phe} is not sufficiently overexpressed at this temperature to overcome the defect in decoding, or if part of the defect in tRNA^{Phe} occurs at a step after binding of the tRNA to the A-site of the ribosome, since additional copies of hypomodified tRNAs should not affect translation after this step. Because overexpression of tRNA^{Phe} nearly completely suppressed the slow growth phenotype of *Sp trm734Δ* mutants at all temperatures, the defect is almost certainly caused by loss of Gm₃₄ on tRNA^{Phe} (Fig. 4E).

Although there is conservation of the Trm7 circuitry for tRNA^{Phe} anticodon loop modification and for the importance of these modifications in *S. pombe* and *S. cerevisiae*, there are two crucial differences in the biological consequences of mutations in the corresponding genes in the two organisms. First, a *trm7Δ* mutation was more deleterious to growth in *S. pombe* than in *S. cerevisiae*; thus the *Sp trm7Δ* [*P_{mtl1}** Sp trm7⁺ LEU2*] strain (although itself much healthier than the *Sp trm7Δ* haploid) still had a generation time 2.16-fold higher than the wild-type strain (Table 1), which was slightly more than the difference of 1.94-fold observed for an *S. cerevisiae trm7Δ* mutant compared with its wild-type control. Second, an *S. pombe trm734Δ* mutant had a severe growth defect, with a 1.7-fold increased generation time relative to




FIGURE 6. *S. cerevisiae* Trm7 works with human THADA or *S. pombe* Trm732 for function. (A) THADA is the human ortholog of *S. cerevisiae* TRM7. Strains with plasmids as indicated were analyzed as in Figure 5A. (B) *S. pombe trm732⁺* suppresses the slow growth of *S. cerevisiae trm732Δ trm734Δ* mutants. Strains were analyzed as in Figure 5A.

the wild type strain, whereas an *S. cerevisiae trm734Δ* mutant had 1.06-fold difference in generation time. Furthermore, our identification of *FTSJ1* as the human *Sc TRM7* ortholog suggests that defects in the human gene result in a relatively mild, albeit medically serious, condition, since *FTSJ1* splice site, nonsense, and deletion mutations are consistently associated with NSXLID (Freude et al. 2004; Ramser et al. 2004; Froyen et al. 2007; Takano et al. 2008). NSXLID may occur in these patients because specific human tRNAs have a greater requirement for Trm7 modification in development of the central nervous system (CNS) than in other tissues; indeed, recent results suggest that mutation of a tRNA isodecoder expressed specifically in the CNS can lead to ribosome stalling and contribute to neurodegeneration in certain mutant mouse strains (Ishimura et al. 2014).

Our results demonstrating that THADA is the human Trm732 ortholog further suggest that 2'-O-methylation of the tRNA anticodon loop may be associated with human health. THADA is associated with epithelial thyroid adenomas (Rippe et al. 2003); and genome wide association studies have implicated THADA alleles in type 2 diabetes (Zeggini et al. 2008) and polycystic ovary syndrome (Chen et al. 2011c). However, the biological significance of these associations is not known, and the linkage to Nm₃₂ modification remains to be determined.

NSXLID associated with defective *FTSJ1* (Freude et al. 2004; Ramser et al. 2004; Froyen et al. 2007; Takano et al. 2008) adds to a growing list of neurological disorders associated with defective tRNA modification. This list includes intellectual disability associated with a point mutation in *hADAT3*, the predicted homolog of a subunit of the yeast tRNA A₃₄ deaminase (Alazami et al. 2013); a frameshift mutation in *hTRMT1* (Najmabadi et al. 2011), which has tRNA m^{2,2}G₂₆ (N²,N²-dimethylguanosine) methyltransferase activity (Liu and Stråby 2000); mutations in *NSUN2* (Abbasi-Moheb et al. 2012; Khan et al. 2012; Martinez et al. 2012), which modifies C₃₄, C₄₈, C₄₉, and C₅₀ on target tRNAs to m⁵C; and mutations in *hELP2* (Najmabadi et al. 2011), a member of the ELP complex responsible for formation of the cm⁵U moiety found on mcm⁵U₃₄, ncm⁵U₃₄, mcm⁵s²U₃₄ and related modifications. In addition, familial disautonomia is associated with mutations in *hELP1* (*IKBAKP*) (Anderson et al. 2001). The ELP complex has been reported to have functions in addition to tRNA modification in humans (Creppe et al. 2009; Okada et al. 2010), suggesting the possibility that these disease associations could be due to non-tRNA related defects. Regardless, it is notable that many of the associations between defective tRNA modification and neurological disorders are linked to wobble residue 34, which is required for efficient and accurate decoding of mRNA (Agris et al. 2007; Johansson et al. 2008), suggesting that neurological development is extremely sensitive to defects in translation. It remains to be seen whether *FTSJ1*-associated NSXLID is specifically due to Nm₃₄ defects, to Nm₃₂ defects, or to both.

MATERIALS AND METHODS

Identification of Trm7, Trm732, and Trm734 sequence homologs from diverse eukaryotic genomes

BLAST searches (<http://blast.ncbi.nlm.nih.gov/>) for Trm7, Trm732, and Trm734 homologs were performed against sequenced genomes from a diverse set of eukaryotes including representatives from all five eukaryotic supergroups (Adl et al. 2012), including Amoebozoa (*Dictyostelium discoideum*), Archaeplastida (*Arabidopsis thaliana*, *Cyanidioschyzon merolae*, *Ostreococcus tauri*, *Oryza sativa*, and *Zea mays*), Excavata (*Giardia intestinalis*, *Naegleria gruberi*), Opisthokonta (*Bombyx mori*, *Caenorhabditis elegans*, *Danio rerio*, *Drosophila melanogaster*, *Homo sapiens*, *Monosiga brevicollis*, *Mus musculus*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Xenopus tropicalis*), and SAR (Stramenopiles + Alveolates + Rhizaria) (*Cryptomonas paramecium*, *Guillardia theta*, *Phytophthora infestans*, *Tetrahymena thermophila*, *Thalassiosira pseudonana*, *Toxoplasma gondii*, and *Trypanosoma brucei*).

Yeast strains

Yeast strains are listed in Table 6. The *S. pombe* haploid *Sp trm7Δ::kanMX* [*P_{trm7}* *Sp trm7⁺ ura4⁺*] (yMG1052A) strain was generated by transformation of the *Sp trm7Δ::kanMX* heterozygous diploid with pMG360A [*P_{trm7}* *Sp trm7⁺ ura4⁺*], followed by sporulation on EMM lacking uracil supplemented with 500 mg/L G418, selection of haploids, and PCR confirmation of the knockout. The haploid *S. pombe trm732Δ::kanMX* mutant strain (yMG958B) was generated by sporulation of the heterozygous diploid on yeast extract with 3% dextrose (YE medium) with 200 mg/L G418, selection of haploids, and PCR confirmation. The haploid *S. pombe trm734Δ* [*P_{trm734}* *Sp trm734⁺ ura4⁺*] mutant strain (yMG1289-1) was generated by transformation of the *Sp trm734Δ::kanMX* heterozygous diploid with a *LEU2* tRNA^{Phe} plasmid (pMG308C), followed by sporulation on YE medium with 200 mg/L G418, selection of haploids and PCR verification. This strain was then transformed with pMG426G (*P_{trm734}* *Sp trm734⁺ ura4⁺*), followed by selection of *leu⁻/ura⁺* colonies, and PCR confirmation. The *S. cerevisiae trm7Δ::ble^R* [*URA3 Sc TRM7*] (yMG348-1), *Sc trm732Δ::ble^R* (yMG814-1), *Sc trm734Δ::ble^R* (yMG724-5), and *Sc trm734Δ::ble^R trm732Δ::kanMX* [*Sc TRM734 URA3 CEN*] (yMG818-1) strains were described previously (Guy et al. 2012). Double mutant *S. cerevisiae trm7Δ* strains were constructed by PCR amplification of DNA from the appropriate YKO collection *kanMX* strain (Open Biosystems), followed by transformation of the DNA into yMG348-1.

Plasmids

Plasmids used in this study are listed in Table 7. The *S. pombe* vector expressing *S. pombe trm7⁺* under control of the *P_{trm7}* (no message in thiamine) (pMG360A) and *P_{trm7}Δ* (low strength no message in thiamine) promoters (pMG527B) were constructed by PCR of the *Sp trm7⁺* cDNA ORF from an *S. pombe* cDNA library (Fikes et al. 1990) and insertion into the *XhoI* and *SmaI* sites of pREP4X or pREP81X vectors, respectively. The *S. pombe* vector expressing *Sp trm734⁺* under control of the native *Sp trm734⁺* promoter (pMG426G) was constructed by PCR of *Sp trm734⁺* from *S. pombe* genomic DNA and insertion at the *PstI* and *XhoI* sites of pREP4X to

TABLE 6. Strains used in this study

Strain	Genotype	Source
SP286 (wild-type <i>S. pombe</i> diploid)	<i>ade6-M210/ade6-M216 ura4-D18/ura4-D18 leu1-32/leu1-32</i>	Bioneer
yMG957B (wild-type haploid)	SP286	This study
<i>spac4F10.03cΔ</i> (<i>Sp trm7Δ</i> heterozygous diploid)	SP286, <i>spac4F10.03cΔ::kanMX/SPAC4F10.03c</i>	Bioneer
yMG1052A	SP286, <i>spac4F10.03cΔ::kanMX</i> [<i>P_{nmt1} SPAC4F10.03c ura4⁺</i>]	This study
<i>spcc1306.02Δ</i> (<i>Sp trm734Δ</i> heterozygous diploid)	SP286, <i>spbc1306.02Δ::kanMX/SPBC1306.02</i>	Bioneer
yMG1289-1	SP286, <i>spbc1306.02Δ::kanMX</i> [<i>P_{SPBC1306.02} SPBC1306.02⁺ ura4⁺</i>]	This study
yMG1291	SP286, <i>spbc1306.02Δ::kanMX</i>	This study
<i>spcc1494.07Δ</i> (<i>Sp trm732Δ</i> heterozygous diploid)	SP286, <i>spcc1494.07Δ::kanMX/SPCC1494.07</i>	Bioneer
yMG958B	SP286, <i>spcc1494.07Δ::kanMX</i>	This study
BY4741 (wild <i>S. cerevisiae</i> haploid)	<i>MATa his3-Δ1 leu2Δ0 met15-Δ0 ura3-Δ0</i>	Open Biosystems
yMG348-1	BY4741, <i>Sc trm7Δ::ble^R</i> [<i>Sc TRM7 URA3 CEN</i>]	Guy et al. (2012)
yMG105	BY4741, <i>Sc trm7Δ::ble^R</i>	Guy et al. (2012)
yMG724-5	BY4741, <i>Sc trm734Δ::ble^R</i>	Guy et al. (2012)
yMG814-1	BY4741, <i>Sc trm732Δ::ble^R</i>	Guy et al. (2012)
yMD1-4	yMG348-1, <i>Sc trm734Δ::kanMX</i>	This study
yMD2-4	yMG348-1, <i>Sc trm732Δ::kanMX</i>	This study
yMG818-1	BY4741, <i>Sc trm734Δ::ble^R, Sc trm732Δ::kanMX</i> [<i>Sc TRM734 URA3 CEN</i>]	Guy et al. (2012)

remove the *P_{nmt1}* promoter sequence. tRNA expression vectors were generated by PCR of the appropriate tRNA fragment from genomic DNA and insertion at the PstI and XhoI sites of pREP3X. For expression in *S. cerevisiae*, human *FTSJ1* and *THADA* ORFs were cloned from cDNA plasmids (Open Biosystems), as was *D. melanogaster* ORF *CG5220* (*Dm TRM7*) (Drosophila Genomics Resource Center). *S. pombe trm732⁺* ORF was cloned from an

S. pombe cDNA library (Fikes et al. 1990). ORFs were then inserted by ligation independent cloning (LIC) into [2μ *P_{GALL10}*] *S. cerevisiae* dual ORF expression vectors which express ORFs under *P_{GALL1}* control with a C-terminal PT tag (ORF-3C site-HA epitope-His6- ZZ domain of protein A), and ORFs under *P_{GALL10}* control with no tag, essentially as described previously (Quartley et al. 2009; Guy et al. 2012). All plasmids were confirmed by sequencing before use.

TABLE 7. Plasmids used in this study

Plasmid	Parent	Description	Source
pREP4X		<i>P_{nmt1} ura4⁺</i>	Forsburg (1993)
pMG360A	pREP4X	<i>P_{nmt1} SPAC4F10.03⁺ cDNA ura4⁺</i>	This study
pREP81X		<i>P_{nmt1}⁺ LEU2</i> (low strength <i>P_{nmt1}</i>)	Forsburg (1993)
pMG527B	pREP81X	<i>P_{nmt1}⁺ SPAC4F10.03⁺ cDNA LEU2</i>	This study
pREP3X		<i>P_{nmt1} LEU2</i>	Forsburg (1993)
pMG308C	pREP3X	<i>LEU2 tRNA^{Phe}</i>	This study
pMG309A	pREP3X	<i>LEU2 tRNA^{Trp}</i>	This study
pMG310A	pREP3X	<i>LEU2 tRNA^{Leu(UAA)}</i>	This study
pMG426G	pREP4X	<i>P_{SPBC1306.02} SPBC1306.02⁺ ura4⁺</i>	This study
pAVA579		<i>CEN URA3 LIC</i>	Quartley et al. (2009)
pMG13	pAVA579	<i>CEN URA3 Sc TRM7</i>	Guy et al. (2012)
pBG2619		2μ <i>LEU2 P_{GALL10} LIC</i>	Quartley et al. (2009)
pMG240A	pBG2619	2μ <i>LEU2 P_{GALL10} Sc TRM7</i>	Guy et al. (2012)
pMG298B	pBG2619	2μ <i>LEU2 P_{GALL10} SPAC4F10.03⁺-PT cDNA</i>	This study
pOT2-CG5220		pOT2-CG5220 cDNA	Drosophila Genomics Resource Center
pMG291A	pBG2619	2μ <i>LEU2 P_{GALL10} CG5220</i>	This study
pOTB7 <i>FTSJ1</i>		pOTB7 <i>FTSJ1</i> cDNA	Open Biosystems
pCR-BluntII-TOPO <i>FTSJ2</i>		pCR-BluntII-TOPO <i>FTSJ2</i> cDNA	Open Biosystems
pMG133A	pBG2619	2μ <i>LEU2 P_{GALL10} FTSJ1</i>	This study
pMG135A	pBG2619	2μ <i>LEU2 P_{GALL10} FTSJ2</i>	This study
pBP2A	pAVA579	<i>CEN URA3 Sc TRM734</i>	Guy et al. (2012)
pMG140A	pBG2619	2μ <i>LEU2 P_{GALL1} FTSJ1-PT</i>	This study
pENTR223.1 <i>THADA</i>		pENTR223.1 <i>THADA</i> cDNA	Open Biosystems
pMG244A	pMG140A	2μ <i>LEU2 P_{GALL1} FTSJ1-PT P_{GALL10}-THADA</i>	This study
pMG245A	pBG2619	2μ <i>LEU2 P_{GALL10} THADA</i>	This study
pMG299A	pBG2619	2μ <i>LEU2 P_{GALL10} SPCC1494.07⁺-PT cDNA</i>	This study

Isolation and purification of tRNA

S. pombe strains were grown at 30°C to mid-log phase in YE medium supplemented with 225 mg/L adenine, lysine, histidine, leucine, and uracil (YES), or in EMM with appropriate supplements at 225 mg/L. For analysis of tRNA from the *Sp trm7Δ* [*P_{trm1}+* *Sp trm7⁺* *LEU2*] and wild-type strains under repressive conditions, thiamine was added to EMM-Leu at 5 mg/L. *S. cerevisiae* strains were grown at 30°C to mid-log phase in S dropout medium containing 2% raffinose and 2% galactose. Bulk low molecular weight RNA was extracted from 300 OD-mL pellets, and appropriate 5' biotinylated oligonucleotides were used to purify tRNA as previously described (Jackman et al. 2003).

HPLC and analysis of tRNA

Purified tRNA was digested with P1 nuclease and phosphatase as previously described, and nucleosides were subjected to HPLC analysis essentially as previously described (Jackman et al. 2003). For tRNA^{Phe}, HPLC was done at pH 7.0 to maximize separation of Gm and m¹G as previously described (Guy et al. 2012). For detection of yW, HPLC was performed with buffers and gradients essentially as previously described (Noma et al. 2006).

ACKNOWLEDGMENTS

We thank E. Grayhack for discussions and insights, J. Gecz and M. Shaw (University of Adelaide) for discussions, and S. Forsburg (University of Southern California), S. Shuman (Sloan-Kettering Institute), and B. Schwer (Weill Cornell Medical College) for the gift of plasmids. We thank M. Daly for help in constructing yeast strains. This research was supported by National Institutes of Health (NIH) grant GM052347 to E.M.P. M.P.G. was supported by a NIH postdoctoral training grant NCI T32 CA09363.

Received August 6, 2014; accepted October 9, 2014.

REFERENCES

- Abbasi-Moheb L, Mertel S, Gonsior M, Nouri-Vahid L, Kahrizi K, Cirak S, Wieczorek D, Motazacker MM, Esmaeili-Nieh S, Cremer K, et al. 2012. Mutations in *NSUN2* cause autosomal-recessive intellectual disability. *Am J Hum Genet* **90**: 847–855.
- Adl SM, Simpson AG, Lane CE, Lukeš J, Bass D, Bowser SS, Brown MW, Burki F, Dunthorn M, Hampl V, et al. 2012. The revised classification of eukaryotes. *J Eukaryot Microbiol* **59**: 429–493.
- Agris PF, Vendeix FA, Graham WD. 2007. tRNA's wobble decoding of the genome: 40 years of modification. *J Mol Biol* **366**: 1–13.
- Alazami AM, Hijazi H, Al-Dosari MS, Shaheen R, Hashem A, Aldahmesh MA, Mohamed JY, Kentab A, Salih MA, Awaji A, et al. 2013. Mutation in *ADAT3*, encoding adenosine deaminase acting on transfer RNA, causes intellectual disability and strabismus. *J Med Genet* **50**: 425–430.
- Alexandrov A, Chernyakov I, Gu W, Hiley SL, Hughes TR, Grayhack EJ, Phizicky EM. 2006. Rapid tRNA decay can result from lack of non-essential modifications. *Mol Cell* **21**: 87–96.
- Anderson J, Phan L, Cuesta R, Carlson BA, Pak M, Asano K, Björk GR, Tamame M, Hinnebusch AG. 1998. The essential Gcd10p–Gcd14p nuclear complex is required for 1- methyladenosine modification and maturation of initiator methionyl-tRNA. *Genes Dev* **12**: 3650–3662.
- Anderson SL, Coli R, Daly IW, Kichula EA, Rork MJ, Volpi SA, Ekstein J, Rubin BY. 2001. Familial dysautonomia is caused by mutations of the *IKAP* gene. *Am J Hum Genet* **68**: 753–758.
- Begley U, Dyavaiah M, Patil A, Rooney JP, DiRenzo D, Young CM, Conklin DS, Zitomer RS, Begley TJ. 2007. Trm9-catalyzed tRNA modifications link translation to the DNA damage response. *Mol Cell* **28**: 860–870.
- Chan CT, Dyavaiah M, DeMott MS, Taghizadeh K, Dedon PC, Begley TJ. 2010. A quantitative systems approach reveals dynamic control of tRNA modifications during cellular stress. *PLoS Genet* **6**: e1001247.
- Chan CT, Pang YL, Deng W, Babu IR, Dyavaiah M, Begley TJ, Dedon PC. 2012. Reprogramming of tRNA modifications controls the oxidative stress response by codon-biased translation of proteins. *Nat Commun* **3**: 937.
- Chen C, Huang B, Anderson JT, Byström AS. 2011a. Unexpected accumulation of mcm³U and mcm⁵S²U in a *trm9* mutant suggests an additional step in the synthesis of mcm⁵U and mcm⁵S²U. *PLoS One* **6**: e20783.
- Chen C, Huang B, Eliasson M, Rydén P, Byström AS. 2011b. Elongator complex influences telomeric gene silencing and DNA damage response by its role in wobble uridine tRNA modification. *PLoS Genet* **7**: e1002258.
- Chen ZJ, Zhao H, He L, Shi Y, Qin Y, Shi Y, Li Z, You L, Zhao J, Liu J, et al. 2011c. Genome-wide association study identifies susceptibility loci for polycystic ovary syndrome on chromosome 2p16.3, 2p21 and 9q33.3. *Nat Genet* **43**: 55–59.
- Chernyakov I, Whipple JM, Kotlawa L, Grayhack EJ, Phizicky EM. 2008. Degradation of several hypomodified mature tRNA species in *Saccharomyces cerevisiae* is mediated by Met22 and the 5'-3' exonucleases Rat1 and Xrn1. *Genes Dev* **22**: 1369–1380.
- Creppe C, Malinouskaya L, Volvert ML, Gillard M, Close P, Malaise O, Laguesse S, Cornez I, Rahmouni S, Ormenese S, et al. 2009. Elongator controls the migration and differentiation of cortical neurons through acetylation of α -tubulin. *Cell* **136**: 551–564.
- Dihanich ME, Najarian D, Clark R, Gillman EC, Martin NC, Hopper AK. 1987. Isolation and characterization of *MOD5*, a gene required for isopentenylolation of cytoplasmic and mitochondrial tRNAs of *Saccharomyces cerevisiae* [published erratum appears in *Mol Cell Biol* 1987; **7**: 2035]. *Mol Cell Biol* **7**: 177–184.
- El Yacoubi B, Hatini I, Deutsch C, Kahveci T, Rousset JP, Iwata-Reuyl D, Murzin AG, de Crecy-Lagard V. 2011. A role for the universal Kae1/Qri7/YgjD (COG0533) family in tRNA modification. *EMBO J* **30**: 882–893.
- Esberg A, Huang B, Johansson MJ, Byström AS. 2006. Elevated levels of two tRNA species bypass the requirement for elongator complex in transcription and exocytosis. *Mol Cell* **24**: 139–148.
- Fernandez AG, Gunsalus KC, Huang J, Chuang LS, Ying N, Liang HL, Tang C, Schetter AJ, Zegar C, Rual JF, et al. 2005. New genes with roles in the *C. elegans* embryo revealed using RNAi of ovary-enriched ORFeome clones. *Genome Res* **15**: 250–259.
- Fernández-Vázquez J, Vargas-Pérez I, Sanso M, Buhne K, Carmona M, Paulo E, Hermand D, Rodríguez-Gabriel M, Ayte J, Leidel S, et al. 2013. Modification of tRNA^{Lys}_{UUU} by elongator is essential for efficient translation of stress mRNAs. *PLoS Genet* **9**: e1003647.
- Fikes JD, Becker DM, Winston F, Guarente L. 1990. Striking conservation of TFIID in *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae*. *Nature* **346**: 291–294.
- Forsburg SL. 1993. Comparison of *Schizosaccharomyces pombe* expression systems. *Nucleic Acids Res* **21**: 2955–2956.
- Freude K, Hoffmann K, Jensen LR, Delatycki MB, des Portes V, Moser B, Hamel B, van Bokhoven H, Moraine C, Fryns JP, et al. 2004. Mutations in the *FTSJ1* gene coding for a novel S-adenosylmethionine-binding protein cause nonsyndromic X-linked mental retardation. *Am J Hum Genet* **75**: 305–309.
- Froyen G, Bauters M, Boyle J, Van Esch H, Govaerts K, van Bokhoven H, Ropers HH, Moraine C, Chelly J, Fryns JP, et al. 2007. Loss of *SLC38A5* and *FTSJ1* at Xp11.23 in three brothers with non-

- syndromic mental retardation due to a microdeletion in an unstable genomic region. *Hum Genet* **121**: 539–547.
- Fu D, Brophy JA, Chan CT, Atmore KA, Begley U, Paules RS, Dedon PC, Begley TJ, Samson LD. 2010. Human AlkB homolog ABH8 is a tRNA methyltransferase required for wobble uridine modification and DNA damage survival. *Mol Cell Biol* **30**: 2449–2459.
- Gerber AP, Keller W. 1999. An adenosine deaminase that generates inosine at the wobble position of tRNAs. *Science* **286**: 1146–1149.
- Grosjean H. 2009. Nucleic acids are not boring long polymers of only four types of nucleotides: a guided tour. In *DNA and RNA modification enzymes: structure mechanism, function and evolution* (ed. Grosjean H), pp. 1–12. Landes Bioscience, Austin, TX.
- Guy MP, Podyma BM, Preston MA, Shaheen HH, Krivos KL, Limbach PA, Hopper AK, Phizicky EM. 2012. Yeast Trm7 interacts with distinct proteins for critical modifications of the tRNA^{Phe} anticodon loop. *RNA* **18**: 1921–1933.
- Hedges SB. 2002. The origin and evolution of model organisms. *Nat Rev Genet* **3**: 838–849.
- Ishimura R, Nagy G, Dotu I, Zhou H, Yang XL, Schimmel P, Senju S, Nishimura Y, Chuang JH, Ackerman SL. 2014. RNA function. Ribosome stalling induced by mutation of a CNS-specific tRNA causes neurodegeneration. *Science* **345**: 455–459.
- Jackman JE, Montange RK, Malik HS, Phizicky EM. 2003. Identification of the yeast gene encoding the tRNA m¹G methyltransferase responsible for modification at position 9. *RNA* **9**: 574–585.
- Joardar A, Malliahgari SR, Skariah G, Gupta R. 2011. 2'-O-methylation of the wobble residue of elongator pre-tRNA^{Met} in *Haloferax volcanii* is guided by a box C/D RNA containing unique features. *RNA Biol* **8**: 782–791.
- Johansson MJ, Esberg A, Huang B, Björk GR, Byström AS. 2008. Eukaryotic wobble uridine modifications promote a functionally redundant decoding system. *Mol Cell Biol* **28**: 3301–3312.
- Kadaba S, Krueger A, Trice T, Krecic AM, Hinnebusch AG, Anderson J. 2004. Nuclear surveillance and degradation of hypomodified initiator tRNA^{Met} in *S. cerevisiae*. *Genes Dev* **18**: 1227–1240.
- Kalhor HR, Clarke S. 2003. Novel methyltransferase for modified uridine residues at the wobble position of tRNA. *Mol Cell Biol* **23**: 9283–9292.
- Khan MA, Rafiq MA, Noor A, Hussain S, Flores JV, Rupp V, Vincent AK, Malli R, Ali G, Khan FS, et al. 2012. Mutation in *NSUN2*, which encodes an RNA methyltransferase, causes autosomal-recessive intellectual disability. *Am J Hum Genet* **90**: 856–863.
- Kim DU, Hayles J, Kim D, Wood V, Park HO, Won M, Yoo HS, Duhig T, Nam M, Palmer G, et al. 2010. Analysis of a genome-wide set of gene deletions in the fission yeast *Schizosaccharomyces pombe*. *Nat Biotechnol* **28**: 617–623.
- Kuchino Y, Borek E, Grunberger D, Mushinski JF, Nishimura S. 1982. Changes of post-transcriptional modification of wye base in tumor-specific tRNA^{Phe}. *Nucleic Acids Res* **10**: 6421–6432.
- Lamichane TN, Blewett NH, Crawford AK, Cherkasova VA, Iben JR, Begley TJ, Farabaugh PJ, Maraia RJ. 2013. Lack of tRNA modification isopentenyl-A37 alters mRNA decoding and causes metabolic deficiencies in fission yeast. *Mol Cell Biol* **33**: 2918–2929.
- Laten H, Gorman J, Bock RM. 1978. Isopentenyladenosine deficient tRNA from an antisuppressor mutant of *Saccharomyces cerevisiae*. *Nucleic Acids Res* **5**: 4329–4342.
- Lemieux J, Lakowski B, Webb A, Meng Y, Ubach A, Bussi re F, Barnes T, Hekimi S. 2001. Regulation of physiological rates in *Caenorhabditis elegans* by a tRNA-modifying enzyme in the mitochondria. *Genetics* **159**: 147–157.
- Liu J, Str by KB. 2000. The human tRNA(m²G²⁶)dimethyltransferase: functional expression and characterization of a cloned *hTRM1* gene. *Nucleic Acids Res* **28**: 3445–3451.
- Machnicka MA, Milanowska K, Osman Oglou O, Purta E, Kurkowska M, Olchowik A, Januszewski W, Kalinowski S, Dunin-Horkawicz S, Rother KM, et al. 2013. MODOMICS: a database of RNA modification pathways—2013 update. *Nucleic Acids Res* **41**: D262–D267.
- Martinez FJ, Lee JH, Lee JE, Blanco S, Nickerson E, Gabriel S, Frye M, Al-Gazali L, Gleeson JG. 2012. Whole exome sequencing identifies a splicing mutation in *NSUN2* as a cause of a Dubowitz-like syndrome. *J Med Genet* **49**: 380–385.
- Muramatsu T, Nishikawa K, Nemoto F, Kuchino Y, Nishimura S, Miyazawa T, Yokoyama S. 1988. Codon and amino-acid specificities of a transfer RNA are both converted by a single post-transcriptional modification. *Nature* **336**: 179–181.
- Najmabadi H, Hu H, Garshasbi M, Zemojtel T, Abedini SS, Chen W, Hosseini M, Behjati F, Haas S, Jamali P, et al. 2011. Deep sequencing reveals 50 novel genes for recessive cognitive disorders. *Nature* **478**: 57–63.
- Naor A, Thiaville PC, Altman-Price N, Cohen-Or I, Allers T, de Cr cy-Lagard V, Gophna U. 2012. A genetic investigation of the KEOPS complex in halophilic Archaea. *PLoS One* **7**: e43013.
- Noma A, Kirino Y, Ikeuchi Y, Suzuki T. 2006. Biosynthesis of wybutosine, a hyper-modified nucleoside in eukaryotic phenylalanine tRNA. *EMBO J* **25**: 2142–2154.
- Nyswaner KM, Checkley MA, Yi M, Stephens RM, Garfinkel DJ. 2008. Chromatin-associated genes protect the yeast genome from Ty1 insertional mutagenesis. *Genetics* **178**: 197–214.
- Okada Y, Yamagata K, Hong K, Wakayama T, Zhang Y. 2010. A role for the elongator complex in zygotic paternal genome demethylation. *Nature* **463**: 554–558.
- Phizicky EM, Alfonzo JD. 2010. Do all modifications benefit all tRNAs? *FEBS Lett* **584**: 265–271.
- Pintard L, Leco nte F, Bujnicki JM, Bonnerot C, Grosjean H, Lapeyre B. 2002. Trm7p catalyses the formation of two 2'-O-methylriboses in yeast tRNA anticodon loop. *EMBO J* **21**: 1811–1820.
- Preston MA, D'Silva S, Kon Y, Phizicky EM. 2013. tRNA^{His} 5-methylcytidine levels increase in response to several growth arrest conditions in *Saccharomyces cerevisiae*. *RNA* **19**: 243–256.
- P tz J, Florentz C, Benseler F, Gieg  R. 1994. A single methyl group prevents the mischarging of a tRNA. *Nat Struct Biol* **1**: 580–582.
- Quartley E, Alexandrov A, Mikucki M, Buckner FS, Hol WG, DeTitta GT, Phizicky EM, Grayhack EJ. 2009. Heterologous expression of *L. major* proteins in *S. cerevisiae*: a test of solubility, purity, and gene recoding. *J Struct Funct Genomics* **10**: 233–247.
- Ramser J, Winne penninckx B, Lenski C, Errijgers V, Platzer M, Schwartz CE, Meindl A, Kooy RF. 2004. A splice site mutation in the methyltransferase gene *FTSJ1* in Xp11.23 is associated with non-syndromic mental retardation in a large Belgian family (MRX9). *J Med Genet* **41**: 679–683.
- Rippe V, Drieschner N, Meiboom M, Murua Escobar H, Bonk U, Belge G, Bullerdiek J. 2003. Identification of a gene rearranged by 2p21 aberrations in thyroid adenomas. *Oncogene* **22**: 6111–6114.
- Shi Y, Stefan CJ, Rue SM, Teis D, Emr SD. 2011. Two novel WD40 domain-containing proteins, Ere1 and Ere2, function in the retromer-mediated endosomal recycling pathway. *Mol Biol Cell* **22**: 4093–4107.
- Songe-M ller L, van den Born E, Leihne V, V gb  CB, Kristoffersen T, Krokan HE, Kirpekar F, Falnes P , Klungland A. 2010. Mammalian ALKBH8 possesses tRNA methyltransferase activity required for the biogenesis of multiple wobble uridine modifications implicated in translational decoding. *Mol Cell Biol* **30**: 1814–1827.
- Srinivasan M, Mehta P, Yu Y, Prugar E, Koonin EV, Karzai AW, Sternglanz R. 2011. The highly conserved KEOPS/EKC complex is essential for a universal tRNA modification, t6A. *EMBO J* **30**: 873–881.
- Takano K, Nakagawa E, Inoue K, Kamada F, Kure S, Goto Y; Japanese Mental Retardation Consortium. 2008. A loss-of-function mutation in the *FTSJ1* gene causes nonsyndromic X-linked mental retardation in a Japanese family. *Am J Med Genet B Neuropsychiatr Genet* **147B**: 479–484.
- Tewari R, Bailes E, Bunting KA, Coates JC. 2010. Armadillo-repeat protein functions: questions for little creatures. *Trends Cell Biol* **20**: 470–481.

- Tkaczuk KL, Dunin-Horkawicz S, Purta E, Bujnicki JM. 2007. Structural and evolutionary bioinformatics of the SPOUT superfamily of methyltransferases. *BMC Bioinformatics* **8**: 73.
- Urbonavicius J, Qian Q, Durand JM, Hagervall TG, Björk GR. 2001. Improvement of reading frame maintenance is a common function for several tRNA modifications. *EMBO J* **20**: 4863–4873.
- Wolf J, Gerber AP, Keller W. 2002. tadA, an essential tRNA-specific adenosine deaminase from *Escherichia coli*. *EMBO J* **21**: 3841–3851.
- Yarham JW, Lamichhane TN, Pyle A, Mattijssen S, Baruffini E, Bruni F, Donnini C, Vassilev A, He L, Blakely EL, et al. 2014. Defective i⁶A37 modification of mitochondrial and cytosolic tRNAs results from pathogenic mutations in TRIT1 and its substrate tRNA. *PLoS Genet* **10**: e1004424.
- Zeggini E, Scott LJ, Saxena R, Voight BF, Marchini JL, Hu T, de Bakker PI, Abecasis GR, Almgren P, Andersen G, et al. 2008. Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat Genet* **40**: 638–645.
- Zuin A, Gabrielli N, Calvo IA, Garcia-Santamarina S, Hoe KL, Kim DU, Park HO, Hayles J, Ayté J, Hidalgo E. 2008. Mitochondrial dysfunction increases oxidative stress and decreases chronological life span in fission yeast. *PLoS One* **3**: e2842.



RNA

A PUBLICATION OF THE RNA SOCIETY

Conservation of an intricate circuit for crucial modifications of the tRNA^{Phe} anticodon loop in eukaryotes

Michael P. Guy and Eric M. Phizicky

RNA 2015 21: 61-74 originally published online November 17, 2014

Access the most recent version at doi:[10.1261/rna.047639.114](https://doi.org/10.1261/rna.047639.114)

References

This article cites 67 articles, 27 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/21/1/61.full.html#ref-list-1>

Creative Commons License

This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>
